

Echocardiographic Study in Preterm Infant with Hemodynamic Significant Patent Ductus Arteriosus

by Sunny Mariana Samosir

Submission date: 15-Feb-2022 08:48PM (UTC+0800)

Submission ID: 1762915725

File name: Infant_with_Hemodynamic_Significant_Patent_Ductus_Arteriosus.pdf (351.15K)

Word count: 2605

Character count: 13397

Article Type: Research Article

30

Echocardiographic Study in Preterm Infant with Hemodynamic Significant Patent Ductus Arteriosus

Sunny Mariana Samosir¹, Martono Tri Utomo², Mahrus A. Rahman³, Risa Etika⁴, Dina Angelika², Kartika Darma Handayani², Agus Harianto²

¹Researcher, Pediatric Resident, ²Senior staff in Neonatology Division, ³Senior staff, Head Division of Cardiology,

⁴Senior staff, Head Division of Neonatology Faculty of Medicine, Airlangga University, Dr. Soetomo General Hospital, Surabaya, Indonesia

Abstract

Background: Potential complications of hemodynamic significant patent ductus arteriosus (hsPDA) after birth include heart failure, need for respiratory support, renal dysfunction, intraventricular hemorrhage, as well as long term altered growth and development. Nevertheless, clinical signs of patent ductus arteriosus (PDA) are not sensitive and specific enough. Therefore, echocardiography still remains the preferred method to evaluate the ductal patency in preterm infant. The present study aimed to evaluate the echocardiography characteristic in preterm infant with hsPDA.

Methods: A cross-sectional study was conducted on preterm infants aged 3-7 days with 24-33^{6/7} weeks of gestation. Data taken were demographic, clinical and echocardiography. Diagnosis of hsPDA was carried out by echocardiography; defined as >1.5mm diameter of ductus and >1.4 left pulmonary artery and aorta (La/Ao) ratio. The statistical analysis was undertaken using SPSS 21.0.

Results: There were 11 out of 52 preterm infants diagnosed hsPDA. Mean birth weight was 1213±293 gram; Mean gestational age was 30.72±2.01 weeks. In hsPDA group, mean ductus diameter was 2.84±0.93 mm, mean La/Ao ratio was 1.56±0.26, and mean ejection fraction (EF) was 71.55±5.72%.

Conclusion: Echocardiographic evaluation is important for addressing hsPDA in preterm infants.

Keywords: echocardiographic, preterm infant, hemodynamic significant patent ductus arteriosus

Introduction

The birth rate of premature babies continues to increase year after year. Complications of preterm birth are still the leading cause of infant mortality,

reaching one million deaths in 2015.¹ Patent ductus arteriosus (PDA) is the most common cardiovascular disorder in premature infants.² The ducts of arteriosus will normally shrink after birth in a reasonably month-old baby and are functionally closed at the age of 72 hours. The likelihood of spontaneous closure of the ductus arteriosus (DA) in full-term infants without congenital heart disease is very high; but in the preterm infant, closure rates are poorer.^{3,4}

Some studies linking uncorrected hsPDA to infant morbidity for less than a month include

Corresponding author:

Martono Tri Utomo, MD, PhD
Neonatologist in Neonatology Division Child Health Department/Faculty of Medicine Universitas Airlangga, Dr. Soetomo General Academic Hospital, Surabaya, Indonesia, Email: mrmartono73@gmail.com

intraventricular bleeding, necrotizing enterocolitis (NEC) and renal insufficiency that can be caused by decreased systemic blood flow and tissue ischemic consequences.⁵ Decreased brain oxygenation in infants less than a month can occur in infants with hsPDA resulting in brain damage and developmental disorders.⁶ Clinical assessment and echocardiography are reliable methods of diagnosing an hsPDA. Therefore a timely echocardiographic evaluation is needed. The aim of this study was to define the basic functional echocardiographic characteristic of preterm infant with hsPDA.

Methods

A cross-sectional study was conducted in all preterm infants born between November 2019 and May 2020 at the tertiary level neonatal intensive care unit of Dr. Soetomo General Hospital whose oxygen support devices (i.e. high flow nasal canula, continuous positive airway pressure, invasive and non-invasive ventilator) were eligible for inclusion. Patients with multiple congenital anomaly, ductal dependent cyanotic heart defect, early onset of septicemia, and incomplete consent from parents were excluded.

Echocardiography screening was performed between 3rd and 7th postnatal day by pediatric cardiology consultant using Sonoscape Portable Digital Color Doppler Ultrasound System Model S9 (SonoScape, Shenzhen). hsPDA was considered existing if there were a ductus arteriosus with diameter > 1.5 mm on constriction phase, pulmonal perfusion

seen in the left pulmonal artery diameter and aorta diameter ratio (LA/Ao) was > 1.4, and left to right shunt were present. While DA that was already closed and did not fulfill the requirement for hsPDA was considered non-hsPDA.

Echocardiographic characteristics of babies with hsPDA were described as mean (M) and standard deviation (SD). Differences in clinical characteristics of the two sample groups of hsPDA and non hsPDA were analyzed using Chi square test, Fisher exact test, and exact probability test. IBM SPSS 21.0 was used for all statistical analyses.

The ethical clearance was issued by the Ethical Committee of Dr. Soetomo General Hospital (No.1766/105/XI/2019). Before the subject recruitment, the researchers had explained to the parents about the general research information and the consent.

Result

Echocardiography evaluation were performed in 52 out of 191 preterm infants treated in NICU during study. Four babies with early onset of septicemia, three babies with multiple congenital anomalies, and babies without complete data and consent were excluded from the study. There were 25 (48%) and 27 (52%) preterm male and female babies, respectively, as shown in Table 1. There were 11 infants meet the criteria of hsPDA based on echocardiography.

Table 1. Subject characteristic

	hsPDA n (11)	non-hsPDA n (41)	Total n (52)	P
Sex				
Male	5 (9.6)	20 (38.5)	25 (48.1)	1.000*
Female	6 (11.5)	21 (40.4)	27 (51.9)	
Gestational age				
Extremely Preterm (<28 weeks)	1 (1.9)	3 (5.8)	4 (7.7)	1.000**
Very Preterm (28 - <32 weeks)	6 (11.5)	23 (44.3)	29 (55.8)	
Moderate Late Preterm (32 - <37 weeks)	4 (7.7)	15 (28.8)	19 (36.5)	

Cont... Table 1. Subject characteristic

19 Birth weight				
ELBW (<1000 g)	2 (3.9)	6 (11.5)	8 (15.4)	0.902**
VLBW (<1500 g)	7 (13.5)	24 (46.1)	31 (59.6)	
LBW (<2500 g)	2 (3.9)	11 (21.1)	13 (25)	
Age during recruitment				0.118
3 day	3 (5.7)	21 (40.5)	24 (46.2)	
4 day	6 (11.5)	10 (19.3)	16 (30.8)	
5 day	1 (1.9)	5 (9.6)	6 (11.5)	
6 day	1 (1.9)	4 (7.7)	5 (9.6)	
7 day	0 (0.0)	1 (1.9)	1 (1.9)	
Infant's comorbidity				0.086
Respiratory distress synd.	6 (11.5)	12 (23.1)	18 (34.6)	
Perinatal Asphyxia	2 (3.8)	11 (21.1)	13 (24.9)	
PPHN	3 (5.7)	3 (5.7)	6 (11.4)	
Hyperbilirubinemia	4 (7.7)	6 (11.5)	10 (19.2)	
Others	1 (1.9)	9 (17.3)	10 (19.2)	

Note: Sample characteristic using Chi square test; *Chi square corrected; **Fisher's exact test

The echocardiography characteristics of hsPDA group research subjects presented in table 2 include ejection fraction (EF), arteriosus duct diameter (DA), left pulmonary artery diameter (La), aortic diameter (Ao), as well as La compared Ao ratio. In determining the status of hsPDA used parameters in the form of DA diameter and La/Ao ratio as a marker of the presence of pulmonary hyperperfusion. In this

study obtained the smallest DA diameter of 1.5mm and the largest 4.8mm in the hsPDA group. Of the eleven hsPDA group samples, a diameter of duktus was obtained with an average of 2.84±0.93. The comparison between La and Ao obtained an average score of 1.56±0.26. Ejection fraction in hsPDA group was 71.55±5.72 and the EF in non hsPDA group was 73.94±0.26 (range 56.5-93.12%).

Table 2. Echocardiographic characteristic in hsPDA group

Echocardiographic Characteristic	Patients (n)	hsPDA Mean ± SD
Ejection fraction (%)	11	71.55 ± 5.72
DA diameter (mm)	11	2.84 ± 0.93
Left pulmonary artery diameter (La)	11	11.30 ± 1.97
Aorta diameter (Ao)	11	7.19 ± 1.35
La/Ao ratio	11	1.56 ± 0.26

Discussion

This study showed no significance between gestational age ($p = 0.981$), birth weight ($p = 0.832$), single birth of twin mapun ($p = 0.322$) between hsPDA and non hsPDA groups. Based on gestational age the most to smallest age group was the 28-<32-week group (55.8%), followed by 32-33^{6/7} weeks (36.5%) and gestational age <28 weeks (7.7%). In contrast to other studies where there was an increase in the incidence of hsPDA inversely proportional to the gestational age. Studies in Italy by the Italian Neonatal Network showed higher incidence of PDAs at younger gestational ages: 88.9% and 81.5% at gestational ages of 23 and 24 weeks. PDA incidence decreased at gestation age of 25 weeks by 70.3% and 29 weeks by 32%.⁷

In this study, there were 2/11 neonates with hsPDA having a birth weight of 1000-<1500 grams and as much as 6/41 in the non hsPDA group. The very low birth weight (VLBW) group with a birth weight of 1000-1499 grams in both the hsPDA (7/11) and non hsPDA (24/41) groups had the highest frequency compared to the extremely low birth weight (ELBW, < 1000grams) and low birth weight (LBW, 1500-<2500 grams) groups. These findings are in accordance with previous studies in Palembang, where the majority of babies with PDAs have a birth weight of ≤ 1500 gram and are a risk factor for the occurrence of PDA.⁸ Different results were obtained in the study by Su et al., where the incidence of PDA was higher in ELBW (79%) compared to VLBW (65%) neonatal age of 4 days.⁹ This difference in proportion can be due to the spread of different samples between birth weight groups where the number of ELBW samples is only 8 infants (15.4%) from the total sample.

The duct of arteriosus normally shrinks after birth in a term infant and is functionally closed at the age of 72 hours. The closure of the ducts was delayed to 4 days in 10% of the babies born with 30 to 37 weeks of gestational age, 80% of babies born gestational age 25 to 28 weeks, and 90% in infants born less than 24

weeks.^{3,10} The persistent ductus arteriosus can cause hemodynamic disturbance where the systemic shunt to the pulmonary resulting in pulmonary hyperperfusion and systemic hypoperfusion.¹¹

Although ductus arteriosus evaluation is focused on the functional echocardiography, the baby should always undergo a complete echocardiogram first. An initial echocardiogram enables structural congenital heart disease to be excluded. When performing functional echocardiography, it is also important to bear in mind the difficulties and challenges associated with scanning a premature infant. Some babies are very small, they can be agitated or unstable, have a high heart rate, and sometimes a poor echo window, particularly if they are ventilated and have severe underlying lung disease.⁴

In the early hours of life, relatively high pulmonary pressures result in a balanced pulmonary to systemic circulation. A heart murmur, hyperactive precordium, bounding pulses, and a widened systolic to diastolic pulse pressure amplitude are recognized clinical signs attributed to the presence of a hsPDA. Most clinical signs lack sensitivity in the first days of life and hence during this period, a PDA is usually diagnosed by echocardiography.¹² In this study, the pulsation rate between hsPDA and non hsPDA babies is not differ significantly.

This study obtained an average diameter of ductus of 2.84 ± 0.93 mm. This result is higher than the study by Visconti et al, in which a spontaneous closed duct with diameter of 1.63mm, ductus that required pharmacological therapy with diameter of 2.24mm, and ductus diameter which require operative therapy of 2.39mm.¹³ Determination of hsPDA is generally given when transductal diameter of >1.5mm is obtained. Based on previous studies the value of this intersection indicates the presence of perfusion disorders of the target organ. The La/Ao ratio indicating pulmonary hyper perfusion showed an average value of 1.56 ± 0.26 . This figure meets the criteria made by Sehgal and McNamara where the

ducts with an L/Ao ratio of $> 1.4 \pm 1.6:1$ give the effect of moderate hemodynamic disorders on neonates.¹⁴ One baby gave a manifestation of heart failure that was exacerbated by late onset of septicemia that ended in mortality. Three babies in hsPDA group had persistent pulmonary hypertension of the newborn, thus requiring vasodilator treatment prior to ductal closure.

The limitation of this study is focused solely on evaluating the presence of the ductal and left ventricle function. The sample limitation is also a shortcoming in this study. Therefore, future study is needed.

Conclusion

In conclusions, doppler echocardiographic assessment of ductal flow in preterm infant is useful to predict the development of hsPDA. Clinicians need to perform simple echocardiography screening after 3 days of life in preterm infant.

Acknowledgement: The authors thank the Director of Dr. Soetomo General Hospital, Surabaya Indonesia Airlangga, Surabaya, Indonesia for supporting this research.

Ethical Clearance

This study had got permission from the ethics committee of The Faculty of Medicine, Airlangga University Before the subject recruitment, the explanation was done to the parents.

Source of Funding - Self

Conflict of Interest – Nil

References

1. WHO. Preterm birth. 2018. Available from: <http://www.who.int/mediacentre/factsheets/fs363/en/>.
2. Van Overmeire B, Chemtob S. The pharmacologic closure of the patent ductus arteriosus. *Semin Fetal Neonatal Med.* 2005;10:177–84.
3. Benitz, WE. Patent ductus arteriosus in preterm infants. *Pediatrics.* 2016;137:e20153730.
4. Arlettaz R. Echocardiographic evaluation of patent ductus arteriosus in preterm infants. *Front Pediatr.* 2017;5:1–9.
5. Van Der Laan ME, Roofthoof MTR, Fries MWA, Berger RMF, Schat TE, Van Zoonen AGJF, et al. A hemodynamically significant patent ductus arteriosus does not affect cerebral or renal tissue oxygenation in preterm infants. *Neonatology* 2016;110:141–7.
6. Cohen E, Dix L, Baerts W, Alderliesten T, Lemmers P, Van Bel F. Reduction in cerebral oxygenation due to patent ductus arteriosus is pronounced in small-for-gestational-age neonates. *Neonatology.* 2017;111:126–32.
7. Dani C, Poggi C, Cianchi I, Corsini I, Vangi V, and Pratesi S. Effect on cerebral oxygenation of paracetamol for patent ductus arteriosus in preterm infants. *Eur J Pediatr.* 2018;1–7.
8. Bernati N, Nova R, Tasli JM. T. Risk factors for patent ductus arteriosus in preterm neonates. *Sari Pediatr.* 2014;54:132–6.
9. Su BH, Lin HY, Chiu HY, Tsai ML, Chen YT, Lu IC. Therapeutic strategy of patent ductus arteriosus in extremely preterm infants. *Pediatr Neonatol.* 2020;61:133–41.
10. Hung Y, Yeh J. Molecular mechanisms for regulating postnatal ductus arteriosus closure. *Int J Mol Sci.* 19:1861.
11. Härkin P, Marttila R, Pokka T, Saarela T, Hallman M. Morbidities associated with patent ductus arteriosus in preterm infants. Nationwide cohort study. *J Matern Neonatal Med.* 2018;31:2576–83.
12. Van Laere D, van Overmeire B, Gupta S, El Khuffash A, Savoia M, McNamara PJ, et al. Application of NPE in the assessment of a patent ductus arteriosus. *Pediatr Res.* 2018;84:46–56.
13. Visconti LF, Morhy SS, Deutsch ADA, Tavares GMP, Wilberg TJM, Rossi F de S. Clinical and echocardiographic characteristics associated

with the evolution of the ductus arteriosus in the neonate with birth weight lower than 1,500g. Einstein (Sao Paulo). 2013;11:317–23.

14. Mcnamara PJ, Sehgal A. Disease Staging. Arch Dis Child Fetal Neonatal. 2007;92:424–7.

Echocardiographic Study in Preterm Infant with Hemodynamic Significant Patent Ductus Arteriosus

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

12%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | www.science.gov
Internet Source | 1% |
| 2 | Muhammad Faizi, Nur Rochmah, Irwanto Irwanto, Irma Soelistiyo, Yuni Hisbiyah, Rayi Kurnia Perwitasari. "Risk Factors for Psychosocial Disorders in Children with Congenital Adrenal Hyperplasia during the COVID-19 Pandemic", Open Access Macedonian Journal of Medical Sciences, 2022
Publication | 1% |
| 3 | ijpediatrics.com
Internet Source | 1% |
| 4 | patient.info
Internet Source | 1% |
| 5 | sloap.org
Internet Source | 1% |
| 6 | M. Fiatt, A.C. Bosio, D. Neves, R. Symanski da Cunha, L.T. Fonseca, R.K. Celeste. "Accuracy of a spontaneous breathing trial for | 1% |

extubation of neonates", Journal of Neonatal-Perinatal Medicine, 2021

Publication

7	www.one-health.panafrican-med-journal.com Internet Source	1 %
8	repository.unair.ac.id Internet Source	1 %
9	Yogen Singh, Alain Fraise, Omer Erdeve, Begum Atasay. "Echocardiographic Diagnosis and Hemodynamic Evaluation of Patent Ductus Arteriosus in Extremely Low Gestational Age Newborn (ELGAN) Infants", Frontiers in Pediatrics, 2020 Publication	1 %
10	www.miniturismo.co.uk Internet Source	1 %
11	Submitted to Universitas Airlangga Student Paper	1 %
12	www.mdpi.com Internet Source	1 %
13	cardiothoracicsurgery.biomedcentral.com Internet Source	<1 %
14	pubmed.ncbi.nlm.nih.gov Internet Source	<1 %
15	Lorri Beatty, Robert Green, Kirk Magee, Peter Zed. "A Systematic Review of Ethanol and	<1 %

Fomepizole Use in Toxic Alcohol Ingestions", Emergency Medicine International, 2013

Publication

16

Wei Yue, Xinrui Han, Jianghe Luo, Zhumei Zeng, Ming Yang. "Effect of music therapy on preterm infants in neonatal intensive care unit: Systematic review and meta - analysis of randomized controlled trials", Journal of Advanced Nursing, 2020

Publication

<1 %

17

[academictree.org](https://www.academictree.org)

Internet Source

<1 %

18

[bmcpediatr.biomedcentral.com](https://www.bmcpediatr.biomedcentral.com)

Internet Source

<1 %

19

[es.scribd.com](https://www.es.scribd.com)

Internet Source

<1 %

20

Jan Hau Lee, Rachel G. Greenberg, Bin H. Quek, Reese H. Clark, Matthew M. Laughon, P. Brian Smith, Christoph P. Hornik. "Association between early echocardiography, therapy for patent ductus arteriosus, and outcomes in very low birth weight infants", Cardiology in the Young, 2017

Publication

<1 %

21

[academic.oup.com](https://www.academic.oup.com)

Internet Source

<1 %

[annalspc.com](https://www.annalspc.com)

22

Internet Source

<1 %

23

departamentos.cardiol.br

Internet Source

<1 %

24

f1000research.com

Internet Source

<1 %

25

www.clinical-medicine.panafrican-med-journal.com

Internet Source

<1 %

26

www.ijphrd.com

Internet Source

<1 %

27

www.openveterinaryjournal.com

Internet Source

<1 %

28

"Emerging Topics and Controversies in Neonatology", Springer Science and Business Media LLC, 2020

Publication

<1 %

29

Birol Karabulut, Baran Cengiz Arcagök, Ayse Simsek. "Utility of the Platelet-to-Lymphocyte Ratio in Diagnosing and Predicting Treatment Success in Preterm Neonates with Patent Ductus Arteriosus", Fetal and Pediatric Pathology, 2019

Publication

<1 %

30

Zhen Han Tan, Vijay R. Baral. "Principles of Clinical Management of Patent Ductus

<1 %

Arteriosus in Extremely Preterm Neonates", Current Pediatric Reviews, 2016

Publication

31

Roofthoof, Daniëlla W. E., Ingrid M. van Beynum, Johan C. A. de Klerk, Monique van Dijk, John N. van den Anker, Irwin K. M. Reiss, Dick Tibboel, and Sinno H. P. Simons. "Limited effects of intravenous paracetamol on patent ductus arteriosus in very low birth weight infants with contraindications for ibuprofen or after ibuprofen failure", *European Journal of Pediatrics*, 2015.

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Echocardiographic Study in Preterm Infant with Hemodynamic Significant Patent Ductus Arteriosus

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
