

Neutrophil-to-lymphocyte ratio as an alternative marker of neonatal sepsis in developing countries

by Khadijah Rizky Sumitro

Submission date: 04-Mar-2021 01:57PM (UTC+0800)

Submission ID: 1523870510

File name: lternative_marker_of_neonatal_sepsis_in_developing_countries.pdf (350.44K)

Word count: 4154

Character count: 20656

Neutrophil-to-Lymphocyte Ratio as an Alternative Marker of Neonatal Sepsis in Developing Countries

Khadijah Rizky Sumitro¹, Martono Tri Utomo^{1*} and Agung Dwi Wahyu Widodo²

¹Department of Child Health, Faculty of Medicine, Universitas Airlangga- Dr. Soetomo Hospital, Surabaya, Indonesia

²Department of Clinical Microbiology, Faculty of Medicine, Universitas Airlangga- Dr. Soetomo Hospital, Surabaya, Indonesia

5

ARTICLE INFO

Article history:

Received: 19 February 2020

Accepted: 25 June 2020

Online:

DOI 10.5001/omj.2021.05

Keywords:

Infant, Newborn; Neonatal Sepsis; Neutrophils; Lymphocytes; Intensive Care Units, Neonatal; C-Reactive Protein.

ABSTRACT

Objectives: We sought to analyze the neutrophil-to-lymphocyte ratio (NLR) as an alternative marker of neonatal sepsis. **Methods:** In this cross-sectional study, we undertook consecutive sampling in all inborn neonates admitted to the Neonatal Intensive Care Unit with clinical manifestations of neonatal sepsis. Neonates with congenital anomalies and referred neonates were excluded. Complete blood count, C-reactive protein (CRP), and blood culture were carried out as the septic workup examinations based on the local Clinical Practical Guidelines. NLR is obtained by dividing the absolute count of neutrophils from lymphocytes manually. A cut-off value of NLR is obtained using a receiver operating characteristic curve. **Results:** The median NLR value of the 104 neonates who met the inclusion and exclusion criteria was 3.63 (2.39–6.12). Neonates with NLR of 2.12 have the area under the curve of 0.630 (95% confidence interval (CI): 0.528–0.741) and 0.725 (95% CI: 0.636–0.814) when combined with CRP = 2.70 mg/dL. Neonates with NLR \geq 2.12 in clinical neonatal sepsis had almost double the risk of providing positive blood culture results (relative risk = 1.867, 95% CI: 1.077–3.235; $p = 0.011$). **Conclusions:** NLR, calculated from complete blood count, can be used as an alternative marker of easy and relatively inexpensive neonatal sepsis, especially in developing countries, and detection of proven neonatal sepsis to be better when combined with CRP.

Neonatal sepsis is a major problem for neonates worldwide and contributes to significant morbidity and mortality (term and preterm), especially in developing countries.^{1,2} It is a clinical syndrome characterized by systemic infections and the isolation of pathogens in the blood (bacteremia) and occurs in infants in the first month of life.^{3–6} As many as 1.6 million neonates die each year due to infection with 60% of deaths occurring in developing countries.⁷ Neonatal sepsis is reported as 1–5 per 1000 live births in developed countries, and a higher incidence is reported in developing countries (10–50 per 1000 live births).^{5,8,9}

Routine parameters used for neonatal sepsis have varying diagnostic values such as total leukocyte count, absolute neutrophil count (ANC), immature/total neutrophil (I/T) ratio, and C-reactive protein (CRP). Procalcitonin, a specific bacterial infection marker, has a high price and not always available in various health service facilities, especially in rural

areas.¹⁰ Blood culture as a gold standard marker, takes a long time, and often gave negative results.¹ The neutrophil-lymphocyte ratio (NLR) is inexpensive and included as part of complete blood counts and does not require additional examination.¹⁰ Research on NLR in bacterial infections in children, especially neonates, was still limited. Normal NLR values in healthy neonatal or pediatric populations have only been reported once with an average NLR value of 0.52–0.91.¹¹ To date, no studies in neonates have been conducted in our hospital previously, so the purpose of this study was to analyze the NLR as an initial marker of neonatal sepsis that can be used in limited-resource areas.

METHODS

We conducted a six-month (April to September 2019) observational analytical study using a cross-sectional design with consecutive sampling in all inborn neonates treated in Neonatal Intensive

*Corresponding author: nmartono73@gmail.com

Care Unit (NICU) with clinically neonatal sepsis. Complete blood count, CRP, and blood culture were carried out before giving antibiotics according to the local Clinical Practice Guidelines. As mentioned by Haque, suspected neonatal sepsis was characterized by the presence of one or more fetal inflammatory response syndrome (FIRS) criteria in conjunction with signs and symptoms of infection. FIRS manifested by at least two of the following criteria, tachypnea (respiratory rate/ > 60 bpm) plus either grunting/retraction or desaturation, body temperature abnormalities (> 37.9 °C or < 36 °C), capillary refill time (CRT) > 3 seconds, abnormal leukocyte count (< 4000 or > 34000/mm³), CRP > 10 mg/dL, IL-6 or IL-8 > 70 pg/mL, and positive 16S rRNA genes PCR. Sign and symptoms of infection can be grouped into clinical variables (temperature instability, heart rate (HR) ≥ 180 or ≤ 100, respiratory rate > 60 bpm plus grunting/retraction or desaturation, lethargy/ altered mental status, glucose intolerance, plasma glucose > 10 mmol/L, and drinking intolerance), perfusion variable (CRT > 3 seconds, plasma lactic acid > 3 mmol/L), hemodynamic variables (blood pressure < 2 SD below normal age, systolic blood pressure < 50 mmHg for one day old, and < 65 mmHg for ≤ 1 month old), and inflammatory variables (leukocytosis > 34000/mm³; leukopenia < 4000/mm³; immature neutrophils > 10%; I/T ratio > 0.2; thrombocytopenia < 100 000/mm³; CRP > 10 mg/dL or > 2 standard deviation (SD) above normal value; procalcitonin > 8.1 mg/dL or > 2 SD above normal value; IL-6 or IL-8 > 70 pg/mL; and positive 16S rRNA genes PCR.⁷ Basic characteristics such as sex, gestational age, birth weight, Lubschenko score, mode of delivery, history of premature rupture of membranes, history of mother with preeclampsia/ eclampsia, history of prenatal steroid use, and neonatal sepsis onset were evaluated in this study.

Ethical clearance in this study was approved by the Ethical Committee in Health Research of Dr. Soetomo General Academic Surabaya (ref. no. 1047/KEPK/III/2019).

Complete blood count performed by automated hematology analyzer and include white blood count (WBC) differential to evaluate the WBC based on light scattering characteristics. The ANC and absolute lymphocyte count were identified and counted in the WBC differential. The NLR is obtained by dividing the ANC by the absolute

lymphocytes count recorded in the medical record manually. The diagnosis of neonatal sepsis is categorized into two groups based on blood culture results; proven neonatal sepsis if the blood culture results are positive and suspected neonatal sepsis if negative.

Data were analyzed using SPSS and presented as median and interquartile range. Sex, gestational age, birth weight, Lubchenko score, mode of delivery, history of premature rupture of membranes, history of mothers with preeclampsia/eclampsia, history of prenatal steroids, and neonatal sepsis onset were analyzed using the chi-square test. Total leukocyte count, ANC, absolute lymphocytes count, NLR, and CRP were analyzed using Mann-Whitney U test with significance defined as a *p*-value < 0.050. The NLR cut-off value was established using a receiver operating characteristic curve.

RESULTS

The number of inborn neonates admitted to the NICU during the six-month study period was 492, and 260 of them were suspected neonatal sepsis. A total 156 neonates were excluded from the study due to congenital abnormalities (n = 36) and were referred from other hospitals (outborn neonates, n = 120), so the total subjects who met the inclusion criteria were 104 neonates [Figure 1]. Neonatal sepsis in this study had an incidence of 10.6% with 52 (50.0%) neonates categorized as proven neonatal sepsis. Gram-negative bacteria dominated as the cause of neonatal sepsis (75.0%), with extended-spectrum β -lactamase producing *Klebsiella pneumoniae* (+) isolates making up 61.5%. Table 1 informs the subjects and laboratory characteristics of the subjects in this study. The median NLR (*p* = 0.018) and CRP (*p* = 0.001) in the proven neonatal sepsis neonates




Figure 1: Flow diagram the research design.

Table 1: Subjects characteristics.

Variables	Proven neonatal sepsis n (%)	Suspected neonatal sepsis n (%)	Total n (%)	p-value
Sex				0.239*
Male	30 (57.7)	24 (46.2)	54 (51.9)	
Female	22 (42.3)	28 (53.8)	50 (48.1)	
Gestational age				0.184*
Extremely preterm (< 28 weeks)	5 (9.6)	3 (5.8)	8 (7.7)	
Very preterm (≥ 28–32 weeks)	8 (15.4)	18 (34.6)	26 (25.0)	
Moderate late preterm (≥ 32–37 weeks)	33 (63.5)	24 (46.2)	57 (54.8)	
Term (≥ 37 weeks)	6 (11.5)	7 (13.5)	13 (12.5)	
Birth weight				0.008*
Normal	3 (5.8)	7 (13.5)	10 (9.6)	
Low birth weight (< 2500 g)	27 (51.9)	16 (30.8)	43 (41.3)	
Very low birth weight (< 1500 g)	20 (38.5)	17 (32.7)	37 (35.6)	
Extremely low birth weight (< 1000 g)	2 (3.8)	12 (23.1)	14 (13.5)	
Lubchenco score				0.512*
Appropriate for gestational age	39 (75.0)	36 (69.2)	75 (72.1)	
Small for gestational age	13 (25.0)	16 (30.8)	29 (27.9)	
Mode of delivery				0.222*
Spontaneous	16 (30.8)	22 (42.3)	38 (36.5)	
Scetio caesaria	35 (67.3)	28 (53.8)	63 (60.6)	
Others	1 (1.9)	2 (3.8)	3 (2.9)	
History of premature rupture of membrane				1.000*
Yes	15 (28.8)	15 (28.8)	30 (28.8)	
No	37 (71.2)	37 (71.2)	74 (71.2)	
History of preeclampsia/eclampsia				0.543*
Yes	21 (40.4)	18 (34.6)	39 (37.5)	
No	31 (59.6)	34 (65.4)	65 (62.5)	
History of prenatal steroid				0.464*
Yes	12 (23.1)	9 (17.3)	21 (20.2)	
No	40 (76.9)	43 (82.7)	83 (79.8)	
Onset neonatal sepsis				0.003*
Early-onset neonatal sepsis	13 (25.0)	28 (53.8)	41 (39.4)	
Late-onset neonatal sepsis	39 (75.0)	24 (46.2)	63 (60.6)	
Total leucocyte count, /mm³, median (IQR)	11 700 (6405–17 590)	10 215 (4920–16 010)	11 265 (5745–17 265)	0.465**
Absolute neutrophil count, /mm³, median (IQR)	7310 (4920–11 580)	5390 (2995–10 495)	6375 (3260–11 205)	0.155**
Absolute lymphocyte count, /mm³, median (IQR)	1840 (1120–3145)	2155 (1270–3540)	2010 (1160–3245)	0.359**

*Chi-Square test. **Mann-Whitney U test.

IQR: interquartile range.

was significantly higher than in the suspected group [Figure 2].

Neonates with NLR of 2.12 have an area under the curve (AUC) of 0.630 (95% confidence interval (CI): 0.528–0.741) with a cut-off 2.12. The AUC of NLR did not differ statistically when compared to CRP at a cut-off 2.70 mg/dL, which was 0.690 (95% CI: 0.592–0.793; $p = 0.454$) [Figure 3a].

A better AUC of 0.725 (95% CI: 0.636–0.814) can be achieved if NLR is combined with CRP [Figure 3b]. The NLR has high sensitivity of 80.8% but low specificity of 42.3%. The positive predictive value (PPV) and negative predictive value (NPV) were 58.3% and 68.8% with 62.5% accuracy. The combination of NLR and CRP have 75.0% sensitivity, 67.3% specificity, 69.6% PPV, 72.9%




Figure 2: Box plot of (a) neutrophil-lymphocyte ratio and (b) C-reactive protein based on blood culture (Mann-Whitney U test).

NPV and 71.2% accuracy. Neonates with an NLR ≥ 2.12 have almost twice the risk of giving a positive blood culture result (relative risk (RR) = 1.867, 95% CI: 1.077–3.235; $p = 0.011$), whereas when combined with CRP > 2.70 mg/dL, the risk of giving positive blood culture results is more than doubled (RR = 2.106, 95% CI: 1.396–3.179; $p < 0.001$).

DISCUSSION

Preterm and low birth weight (LBW) infants were the highest subjects to have neonatal sepsis in this study. This results were similar to the previous study reported in Denpasar in 2008 (50% preterm and 53.6% LBW neonates) and Surabaya in 2010 (77.78% preterm and 32% LBW neonates).^{8,12} The incidence of neonatal sepsis is inversely related to

birth weight and gestational age. It is mostly found in infants with very LBW < 1500 g and gestational age < 28 weeks.^{5,13} Prematurity with sepsis as a complication was the leading cause of neonatal mortality. Higher neonatal death was also found at lower gestational age, with the majority born at < 26 weeks. The majority of neonatal death was noted in infants with a birth weight < 1000 g.¹⁴ Our study included more males (51.9%). Similar results were reported by several previous studies ranging from 55.2 to 59.9% although there was no difference in the risk of neonatal sepsis by sex.^{15–19}

As many as 75.0% of positive blood cultures were obtained in late-onset neonatal sepsis (LOS) and dominated by gram-negative bacteria. It is significantly 3.5-times more common than early-onset neonatal sepsis (EOS). LOS with positive




Figure 3: Area under curve (a) neutrophil-lymphocyte ratio (NLR) and C-reactive protein (CRP) and (b) NLR combined with CRP as initial marker of neonatal sepsis.

blood cultures was also reported in previous studies by Shehab El-Din et al,¹⁶ at 65% and Yusef et al,²⁰ at 72%. LOS was associated with nosocomial bacteria and more common in preterm neonates due to long term treatment in NICU.^{1,21,22} Immature immune responses and high invasive life support such as central venous catheters (CVC) or endotracheal tubes and total parenteral nutrition (TPN) support make preterm neonates more susceptible to infection than term neonates.^{1,5,21} The domination of gram-negative bacteria was also reported in a previous study, and it was found that intrapartum antibiotics, TPN duration, CVC duration, and mechanical ventilation duration were potential maternal and neonatal risk factors for late-onset gram-negative sepsis.¹⁹ We did not evaluate the use of CVC, endotracheal tubes, and TPN in this study. Low humoral immune system in preterm neonates occurs because the transfer of transplacental immunoglobulin G from mother to fetus begins in the second trimester and peaks in the third trimester of pregnancy.^{21,22}

Nearly 90% of neonatal sepsis with positive blood cultures in this study had normal leukocyte counts (4000–34 000/mm³). Abnormal total leukocyte counts were only found in eight neonates (four EOS and four LOS) with total leukocyte counts < 4000/mm³ and no neonates with total leukocyte counts > 34 000/mm³. Sucilathangam et al,²³ also reported that normal total leukocyte counts were obtained in 12/14 (85.72%) neonates with positive blood culture results. A normal total leukocyte count cannot rule out the presence of neonatal sepsis because 50.0% of neonatal sepsis with positive blood cultures have a normal total leukocyte count.⁹

Higher median NLR was found in proven neonatal sepsis or neonates with positive blood cultures, and higher mean NLR in positive blood cultures (3.69±3.0) compared to negative blood cultures (1.56±1.83, $p < 0.001$).¹ Significantly, higher NLR compared to healthy neonates were also reported in several studies. The term neonatal sepsis group had a significantly higher NLR than healthy term neonates reported by Can et al,²⁴ with 2.88±0.16 and 0.21±0.12, respectively ($p = 0.02$), and Omran et al,²⁵ with 2.9±1.7 and 1.6±0.4, respectively ($p < 0.001$). Overall studies in neonatal sepsis groups have higher NLR values compared to NLR values previously reported by Hamiel et al,¹¹ in healthy neonates or the pediatric population with an average NLR 0.52–0.91. The

high NLR value in the sepsis group is due to an imbalance between the levels of neutrophils and lymphocytes. Increasing neutrophils is a first-line defense mechanism in the primary role of neutrophils as an innate immune system response to fight bacterial infections and stimulates the process of emergency granulopoiesis. An increase in neutrophil counts is accompanied by a decrease in lymphopoiesis and monocytopoiesis and also a decrease in neutrophil apoptosis but an increase in lymphocyte apoptosis resulting in neutrophilia and lymphocytopenia.^{1,26–28}

The NLR cut-off in this study was higher than in previous studies. The NLR 1.81 cut-off value in neonates with risk factors for sepsis and neonatal sepsis in Dr. Moewardi Hospital, Surakarta, has 86.1% sensitivity, 85.1% specificity, 68.9% PPV, and 94.1% NPV.²⁹ The NLR cut-off in this study was lower than in some previous studies but with higher sensitivity. Omran et al,²⁵ obtained an average NLR value in 35 term neonates with positive and negative blood cultures compared to healthy term neonates of 2.9±1.7 with NLR 2.7 has 80% sensitivity and 57.1% specificity with AUC of 0.791±0.057. Ruslie et al,³⁰ obtained an NLR 9.4 cut-off value having a sensitivity of 61.5% and a specificity of 66.7% in 94 neonates with clinical manifestations of sepsis (positive blood culture versus negative blood culture), which was dominated by term neonates as the study population.

The AUC of NLR and CRP are not statistically significantly different, but the best AUC is obtained if both are combined. Only one previous study also reported the best AUC of 0.79 (95% CI: 0.70–0.88) was found when NLR and CRP were combined compared to NLR or CRP alone. The study was conducted in term neonates aged 7–28 days old until 90 days old who experience serious bacterial infection.¹¹

This is the first study in Surabaya, Indonesia, that assesses NLR as a neonatal sepsis marker. The use of NLR is expected to reduce the use of antibiotics that are not appropriate, complications, and neonatal mortality due to infections, especially bacterial infections.

This study's limitation is that it was collected at only one research center and we excluded neonates transferred from other hospitals. This study also did not include healthy neonates as a control group so the normal range value of NLR could not be known.

23

CONCLUSION

The NLR, by dividing the absolute count of neutrophils from lymphocytes from complete blood count, can be used as an alternative marker of neonatal sepsis, especially in developing countries. It is an easy and relatively inexpensive examination. Better detection of proven neonatal sepsis can be achieved using NLR when combined with CRP.

40

Disclosure

The authors declared no conflicts of interest. No funding was received for this study.

Acknowledgements

The authors thank all patients who have been involved in this study, and all team members and colleagues for assisting this research.

REFERENCES

- Alkan Ozdemir S, Arun Ozer E, Ilhan O, Sutcuoglu S. Can neutrophil to lymphocyte ratio predict late-onset sepsis in preterm infants? *J Clin Lab Anal* 2018 May;32(4):e22338.
- Hasibuan BS. Comparison of microbial pattern in early and late onset neonatal sepsis in referral center Haji Adam Malik hospital Medan Indonesia. *IOP Conf Ser Earth Environmental Sci* 2018;125(1):012053.
- Edwards MS, Baker CJ. Sepsis in the newborn. In: Gershon A, Hotez P, Katz S, editors. *Krugman's Infectious Disease of Children*. 11th ed. Philadelphia, 2003. p. 545–563.
- Polin RA; Committee on Fetus and Newborn. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics* 2012 May;129(5):1006-1015.
- Bany-Mohammed F. Sepsis. In: Gomella TL, Cunningham MD, Eyal FG, editors. *Neonatology: management, procedures, on-call problems, diseases and drugs*. 7th ed. New York, 2013. p. 865-874.
- Coetzee M, Mbowane N, De Witt T. Neonatal sepsis: Highlighting the principles of diagnosis and management. *S Afr J Child Heal* 2017 Jul;11(2):99-103.
- Haque KN. Definitions of bloodstream infection in the newborn. *Pediatr Crit Care Med* 2005 May;6(3) (Suppl):S45-S49.
- Kardana IM. Incidence and factors associated with mortality of neonatal sepsis. *Paediatr Indones* 2011 May;51(3):144-148.
- Gomella TL. Postdelivery antibiotic. In: Gomella TL, Cunningham MD, Eyal FG, editors. *Neonatology: management, procedures, on-call problems, diseases and drugs*. 7th ed. New York, 2013. p. 492-501.
- Holub M, Beran O, Kaspiřiková N, Chalupa P. Neutrophil to lymphocyte count ratio as a biomarker of bacterial infections. *Cent Eur J Med* 2012 Jan;7(2):258-261.
- Hamiel U, Bahat H, Kozar E, Hamiel Y, Ziv-Baran T, Goldman M. Diagnostic markers of acute infections in infants aged 1 week to 3 months: a retrospective cohort study. *BMJ Open* 2018 Jan;8(1):e018092.
- Utomo MT. Neonatal sepsis in low birth weight infants in Dr. Soetomo General Hospital. *Indones J Trop Infect Dis* 2010 May-Aug;1(2):86-89.
- Dong Y, Speer CP. Late-onset neonatal sepsis: recent developments. *Arch Dis Child Fetal Neonatal Ed* 2015 May;100(3):F257-F263.
- Abdellatif M, Ahmed M, Bataclan MF, Khan AA, Al Battashi A, Al Maniri A. The patterns and causes of neonatal mortality at a tertiary hospital in Oman. *Oman Med J* 2013 Nov;28(6):422-426.
- Ozkan H, Cetinkaya M, Koksak N, Celebi S, Hacimustafaoglu M. Culture-proven neonatal sepsis in preterm infants in a neonatal intensive care unit over a 7 year period: coagulase-negative Staphylococcus as the predominant pathogen. *Pediatr Int* 2014 Feb;56(1):60-66.
- Shehab El-Din EM, El-Sokkary MM, Bassiouny MR, Hassan R. Epidemiology of neonatal sepsis and implicated pathogens: a study from Egypt. *Biomed Res Int* 2015;2015(1):509484.
- Mahallei M, Rezaee MA, Mehramuz B, Beheshtirooy S, Abdinia B. Clinical symptoms, laboratory, and microbial patterns of suspected neonatal sepsis cases in a children's referral hospital in northwestern Iran. *Medicine (Baltimore)* 2018 Jun;97(25):e10630.
- Murthy S, Godinho MA, Guddattu V, Lewis LE, Nair NS. Risk factors of neonatal sepsis in India: A systematic review and meta-analysis. *PLoS One* 2019 Apr;14(4):e0215683.
- Abdellatif M, Al-Khabori M, Rahman AU, Khan AA, Al-Farsi A, Ali K. Outcome of late-onset neonatal sepsis at a tertiary hospital in Oman. *Oman Med J* 2019 Jul;34(4):302-307.
- Yusef D, Shalakhiti T, Awad S, Algharaibeh H, Khasawneh W. Clinical characteristics and epidemiology of sepsis in the neonatal intensive care unit in the era of multi-drug resistant organisms: A retrospective review. *Pediatr Neonatol* 2018 Feb;59(1):35-41.
- Kaufman D, Fairchild KD. Clinical microbiology of bacterial and fungal sepsis in very-low-birth-weight infants. *Clin Microbiol Rev* 2004 Jul;17(3):638-680.
- Zea-Vera A, Ochoa TJ. Challenges in the diagnosis and management of neonatal sepsis. *J Trop Pediatr* 2015 Feb;61(1):1-13.
- Sucilathangam G, Amuthavalli K, Velvizhi G, Ashihabegum MA, Jeyamurugan T, Palaniappan N. Early diagnostic markers for neonatal sepsis: comparing procalcitonin (pct) and c-reactive protein (crp). *J Clin Diagn Res* 2012 May;6(4):627-631.
- Can E, Hamilcikan Ş, Can C. The value of neutrophil to lymphocyte ratio and platelet to lymphocyte ratio for detecting early-onset neonatal sepsis. *J Pediatr Hematol Oncol* 2018 May;40(4):e229-e232.
- Omran A, Maarouf A, Saleh MH, Abdelwahab A. Salivary C-reactive protein, mean platelet volume and neutrophil lymphocyte ratio as diagnostic markers for neonatal sepsis. *J Pediatr (Rio J)* 2018 Jan - Feb;94(1):82-87.
- Wesche DE, Lomas-Neira JL, Perl M, Chung CS, Ayala A. Leukocyte apoptosis and its significance in sepsis and shock. *J Leukoc Biol* 2005 Aug;78(2):325-337.
- Manz MG, Boettcher S. Emergency granulopoiesis. *Nat Rev Immunol* 2014 May;14(5):302-314.
- Hwang SY, Shin TG, Jo IJ, Jeon K, Suh GY, Lee TR, et al. Neutrophil-to-lymphocyte ratio as a prognostic marker in critically-ill septic patients. *Am J Emerg Med* 2017 Feb;35(2):234-239.
- Nugroho ID. Nilai diagnostik neutrophil to lymphocyte count ratio sebagai penanda awal pada neonatus terinfeksi [Thesis]. Universitas Sebelas Maret; 2018. Available from: <https://eprints.uns.ac.id/40470/>.
- Ruslie RH, Tjipta DG, Samosir CT, Hasibuan BS. Bacterial pattern and role of laboratory parameters as marker for neonatal sepsis. *IOP Conf Ser Earth Environ Sci* 2018;125(1):012057.

Neutrophil-to-lymphocyte ratio as an alternative marker of neonatal sepsis in developing countries

ORIGINALITY REPORT

23%

SIMILARITY INDEX

16%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Santosh K Panda, Manas K Nayak, Soumini Rath, Palash Das. "The Utility of the Neutrophil-Lymphocyte Ratio as an Early Diagnostic Marker in Neonatal Sepsis", Cureus, 2021
Publication 1%
- 2** Mohammad N. Almohammady, Eman M. Eltahlawy, Nashwa M. Reda. "Pattern of bacterial profile and antibiotic susceptibility among neonatal sepsis cases at Cairo University Children Hospital", Journal of Taibah University Medical Sciences, 2020
Publication 1%
- 3** cdn.intechopen.com
Internet Source 1%
- 4** www.heanoti.com
Internet Source 1%
- 5** modified-argon-95704.appspot.com
Internet Source 1%

6

Neeraj Kumar Neeraj Kumar, Dayal R, Agrawal D, Kumar P, Bhatia R, Goyal A, Verma D. "IS PROCALCITONIN A BETTER DIAGNOSTIC MARKER THAN CRP IN NEONATAL SEPSIS?", Journal of Evolution of Medical and Dental Sciences, 2014

Publication

1 %

7

Yearbook of Intensive Care and Emergency Medicine, 2007.

Publication

1 %

8

Melek Büyükeren, Hasan Tolga Çelik, Şule Yiğit, Oytun Portakal, Banu Sancak, Murat Yurdakök. "The role of the delta neutrophil index in determining the etiology of neonatal sepsis", Turkish Journal of Biochemistry, 2020

Publication

1 %

9

repository-tnmgrmu.ac.in

Internet Source

1 %

10

M J Schindl. "The value of residual liver volume as a predictor of hepatic dysfunction and infection after major liver resection", Gut, 2005

Publication

1 %

11

www.slideshare.net

Internet Source

1 %

12

paediatricaindonesiana.org

Internet Source

1 %

13	jamanetwork.com Internet Source	1 %
14	app.trdizin.gov.tr Internet Source	1 %
15	Herzog, Maruša, Lilijana Kornhauser Cerar, Tanja Premru Sršen, Ivan Verdenik, and Miha Lučovnik. "Impact of risk factors other than prematurity on periventricular leukomalacia. A population-based matched case control study", <i>European Journal of Obstetrics & Gynecology and Reproductive Biology</i> , 2015. Publication	1 %
16	"Monday, 1 September 2008", <i>European Heart Journal</i> , 09/02/2008 Publication	1 %
17	journals.lww.com Internet Source	1 %
18	epdf.tips Internet Source	1 %
19	balimedicaljournal.org Internet Source	<1 %
20	www.ijss-sn.com Internet Source	<1 %
21	www.nature.com Internet Source	<1 %

22 Jie Xu, Ziling Nie, Junjie Zhu. "Characterization and Selection of Probability Statistical Parameters in Random Slope PWM Based on Uniform Distribution", IEEE Transactions on Power Electronics, 2021
Publication <1 %

23 www.turingmachine.org
Internet Source <1 %

24 www.researchsquare.com
Internet Source <1 %

25 www.hindawi.com
Internet Source <1 %

26 www.wjgnet.com
Internet Source <1 %

27 C. Seyler. "Risk Factors for Active Tuberculosis after Antiretroviral Treatment Initiation in Abidjan", American Journal of Respiratory and Critical Care Medicine, 04/07/2005
Publication <1 %

28 www.msmanuals.com
Internet Source <1 %

29 Haque, K.N.. "Defining common infections in children and neonates", Journal of Hospital Infection, 200706
Publication <1 %

www.mdpi.com

30

Internet Source

<1 %

31

www.tandfonline.com

Internet Source

<1 %

32

Tom-Revzon, Catherine. "Strategic Use of Antibiotics in the Neonatal Intensive Care Unit :", The Journal of Perinatal & Neonatal Nursing, 2004.

Publication

<1 %

33

research.library.mun.ca

Internet Source

<1 %

34

Ida Sue Baron, Fern R. Litman, Margot D. Ahronovich, Jennifer C. Gidley Larson. "Neuropsychological Outcomes of Preterm Triplets Discordant for Birthweight: A Case Report", The Clinical Neuropsychologist, 2007

Publication

<1 %

35

jped.elsevier.es

Internet Source

<1 %

36

www.thieme-connect.com

Internet Source

<1 %

37

Yaping Li, Muqi Wang, Wenjun Wang, Dandan Feng, Huiling Deng, Yufeng Zhang, Shuangso Dang, Song Zhai. "

<1 %

Prognostic Value of Neutrophil-to-Lymphocyte Ratio in Predicting Death Risk in Patients with Severe Hand, Foot and Mouth Disease

", Therapeutics and Clinical Risk Management, 2020

Publication

38

M A Bhat. "Organism-specific platelet response and factors affecting survival in thrombocytopenic very low birth weight babies with sepsis", Journal of Perinatology, 06/25/2009

Publication

<1 %

39

www.ajol.info

Internet Source

<1 %

40

mafiadoc.com

Internet Source

<1 %

41

www.ijphrd.com

Internet Source

<1 %

42

Nadire Seval Gündem, Nuriye Tarakçı Emiroglu, Erkan Atas. "Causative Agents, Antibiotic Resistance Patterns and Risk Factors in Early and Late Onset Neonatal Sepsis in a Neonatal Intensive Care Unitof", INDIAN JOURNAL OF NEONATAL MEDICINE AND RESEARCH, 2019

Publication

<1 %

43

ijpsr.com
Internet Source

<1 %

44

Aytenew Getabelew, Mihret Aman, Endashaw Fantaye, Tomas Yeheyis. "Prevalence of Neonatal Sepsis and Associated Factors among Neonates in Neonatal Intensive Care Unit at Selected Governmental Hospitals in Shashemene Town, Oromia Regional State, Ethiopia, 2017", International Journal of Pediatrics, 2018

Publication

<1 %

45

Gulali Aktas, Tuba Duman, Burcin Atak, Ozge Kurtkulagi, Satilmis Bilgin, Elif Basaran, Muhammed Demirkol, Mehmet Kosekli. "Irritable bowel syndrome is associated with novel inflammatory markers derived from hemogram parameters", Family Medicine & Primary Care Review, 2020

Publication

<1 %

46

mlsb.kmu.edu.tw
Internet Source

<1 %

47

Xuwei Cui, Yongyan Shi, Xinyi Zhao, Xindong Xue, Jianhua Fu. "The clinical significance of interleukin-6, C-reactive protein, and procalcitonin in the early recognition of nosocomial infections in preterm infants", Research Square, 2020

Publication

<1 %

48	f1000research.com Internet Source	<1 %
49	www.ncbi.nlm.nih.gov Internet Source	<1 %
50	Stephanie Royer Moss, Hallie C. Prescott. "Are we getting any better at diagnosing sepsis?", Elsevier BV, 2020 Publication	<1 %
51	www.paediatricaindonesiana.org Internet Source	<1 %
52	Ismail Alay, Cihan Kaya, Ibrahim Karaca, Ecem Eren, Murat Hosgoren, Fidan Aslanova, Huseyin Cengiz, Murat Ekin, Levent YaSar. "The effectiveness of neutrophil to lymphocyte ratio in prediction of medical treatment failure for tubo_ovarian abscess", Journal of Obstetrics and Gynaecology Research, 2019 Publication	<1 %
53	www.scientiaricerca.com Internet Source	<1 %
54	www.e-cep.org Internet Source	<1 %
55	test.dovepress.com Internet Source	<1 %

advancesindifferenceequations.springeropen.com

56

<1 %

57

"1-A6: Cell and Molecular Biology 2 : Poster Sessions", *Respirology*, 2013.

Publication

<1 %

58

Dildar Duman, Emine Aksoy, Meltem Coban Agca, Nagihan Durmus Kocak et al. "The utility of inflammatory markers to predict readmissions and mortality in COPD cases with or without eosinophilia", *International Journal of Chronic Obstructive Pulmonary Disease*, 2015

Publication

<1 %

59

Hebatallah A. Shaaban, Nesma Safwat. "Mean platelet volume in preterm: A predictor of early onset neonatal sepsis", *The Journal of Maternal-Fetal & Neonatal Medicine*, 2018

Publication

<1 %

60

Deepak Sharma, Nazanin Farahbakhsh, Sweta Shastri, Pradeep Sharma. "Biomarkers for diagnosis of neonatal sepsis: a literature review", *The Journal of Maternal-Fetal & Neonatal Medicine*, 2017

Publication

<1 %

61

Ahmed Omran, Abdallah Maarooof, Mai H. Saleh, Amina Abdelwahab. "Salivary C-reactive protein, mean platelet volume and neutrophil

<1 %

lymphocyte ratio as diagnostic markers for neonatal sepsis", Jornal de Pediatria, 2018

Publication

62

Ting You, Han Zhang, Lu Guo, Ke-Ran Ling, Xiao-Yu Hu, Lu-Quan Li. " Differences in clinical characteristics of early- and late-onset neonatal sepsis caused by pneumoniae ", International Journal of Immunopathology and Pharmacology, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Neutrophil-to-lymphocyte ratio as an alternative marker of neonatal sepsis in developing countries

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
