

Zinc supplementation in cytokine regulation during LPS- induced sepsis in rodent

by Martono Tri Utomo

Submission date: 04-Mar-2021 02:37PM (UTC+0800)

Submission ID: 1523893982

File name: n_in_cytokine_regulation_during_LPS-induced_sepsis_in_rodent.pdf (416.16K)

Word count: 3638

Character count: 18716

Zinc Supplementation in Cytokine Regulation During LPS-induced Sepsis in Rodent

Martono T Utomo^{1*}, Subijanto M Sudarmo¹, Ketut Sudiana²

1. Department of Child Health Faculty of Medicine, Universitas Airlangga.

2. Department of Patology Anatomy Faculty of Medicine, Universitas Airlangga.

Abstract

Severe sepsis increased pro-inflammatory cytokines and low Zinc levels were also found in patients with sepsis. To explain the mechanisms of sepsis improvement after Zinc administration through cytokine regulations

Total sample of 40 rodents was randomized into 4 groups with treatment of LPS in LPS group and LPS-Zinc and placebo in Control and Zinc group, then followed by blood sampling at the second hour to measure Zinc level by AAS. Furthermore, in the LPS-Zinc and Zinc group, Zinc was administered orally, while the control and LPS group were given placebo orally. The blood samples were then collected at the 8th, 24th, and 72nd hour to measure TNF- α , IL-6, IL-10, and TGF- β by sandwich-ELISA method and the Zinc content was also measured at the 72nd hour using AAS. One-way Anova, Kruskal Wallis, Mann-Whitney, paired t-tests, and Wilcoxon Signed Rank Test were employed as statistical analyses.

There were some decreases in the level of TNF- α and IL-6 plasma and an elevated levels of IL-10 and TGF- β in the LPS-Zinc group compared to the LPS group.

Administration of Zinc in sepsis improve sepsis condition through decreased pro-inflammatory cytokines (TNF- α and IL-6) and increased anti-inflammatory cytokines (IL-10 and TGF- β).

Experimental article (J Int Dent Med Res 2020; 13(1): 46-50)

Keywords: Zinc, sepsis, cytokine

Received date: 25 October 2018

Accept date: 19 August 2019

Introduction

Sepsis remains a world health problem due to its high mortality rate despite optimal therapy using antibiotics, fluids, and inotropes.^{1,2} Mortality rate of sepsis in hospitals is approximately 15% among adults and 25% among children.^{1,3} During sepsis, an increase in pro-inflammatory cytokines of IL-1 β , IL-6, IL-8, IL-12, IFN, and TNF- α are associated with mortality in sepsis.⁴ Low serum Zinc levels are also found in severe and critical sepsis among children and adults.⁵⁻⁷ Zinc supplementation can reduce pro-inflammatory cytokines of IL-6, IL-8, and TNF- α .^{5,8,9} In addition, during Zinc deficiency, there is also a low level of anti-inflammatory cytokines of I-4, IL-10 and TGF- β .¹⁰ Several studies have shown that administration

of Zinc for sepsis can be useful to reduce mortality rate.^{11,12} This study aims to investigate the effect of Zinc on cytokine regulation under septic conditions.

Materials and methods

Animal studies

This study used a sample of 40 Sprague-Dawley rodents (10-12 weeks old) which were acclimated 2 weeks before treatment, given standard food, and grouped randomly into Control, LPS, LPS-Zinc, and Zinc groups (10 rodents in each group). The administration of LPS *E coli* serotype O111: B4 from Sigma with a dose of 10 mg / kg (500 mg was prepared, yet only 100 mg was taken). Then it was diluted with 10 mL of aquabidest and injected intravenously in the tail of rodents in LPS group and LPS-Zinc group as much as 0.2 mL, whereas in the Control and Zinc groups, rodents were given normal saline of 0.2 mL intravenously. In addition, in order to facilitate intravenous injection, 0.1 mL of ketamine was administered.

*Corresponding author:

Martono T Utomo

Department of Child Health Faculty of Medicine, Universitas Airlangga.

E-mail: martono-t-u@fk.unair.ac.id

Zinc drop (10 mg/mL) (Zincpro® drop 15 mL, Combiphar Indonesia) was diluted to 1 mg/mL Zinc sulfate solution by adding 9 mL of aquabidest to every 1 mL Zinc drop. Two mg/kg of Zinc dose was to human body, or equivalent to 4.65 mg / kg of rodents' body weight. It was given to rodents via gastric tube to the LPS-Zinc and Zinc group with a dose of 1 mL for 3 days, while in the control and LPS groups, rodents were given a placebo aquabidest of 1 mL.

Blood sampling was conducted through rodents' tail for 5 rodents in each group at the second hour to analyze Zinc levels and at the 8th and 24th hours for the next 5 rodents in each group. Then at the 72nd hour, we took blood from rodents' heart and aorta to analyze cytokines of TNF- α , IL6, IL10, and TGF- β . Examination of Zinc levels was also carried out at the 72nd hour. Blood samples without EDTA were centrifuged at 6,000 G for 10 minutes and at 25 °C within 30 minutes, and then stored in a refrigerator at -20 °C until they were analyzed.

Plasma Zinc

Plasma Zinc was assayed by Atomic Absorption Spectrophotometry (AAS) methods that was used in the laboratory of Science and Technology, Brawijaya University. This methods had been described in the previous study (13).

Cytokine analysis

Measurements of cytokines. The concentrations of TNF- α , IL6, IL10, and TGF- β in the supernatant were measured by sandwich-ELISA, using a FineTest ELISA Kit manual instruction's from Wuhan Fine Biological Technology Co.,Ltd. This study was approved by the Ethical Committee of the Brawijaya University Malang.²⁴

Statistical analysis

The data are expressed as means \pm SDs. All data in 4 groups were compared using one-way Anova and post hoc test. If the data were not normally distributed, further Kruskal-Wallis test was employed. The data in a group at the 8th, 24th, and 72nd hour were compared with same subject anova when the data distribution was normal, or Friedman anova if they were not normally distributed. The data of Zinc level at the 2nd and 72nd hours were analyzed with t-test dependent.

Results

During study, two rodents were dead in LPS group at the 8th and 24th hour. In this study the minimal sample for each group was 5 and duplicated to 10, because there was blood collection more than 2 times in 24 hours. Sample characteristics between groups were comparable, as depicted in Table 1 which shows that there is no differences in rodents' weight in 4 groups.

Group	n	Body weight (g)				p
		Mean	SD	Minimum	Maximum	
Control	5	221.40	6.50	213	229	0.34
LPS	5	240.80	10.50	225	253	
LPS-Zinc	5	226.60	19.46	208	256	
Zinc	5	232.80	24.81	201	263	

Table 1. Rodents' body weights in 4 groups

LPS E Coli administration influences plasma Zinc level

The effects of intravenous LPS *E Coli* on Zinc levels are illustrated in Table 2. Giving intravenous LPS *E Coli* to the LPS and LPS-Zinc groups can reduce Zinc levels below normal levels with normal values of Zinc levels 0.84-1.59 mg / L, though there are no differences between the four groups. Increased Zinc levels were found in the LPS, LPS-Zinc, and Zinc groups at the 72nd hour. The increase Zinc levels in the LPS and LPS-Zinc groups illustrated some fluctuations in Zinc levels during sepsis condition. While for the case of Zinc group, it proved that giving Zinc supplementation for 3 days could increase Zinc levels. Zinc levels at the 72nd hour of the control group, LPS, and LPS-Zinc were not significantly different, which showed fluctuations in Zinc levels during sepsis.

Group	n	Plasma Zinc (mg/L)		p
		2 hours	72 hours	
Control	5	0.87 \pm 0.09 (0.75-0.97)	0.91 \pm 0.22 ^a (0.57-1.19)	0.67
LPS	5	0.62 \pm 0.21 (0.42-0.97)	1.14 \pm 0.24 ^a (0.94-1.54)	0.04†
LPS-Zinc	5	0.75 \pm 0.06 (0.71-0.86)	0.99 \pm 0.07 ^a (0.94-1.11)	0.04†
Zinc	5	0.87 \pm 0.18 (0.62-1.12)	1.49 \pm 0.17 ^b (1.33-1.74)	0.001†
P		0.07	0.001*	

Table 2. Plasma Zinc at the 2nd and 72nd hour after LPS E coli administration.

* Significant at $\alpha=0.05$ (One way Anova)
 † Significant at $\alpha=0.05$ (Paired t-test / Wilcoxon Signed Rank Test)
^{a,b} Same superscript in the column revealed no difference between groups (multiple comparisons LSD)

Zinc supplementation reduces the pro-inflammatory cytokine

Modeling of sepsis experimental animals in this study using intravenous injection of LPS *E coli* was successful, which was characterized by an inflammatory response in the form of high levels of pro-inflammatory cytokines of TNF- α and IL-6 in the LPS and LPS-Zinc groups compared to the control group. Zinc supplementation reduced levels of pro-inflammatory cytokines of TNF- α and IL-6 in septic experimental animals, which was seen from the lower levels of pro-inflammatory cytokines in the LPS-Zinc group compared to LPS at the 8th, 24th, and 72nd hour. TNF cytokine values - α of the LPS group fluctuated at 8th, 24th, and 72nd hour. Similar occurrence also happened in the control group. While the cytokine values of IL-6 in LPS and LPS-Zinc groups fluctuated at the 8th, 24th, and 72nd hour, peaking at the 24th hour and sloping down at the 72nd hour. These results are illustrated in Tables 3 and 4.

TNF- α plasma concentration (pg/mL)					
Group	n	8 th hour	24 th hour	72 nd hour	p
Control	5	79.95 \pm 2.04 ^a (76.55-81.55)	81.55 \pm 1.30 ^a (80.55-84.05)	76.35 \pm 3.19 ^a (71.55-80.55)	0.029†
LPS	5	122.55 \pm 11.40 ^c (111.55-136.55)	110.05 \pm 5.48 ^b (106.55-119.05)	108.05 \pm 4.87 ^b (101.55-114.05)	0.034†
LPS-Zinc	5	92.75 \pm 1.25 ^b (91.55-94.55)	95.35 \pm 2.17 ^b (91.55-96.55)	93.35 \pm 2.49 ^c (91.55-96.55)	0.211
Zinc	5	77.85 \pm 4.15 ^a (71.55-81.55)	76.55 \pm 6.12 ^a (71.55-86.55)	82.35 \pm 4.27 ^b (76.55-86.55)	0.101
p		0.000*	0.001*	0.001*	

Table 3. TNF- α plasma concentration at 8th, 24th, and 72nd hours.

* Significant at $\alpha=0.05$ (Brown-Forsythe/Kruskal-Wallis)
 † Significant at $\alpha=0.05$ (Same subject Anova/Anova Friedman)
^{a,b,c,d} Same superscript in the column revealed no difference between groups (multiple comparisons Games-Howell/Mann-Whitney)

IL-6 plasma concentration (pg/mL)					
Group	n	8 th hour	24 th hour	72 nd hour	p
Control	5	1.58 \pm 0.09 ^b (1.42-1.64)	1.39 \pm 0.11 ^a (1.31-1.55)	1.57 \pm 0.05 ^a (1.53-1.63)	0.091
LPS	5	4.91 \pm 0.40 ^c (4.25-5.21)	5.57 \pm 0.16 ^d (5.54-5.93)	3.39 \pm 0.38 ^c (3.09-4.05)	0.000†
LPS-Zinc	5	2.15 \pm 0.09 ^b (2.06-2.25)	3.83 \pm 0.19 ^c (3.64-4.11)	2.51 \pm 0.15 ^b (2.34-2.74)	0.000†
Zinc	5	2.10 \pm 0.17 ^b (1.91-2.27)	2.21 \pm 0.44 ^b (1.81-2.72)	1.84 \pm 0.26 ^b (1.55-2.14)	0.304
p		0.000*	0.000*	0.001*	

Table 4. IL-6 plasma concentration at the 8th, 24th, and 72nd hour.

* Significant at $\alpha=0.05$ (Brown-Forsythe/Kruskal-Wallis)
 † Significant at $\alpha=0.05$ (Same subject Anova)
^{a,b,c,d} Same superscript in the column revealed no difference between groups (multiple comparisons Games-Howell/Mann-Whitney)

Zinc supplementation increases anti-inflammatory cytokine

In contrast to pro-inflammatory cytokines, Zinc increases anti-inflammatory cytokines of IL-10 and TGF β . There were higher levels of IL-10 and TGF- β in the LPS-Zinc group than in the LPS group at the 8th, 24th, and 72nd hour. The IL-10 levels of LPS and LPS-Zinc groups fluctuated at the highest levels at the 24th hour. TGF- β level of the LPS group was also fluctuated with the same pattern, while the TGF- β level of the LPS-Zinc group found a different pattern, namely there was a gradual increase from the 8th to the 72nd hour.

IL-10 plasma concentration (pg/mL)					
Group	n	8 th hour	24 th hour	72 nd hour	p
Control	5	1.27 \pm 0.04 ^a (1.24-1.34)	1.37 \pm 0.08 ^a (1.31-1.46)	1.38 \pm 0.05 ^a (1.33-1.44)	0.07
LPS	5	2.12 \pm 0.21 ^b (1.83-2.33)	3.83 \pm 0.32 ^c (3.37-4.11)	2.17 \pm 0.20 ^b (1.95-2.48)	0.000†
LPS-Zinc	5	2.96 \pm 0.52 ^b (2.38-3.78)	5.16 \pm 0.56 ^d (4.52-5.74)	4.06 \pm 0.57 ^d (3.26-4.51)	0.002†
Zinc	5	2.11 \pm 0.29 ^b (1.85-2.54)	1.99 \pm 0.39 ^b (1.65-2.64)	1.63 \pm 0.14 ^b (1.51-1.87)	0.06
p		0.000*	0.000*	0.000*	

Table 5. IL-10 plasma concentration at 8th, 24th, and 72nd hours

* Significant at $\alpha=0.05$ (Brown-Forsythe/Kruskal-Wallis)
 † Significant at $\alpha=0.05$ (Same subject Anova)
^{a,b,c,d} Same superscript in the column revealed no difference between groups (multiple comparisons Games-Howell/Mann-Whitney)

TGF- β plasma concentration (pg/mL)					
Group	n	8 th hour	24 th hour	72 nd hour	p
Control	5	0.80 \pm 0.03 ^a (0.76-0.84)	0.85 \pm 0.06 ^a (0.76-0.92)	0.77 \pm 0.03 ^a (0.74-0.83)	0.08
LPS	5	1.75 \pm 0.09 ^b (1.64-1.87)	2.02 \pm 0.38 ^c (1.68-2.59)	1.98 \pm 0.29 ^c (1.60-2.39)	0.35
LPS-Zinc	5	2.59 \pm 0.24 ^c (2.37-2.97)	3.95 \pm 0.27 ^d (3.73-4.42)	4.74 \pm 0.25 ^d (4.40-4.98)	0.000†
Zinc	5	1.73 \pm 0.17 ^b (1.57-1.97)	1.65 \pm 0.20 ^b (1.48-1.90)	1.37 \pm 0.16 ^b (1.21-1.55)	0.05
p		0.000*	0.000*	0.000*	

Table 6. TGF- β plasma concentration at the 8th, 24th, and 72nd hour

* Significant at $\alpha=0.05$ (Brown-Forsythe/Oneway Anova)
 † Significant at $\alpha=0.05$ (Same subject Anova)
^{a,b,c,d} Same superscript in the column revealed no difference between groups (multiple comparisons Games-Howell/LSD)

Discussion

This study showed some improvements in sepsis in terms of decreased inflammatory response in the form of decreasing pro-inflammatory cytokines (IL-6 and TNF- α) and increasing anti-inflammatory cytokines (IL-10 and TGF- β). This study is different from previous studies for rodent which were given Zinc before being induced to sepsis by CLP. In the previous study, similar results were obtained in the form of a decrease in pro-inflammatory cytokines of IL-6 and IL-1 β .¹⁴ In this study, sepsis was modeled in experimental animals by administering intravenous LPS *E. Coli*. The use of intra-venous LPS *E Coli* as a sepsis experiment in animal model was similarly used in previous studies, resulting in the increase of pro-inflammatory cytokines.^{15,16} The successful use of LPS *E coli* for the inflammatory reaction caused by sepsis in animal models in this study was demonstrated by an increase in pro-inflammatory cytokines (TNF- α and IL-6) and anti-inflammatory cytokines (IL-10 and TGF- β). An increase in pro-inflammatory cytokines (TNF- α and IL-6) and anti-inflammatory cytokines (IL-10 and TGF- β) are associated with the worsening or progression of sepsis.¹⁷ Increasing pro-inflammatory cytokine TNF- α and IL-6 are also occurred in local infection such as periodontitis.¹⁸

The decrease in plasma Zinc level in two hours after the administration of LPS *E coli* in the LPS and LPS-Zinc groups in this study was not as large as the decrease in plasma Zinc levels in sepsis and critical conditions in the previous study (45-48 $\mu\text{g} / \text{dL}$).⁵⁻⁷ In this study, the mean Zinc level in the LPS group was 0.62 mg/L or 62 $\mu\text{g} / \text{dL}$ and the mean LPS-Zinc group's Zinc level was 0.75 mg /L or 75 $\mu\text{g}/\text{dL}$ which was included in Zinc deficiency criteria according to the normal Zinc levels of 84 - 159 $\mu\text{g} / \text{dL}$.¹⁸ While in the control and Zinc groups, there were normal Zinc levels, though there were no differences with the LPS and LPS-Zinc groups. For this study, symptoms of sepsis might be triggered by intravenous LPS *E coli* and did not cause severe sepsis symptoms. Low Zinc levels in sepsis, which were found in patients with severe and critical sepsis in children and adults, were associated with mortality.⁵⁻⁷ Blood's low Zinc levels in sepsis are caused by the influence of pro-inflammatory cytokines (namely IL1 β and IL-6) which activate STAT mediated signals and

upregulation of ZIP14 and ZIP 6; then, they also trigger intracellular blood Zinc influx (20). Low Zinc levels and increase in cytokines of IL-6, IL-8, IL-1 TN, and TNF- α occur more frequently in sepsis, compared to healthy individuals.⁵

Increased levels of TNF- α are found in sepsis and associated with shock, coagulopathy, death, and survival. TNF- α is also used as a sepsis diagnosis and evaluation of therapy in sepsis.²⁰ TNF- α has an important role in inflammation. Moreover, rhTNF's administration in experimental animals can cause symptoms of hypotension, metabolic acidosis, massive pulmonary hemorrhage, acute tubular necrosis in kidney, and gastrointestinal bleeding lesions.^{21,22} TNF- α levels in experimental animals injected with LPS *E coli* and oral Zinc administration was lower than the experimental group injected with LPS *E coli* and given placebo in this study. The decrease in TNF- α in the group given Zinc was also found in previous studies, such as in cases of diarrhea in children,²³ sepsis in rodents as animal model,⁹ sickle-cell disease,²⁴ and incidence of infection among elderly.²⁵

Increase in IL-6 occurs in SIRS conditions associated with infection and mortality.^{26,27} Increase in IL-6 in severe sepsis causes an increase in capillary leak and a decrease in intestinal contractions.^{28,29} The administration of Zinc in this study reduced IL-6 level in experimental animal groups which were injected with intravenous LPS *E coli* and Zinc, compared to the experimental group which was only injected with LPS *E coli*.

IL-10 and TGF- β are anti-inflammatory cytokines secreted by Treg. Previous study showed that giving Zinc to healthy people for 10 days can induce Treg.³⁰ IL-10, which was secreted by Treg, inhibits Th1, so pro-inflammatory cytokines from Th1 are reduced.³¹ In this study, an increase in IL-10 in the LPS-Zinc group was higher than the LPS group. It indicated that Zinc's effect on the increase in IL-10 cytokines already occurred at the 8th hour. In the previous study, there was an increase after 10 days of Zinc administration, because no analysis was performed at an earlier hour.³⁰ Increased TGF- β secretion, which is an anti-inflammatory cytokine secreted by Treg, occurs in the LPS-Zinc group compared to the LPS group. Giving Zinc can increase TGF- β 1, elevation of intestinal villi, and ratio villi crypt in the intestine, so that it can improve intestinal

permeability.³² TGF- β can inhibit TNF- α cytokines through translated inhibition of TNF mRNA.³³

The typical decreased vertical facial height of this patient resulted from the congenital.

Conclusions

Administration of Zinc in sepsis can improve sepsis condition through cytokine regulation in the form of decreased pro-inflammatory cytokines (TNF- α and IL-6) and increased anti-inflammatory cytokines (IL-10 and TGF- β).

Declaration of Interest

The authors report no conflict of interest and the article is not funded or supported by any research grant.

References

1. Weiss SL, Fitzgerald JC, Pappachan J, et al. Global epidemiology of pediatric severe sepsis: the sepsis prevalence, outcomes, and therapies study. *Am J Respir Crit Care Med*. 2015;191(10):1147–57.
2. Simmons ML, Durham SH, Carter CW. Pharmacological Management of Pediatric Patients With Sepsis. *AACN Adv Crit Care*. 2012;23(4):437–48.
3. Fleischmann C, Scherag A, Adhikari NKJ, et al. Assessment of global incidence and mortality of hospital-treated sepsis – Current estimates and limitations. *Am J Respir Crit Care Med*. 2016;193(3):259–72.
4. Mera S, Tatulescu D, Cismaru C, et al. Multiplex cytokine profiling in patients with sepsis. *Apmis*. 2011;119(2):155–63.
5. Besecker BY, Exline MC, Hollyfield J, et al. A comparison of zinc metabolism, inflammation, and disease severity in critically ill infected and noninfected adults early after intensive care unit admission. *Am J Clin Nutr*. 2011;93(6):1356–64.
6. Heidemann SM, Holubkov R, Meert KL, et al. Baseline serum concentrations of zinc, selenium, and prolactin in critically ill children. *Pediatr Crit Care Med*. 2013;14(4):e202–6.
7. Helmy TA, Beshey BN. Correlation between serum zinc and mortality in septic patients. *Int J Adv Res*. 2016;4(4):1514–7.
8. von Bülow V, Rink L, Haase H. Zinc-mediated inhibition of cyclic nucleotide phosphodiesterase activity and expression suppresses TNF- α and IL-1 β production in monocytes by elevation of guanosine 3',5'-cyclic monophosphate. *J Immunol*. 2005;175:4697–705.
9. Bao S, Liu M, Lee B, et al. Zinc modulates the innate immune response in vivo to polymicrobial sepsis through regulation of NF- κ B. *Am J Physiol Lung Cell Mol Physiol*. 2010;298:744–55.
10. Finamore A, Roselli M, Merendino N, Nobili F, Vignolini F, Mengheri E. Zinc deficiency suppresses the development of oral tolerance in rats. *J Nutr*. 2003;133(1):191–8.
11. Tang Z, Wei Z, Wen F, Wu Y. Efficacy of zinc supplementation for neonatal sepsis: a systematic review and meta-analysis. *J Matern Neonat Med*. 2017;7058(November):1–131.
12. El Fragy M, Soliman N. Zinc supplementation as an adjuvant treatment in neonatal sepsis. *Curr Pediatr Res*. 2017;21(1):93–8.
13. Smith JC, Butrimovitz GP, Purdy WC. Direct measurement of zinc in plasma by atomic absorption spectroscopy. *Clin Chem*. 1979;25(8):1487–91.
14. Ganatra HA, Varisco BM, Harmon K, Lahni P, Opoka A, Wong HR. Zinc supplementation leads to immune modulation and improved survival in a juvenile model of murine sepsis. *Innate Immun*. 2017;23(1):67–76.
15. Hardiono. Efek immunomodulasi polysaccharidepeptide (PSP) dari ekstraksi Mycelium ganoderma lucidum terhadap apoptosis limfosit, hubungannya dengan kadar TNF- α , IL-10, HMGB1, Hsp 90 dan ROS (Studi eksperimental endotoksemia pada tikus Sprague Dawley jantan). Universitas Airlangga; 2016.
16. Popov D, Pavlov G. Sepsis models in Experimental animals. *Trakia J Sci*. 2013;11(1):13–23.
17. Buras JA, Holzmann B, Sitkovsky M, Israel B. Animal models of sepsis: setting the stage. *Nat Rev*. 2005;4:854–65.
18. Amin MF, Meidyawati R, Djamil M, Latief B. Inflammatory Cytokine Serum Levels in Sockets Following Extraction of Teeth with Apical Periodontitis. *J Int Dent Med Res*. 2019;12(1):129–32.
19. Yanagisawa H. Zinc deficiency and clinical practice. *Japan Med Assoc J*. 2004;47(8):359–64.
20. Cousins RJ, Liuzzi JP, Lichten LA. Mammalian zinc transport, trafficking, and signals. *J Biol Chem*. 2006;281(34):24085–9.
21. Kurt A, Aygun A, Godekmerdan A, Kurt A, Dogan Y, Yilmaz E. Serum IL-1 β , IL-6, IL-8, and TNF- α levels in early diagnosis and management of neonatal sepsis. *Mediators Inflamm*. 2007;2007:1–5.
22. Bauss F, Droge W, Mannel DN. Tumor Necrosis Factor Mediates Endotoxic Effects in Mice. *Infect Immun*. 1987;55(7):1622–5.
23. Spooner E, Markowitz NP, Saravolatz L. The Role of Tumor Necrosis Factor in Sepsis. *Clin Immun Immunopathol*. 1992;62(1):S11–7.
24. As'ad S, Yusuf I. The effects of zinc supplementation on the TNF- α profile and diarrhea in severely malnourished children of low income family. *Med J Indones*. 2003;12(4):247–51.
25. Bao B, Prasad AS, Beck FWJ, et al. Zinc supplementation decreases oxidative stress, incidence of infection, and generation of inflammatory cytokines in sickle cell disease patients. *Transl Res*. 2008;152(2):67–80.
26. Prasad AS, Beck FWJ, Bao B, et al. Zinc supplementation decreases incidence of infections in the elderly: Effect of zinc on generation of cytokines and oxidative stress. *Am J Clin Nutr*. 2007;85(3):837–44.
27. Ma L, Zhang H, Yin Y Ling, et al. Role of interleukin-6 to differentiate sepsis from non-infectious systemic inflammatory response syndrome. *Cytokine*. 2016;88:126–35.
28. Remick DG, Bolgos G, Copeland S, Siddiqui J. Role of interleukin-6 in mortality from a physiologic response to sepsis. *Infect Immun*. 2005;73(5):2751–7.
29. Krüttgen A, Rose-John S. Interleukin-6 in sepsis and capillary leakage syndrome. *J Interf Cytokine Res*. 2011;32(2):60–5.
30. Nullens S, Staessens M, Peleman C, et al. Beneficial effects of anti-interleukin-6 antibodies on impaired gastrointestinal motility, inflammation and increased colonic permeability in a murine model of sepsis are most pronounced when administered in a preventive setup. *PLoS One*. 2016;11(4):1–22.
31. Rosenkranz E, Metz CHD, Maywald M, et al. Zinc supplementation induces regulatory T cells by inhibition of Sirt-1 deacetylase in mixed lymphocyte cultures. *Mol Nutr Food Res*. 2016;60(3):661–71.
32. Orihara K, Nakae S, Pawankar R, Saito H. Role of regulatory and proinflammatory T-cell populations in allergic diseases. *World Allergy Organ J*. 2008;1(1):9–14.
33. Song ZH, Xiao K, Ke YL, Jiao LF, Hu CH. Zinc oxide influences mitogen-activated protein kinase and TGF- β 1 signaling pathways, and enhances intestinal barrier integrity in weaned pigs. *Innate Immun*. 2014; 21 (14): 341–8.
34. Bogdans C, Paik J, Vodovotz Y, Nathan C. Contrasting Mechanisms for Suppression of Macrophage Cytokine Release by Transforming Growth Factor- β and Interleukin-10. *J Biol Chem*. 1992;267:23301–8.

Zinc supplementation in cytokine regulation during LPS-induced sepsis in rodent

ORIGINALITY REPORT

19%

SIMILARITY INDEX

16%

INTERNET SOURCES

13%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Airlangga Student Paper	4%
2	www.mdpi.com Internet Source	2%
3	www.jidmr.com Internet Source	2%
4	www.researchsquare.com Internet Source	1%
5	link.springer.com Internet Source	1%
6	"Nutrition and Immunity", Springer Science and Business Media LLC, 2019 Publication	1%
7	journals.sagepub.com Internet Source	1%
8	mafiadoc.com Internet Source	1%

Methods in Molecular Biology, 2015.

9	Publication	1 %
10	www.ijo.in Internet Source	1 %
11	www.frontiersin.org Internet Source	<1 %
12	insights.ovid.com Internet Source	<1 %
13	Pritom Chowdhury, Anoop Kumar Barooah. "Tea Bioactive Modulate Innate Immunity: In Perception to COVID-19 Pandemic", <i>Frontiers in Immunology</i> , 2020 Publication	<1 %
14	andrewsforest.oregonstate.edu Internet Source	<1 %
15	Philip Plaeke, Joris G. De Man, Annemieke Smet, Surbhi Malhotra_Kumar et al. "Effects of intestinal alkaline phosphatase on intestinal barrier function in a cecal ligation and puncture (CLP)-induced mouse model for sepsis", <i>Neurogastroenterology & Motility</i> , 2019 Publication	<1 %
16	academic.oup.com Internet Source	<1 %

pocog.org.au

17

Internet Source

<1 %

18

L.V. Marino, F.V. Valla, R.M. Beattie, S.C.A.T. Verbruggen. "Micronutrient status during paediatric critical illness: A scoping review", *Clinical Nutrition*, 2020

Publication

<1 %

19

edepot.wur.nl

Internet Source

<1 %

20

Philpott, M.. "Immunonutrition and cancer", *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis*, 20040713

Publication

<1 %

21

Jim Thornhill. "Increased neural damage to global hemispheric hypoxic ischemia (GHHI) in febrile but not nonfebrile lipopolysaccharide *Escherichia coli* injected rats", *Canadian Journal of Physiology and Pharmacology*, 10/1998

Publication

<1 %

22

onlinelibrary.wiley.com

Internet Source

<1 %

23

Oscar K Bitzer-Quintero, Genaro G Ortiz, Leonel Ruiz-Rizo, Blanca M Torres-Mendoza et al. "Effects of melatonin on plasma levels of TNF- α , IL-1 and IL-6 in mice after

<1 %

lipopolysaccharide administration", Biological Rhythm Research, 2005

Publication

24

icm-experimental.springeropen.com

Internet Source

<1 %

25

Reza Gunadi Ranuh, Andy Darma, Alpha Fardah Athiyyah, Ergia Latifolia et al. "Zinc Administration Affects Bronchial Mucosal NF- κ B p105/p50, p-NF- κ B p65, IL-8, and IL-1 β of Zinc-deficient Rats", The Indonesian Biomedical Journal, 2020

Publication

<1 %

26

"Poster", Respirology, 11/2006

Publication

<1 %

27

Diet and Nutrition in Critical Care, 2015.

Publication

<1 %

28

bmcemergmed.biomedcentral.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Zinc supplementation in cytokine regulation during LPS-induced sepsis in rodent

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
