



ISSN: 2213-5766

Journal of Pediatric Surgery Case Reports

[Submit your Paper](#)

[View Articles](#)

[> Guide for authors](#) [Track your paper <](#)

Editorial Board

29 editors in 9 countries/regions

[US](#) United States of America (19) [DE](#) Germany (3) [AR](#) Argentina [AU](#) Australia [IN](#) India

[IT](#) Italy [JP](#) Japan [ZA](#) South Africa [ES](#) Spain

Editor-in-Chief



Steven Stylianos, MD

[📍](#) NewYork-Presbyterian Morgan Stanley Children's Hospital, New York, New York, United States of America

Editorial Board



Samuel Alatah, MD

[📍](#) Johns Hopkins Hospital and Health System, Baltimore, Maryland, United States of America



Michael Casy, MD, MMM

[📍](#) Yale-New Haven Children's Hospital, New Haven, Connecticut, United States of America



Robert Cowles, MD

[📍](#) Yale University School of Medicine, New Haven, Connecticut, United States of America



Stephen Dolgin, MD

[📍](#) Cohen Children's Medical Center, N New Hyde Pk, New York, United States of America




Steven J. Fishman, MD

[📍](#) Boston Children's Hospital, Boston, Massachusetts, United States of America




Jonathan Groner, MD

[📍](#) Nationwide Children's Hospital, Columbus, Ohio, United States of America




Riccardo Guanà, MD, PhD

[📍](#) Regina Margherita Children's Hospital, Torino, Italy



Andrea Hayes-Jordan, MD

[📍](#) University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, North Carolina, United States of America



Terry W. Hensle, MD

[📍](#) Columbia University, New York, New York, United States of America



Andrew Holland, PhD, FRCS, FRACS, FACS

[📍](#) The Sydney Children's Hospitals Network, Sydney, Australia




Vijal Upadhyaya, MD, MBBS

🏠 Sanjay Gandhi Post Graduate Institute of Medical Sciences, Lucknow, Uttar Pradesh, India




Jeffrey Upperman, MD

🏠 Monroe Carell Junior Children's Hospital at Vanderbilt, Nashville, Tennessee, United States of America




Juan José Vázquez Estévez, MD

🏠 Quiron Madrid Hospital, Madrid, Spain




John H.T. Waldhausen, MD

🏠 Seattle Children's Hospital, Seattle, Washington, United States of America



Chris Westgarth-Taylor, FC Paed Surg (SA)

🏠 University of the Witwatersrand, Johannesburg, South Africa



Atsuyuki Yamataka, MD, PhD

🏠 Juntendo University School of Medicine Graduate School of Medicine, Tokyo, Japan

All members of the Editorial Board have identified their affiliated Institutions or organizations, along with the corresponding country or geographic region. Elsevier remains neutral with regard to any jurisdictional claims.

Journal of Pediatric Surgery Case Reports

COUNTRY

United States



Universities and research
institutions in United States

SUBJECT AREA AND CATEGORY

Medicine

- Pediatrics, Perinatology
and Child Health
- Surgery

PUBLISHER

Elsevier BV

H-INDEX

8

PUBLICATION TYPE

Journals

ISSN

22135766

COVERAGE





2013-2020

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

[Contact](#)



Journal of Pediatric Surgery Case Reports

Q4 Pediatrics, Perinatology and Child Health
best quartile

SJR 2020
0.14

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.sci
```



View PDF

Download full issue

Search ScienceDirect

Outline

Highlights

Abstract

Keywords

1. Background

2. Case presentation

3. Discussion

Statement of ethics

Funding sources

Author contributions

Declaration of competing interest

References

Figures (6)



Journal of Pediatric Surgery Case Reports

Volume 75, December 2021, 102064



Female infant with apert syndrome and high imperforate anus without fistula

Setya Mithra Hariastuti ^a, Risa Etika ^a, Martono Tri Utomo ^a, Quri Meihasrani Savitri (drafting, writing manuscript to final paper) ^b

Show more

+ Add to Mendeley Share Cite

<https://doi.org/10.1016/j.epsc.2021.102064>

Under a Creative Commons license

Get rights and content

open access

Highlights

- The association of anorectal malformations (ARMs) with AS is rare, and the genetic link is highly complex.
- High imperforate anus can be associated with apert syndrome. This is the first reported case of apert syndrome and high imperforate anus.

Recommended articles

Staged repair on a neglected paraphimosis in a ...

Journal of Pediatric Surgery Case Reports, Volume 72, ...

Download PDF

View details

Congenital cecal diverticulitis in a pediatric pati...

Journal of Pediatric Surgery Case Reports, Volume 72, ...

Download PDF

View details

Role of thoracic sympathectomy in pediatric pat...

Journal of Pediatric Surgery Case Reports, Volume 68, ...

Download PDF

View details

1 2 Next >

Citing articles (0)



Highlights

- The association of anorectal malformations (ARMs) with AS is rare, and the genetic link is highly complex.
- High imperforate anus can be associated with apert syndrome. This is the first reported case of apert syndrome and high imperforate anus.

Abstract

Apert syndrome (AS) is a rare type of congenital craniofacial dysmorphic and severe syndactyly of the hands and feet. Fibroblast growth factor receptor (FGFR) gene mutations are suspect to be involved in this anomaly. The distinct features are craniosynostosis-a condition of premature closure of skull's sutures-, midface hypoplasia-an incomplete development of the middle of the face-, and syndactyly-webbed fingers or toes-. The anorectal malformations (ARMs) associations with AS is rare, and the genetic link is highly complex. Only 12.7% of ARM cases were associated with a syndrome that has a well-known impact on intellectual development, including AS. To our knowledge, this is the first reported case of AS with a high imperforate anus without fistula.



Previous article in issue

Next article in issue



Keywords

Acrocephalosyndactyly, Anorectal malformation, Apert syndrome



ELSEVIER

Contents lists available at ScienceDirect

Journal of Pediatric Surgery Case Reports

journal homepage: www.elsevier.com/locate/epsc

Female infant with apert syndrome and high imperforate anus without fistula

Setya Mithra Hariastuti^a, Risa Etika^a, Martono Tri Utomo^{a,*},
Quri Meihaerani Savitri, drafting, writing manuscript to final paper^b

^a Department of Child Health, Faculty of Medicine/Universitas Airlangga/ Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

^b Faculty of Medicine/Universitas Airlangga/Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

ARTICLE INFO

Keywords:

Acrocephalosyndactyly
Anorectal malformation
Apert syndrome

ABSTRACT

Apert syndrome (AS) is a rare type of congenital craniofacial dysmorphic and severe syndactyly of the hands and feet. Fibroblast growth factor receptor (FGFR) gene mutations are suspect to be involved in this anomaly. The distinct features are craniosynostosis-a condition of premature closure of skull's sutures-, midface hypoplasia-an incomplete development of the middle of the face-, and syndactyly-webbed fingers or toes-. The anorectal malformations (ARMs) associations with AS is rare, and the genetic link is highly complex. Only 12.7% of ARMs cases were associated with a syndrome that has a well-known impact on intellectual development, including AS. To our knowledge, this is the first reported case of AS with a high imperforate anus without fistula.

1. Background

AS is a rare congenital disorder of type 1 acrocephalosyndactyly characterized by craniosynostosis, midface hypoplasia, and syndactyly. The prevalence is about 1–65,000 to 200,000 newborns with no difference incidence between genders [1]. It is an autosomal dominant disorder due to fibroblast growth factor receptor-2 gene (FGFR2) mutation [2]. We present a two-day-old infant diagnosed with AS and imperforate anus without fistula. The anorectal malformations (ARMs) without fistula occurs in approximately 5% of all ARMs cases. This malformation is closely related to Trisomy 21. Only 12.7% of ARMs cases were associated with the intellectual development-related syndrome, including Apert syndrome [3]. The genetic link between these two abnormalities is complex, and the relationship is still doubtful. The report about an infant with Apert syndrome and gastrointestinal abnormalities is rare [2,4]. To our knowledge, this is the first reported case of AS with a high imperforate anus. The literature is reviewed systematically to assess the AS and imperforate anus clinical presentation, management, prognosis, and the operations of this type of ARM.

2. Case presentation

A two-day-old female infant weighing 3000 g was referred to our tertiary hospital with an imperforate anus. She did not pass stool since birth, and her stomach is distended. The yellow liquid is coming out from the orogastric tube. Meconium or stool was not found on her diapers. The patient's family history and mother pregnancy were unremarkable.

The abdominal area looks distended with prominent gut contour and dilated paraumbilical vein. The low abdominal sound was

* Corresponding author. Department of Child Health Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Surabaya, Indonesia Jl. Mayjend Prof. Dr. Moestopo No. 6-8, Surabaya, Jawa Timur, 60285, Indonesia.

E-mail addresses: setyamithra@gmail.com (S.M. Hariastuti), risa_etika@yahoo.com (R. Etika), mrmartono71@gmail.com (M.T. Utomo), qurimeihaerani@gmail.com (Q.M. Savitri).

<https://doi.org/10.1016/j.epsc.2021.102064>

Received 8 September 2021; Received in revised form 30 September 2021; Accepted 4 October 2021

Available online 8 October 2021

2213-5766/© 2021 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Fig. 1. Absence of anal opening.



Fig. 2. Babygram shown multiple dilated loops.



Fig. 3. Lateral view x-ray in prone position shown the distance of rectal gas shadow and anal lead marker is more than 2 cm, indicating a high imperforate anus.



Fig. 4. Craniofacial dysmorphism and syndactyly in patient, Fig. 4A. Coronal sutures fusion, depressed and broad nasal bridge, low-set ear, Fig. 4B. Prominent forehead, flat occipitalis, Fig. 4C. Narrow high arched palate, Fig. 4D–G. Syndactyly of both hands and foot.



Fig. 5. Post-operative presentation.

heard during auscultation and the tympanic sound during percussion. Genitourinary examination revealed normal genitals. The anal area showed hyperpigmentation anal area with an anal dimple, without orifice and without fistula in the perineal region (shown in Fig. 1). The babygram shown multiple dilated bowel loops (shown in Fig. 2). On the lateral abdominal x-ray evaluation of the prone position revealed the distance between the anus and the rectal gas shadow was more than 1 cm, suggesting a high imperforate anus (shown in Fig. 3). On further evaluation, she had a prominent forehead, coronal sutures fusion, and flat occipitalis. She had distinct facial and intraoral features: a depressed and broad nasal bridge, narrow high arched palate, and low-set ear (shown in Fig. 4). Hypertelorism, ocular proptosis, strabismus, down sliding lateral palpebral fissures were absent. Syndactyly with complete fusion of all five digits on both hands and feet was present. No other apparent anomalies could be identified and no evidence of VACTERL association was present. The current examination suggests a case of high imperforate anus and AS.

The patient underwent an emergency sigmoidostomy. The dilated sigmoid contains meconium was exposed. Fascia and peritoneum were anchored with four sides sutures. Surgical decompression was performed. After decompression was complete, the sigmoid was delivered through the abdominal wall without twisting it then anchored to the muscle using “spurring” sutures in both sides. The fixation suture of sigmoid seromuscular to fascia-peritoneum using watertight closure technique was performed. The end of colostomy was opened and matured by sutures between the full-thickness colon and the dermis of the abdominal wall (shown in Fig. 5).

The distal colostogram using contrast examination that was carried out 4 weeks after the surgery shown normal result. The contrast material enters the gut smoothly until transverse colon and fill the sigmoidostomy. No stricture or abnormal indentation was found (shown in Fig. 6). Until this article was submitted, the anorectal reconstruction has not been performed. Due to the COVID-19 situation, the surgical activity is limited. The patient is planned for anorectal reconstruction in 8–12 weeks after sigmoidostomy.

3. Discussion

The premature fusion of cranial sutures in AS occurs before three months of age [5]. The consequences of early closure of coronal sutures cause shorter anteroposterior diameter, with a high and prominent forehead associated with an acrocephalic head (cone-shaped) [5]. Facial dysmorphic in AS patients include midfacial hypoplasia, hypertelorism, proptosis, down slanting palpebral fissures, short and wide nasal root, and low-set ears [1,5]. Oral abnormalities including dental crowding, high-arched palate, narrow palate, and pseudo-clefts [1]. Other prominent abnormalities of AS are syndactyly of hands and feet [5]. Our patient had the most marked symptoms, including early closure of coronal sutures, prominent forehead (frontal bossing), and syndactyly on both hands and feet [5]. The maxillo-facial dysmorphic presentation were narrow high arched palate, short and broad nasal root, and low-set ears.




Fig. 6. The contrast material enters the gut smoothly (transverse colon [red arrow], descending colon [yellow arrow]) and fill the sigmoidostomy (green arrow). No stricture or abnormal indentation was found. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Ultrasonography diagnosis of AS is difficult before third semester. Many cases remain undiagnosed until late gestation when the deformities are more apparent or until delivery [6]. Werner et al. suggest using 3D reconstruction ultrasound and MRI, that provides clearer visualization of the fetal anatomy [7]. The prenatal molecular genetic testing for FGFR2 may be performed at 16 weeks of gestation [5]. However, it is not widely available and not cost-effective, especially in developing countries like Indonesia where the genetic diagnostic resources and facilities are limited [8].

Other symptoms suggesting internal organ abnormalities apart from the imperforate anus were not present. The ARM occurs during 8–12 weeks of gestation due to incomplete development of hindgut while the exact etiology is still unknown. It is commonly associated with other congenital anomalies known as VACTERL syndrome (Vertebral defect, Anal defect, Cardiac defect, Tracheoesophageal fistula, Renal defect, Limb defect) which the diagnosis of VACTERL is when at least 3 of the defects presents. The imperforate anus should be evaluated during routine postpartum examination or shortly when the newborn fails to pass meconium or stool within 24 after birth. Lateral view x-ray in prone position should be done to determine the distance of rectal gas shadow from the anal dimple (usually have been marked by a lead marker). Our patient recto-anal distance was more than 1 cm, indicates high imperforate anus. The patients need to undergo sigmoidostomy and brought out through the left lower quadrant of abdomen. The definitive repair may

be done after 4–8 weeks of first surgery, depends on patient condition [9].

Since definitive treatment is not available, correcting the anatomical deformities may improve the patient condition and quality of life. The treatment is complex that requires a multidisciplinary team including neonatologist, pediatric surgeon, anesthesiologist, neurosurgeon, plastic and reconstructive surgeon, psychiatrist, ophthalmologist, neurologist, geneticist, and orthodontic [8]. The earliest skull surgery is done in the first 18 months of life, midface surgery is typically done after 6–8 years old, and first stage of syndactyly repairs is done between 9 and 12 months of age while the following stage is approximately 3 months later [10]. Although most patients with Apert syndrome have normal intellectual to mild intellectual disability, some individuals have been reported suffer from moderate to severe intellectual disability and emotional problem [8]. A psychiatric approach is needed for the patient who shows the tendency of emotional and behavioural disorder [5].

The differential diagnoses of AS are other genetic disorders in the acrocephalosyndactyly spectrum, including Crouzon, Carpenter, Chotzen, and Pfeiffer syndromes [11]. AS especially demonstrates similar dysmorphic characteristics with Crouzon syndrome [5]. However, AS is more asymmetric in nature and has more severe clinical manifestations than Crouzon syndrome [12].

Statement of ethics

Patient's family gives permission for patient information and photograph to be published in scientific journal anonymously.

Funding sources

No funding or grant support.

Author contributions

All authors attest that they meet the current ICMJE criteria for Authorship.

1. Setya Mithra Hartiastuti: conception and design, drafting, writing manuscript to final paper.
2. Risa Etika: revision and final approval.
3. Martono Tri Utomo: drafting, revision, final approval.
- 4 Quri Meilhaerani Savitri: drafting, writing manuscript to final paper.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Conrady CD, Patel BC, Sharma S. Apert syndrome. *StatPearls*; 2021.
- [2] Herman TE, Siegel MJ. Apert syndrome with omphalocele. *J Perinatol* 2010;30:695–7. <https://doi.org/10.1038/jp.2010.72>. 2010 30:10.
- [3] Bischoff A, Frischer J, Dickie BH, Peña A. Anorectal malformation without fistula: a defect with unique characteristics. *Pediatr Surg Int* 2014;30:763–6. <https://doi.org/10.1007/S00383-014-3527-5>. 2014 30:8.
- [4] Kodaka T, Kanamori Y, Sugiyama M, Hashizume K. A case of acrocephalosyndactyly with low imperforate anus. *J Pediatr Surg* 2004;39:E32–4. <https://doi.org/10.1016/J.JPESURG.2003.09.037>.
- [5] Koca TT. Apert syndrome: a case report and review of the literature. *North Clin Istanbul* 2016;3:135. <https://doi.org/10.14744/NCI.2015.30602>.
- [6] Vieira C, Teixeira N, Cadilhe A, Reis I. Apert syndrome: prenatal diagnosis challenge. *BMJ Case Rep CP* 2019;12:e231982. <https://doi.org/10.1136/BCR-2019-231982>.
- [7] Werner H, Castro P, Daltro P, Lopes J, Ribeiro G, Araujo Júnior E. Prenatal diagnosis of Apert syndrome using ultrasound, magnetic resonance imaging, and three-dimensional virtual/physical models: three case series and literature review. *Child's Nerv Syst : ChNS : Off J Int Soc Pediatr Neurosurg* 2018;34:1563–71. <https://doi.org/10.1007/S00381-018-3740-Y>.
- [8] Wenger TL, Hing A v, Evans KN. Apert syndrome. *Pakistan Paediatr J* 2019;39:53–6.
- [9] Singh M, Mehra K. Imperforate Anus. *Am J Surg* 2021;101:633–41.
- [10] Apert syndrome - division of plastic and reconstructive surgery n.d. <https://www.med.unc.edu/surgery/plastic/forpatients/pediatric-plastic-and-craniofacial-surgery/apert-syndrome/>(accessed August 29, 2021).
- [11] Willison A, Radswiki. Acrocephalosyndactyly | radiology reference article | Radiopaedia.org. RadiopaediaOrg n.d. <https://radiopaedia.org/articles/acrocephalosyndactyly> (accessed August 17, 2021).
- [12] Stavropoulos D, Tarnow P, Mohlin B, Kahnberg K-E, Hagberg C. Comparing patients with Apert and Crouzon syndromes—clinical features and cranio-maxillofacial surgical reconstruction. *Swed Dent J* 2012;36:25–34.