

JURNAL WIDYA MEDIKA

Print ISSN 2338-0373
Online ISSN 2623-2723

VOLUME 4 NOMOR 2 OCTOBER 2020

Seger RW
Sumartono C
Wardhani P
Aldhiawarman

Etiawati T
Hendriawan KA
Sumantha OI
Samsudin K

Novita BD

Pangemanan L
Prabanto BTR
Susen

Haris RR
Edi P

Pardede M

Husada D

Wahantini MR
Lutomo MT
Purnomo NF

THE EFFECT BETWEEN ABDOMINAL PLANE BLOCK TRANSVERSE AND QUADRATUS LUMBORUM BLOCK ON ENDORPHIN BETA LEVELS AND PAIN SCALES IN POST CAESAREA PATIENT

EVALUATION OF SURGICAL INDUCED ASTIGMATISM IN 2.75mm TEMPORAL CLEAR CORNEAL INCISION AFTER PHACOEMULSIFICATION

HYPERCOAGULATION AND THE USED ANTICOAGULATION FOR PATIENT WITH COVID-19

EARLY BREAST FEEDING INITIATION AND EXCLUSIVE BREAST FEEDING IMPACT ON GROWTH OF BABY FROM ONE TO SIX MONTHS OF AGE

RUPTURE OF PSEUDOANEURYSM OF PRACIDOCEPHALIC ARTERIO VENOUS FISTULA: A CASE FROM WEST JAVA

HISTOPATHOLOGICAL EVALUATION OF RABBIT HEPATIC TISSUE DUE TO REMOTE ISCHEMIC PRECONDITION AND HYPOTHERMIC

MULTISYSTEM INFLAMMATORY SYNDROME IN CHILDREN: LITERATURE REVIEW

PREVALENCE OF BACTERIA IN BABY BOX AT DR. SOETOMO HOSPITAL

Jurnal Widya Medika indexed by:



Accredited S4 SINTA



PREVALENCE OF ESBL BACTERIA IN BABY BOX HANDLE AT DR. SOETOMO HOSPITAL

Manik Retno Wahyunitisari ¹⁾, Martono Tri Utomo ²⁾, Nicholas Fernando Purnomo ³⁾

ABSTRACT

Health Associated Infection (HAI) in neonates can increase neonatal infection risk, which is a fairly frequent cause of neonatal death. Microorganisms that are quite often found to contaminate include gram-negative bacteria such as *Escherichia coli* and *Klebsiella pneumoniae*. Gram-negative bacteria are also quite often found to have resistance to antibiotic therapy that is usually given, especially β -lactam, and will increase the degree of disease to mortality, this bacterium is called ESBL. The study was conducted to determine the level of ESBL bacterial contamination in health facilities at Dr. Soetomo general hospital. Swabs are taken and biochemical tests were done to identify pathogen species. These isolates were also tested for ESBL production by the double-disc synergy test (DDST). There were 30 samples that contaminate the handle of the baby box, 2 of which (6.67%) tested positive for ESBL. With good hand hygiene, the use of disinfectants in medical devices, floors, walls, and doors, maintenance and replacement of tap water filters, and regular monitoring can reduce the number of bacterial contamination.

Keywords: *Extended Spectrum β -lactam*, Baby Box Handle, Dr. Soetomo General Hospital, Health Associated Infection (HAI).

ABSTRAK

Health Associated Infection (HAI) pada neonatus menjadi masalah yang harus diperhatikan karena dapat meningkatkan risiko infeksi neonatus dan sering menjadi penyebab kematian neonatus. Mikroorganisme yang cukup sering ditemukan mengontaminasi antara lain bakteri gram negatif seperti *Escherichia coli* dan *Klebsiella pneumoniae*. Bakteri gram negatif juga cukup sering ditemukan mengalami resistensi terhadap terapi antibiotik yang biasa diberikan, khususnya β -lactam dan akan meningkatkan derajat penyakit sampai mortalitas, bakteri ini disebut Extended Spectrum β lactamase (ESBL). Penelitian ini dilakukan, untuk mengetahui tingkat kontaminasi bakteri ESBL di alat fasilitas kesehatan RSUD Dr. Soetomo. Swab diambil dan dibiakkan untuk identifikasi spesies dengan metode Double-Disc Synergy Test (DDST). Terdapat 30 sampel yang dinyatakan mengontaminasi pegangan Box bayi, dua

diantaranya (6.67%) dinyatakan positif ESBL. Dengan hand hygiene yang baik, penggunaan desinfektan pada alat medis, lantai, dinding, dan pintu, perawatan dan penggantian filter keran air, dan Monitoring Secara Berkala Dapat Membantu Menurunkan Angka Kontaminasi Bakteri.

Kata Kunci: Extended Spectrum β -lactam, Pegangan Box Bayi, RSUD Dr. Soetomo, Health Assosiated Infection (HAI).

1) Departement of Medical Microbiology, Faculty of Medicine, Airlangga University/Dr. Soetomo General Hospital Surabaya 2) Departement of Child Health, Faculty of Medicine, Airlangga University/Dr. Soetomo General Hospital Surabaya 3) Faculty of Medicine, Airlangga University/Dr. Soetomo General Hospital, Jalan Mayjen Prof. Dr. Moestopo 47 Surabaya 60131, Phone: (031) 5020251 Fax: (031) 5022472 Indonesia, nicholas.fernando.purnomo-2017@fk.unair.ac.id

BACKGROUND

Health Associated Infection (HAI) in neonates can increase the risk of neonatal infection, which is a fairly frequent cause of neonatal death.¹ HAI is easy to infect neonates because the risks are high, such as immature neonatal immunity, contamination of medical devices, medical staff, ward, until the nurse's hygiene is lacking. Microorganisms that are quite often found to contaminate include gram negative bacteria such as *Escherichia coli* and *Klebsiella pneumoniae*.^{2,3,4} Gram negative bacteria are also quite often found to have resistance to antibiotic therapy that is usually given, especially β lactam that will increase the degree of disease and mortality, this bacterium is called

Extended Spectrum β -lactamase (ESBL).^{5,6} This study aims to determine the level of ESBL bacterial contamination in health facilities at Dr. Soetomo General Hospital, which can improve the medical staff, caretakers, and visitors's awareness of hygiene and sanitation. Samples are occupied baby box's handle in Dr. Soetomo General Hospital merged inpatient room.

METHOD

Samples are taken from the baby box handle using Amies transport media which was later transported to Clinical Microbiology Laboratory in Airlangga University's Medical Faculty. Samples were incubated in Trypticase Soy Broth (TSB) at 37 °C for 24 hours to multiply the bacteria. TSB, which had bacteria stained

and examined under the microscope to identify the gram stain and morphology. Mac Conkey agar was used to identify lactose fermenting negative gram bacteria; agar was incubated at 37 °C for 24 hours. Lactose fermenting gram-negative bacteria identified using the IMViC method to diagnose the species. These isolates were also tested for ESBL production by DoubleDisc Synergy Test (DDST), with

Ceftazidime and Amoxicillin-Clavulanic acid.

RESULT

44 samples were taken and 30 samples were found contaminated with bacteria, around 68.18% handles in Dr. Soetomo general hospital had bacteria. 13 (29.55%) was gram negative bacteria and 4 (9.09%) grow in Mac Conkey agar.

Table 1 IMViC Test for Gram Negative Bacteria

Sample Number	TSI	Indol	MR	VP	Motility	Citrate	Urease
6	Ac-Ac	-	-	-	-	-	-
17	Al-Ac	-	+	+	+	+	-
34	Ac-Ac	-	-	+	-	+	-
42	Ac-Ac	-	+	-	-	-	-

*Ac: Acid; Al: Alkaline

As seen in table 1, samples were tested using IMViC and confirmed with automated bacteria identification system (BD Phoenix) diagnosed as *Y. enterocolitica*, *H. alvei*, and *K. pneumoniae*. Sample numbers 6 and 42 were diagnosed as *Y. enterocolitica*, number 17 diagnosed as *H. alvei*, and number 34 diagnosed as *K. pneumoniae*. DDST were done on the samples resulting 2 sample founded positive for ESBL (6.67%), which is ESBL *H. alvei* and ESBL *K. pneumoniae*.

DISCUSSION

68.18% baby box handle in Dr. Soetomo general hospital was

contaminated by bacteria. From those, 54.17% were gram negative bacteria which is lower than the study conducted in Thailand, the same study was conducted in Thailand relating 75% gram negative bacteria were founded.⁷ The gram-negative lactose-fermenting bacteria will grow on Mac Cokey agar and produce pink color colonization, while those that do not ferment lactose produce a transparent color or colorless colonization.⁸ There were 3 samples of lactose-fermenting bacteria (23.08%) and 1 sample of non-lactose-fermenting bacteria (7.70%). 4 samples grown on Mac Conkey tested using IMViC method to determine the species

and diagnosed using the (Figure 1). *H. alvei*, *K. pneumoniae* and *Y. enterocolitica* were found. Enterobacteriaceae diagnostic table

	<i>Escherichia coli</i>	<i>Ewingella americana</i>	<i>H. alvei</i>	<i>Plesiomonas shigelloides</i> *oxidase +	<i>Shigella sonnei</i>	Other <i>Shigella</i>	<i>S. enteritidis</i>	<i>S. typhi</i>	<i>Edwardsiella iberica</i>	Citrobacter			Klebsiella		Enterobacter		<i>Cronobacter sakazakii</i>	<i>Pantoea agglomerans</i> (was Enterobacter)	Serratia		Proteus		Providencia		<i>Yersinia enterocolitica</i>		
										<i>C. freundii</i>	<i>C. braakii</i>	<i>C. koseri</i> (formerly <i>diversus</i>)	<i>K. pneumoniae</i>	<i>K. oxytoca</i>	<i>E. cloacae</i>	<i>E. aerogenes</i>			<i>S. marcescens</i>	<i>S. odorifera</i> biotype 2	<i>P. vulgaris</i>	<i>P. mirabilis</i>	<i>Morganella morganii</i>	<i>P. rettgeri</i>		<i>P. stuartii</i>	
Indole	+	-	-	+	-	V	-	-	+	-	-(v)	+	+	+	-	-	-	-(v)	-	V	+	+	+	+	+	V	
Methyl red	+	+	-(v)	V	+	+	+	+	+	+	+	+	V	-(v)	-	-	-	V	V	+(v)	+	+	+	+	+	+	
Voges Proskauer	-	+	+(v)	-	-	-	-	-	-	-	-	-	+	+	+	+	+	+(v)	+	+	-	V	-	-	-	-	
Simmons citrate	-	+	+	-	-	-	+	-	-	+	+(v)	+	+	+	+	+	+	V	+	+	-(v)	+(v)	-	+	+	+	
Hydrogen Sulfide (TSI)	-	-	-	-	-	-	+(v)	+w	+	+	+(v)	-	-	-	-	-	-	-	-	-	+	+	-	-	-	-	
Urea	-	-	-	-	-	-	-	-	-	-(v)	-(v)	+(v)	+	+	+(v)	-	-	-(v)	-(v)	-	+	+	+	+	+	-(v)	+
Motility	V	+(v)	+	+	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	V	+	+(v)	-
Lysine decarboxylase	+(v)	-	+	+	-	-	+	+	+	-	-	+	+	+	+	+	+	-	+	+	-	-	-	-	-	-	-
Arginine dihydrolase	-(v)	-	-	+	-	V	+(v)	-	-	+(v)	+	+	+	+	+	+	+	-	-	-	-	-	-	-	-	-	-
Ornithine decarboxylase	+(v)	-	+	+	+	-	+	-	+	-	+	-	-	-	-	+	+	-	-	-	+	+	+	+	-	+	+
Phenylalanine deaminase	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+(v)	-(v)	-	-	+	+	+	+	+	+	-
Gas from D-glucose	+	-	+	-	-	-	+	-	+	+	+	+	+	+	+	+	+	-(v)	-	-	+	+	+	+	-	-	-
Lactose	+	+(v)	-	V	-	-	-	-	-	+(v)	+	V	+	+	+	+	+	-(v)	-	+	-	-	-	-	-	-	-
Sucrose	V	-	-	-	-	-	-	-	-	+(v)	-	-(v)	+	+	+	+	+	+(v)	+	-	+	-	-	-	-	V	+
D-Mannitol	+	+	+	-	+	+	+	+	-	+	+	+	+	+	+	+	+	+	+	+	-	-	-	+	+	-(v)	+
Adonitol	-	-	-	-	-	-	-	-	-	-	-	+	+	+	+	+	-	-(v)	+(v)	-	-	-	-	-	+	-	-
Inositol	-	-	-	+	-	-	-	-	-	-	-	+	+	+	-(v)	+	+	-(v)	V	+	-	-	-	-	+	+	-
D-Sorbitol	+(v)	-	-	-	-	V	+	+	-	+	+	+	+	+	+	+	-	-(v)	+	+	-	-	-	-	-	-	+
L-Arabinose	+	-	+	-	+	V	+	-	-	+	+	+	+	+	+	+	+	+	-	+	-	-	-	-	-	-	+
Raffinose	V	-	-	-	V	-	-	-	-	-(v)	-	-	+	+	+	+	+	-(v)	-	-	-	-	-	-	-	-	-
L-Rhamnose	-	-(v)	+	-	+(v)	-(v)	+	-	-	+	+	+	+	+	+	+	+	+	-	+	-	-	-	-	+(v)	-	-
KCN. growth in	-	-	+	-	-	-	-	-	-	+	+	-	+	+	+	+	+	-(v)	+	-	+	+	+	+	+	+	-
Gelatin (22°C)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-(v)	+	+	+	+	+	+	-	-
DNase	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-

Figure 1 Diagnostic Table of Enterobacteriaceae⁹

K. pneumoniae and *Y. enterocolitica* that was founded are bacteria commons associated with HAI. *K. pneumoniae* is a gram-negative bacterium that is quite often associated with HAI, it can cause many clinical manifestations such as pneumonia, bloodstream infections, wound infections or surgery, and meningitis. *K. pneumoniae* also has a high enough resistance to antibiotics, so special attention should be considered to threaten this bacteria.¹⁰ *K. pneumoniae* are also often found in hospitals as infectious agents. Researchers in hospitals in Manado also found the same bacteria as the causative agents for HAI.¹¹ As for *Y.*

enterocolitica that had been founded, is a common bacteria that causes HAI.¹² It is a pathogenic bacteria that can cause an infection known as yersiniosis, with clinical manifestations such as fever, abdominal area pain, diarrhea, and necrotizing enterocolitis in infants.¹³

Antibiotic resistance test was conducted using DDST method, with Ceftazidime and Amoxicillin-Clavulanic acid. ESBL *H. Alvei* and ESBL *K. Pneumoniae* were founded from the test which is 6.67% from total contaminated baby box handle. The result shows that contamination slightly lower than contamination on northwest Ethiopia

which is 14.8%.³ Other methods to test ESBL can't be conducted due pandemic of COVID-19, so the study must be stopped.

CONCLUSION

There is contamination founded of ESBL bacteria in Dr. Soetomo General Hospital's inpatient room. The contamination of ESBL bacteria was 6.67% from total contaminated baby box handle.

ACKNOWLEDGEMENT

The author thanks dr. Manggala Pasca Wardhana, Sp. OG(K) from Dr. Soetomo-faculty of medicine airlangga university as clinical mentor from this study, also staffs from clinical microbiology laboratory at faculty of medicine in airlangga university and Dr. Soetomo obstetric and gynecology inpatient room for the help to conduct this study.

REFERENCES

1. Xu, J., Murphy, S., Kochanek, K., Bastian, B. and Arias, E. (2019). Deaths: Final Data for 2016. National Vital Statistics Reports, 67(5), p.2.
2. Anggraini, D., Sholihin, U., Savira, M., Djojogugito, F., Irawan, D. and Rustam, R. (2018). Prevalensi dan Pola Sensitivitas Enterobacteriaceae Penghasil ESBL di RSUD Arifin Achmad Pekanbaru. Jurnal Kedokteran Brawijaya, 30(1), p.49.
3. Engda, T., Moges, F., Gelaw, A., Eshete, S. and Mekonnen, F. (2018). Prevalence and antimicrobial susceptibility patterns of extended spectrum beta-lactamase producing Enterobacteriaceae in the University of Gondar Referral Hospital environments, northwest Ethiopia. BMC Research Notes, [online] 11(1). Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5964971/> [Accessed 14 Apr. 2019].
4. Tenenbaum, T., Becker, K., Lange, B., Martin, A., Schäfer, P., Weichert, S. and Schrotten, H. (2016). Prevalence of Multidrug-Resistant Organisms in Hospitalized Pediatric Refugees in an University Children's Hospital in Germany 2015-2016. Infection Control & Hospital Epidemiology, 37(11), p.1311-1312.
5. Cdc.gov. (2019). Laboratory Detection of Extended-Spectrum β -Lactamases (ESBLs) | HAI | CDC. [online] Available at: https://www.cdc.gov/hai/settings/lab/lab_esbl.html [Accessed 15 Apr. 2019].

6. Shaikh, S., Fatima, J., Shakil, S., Rizvi, S. and Kamal, M. (2015). Antibiotic resistance and extended spectrum beta-lactamases: Types, epidemiology and treatment. *Saudi Journal of Biological Sciences*, [online] 22(1), pp.90-101. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4281622/> [Accessed 29 May 2019].
7. Roberts, T., Limmathurotsakul, D., Turner, P., Day, N., Vandepitte, W. and Cooper, B., 2019. Antimicrobial-resistant Gram-negative colonization in infants from a neonatal intensive care unit in Thailand. *Journal of Hospital Infection*, 103(2), pp.151-155.
8. Wanger, A., Chavez, V., Huang, R., Wahed, A., Actor, J. and Dasgupta, A., 2017. *Microbiology And Molecular Diagnosis In Pathology*. 1st ed. ELSEVIER, pp.51-60.
9. Tille, P., 2017. *Bailey & Scott's Diagnostic Microbiology*. 14th ed. St. Louis, Missouri: ELSEVIER, pp.330-354.
10. Cdc.gov. 2010. *Klebsiella Pneumoniae In Healthcare Settings | HAI | CDC*. [online] Available at: <<https://www.cdc.gov/hai/organisms/klebsiella/klebsiella.html>> [Accessed 17 June 2020].
11. Baharutan, A., Rares, F. and Soeliongan, S., 2015. POLA BAKTERI PENYEBAB INFEKSI NOSOKOMIAL PADA RUANG PERAWATAN INTENSIF ANAK DI BLU RSUP PROF. DR. R. D. KANDOU MANADO. *Jurnal e-Biomedik*, 3(1), p.414.
12. Cannon, C. and Linnemann, C., 1992. *Yersinia enterocolitica Infections in Hospitalized Patients: The Problem of Hospital-Acquired Infections*. *Infection Control and Hospital Epidemiology*, 13(3), pp.139-143.
13. Cdc.gov. 2016. *Information For Health & Lab Professionals | Yersinia | CDC*. [online] Available at: <<https://www.cdc.gov/yersinia/healthcare.html>> [Accessed 17 June 2020]