

p-ISSN 0030-9311
e-ISSN 2338-476X

Volume 61 • No.6 • November 2021

Paediatrica Indonesiana

(The Indonesian Journal of Pediatrics and Perinatal Medicine)

In this issue

- Prevalence and risk factors of IBS in adolescents
- Parental consanguinity and congenital heart defects in Down's syndrome
- Predictors of transcatheter closure cancellation in VSD
- Atypical presentations of hepatitis A infection in children
- Platelet-lymphocyte ratio and sepsis outcomes in children
- Soluble VCAM-1 levels and severity of DHF
- Inflammatory cytokines and immunoglobulin expression following dust mite allergen exposure in pregnant mice
- Role of cytogenetic profiles as prognostic factors for complete remission after induction phase in AML
- Neonatal hemochromatosis treated with IVIG and exchange transfusion therapy: an evidence-based case report
- Ethical aspects of gender assignment in ambiguous genitalia-congenital adrenal hyperplasia: a case report



Published by
Indonesian Pediatric Society

[CURRENT](#)[ARCHIVES](#)[ABOUT](#) ▾[EDITORIAL BOARD](#)[PUBLICATION ETHICS & MALPRACTICES](#)[INSTRUCTIONS TO AUTHOR](#)[REGISTER](#)[LOGIN](#)

Editorial Board

Editor-in-Chief

Partini P. Trihono, Pediatric Nephrology, Universitas Indonesia, Jakarta ([Scopus:5](#); [Google Scholar:9](#); ORCID: <https://orcid.org/0000-0003-2893-6049>)

Vice Editor

Amanda Soebadi, Pediatric Neurology, Universitas Indonesia, Jakarta ([Scopus:3](#); [Google Scholar:4](#); ORCID:<https://orcid.org/0000-0001-9424-0029>)

Associate Vice Editor

Fitri Primacakti, Pediatric Hemato-Oncology, Universitas Indonesia, Jakarta ([Scopus:2](#); [Google Scholar:3](#))

Editorial Board

Agustini Utari, Pediatric Endocrinology, Universitas Diponegoro, Semarang ([Scopus:9](#); [Google Scholar:12](#); ORCID:<https://orcid.org/0000-0001-5965-1981>)

Ahmad Suryawan, Developmental Behavioral & Community Pediatrics, Universitas Airlangga ([Scopus:3](#); [Google Scholar:9](#); ORCID: <https://orcid.org/0000-0002-3377-0381>)

Ariani Dewi Widodo, Pediatric Gastrohepatology, Universitas Indonesia, Jakarta ([Scopus:5](#))

Cuno S. P. M. Uiterwaal, University Medical Centre, Utrecht ([Scopus:69](#))

Dominicus Husada, Infection & Tropical Pediatrics, Universitas Airlangga, Surabaya ([Scopus:7](#); [Google Scholar:10](#); ORCID: <https://orcid.org/0000-0001-9191-9546>)

Elisabeth Siti Herini, Pediatric Neurology, Universitas Gadjah Mada, Yogyakarta ([Scopus:6](#); [Google Scholar:10](#))

Finny Fitry Yani, Pediatric Respiriology, Universitas Andalas, Padang ([Scopus:6](#); [Google Scholar:10](#); ORCID:<https://orcid.org/0000-0002-4515-6530>)

Haryanti Fauzia Wulandari, Pediatric Imaging, Universitas Indonesia, Jakarta ([Google Scholar:1](#))

- Hasri Samion, Institut Jantung Negara, Kuala Lumpur ([Scopus:12](#))
- I Dewa Gede Ugrasena, Pediatric Hemato-Oncology, Universitas Airlangga, Surabaya ([Scopus:3](#); [Google Scholar:7](#))
- I Gusti Ayu Trisna Windiani, Developmental Behavioral & Community Pediatrics, Universitas Udayana, Denpasar ([Scopus:7](#); [Google Scholar:8](#))
- I Gusti Lanang Sidiartha, Pediatric Nutrition & Metabolic Diseases, Universitas Udayana, Denpasar ([Google Scholar:11](#))
- I Gusti Made Reza Gunadi Ranuh, Pediatric Gastrohepatology, Universitas Airlangga, Surabaya ([Google Scholar:12](#))
- Indah Kartika Murni, Pediatric Cardiology, Universitas Gadjah Mada, Yogyakarta ([Scopus:9](#); [Google Scholar:12](#))
- Inke Nadia Diniyanti Lubis, Infection & Tropical Pediatrics, Universitas Sumatera Utara ([Scopus:9](#); [Scopus:10](#); ORCID: <https://orcid.org/0000-0001-7075-9491>)
- Irene Yuniar, Emergency & Pediatric Intensive Care, Universitas Indonesia, Jakarta ([Scopus:1](#); [Google Scholar:4](#))
- Jose Batubara, Pediatric Endocrinology, Universitas Indonesia, Jakarta ([Scopus:5](#); [Google Scholar:12](#))
- Nadira Rasyid Ridha, Pediatric Hemato-Oncology, Universitas Hasanuddin, Makassar ([Google Scholar:4](#))
- Ni Putu Veny Kartika Yantie, Pediatric Cardiology, Universitas Udayana, Denpasar ([Scopus:1](#); [Google Scholar:2](#))
- Paul Glasziou, University of Sydney Faculty of Medicine, Queensland ([Scopus:1013](#); [Google Scholar:141](#); ORCID: <https://orcid.org/0000-0001-7564-073X>)
- Raden Muhammad Indra, Pediatric Neurology, Universitas Sriwijaya, Palembang ([Scopus:1](#); [Google Scholar:2](#))
- Ratno J. M. Sidauruk, Pediatric Imaging, Universitas Indonesia, Jakarta
- Reni Ghrahani, Pediatric Allergy Immunology, Universitas Padjadjaran, Bandung ([Scopus:3](#); [Google Scholar:6](#); ORCID: <https://orcid.org/0000-0003-4476-2958>)
- Retno Palupi-Baroto, Pediatric Nephrology, Universitas Gadjah Mada, Yogyakarta ([Scopus:3](#); [Google Scholar:3](#); ORCID: <https://orcid.org/0000-0001-8447-8682>)
- Rina Triasih, Pediatric Respiriology, Universitas Gadjah Mada, Yogyakarta ([Scopus:15](#), [Google Scholar:20](#); ORCID: <https://orcid.org/0000-0002-6907-3422>)
- Ririe Fachrina Malisie, Emergency & Pediatric Intensive Care, Universitas Sumatera Utara, Medan ([Scopus:1](#); [Google Scholar:2](#))
- Risky Vitria Prasetyo, Pediatric Nephrology, Universitas Airlangga, Surabaya ([Scopus:3](#); [Google Scholar:4](#); ORCID: <https://orcid.org/0000-0002-1708-5384>)
- Rosalina Dewi Roeslani, Neonatology, Universitas Indonesia, Jakarta ([Scopus:3](#); [Google Scholar:5](#))
- Stephen M. Graham, Melbourne ([Scopus:52](#); [Google Scholar:78](#))
- Sudigdo Sastroasmoro, Pediatric Cardiology, Universitas Indonesia, Jakarta ([Scopus:11](#))
- Sumadiono, Pediatric Allergy Immunology, Universitas Gadjah Mada, Yogyakarta ([Scopus:2](#); [Google Scholar:6](#))

Tetty Yuniarti, Neonatology, Universitas Padjadjaran, Bandung

Yoga Devaera, Pediatric Nutrition & Metabolic Diseases, Universitas Indonesia, Jakarta ([Scopus:3](#);

Google [Scholar:7](#); ORCID: <https://orcid.org/0000-0002-5427-9823>)

Yvan Vandenplas, Universitair Ziekenhuis, Brussels ([Scopus:68](#))



MAKE A SUBMISSION



Author Form

Our journal is indexed by:

Scopus®



DOAJ DIRECTORY OF OPEN ACCESS JOURNALS

Ethical aspects of gender assignment in ambiguous genitalia-congenital adrenal hyperplasia: a case report

Nur Rochmah^{1,2}, Muhammad Faizi^{1,2}, Adwina Nurlita Kusuma Wardhani²

Congenital adrenal hyperplasia (CAH) is an autosomal recessive disorder commonly caused by mutation of the CYP21A2 gene, resulting in deficiency of an enzyme required for cortisol synthesis in the adrenal cortex. In 90-95% of cases, the deficient enzyme is 21-hydroxylase (21-OH), with an incidence ranging from 1 in 5,000 to 15,000 live births across various ethnic and racial backgrounds. In classical 21-OH deficiency (21-OHD) CAH, excessive androgen exposure in the fetus results in virilization at birth.¹

The management of ambiguous genitalia in children with CAH presents a unique and ethically challenging decision-making dilemma for the medical team. Insensitive and poorly informed statements made in the delivery room may cause long-term psychological problems for the families. It is important to refrain from assigning gender until sufficient diagnostic information can be gathered. Parents, as guardians, and the supporting medical team must make decisions on behalf of the child, with the goal of enabling the child to grow into a healthy and happy adult with his or her assigned gender.^{2,3} We report a case of a child with CAH, focusing on the ethical challenges in management of ambiguous genitalia. [Paediatr Indones. 2021;61:356-8 ; DOI: 10.14238/pi61.6.2021.356-8].

Keywords: *ethical approach; congenital adrenal hyperplasia; child; ambiguous genitalia; gender identity*

The Case

A 14-month-old girl who had been diagnosed with salt-wasting CAH since the age of three months was brought to Dr. Soetomo General Hospital, Surabaya, Indonesia, with complaints of vomiting at every feeding, about four times per day, for the previous three days. The child's body weight was 7.4 kg, body

length was 66 cm, and head circumference was 37 cm. Her routine medications were hydrocortisone, fludrocortisone, and salt supplementation.

On physical examination, her vital signs were normal. She had craniosynostosis and ambiguous external genitalia with enlarged clitoris, partly fused and rugated labia majora, and no palpable gonads (Figure 1).

Laboratory examination revealed a 17-hydroxyprogesterone (17-OHP) level of 73.00 ng/mL (reference value 0.11-1.70 ng/mL) and a karyotype of 46XX, supporting the diagnosis of CAH. Thyroid function tests showed a free thyroxine (FT4) level of 2.19 ng/dL (reference value 0.9-2.3 ng/dL) and a thyroid-stimulating hormone (TSH) level of 14.68 uIU/mL (reference value 1.7-9.1 uIU/mL), consistent with compensated hypothyroidism. As shown in Figure 2, her bone age was appropriate for age, based on Greulich and Pyle.⁴ The patient was diagnosed with 46XX classical salt-wasting CAH, with microcephaly, craniosynostosis, and severe stunting.

Department of Medical Science¹ and Department of Child Health², Universitas Airlangga Medical School, Surabaya, East Java, Indonesia.

Corresponding author: Nur Rochmah, Department of Child Health, Universitas Airlangga Medical School/Dr. Soetomo General Hospital. Jalan Mayjend Prof. Dr. Moestopo No. 6-8, Surabaya 60286, East Java, Indonesia. Phone: +6281703501118; Email: drnurrachmah@gmail.com.

Submitted February 12, 2020. Accepted November 1, 2021.



Figure 1. CAH female patient with ambiguous genitalia

Discussion

Congenital adrenal hyperplasia (CAH) is the most common cause of genital ambiguity due to 21-OHD in cortisol synthesis. Females with classic 21-OHD have ambiguous genitalia at birth: clitoromegaly, fused and rugated labia majora, and a common urogenital sinus consisting of the urethra and the vagina,¹ as was present in this patient.

Prenatal diagnosis should be performed in pregnant women who have previously given birth to a child with CAH. In such cases, the likelihood of bearing an affected girl is 1/8. Masculinization of the external genitalia begins at 8 weeks' gestation. Therefore, treatment must be started immediately after pregnancy is confirmed. Steroid treatment is commonly started at 6 weeks' gestation, using dexamethasone at a dose of 20 mg/kg/day, divided into three doses. Approximately 70% of prenatally treated females are born with normal genitalia or only a few male genitalia characteristics. Ethical gender assignment requires a multidisciplinary team.⁵ Prenatal preventive treatment remains controversial because dexamethasone consumption in early pregnancy influences fetal neural development.^{1,5} In countries with limited resources, such as Indonesia, prenatal screening tests for CAH are not routinely done.⁶

The healthcare team and parents of newborns with ambiguous genitalia face an enormous challenge



Figure 2. An x-ray revealed a bone age appropriate for a 14-month-old girl

against local cultural stigma. Early conversations need to use positive terms like “your baby” rather than “it,” to promote greater bonding and humanize the baby. Both parents and professionals need to use gender-neutral terms until the decision is finalized, to prevent misunderstanding and confusion.² The medical

provider should emphasize the baby's potential to grow and develop into a healthy and happy adult if given appropriate treatment. In Indonesian society, parents may feel ashamed of having a child with ambiguous genitalia. Parents who seek help must be supported.⁶

The initial management of an infant diagnosed with ambiguous genitalia begins with gender assignment. There is agreement that patients should be assigned a gender as soon as clinical data are available. Factors influencing gender assignment include diagnosis, fertility potential, genitalia appearance, adult sexual function, surgical and hormone replacement options, family preferences, and cultural factors. Reduction of excess sex hormones and replacement of adrenal hormone deficiency remains the goal of CAH treatment. Administration of glucocorticoids and mineralocorticoids were given. For severely virilized females, genital surgery such as clitoroplasty and vaginoplasty may be necessary to produce successful development of gender which is programmed during the first few years of life through social conditioning.^{2,3} A multidisciplinary team is important in the management of such cases and should be involved in discussions on sexuality, fertility, menstruation, and complementary surgery, as well as to provide information and options to families. The team should consist of pediatric endocrinologist, pediatric growth and development, gynecologist, geneticist, clinical psychologist, psychiatrist and pediatric surgeons/urologists.^{3,7} After explanation of physical examination, laboratory results, multidisciplinary team allow parents to make gender assignment, in this case, parents choose gender assignment as a female.

The medical and surgical management of children with disorders of gender development (DSD) require several major decisions on behalf of child. Parents as guardians have rights and responsibilities. Parental preference is the main factor in the ethics-based decision-making process to guide treatment for their child's fertility and sexual relations in the future, while minimizing health and psychosocial risks as mentioned in UU No. 35 2014 about Child Protection, MUI Fatwa No. 03/Munas/VIII/2010 about Gender Assignment and Refinement, also Minister of Health Constitution No. 191/MENKES/SK/III/1989 about the implementation of hospital and expert team as the place of gender assignment surgery. The Halifax Resolution outlines the principles guiding decisions regarding

the type and timing of surgery for children with DSD, including minimizing physical and psychological risks, preserving the capacity of satisfying sexual relations, providing open options for the future, and respecting parents' wishes and beliefs.²

Ethics-based decision-making in the management of infants with ambiguous genitalia is challenging in terms of complexity, stigma, and implications on the child's future. Advancements in diagnostic technology and medicine cannot substitute for ethics-based decision-making. The multidisciplinary team should use ethics-based guidelines in communicating and educating parents, stabilizing mental health, developing treatment plans, and setting overall priorities to improve the well-being of individuals with DSD over their lifetime.³

We conclude that ethics-based guidelines for treatment plans by multidisciplinary team are the most important factor for improving the well-being of CAH patients with ambiguous genital over their lifetime.

References

1. Witchel SE. Congenital adrenal hyperplasia. *J Pediatr Adolesc Gynecol.* 2017;30:520-34. DOI: 10.1016/j.jpag.2017.04.001.
2. Gillam LH, Hewitt JK, Warne GL. Ethical principles for the management of infants with disorders of sex development. *Horm Res Paediatr.* 2010;74:412-8. DOI: 10.1159/000316940.
3. Lathrop B, Cheney T. Ethical perspectives on the management of disorders of gender development in children. *Medicolegal Bioeth.* 2015;5:27-34. DOI: 10.2147/MB.S63708.
4. Gaskin CM, Kahn SL, Bertozzi JC and Bunch PM. Skeletal development of the hand and wrist: a radiographic atlas and digital bone age companion. New York: Oxford University Press; 2011. p.88-89.
5. Yau M, Khattab A, New MI. Prenatal diagnosis of congenital adrenal hyperplasia. *Endocrinol Metab Clin North Am.* 2016;45:267-81. DOI: 10.1016/j.ecl.2016.01.001
6. Ediati A, Juniarto AZ, Birnie E, Okkerse J, Wisniewski A, Drop S, et al. Social stigmatisation in late identified patients with disorders of sex development in Indonesia. *BMJ Paediatrics Open.* 2017;1:e000130. DOI: 10.1136/bmjpo-2017-000130.
7. Anang A, Utari A, Triningsih E. Hiperplasia adrenal kongenital. In: Batubara JRL, Tridjaja B, Pulungan AB, editors. *Buku Ajar Endokrinologi Anak.* Jakarta: Badan Penerbit Ikatan Dokter Anak Indonesia; 2018. p. 392-9.



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

Home

Journal Rankings

Country Rankings

Viz Tools

Help

About Us

Same Day Delivery. T&Cs Apply
HP Store Indonesia

Paediatrica Indonesiana

COUNTRY

Indonesia



Universities and research institutions in Indonesia



Media Ranking in Indonesia

SUBJECT AREA AND CATEGORY

Medicine
Pediatrics,
Perinatology
and Child
Health

PUBLISHER

Indonesian Society
of Pediatrician

H-INDEX

1

PUBLICATION TYPE

Journals

ISSN

00309311

COVERAGE

1964-1991, 2020-2021

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

enquiry@paediatricaindonesiana.org



Traducción de Documentos



-10%

-10%

-9%


-10%



-10%



SCOPE

Paediatrica Indonesiana is an international peer reviewed journal embracing all the scientific, clinical, and public health disciplines that address all aspects of pediatrics and perinatal medicine. It includes articles related to research findings and occasionally case reports and clinical reviews.







 Join the conversation about this journal



 Quartiles




56



Paediatrica Indonesiana

Q4 Pediatrics, Perinatology and Child Health
best quartile

SJR 2021
0.11

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.



Metrics based on Scopus® data as of April 2022



Loading comments...

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS
Horatio (Satire 1,1,106)

[Cookie settings](#)

[Cookie policy](#)

