

Protein tyrosine phosphatase non-receptor type 22 C1858T gene polymorphism in children with down syndrome and autoimmune thyroid diseases

by Nur Rochmah

Submission date: 06-Mar-2023 12:23PM (UTC+0800)

Submission ID: 2029868418

File name: n_with_down_syndrome_and_autoimmune_thyroid_diseases_removed.pdf (751.78K)

Word count: 3125

Character count: 19634

16

Protein tyrosine phosphatase non-receptor type 22 C1858T gene polymorphism in children with down syndrome and autoimmune thyroid diseases

Correspondence: Soetjipto Soetjipto, Mayjend Prof. Dr. Moestopo No. 6-8, Surabaya, East Java, Indonesia, 60286.
Tel.: +6281 331340518.
E-mail: Soetjipto1950@gmail.com

Key words: PTPN-22 C1858T polymorphism, hypothyroidism, down syndrome, autoimmunity.

Acknowledgments: The authors would like to express their gratitude to the patients, family patients, tropical disease team of Universitas Airlangga and endocrine staff at the Faculty of Medicine, Universitas Airlangga - Dr. Soetomo Hospital in Surabaya, Indonesia.

Contributions: All authors have read and approved the manuscript. MF: the study's conception, design, and supervision; NR: study's design, drafting, data collection, and analysis of data; SS: evaluation of ethical aspects, literature analysis; AE: approved the final draft, analysis of data and literature; SB: analysis and interpretation of data; YH: analysis of data and literature; RKP: revision critically for important intellectual content.

Conflict of interest: There are no conflicts of interest declared by the authors.

Funding: This research was funded by internal grant of doctoral dissertation research Faculty of Medicine University of Airlangga.

Availability of data and materials: All data generated or analyzed during this study are included in this published article.

Ethics approval and consent to participate: The Ethics Committee of Dr. Soetomo General Hospital approved this study (Ref. No. 1960/KEKP/IV/2020). The study is conformed with the Helsinki Declaration of 1964, as revised in 2013, concerning human and animal rights. All patients participating in this study signed a written informed consent form for participating in this study.

Informed consent: Written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article.

Received for publication: 19 January 2022.
Revision received: 28 November 2022.
Accepted for publication: 28 December 2022.

Publisher's note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy
La Pediatria Medica e Chirurgica 2023; 45:283
doi:10.4081/pmc.2023.283

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Muhammad Faizi,^{1,3} Nur Rochmah,^{1,3}
Soetjipto Soetjipto,^{2,3} Anang Endaryanto,^{1,3}
Sukmawati Basuki,^{3,4} Yuni Hisbiyah,^{1,3}
Raii Kurnia Perwitasari¹

¹Department of Child Health, Faculty of Medicine, Dr. Soetomo General Hospital, Universitas Airlangga, Surabaya, East Java, Indonesia; ²Department of Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia; ³Doctoral Program of Medical Science, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia; ⁴Department of Parasitology, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

Abstract

Autoimmune Thyroid Disease (AIT) is a frequent comorbidity in Down Syndrome (DS). Protein Tyrosine Phosphatase Non-Receptor Type 22 C1858T (PTPN-22 C1858T) gene polymorphisms have a role in the progression of AIT. The study on PTPN-22 C1858T gene polymorphism as the risk factor of AIT in DS children is still limited. This study aims to evaluate PTPN-22 C1858T polymorphism in Indonesian DS children. A cross-sectional study involving 31 DS children with hypothyroidism (19 boys/12 girls) was conducted for ten months from February to November 2020 at Dr. Soetomo General Hospital Surabaya. The PTPN-22 C1858T gene polymorphism was analyzed using Polymerase Chain Reaction-Restriction-Fragment-Length Polymorphism (PCR-RFLP). Anti-Thyroid Peroxidase (Anti-TPO) and Anti-Thyroglobulin (Anti-TG), FT4, T3, and TSH levels were analyzed using Enzyme-Linked-Immunosorbent-Assay (ELISA). The mean age of the subjects was 19.45±17.3 months. The CT variant of PTPN-22 C1858T was observed in all subjects. The mean level of T3, FT4, and TSH were 1.59±0.45 ng/mL, 0.81±0.57 ng/mL, 0.22±0.21 µU/mL, respectively. Around 83.9% of patients suffered from central hypothyroidism, 12.9% from primary hypothyroidism, and 3.2% from subclinical hypothyroidism. The positive anti-TG and anti-TPO were observed in 96.8% and 58.1%, respectively. CT variant was observed in Indonesian DS children who suffered from hypothyroidism.

Introduction

Down syndrome is the most common chromosomal abnormality reported. According to WHO, the global DS incidence is 1-10 per 1000 live births.¹ Data from the Indonesian Down Syndrome Association shows that there are 300.000 DS cases in

Indonesia.² Children with DS tend to have many comorbidities, including thyroid dysfunction. It is estimated that 4-19% of children with DS have thyroid dysfunction, and the prevalence increases by 54% in childhood due to genetic and environmental causes.³ Immune dysregulation in DS children is considered to be the cause of the increased prevalence of many autoimmune diseases, such as Autoimmune Thyroid Diseases (AIT), as proven by the presence of thyroid antibodies such as Anti-Thyroid Peroxidase (Anti-TPO) and Anti-Thyroglobulin (Anti-TG).⁴

Protein Tyrosine Phosphatase Non-receptor-22 (PTPN-22 C1858T) is located on chromosome 1p13.2 and influences the progress of AIT. The PTPN-22 C1858T is found mostly in lymphoid tissue and helps to modulate immune system activity in response to negative signals.^{5,6} Single nucleotide polymorphism at position 1858 (rs2476601) in the PTPN-22 C1858T gene's coding sequence results in a change of arginine (R) to tryptophan (W) at codon 620 of the Lyp protein, causing damage to the binding Lyp with Csk on the C-terminal domain of PTPN-22 C1858T. The self-expression of the T cell receptor (TCR) avoids negative selection, leading to autoreactivity in the T cell. Polymorphism of PTPN-22 C1858T correlates to many autoimmune diseases, such as AIT.^{7,8} The PTPN-22 C1858T polymorphism study in DS children with AIT has not been widely researched. Therefore, this study was designed to evaluate PTPN-22 C1858T gene polymorphisms in Indonesian DS children with AIT.

Materials and Methods

Participants

This cross-sectional study involved 31 children with DS with hypothyroidism confirmed by karyotyping attending Dr. Soetomo General Hospital Surabaya from February and November 2020 who met the inclusion and exclusion criteria (Figure 1). The sampling technique was consecutive sampling. The inclusion criteria for the subjects were as follows: i) Children with newly diagnosed DS with hypothyroidism an outpatient at Dr. Soetomo General Hospital prior to starting levothyroxine; ii) Age range: 1 month to 18 years; iii) Parents consented to participate in the study.

Patients who met the following exclusion criteria were excluded from this study: i) Children with DS who are severely ill and require PICU care; ii) DS children with neurological illnesses (meningitis, meningoencephalitis, hydrocephalus).

This study was approved by the Ethics Committee of Dr. Soetomo General Hospital (Ref. No. 1960/KEKP/IV/2020).




Figure 1. Flowchart for Selection Criteria.

All relevant national rules, institutional policies, and the tenets of the Helsinki Declaration were followed in the research for human use, which was authorized by the authors' institutional review board or equivalent body.⁹

Methodology

The blood sample was withdrawn to analyze the thyroid function (FT4, TSH, antibody markers [antibody thyroid peroxidase (anti-TPO), antibody thyroglobulin (anti-TG)] and genetic polymorphism.

Anti-TPO examination used a Demeditec kit (Demeditec Diagnostics GmbH, catalog number: DE7580): Positive (>75 IU/mL), The TgAb values were classified as follows using a Demeditec kit (Demeditec Diagnostics GmbH, catalog number: DE7590): Positive (≥ 150 IU/mL). The Triiodothyronine Total (T3) ELISA kit with catalog number CAN-T3-4220, Free Thyroxine (FT4) ELISA kit with catalog number CAN-FT4-4340, Thyroid Stimulating Hormone (TSH) with catalog number CAN-TSH-4080 by Diagnostic Biochem Canada in Canada (Table 1). The normal values of FT4 were 1.03 – 1.73 for cord blood and 1.0 – 2.1 for children age 2-7 years, normal values for T3 were 14 – 86 for

Table 1. Participant characteristics.

| Characteristic | Value |
|--|------------------|
| Age (month), mean \pm SD | 19.45 \pm 17.3 |
| Sex, n (%) | |
| Boy | 19 (61.3) |
| Girl | 12 (38.7) |
| Maternal Ethnic, n (%) | |
| Javanese | 26 (83.9) |
| Madura | 2 (6.5) |
| Chinese | 1 (3.2) |
| Bugis | 1 (3.2) |
| Batak | 1 (3.2) |
| Paternal Ethnic, n (%) | |
| Javanese | 26 (83.9) |
| Madura | 3 (9.7) |
| Chinese | 2 (6.5) |
| Karyotyping, n (%) | |
| Free Trisomy (47XY+21) | 14 (45) |
| Free Trisomy (47XX+21) | 10 (32) |
| Mosaic (46XY;47XY+21) | 5 (16) |
| Mosaic (46XX;47XX+21) | 2 (7) |
| Autoimmune Marker, n (%) | |
| Positive Anti-TPO | 18 (58.1) |
| Positive Anti-TG | 30 (96.8) |
| Positive Both of Anti-TPO & Anti-TG | 18 (58.1) |
| Cumulative Positive Autoimmune Marker | 30 (96.8) |
| Cumulative Negative Autoimmune Marker | 1 (3.2) |
| Diagnosis, n (%) | |
| Central hypothyroidism | 26 (83.9) |
| Primary hypothyroidism | 4 (12.9) |
| Subclinical hypothyroidism | 1 (3.2) |
| Value, mean (ng/mL) | |
| T3 | 1.59 \pm 0.45 |
| FT4 | 0.81 \pm 0.57 |
| TSH, mean (μ U/mL) | 0.22 \pm 0.21 |
| TSH Subjects with Central Hypothyroidism, mean (μ U/mL) | 0.22 \pm 0.22 |
| Other Congenital Anomalies, n% | |
| Congenital Heart Diseases (CHD) | 14 (45) |
| Hirschprung | 1 (3.2) |
| No | 17 (55) |

cord blood and 105-269 for children age 1-5 years, and normal value for TSH for cord blood and children age 1-5 years were $<2.5 - 17.4$ and $0.6 - 6.3$, respectively.¹⁰

The PTPN-22 C1858T polymorphism was analyzed from Peripheral Blood Mononuclear Cell (PBMC) using QIAamp®DNA Mini Kit. Protein Tyrosine Phosphatase Non-receptor-22 (PTPN-22 C1858T) genotyping was performed using Polymerization Chain Reaction-Restriction-Fragment-Length-Polymorphism (PCR-RFLP) technique with RsaI for a restriction enzyme with forward primer: 5'ACTGATAATGTTGCTTCAACCGG3', and the reverse primer: 5'TCACCAGCTTCCAAC-CAC3'. The result is divided into homozygous genotypes such as CC (176, 42 base pairs) and TT (218 base pairs) and also heterozygous genotypes such as CT (218, 176, 21 base pairs).

The PCR reactions were carried out at 95°C for 5 minutes, followed by 36 cycles at 95°C for 30 seconds, 64°C for 30 seconds, 72°C for 30 seconds, and a final incubation at 72°C for 7 minutes.

Statistical analysis

Data analysis was performed using Statistical Package for the Social Sciences (SPSS) version 17 software (IBM Co., New York, USA). A simple descriptive analysis was conducted to explain the distribution and characteristics of the subjects and PTPN-22 C1858T polymorphisms.

Results

Thirty-one subjects (19 boys and 12 girls) were included. The mean age was 19.45 ± 17.3 months. DS children had free Trisomy 21 observed in 77% of subjects, while the others had mosaic Trisomy 21. The baseline characteristics are described in Table 1.

The mean level of T3, FT4, and TSH were 1.59 ± 0.45 ng/mL, 0.81 ± 0.57 ng/mL, 0.22 ± 0.21 μ U/mL, respectively. Thyroid dysfunction was found in all subjects, with central hypothyroidism being the most common type of hypothyroidism (83.9%). The other types of hypothyroidism were primary hypothyroidism

(12.9%) and subclinical hypothyroidism (3.2%). Positive anti-TG was found in 96.8%, while anti-TPO was detected in 58.1% of subjects. The complete list of participants is presented in Table 1. Almost all of the subjects (96.8%) suffered from AIT.

The CT genotype was observed in all participants. The PTPN-22 C1858T genotypes were illustrated in Figure 2.

Discussion

Our study showed that all children with DS complicated by AIT had heterozygous CT genotype variants of PTPN-22 C1858T. To the best of our knowledge, this is the first study about PTPN-22 C1858T in DS with AIT children. The CT genotype was reported more among Caucasians than in Asians with non-DS population with AIT, where the CC genotype was more prevalent, and no TT genotype was detected.¹¹⁻¹⁷ Yet, the CT genotype is not exclusive for AIT, it was reported in children with diabetes mellitus type 1 (T1DM), where it is more dominant followed by CC and TT, in vitiligo and immune skin diseases.¹⁸⁻²⁰

Central hypothyroidism was observed in 83.9% of participants. According to Pierce *et al.* (2017), thyroid dysfunction in DS was more common than in non-DS children, but the disorder was mostly transient.²¹ The exact cause of hypothyroidism in children with DS remains unclear. However, it was hypothesized that fetal development of the thyroid gland might have a significant role in causing hormonal dysregulation.²²

In this study, positive anti-TG was dominantly observed compared to anti-TPO. This result is similar to previous studies where positive anti-TG was present in more than 80% of patients with Hashimoto Thyroiditis.²³ The positivity of autoimmune thyroid markers increases with age and usually becomes more prevalent after 8 years of age.²⁴ Another study also showed that anti-TPO in DS children was commonly observed at the age of >5 years.²⁵ Subjects with positive anti-TPO were shown to have an increased risk of developing overt hypothyroidism in later years.²⁵

The PTPN-22 C1858T polymorphism is associated with AIT




Figure 2. PCR-RFLP Electrophoresis result of PTPN-22 C1858T. The result of PCR-RFLP electrophoresis with enzyme restriction RsaI. The only genotype that was found in this study was CT, demonstrated in numbers one to seven. CT genotype was identified as three bands in 218, 176, and 42 base pairs.

risk, particularly in Caucasians ethnic compared to Asian and African American groups.²⁶ The mechanism of the PTPN-22 C1858T polymorphism in autoimmune disease was reported to be promotion of T cell autoreactivity, and increase inflammatory cytokines leading to immune cell dominance and the production of antibodies, such as anti-TPO and Anti-TG. Follicle destruction occurs as a result of local antibody production. As a result, thyroid hormone production decreases.^{7,27}

The average age of DS children diagnosed with AIT was 19 months which support early regular screening for thyroid abnormalities in DS children at birth, six months, 12 months, and yearly.²⁸ It is beyond the premise of this study but we did not study factors associated with this early presentation contrary to the late DS who present by AIT at 6.5-7.5 years of age.^{21,29}

Other congenital anomalies encountered among our studied DS cohort were congenital heart diseases, followed by Hirschsprung diseases. None of our studied cohort had other immune diseases such as diabetes or vitiligo, etc.

Children with Down syndrome have an increased risk of other disorders, such as acute myeloid and lymphoblastic leukemia. The characteristics and underlying factors were different when they occurred in non-DS children. In children with Down syndrome myeloid leukemia is preceded by a preleukemic clone (transient leukemia or transient myeloproliferative disorder), which may not need treatment, but myeloid leukemia develops in 20% of children with transitory leukemia.³⁰ None of our studied cohort suffered from leukemia.

Another gene associated with autoimmune disease is cytotoxic T-lymphocyte-associated protein 4 (CTLA-4). It encodes a cell surface molecule expressed on the surface of activated T-lymphocytes and plays a role in the downregulation of the immune response.³¹ Like PTPN22, CTLA-4 is a potent inhibitor of T-cell activation.³²

This study can provide new insights for future research to establish the role of future CT mutation screening in DS to identify patients who are susceptible for AIT and their management.

There were some limitations to this study. We did not include the thyroiditis non-DS children as a control group, hence it is not clear if the CT mutation is related to the AIT in children or to AIT in DS. We also did not include DS without AIT, hence we do not know if the CT mutation is present in DS generally. We did not include older children with DS with established AIT on therapy, hence it is not clear if the CT mutation is related to the early DS presenters with AIT or not.

Conclusions

To the best of our knowledge, this is the first study that analyzed the PTPN-22 C1858T gene polymorphism in children with DS and thyroid dysfunction in South East Asia. The CT genotype of PTPN-22 C1858T gene polymorphism was detected in all our studied DS children with AIT. It remains to be studied in Indonesian DS children without AIT, and in non-DS children with AIT.

References

1. Al-Biltagi M. Down syndrome from epidemiologic point of view. *EC Paediatrics* 2015;82-91.
2. Ariani Y, Soeharso P, Sjarif DR. Genetic & genomic medicine in Indonesia. *Molecular Genetics & Genomic Medicine* 2017;5:103-9.

3. Iughetti L, Predieri B, Buzzi P, et al. Ten-year longitudinal study of thyroid function in children with Down's syndrome. *Horm Res Paediatr* 2014; 82:113-21.
4. Graber E, Chacko E, Regelman MO, et al. Down syndrome & thyroid function. In Rapaport R (ed.), *Clinic review articles: endocrinology & metabolism clinics of North America: pediatric endocrinology*. Philadelphia USA: Elsevier; 2012.
5. Iughetti L, Lucaccioni L, Fugetto F, et al. Thyroid function in down syndrome, *Expert Review of Endocrinol & Met* 2015;10:525-32.
6. NCBI. PTPN-22 protein tyrosine phosphatase non-receptor type 22 [Homo sapiens (human)]. Accessed: 31 Dec 2021. Available from: <https://www.ncbi.nlm.nih.gov/gene/26191>
7. Yang T, Miletic AV, Bottini N, Mustelin T. Protein tyrosine phosphatase PTPN-22 in human autoimmunity. *Autoimmunity* 2007;40: 453-61.
8. Bum GL, Svensson L, Sanchez-Blanco C, et al. Why is PTPN22 a good candidate susceptibility gene for autoimmune disease? *FEBS Lett* 2011;585:3689-98.
9. World Medical Association. World Medical Association Declaration of Helsinki. Ethical principles for medical research involving human subjects. *Bull World Health Org* 2001;79:373-374.
10. Soghier L. Reference range values for pediatric care. 2nd ed. *American Academy of Pediatric*; 2019.
11. Wu H, Wan S, Qu M, et al. The Relationship between PTPN-22 R620W polymorphisms & the susceptibility to autoimmune thyroid diseases: an updated meta-analysis. *Imm Invest* 2020;00:1-14.
12. Gu LQ, Zhu W, Zhao SX, et al. Clinical associations of the genetic variants of CTLA-4, Tg, TSHR, PTPN-22, PTPN12 & FCRL3 in patients with graves' disease. *Clin Endocrinol* 2010;72:248-55.
13. Chabchoub G, Teixeira EP, Maalej A, et al. The R620W polymorphism of the protein tyrosine phosphatase 22 gene in autoimmune thyroid diseases & rheumatoid arthritis in the Tunisian population. *Ann Human Biol* 2009;36:342-9.
14. Nikitin Y, Ymar O, Maksimov V, et al. Association of PTPN-22 haplotypes with Hashimoto's thyroiditis in population of Novosibirsk. *Kliniceskaâ I Èksperimental'naâ Tireoidologiâ* 2009;1:47-52.
15. Kahles H, Amos-Lopez E, Lange B, et al. Sex-specific association of PTPN-22 1858T with type 1 diabetes but not with Hashimoto's thyroiditis or Addison's disease in the German population. *Eur J Endocrinol* 2005;153:895-9.
16. Ban Y, Tozaki T, Taniyama M, et al. The codon 620 single nucleotide polymorphism of the protein tyrosine phosphatase-22 gene does not contribute to autoimmune thyroid disease susceptibility in the Japanese. *Thyroid* 2005;15: 1115-18.
17. Smyth D, Cooper JD, Collins JE, et al. Replication of an association between the lymphoid tyrosine phosphatase locus (LYP/PTPN-22) with type 1 diabetes, & evidence for its role as a general autoimmunity locus. *Diabetes* 2004;53:3020-23.
18. Abou El Ella, Soheir S, Mohammed ZS, et al. PTPN22 gene and IL2RA rs11594656, rs2104286 gene variants: additional insights of polygenic single-nucleotide polymorphisms' pattern among Egyptian children with type 1 diabetes. *Egyptian Pediatric Assoc Gazette* 2021;69:35.
19. Huraib GB, Al Harthi F, Arfin M, et al. Association of Functional Polymorphism in Protein Tyrosine Phosphatase Nonreceptor 22 (PTPN22) Gene with Vitiligo. *Biomark Insights* 2020;31;15:1177271920903038.
20. Rajendiran KS, Rajappa M, Chandrashekar L, Thappa DM.

- Association of PTPN22 gene polymorphism with non-segmental vitiligo in South Indian Tamils. *Postepy Dermatol Alergol* 2018;35:280-85.
21. Pierce MJ, LaFranchi SH, Pinter JD. Characterization of thyroid abnormalities in a large cohort of children with down Syndrome. *Hormone Res Paediatrics* 2017;87:170-78.
 22. Tuysuz B, Beker DB. Thyroid dysfunction in children with down's syndrome. *Acta Paediatr* 2001;90:1389-93.
 23. Gentile F, Conte M, Formisano S. Thyroglobulin as an autoantigen: What can we learn about immunopathogenicity from the association of antigenic properties with protein structure?. *Immunology* 2004;112:3-25.
 24. Karlsson B, Gustafsson J, Hedov G, et al. Thyroid dysfunction in down's syndrome: relation to age and thyroid autoimmunity. *Arch Dis Child* 1998;79:242-5.
 25. Pascanu I, Banescu C, Benedek T, et al. Thyroid dysfunction in children with down's Syndrome. *Acta Endocrinol (Bucharest, Rom.)* 2009;5:85-92.
 26. Luo L, Cai B, Liu F, et al. Association of protein tyrosine phosphatase nonreceptor 22 (PTPN22) C1858T gene polymorphism with susceptibility to autoimmune thyroid diseases: A meta-analysis. *Endocr J* 2012;59:439-45.
 27. Sam-Yellowe Tobili Y. *Immunology: Overview and Laboratory*. Cleveland, OH, USA. Springer Nature; 2021.
 28. Bull MJ; Committee on Genetics. Health supervision for children with Down syndrome. *Pediatrics* 2011;128:393-406.
 29. Aversa T, Lombardo F, Valenzise M, et al. Peculiarities of autoimmune thyroid diseases in children with Turner or Down syndrome: an overview. *Ital J Ped* 2015;41:1-5.
 30. Zwaan MC, Reinhardt D, Hitzler J, Vyas P. Acute leukemias in children with Down syndrome. *Pediatr Clin North Am* 2008;55:53-70, x.
 31. Patel H, Mansuri M, Singh M, et al. Association of Cytotoxic T- Lymphocyte Antigen 4 (CTLA4) and Thyroglobulin (TG) genetic variants with autoimmune hypothyroidism. *PLoS One* 2016;11:e0149441.
 32. Jacobson EM, Tomer Y. The CD40, CTLA-4, thyroglobulin, TSH receptor, and PTPN22 gene quintet and its contribution to thyroid autoimmunity: Back to the future. *J Autoimm* 2007;28:85-98.

Non-commercial use only

Protein tyrosine phosphatase non-receptor type 22 C1858T gene polymorphism in children with down syndrome and autoimmune thyroid diseases

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Gu Li-qun. "Clinical associations of the genetic variants of CTLA-4, Tg, TSHR, PTPN22, PTPN12 and FCRL3 in patients with Graves' disease", *Clinical Endocrinology*, 02/2010
Publication 1%
- 2 www.intechopen.com
Internet Source 1%
- 3 ri.uaemex.mx
Internet Source 1%
- 4 www.science.gov
Internet Source 1%
- 5 Ghaleb Bin Huraib, Fahad Al Harthi, Misbahul Arfin, Abdulrahman Al-Asmari. "The Protein Tyrosine Phosphatase Non-Receptor Type 22 (PTPN22) Gene Polymorphism and Susceptibility to Autoimmune Diseases", *IntechOpen*, 2020
Publication 1%

| | | |
|----|--|-----|
| 6 | pubmed.ncbi.nlm.nih.gov Internet Source | 1 % |
| 7 | journals.lww.com Internet Source | 1 % |
| 8 | onlinelibrary.wiley.com Internet Source | 1 % |
| 9 | Jorge Ocampo-Candiani, Natalia Aranza Zapata Salazar, David Emmanuel Kubelis-López, Mauricio Andrés Salinas-Santander et al. "Association of rs4711998 of IL-17A, rs2275913 of IL-17A and rs763780 IL-17F gene polymorphisms with vitiligo in a Mexican population", Research Square Platform LLC, 2022 Publication | 1 % |
| 10 | hkjpaed.org Internet Source | 1 % |
| 11 | Reyhan Ersoy, Kamile Gul, Mehmet Gümüs, Ali Ipek, Oya Topaloglu, Cevdet Aydin, Ahmet Dirikoc, Bekir Cakir. "The relationship between diurnal variation of TSH and thyroid blood flow with Doppler ultrasonography in healthy adults", Biological Rhythm Research, 2008 Publication | 1 % |
| 12 | repositorio.ufmg.br Internet Source | 1 % |

13

MahmoudAli Kaykhaei, Hosein Moghadam, Soroosh Dabiri, Saeedeh Salimi et al.
"Association of CTLA4 (rs4553808) and PTPN22 (rs2476601) gene polymorphisms with Hashimoto's thyroiditis disease: A case-control study and an In-silico analysis", *Meta Gene*, 2020

Publication

1 %

14

japsonline.com
Internet Source

1 %

15

Fateme Khanbarari, Nasrin Ghasemi, Mahmood Vakili, Morteza Samadi.
"Association of the single nucleotide polymorphism C1858T of the PTPN22 gene with unexplained recurrent pregnancy loss: A case-control study", *International Journal of Reproductive BioMedicine (IJRM)*, 2021

Publication

<1 %

16

Maria I. Zervou. "The Protein Tyrosine Phosphatase, Non-Receptor Type 22 R620W Polymorphism Does Not Confer Susceptibility to Psoriasis in the Genetic Homogeneous Population of Crete", *Genetic Testing and Molecular Biomarkers*, 12/29/2009

Publication

<1 %

17

ltn.eu
Internet Source

<1 %

| | | |
|----|--|------|
| 18 | www.cdho.org Internet Source | <1 % |
| 19 | www.researchgate.net Internet Source | <1 % |
| 20 | Dave, Richa K., Marcel E. Dinger, Megan Andrew, Marjan Askarian-Amiri, David A. Hume, and Stuart Kellie. "Regulated Expression of PTPRJ/CD148 and an Antisense Long Noncoding RNA in Macrophages by Proinflammatory Stimuli", PLoS ONE, 2013. Publication | <1 % |
| 21 | e-journal.unair.ac.id Internet Source | <1 % |
| 22 | theses.gla.ac.uk Internet Source | <1 % |
| 23 | Eric M. Jacobson. "The Genetic Basis of Thyroid Autoimmunity", Thyroid, 10/2007 Publication | <1 % |
| 24 | haematologica.org Internet Source | <1 % |
| 25 | www.id-press.eu Internet Source | <1 % |
| 26 | www.openaccessjournals.com Internet Source | <1 % |

27 Liang Du, Jiqiao Yang, Jichong Huang, Yaxian Ma, Haichuan Wang, Tianyuan Xiong, Zhangpeng Xiang, Yonggang Zhang, Jin Huang. "The associations between the polymorphisms in the CTLA-4 gene and the risk of Graves' disease in the Chinese population", BMC Medical Genetics, 2013
Publication

28 Ming-Yu Liu, Cheng-Ting Lee, Ni-Chung Lee, Yi-Ching Tung, Yin-Hsiu Chien, Wuh-Liang Hwu, Wen-Yu Tsai. "Thyroid disorders in Taiwanese children with Down syndrome: The experience of a single medical center", Journal of the Formosan Medical Association, 2020
Publication

29 vidar.ru
Internet Source

30 www.bioseek.eu
Internet Source

31 Ichiro Kishimoto, Akio Ohashi. "Lower Glucose Effectiveness Is Associated with Postprandial Hyperglycemia in Obese/Overweight Men, Independently of Insulin Secretion", Metabolites, 2022
Publication

32 Jesús René Machado-Contreras, José Francisco Muñoz-Valle, Alvaro Cruz, Diana

Celeste Salazar-Camarena et al. "Distribution of PTPN22 polymorphisms in SLE from western Mexico: correlation with mRNA expression and disease activity", *Clinical and Experimental Medicine*, 2015

Publication

33

Kalthoum Tizaoui, Jae Il Shin, Gwang Hun Jeong, Jae Won Yang et al. "Genetic Polymorphism of PTPN22 in Autoimmune Diseases: A Comprehensive Review", *Medicina*, 2022

Publication

34

Katja Zaletel, Simona Gaberscek. "Hashimotos Thyroiditis: From Genes to the Disease", *Current Genomics*, 2011

Publication

35

Luyang Wei, Ning Zhao, Yalun Li, Xiaodan Zheng, Huaxin Zhang, Zhongtao Zhang. "Down syndrome with primary thyroid diffuse large B-cell lymphoma and Hashimoto thyroiditis", *Medicine*, 2019

Publication

36

epag.springeropen.com

Internet Source

37

"The Immunogenetics of Dermatologic Diseases", Springer Science and Business Media LLC, 2022

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

38

"Thyroid Diseases", Springer Science and Business Media LLC, 2018

Publication

<1 %

39

Valeria Calcaterra, Erica Crivicich, Annalisa De Silvestri, Rossella Amariti et al. "Timing, prevalence, and dynamics of thyroid disorders in children and adolescents affected with Down syndrome", Journal of Pediatric Endocrinology and Metabolism, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Protein tyrosine phosphatase non-receptor type 22 C1858T gene polymorphism in children with down syndrome and autoimmune thyroid diseases

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
