

ORIGINAL ARTICLE

Correlation Between Early Age Pregnancy with Low Body Weight (LBW) Newborn at Universitas Airlangga Academic Hospital, Surabaya, Indonesia

Roy Prasajo¹, Budi Prasetyo^{2*}, Widati Fatmaningrum³, Ahmad Hadi Modi⁴

¹Faculty of Medicine, Universitas Airlangga Surabaya, Indonesia

²Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga– Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

³Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga– Dr. Soetomo General Academic Hospital, Surabaya, Indonesia

⁴Faculty of Medicine and Health Science, School of Healthy Aging, Medical Aesthetics and Regenerative Medicine, UCSI University, Kuala Lumpur, Malaysia

ARTICLE INFO

Article history:

Received 28 October 2021

Received in revised form 10 November 2022

Accepted 12 December 2021

Available online 28 April 2022

Keywords:

Cross-Sectional Studies,
Early Age Pregnancy,
Low Birth Weight

*) *Corresponding author:*

budi-p@fk.unair.ac.id

ABSTRACT

Introduction: Birth weight is often used to evaluate a newborn baby's health. Low birth weight (LBW) is one of the leading causes of neonatal mortality. Mothers under the age of 20 (early pregnancy) are at a greater risk of affecting the weight and nutritional status of the unborn child. This research aimed to determine the relationship between early pregnancy and LBW.

Methods: This was a cross-sectional study of women and newborns at Airlangga University academic hospital from January through December of 2017. Data on characteristics such as maternal age and LBW were collected. The correlation between early pregnancy and infant weight was determined using the Chi-square test, with a significance level of $p < 0.05$.

Results: We conducted research on 210 pairs of mothers and their infants. 64 (30.48%) of the pregnancies occurred in women under the age of 20; 61 (29.55%) were born LBW (weighing less than 2,500 grams). Early age pregnancy was linked to a higher proportion of LBW infants ($p = 0.001$). Research indicated that the risk of LBW was more significant in the early groups of pregnancy.

Conclusion: This research establishes a relationship between early age pregnancy and LBW. The findings may aid in identifying vulnerable mothers in need of further assistance and personalized treatments.

Introduction

Birth weight is often used to assess newborn health. Low birth weight (LBW) in newborns is the leading cause of infant mortality. When a newborn weighs less than 2.5 kg, which often happens in preterm births or before to entering gestational age, the newborn is considered LBW.¹ This may occur for various reasons, one of which is the mother's age and nutritional status when pregnant. A proportion of moms under the age of 20 are experiencing a risky pregnancy, which will have an influence on the baby's weight and nutritional status following birth.²

LBW is more prevalent among babies born to young mothers under the age of 20. Teenagers are more likely than not to deliver LBW newborns.^{3,4} This is because they have not attained maturity and hence lack an effective placental transfer mechanism comparable to

that of adult females.⁵ In addition, the reproductive organs of people in their 20s and younger are not in the best shape for having a baby.⁶ However, as women aging, their bodies and health decline, potentially affecting the intrauterine fetus and resulting in LBW in the newborn. LBW is often diagnosed in women between the ages of 20 and 30.⁷⁻⁹

Newborn babies with LBW often face a negative long-term life process. If they do not die after delivery, LBW newborns have a higher chance of growing and developing more slowly than normal-weight newborns.^{10,11} Along with developmental disabilities, those with a history of LBW have an increased chance of developing hypertension, heart disease, and diabetes beyond the age of 40.¹⁰

The factors that contribute to LBW in newborns are projected to be highly connected to the mother's age. The prevalence of early marriage and babies born with LBW



is high around the world, including in Indonesia. The purpose of this research is to determine the relationship between LBW and maternal age in a single-center study in Surabaya.

Methods

A cross-sectional study was conducted from January to December 2017 at the academic hospital of Universitas Airlangga Surabaya, through a birth cohort of hospital deliveries of newborns. The WHO defines adolescent mother as women under the age of 20 who are more likely to have LBW babies, as well as preterm deliveries and severe neonatal disorders.² The research data was collected from medical records of LBW patients at the Department of Obstetrics and Gynecology, Universitas Airlangga Hospital Surabaya. The study population were all normal pregnancies and deliveries, which were divided into 2 groups, namely the group of severe infant cases born <2500 g, which occurred at the Department of Obstetrics and Gynecology, Universitas Airlangga Hospital Surabaya. The total sampling approach was employed.

The study sample consisted of 37-week-old normal pregnancy patients divided into two groups in January 2017 and 2018. Some patients with normal births a 37 weeks and giving birth with a normal delivery, met the inclusion criteria. The criteria for exclusion were patients who had eclampsia, hypertension, diabetes mellitus, postdate, abortion, amelioration, amnion, anemia cases, and pregnant women > 35 years old. The variables in this study were differentiated into 2 groups, independent and dependent variables, as follows: independent variables: age (early age pregnancy and non-early age pregnancy), history of previous pregnancy (primigravida and multigravida). The dependent variable is the baby's weight (LBW and non-LBW). The chi-square test was used to see if early-age pregnancy influenced newborn weight, with $p < 0.05$ considered statistically significant using SPSS version 17.0 (SPSS Inc., Chicago, IL, USA). The ethical committee of Universitas Airlangga Hospital approved this study with ethical number: 172/KEH/2018.

Results

There was 210 participants included in the study, with all subjects having a normal delivery over the study period. Most responses showed that they had become pregnant later (Table 1). The non-early age pregnancy group was nine years older age than the early age pregnancy (15.77±1.34 vs. 24.42±1.99). Primigravida was becoming the majority in early-age pregnancy (27.6%) and the non-early-age pregnancy group (44.3%). The incidence of LBW was more frequent in early age pregnancy group (21.43% vs. 7.62%).

There was a significant correlation between early age pregnancy group and the incidence of LBW ($p=0.001$; Table 2). The earlier the mother ages at giving birth, the higher the tendency of babies born to experience LBW. This study also had a relatively high prevalence ratio (PR=19.243), which showed that the early pregnancy group was at risk of 19.243 times higher in incidence of LBW than the group with pregnancy at an early age (Table 2).

Table 1. Distribution of characteristics of respondents based on maternal age at Airlangga University Hospital Surabaya period January 2017-December 2017

| Characteristics | Early age pregnancy | Non-early age pregnancy |
|---------------------------------------|---------------------|-------------------------|
| Age (mean ± SD) year | 64 (30.48%) | 146 (69.52%) |
| Newborn birth-weight (mean ± SD) gram | 2400.94±418.36 | 2923.29±372.84 |
| Pregnancy | | |
| Primigravida | 58 (27.60%) | 93 (44.30%) |
| Multigravida | 6 (2.90%) | 53 (25.20%) |
| Birth weight | | |
| LBW (%) | 45 (21.43%) | 16 (7.62%) |
| Non-LBW (%) | 19 (9.05%) | 130 (61.90%) |

Table 2. Relationship between maternal age and birth weight at Universitas Airlangga Hospital

| | P value | PR | CI |
|--------------------|---------|--------|--------------|
| Maternal age - LBW | 0.0001 | 19.243 | 9,122-40,594 |

PR = prevalence ratio

Discussion

The findings of this research reveal a significant association between early pregnancy (under the age of 20 years) and low birth weight (LBW) incidence. According to the newest Surabaya study, 30.48 % were categorized early age pregnancy, which is much higher than the rates in developed countries (1.8%-2.1%).¹²⁻¹⁴ Mothers aged 14-19 years did not have fully developed reproductive organs. Nutritional demands were also not fulfilled optimally due to the competition for nourishment between the mother and the developing baby, which might affect fetal growth and development in the form of LBW.^{14,15} The WHO recommended that women between the ages of 20 and 35 years old experience pregnancy and delivery. In 2010, Indonesia had a female mean marriage age of fewer than 25 years (22.17-year-old on average), which increased from 19.28-year-old in 1971.¹⁵ In Indonesia, despite their physical and mental immaturity, child marriage is mainly controlled by social and economic aspects inside and around the girls.¹⁵

Early pregnancy between the ages of 14- and 19-years effect on pregnancy and childbirth problems.¹⁷ Early-age pregnancy also increased the chance of LBW incidence by 4.1 times compared to pregnant women over 20 years. In general, children born to young mothers with LBW had congenital anomalies and physical problems, including epilepsy, mental retardation, blindness, and deafness.¹⁸ Furthermore, even if the infant survives, he or she may be impaired, causing significant issues and experiencing delayed development.¹⁹

Parity is the number of children born alive and born dead by a mother. When a woman becomes pregnant for the first time and her parity exceeds four, she is at risk of giving birth to LBW. Since the mother's uterus has not healed, if she becomes pregnant again, the second pregnancy may be affected.⁸ Because the causes of LBW

are often complex, it was sometimes difficult to establish a preventative intervention. Premature delivery was the most prevalent cause of LBW newborns, and the younger the gestational age, the greater the risk of short- and long-term complications.²⁰⁻²² Generally, maternal variables such as pregnancy disease, maternal age, labor distance, gestational age, parity, environmental factors, fetal factors, and placental factors are connected with LBW.^{8,9} However, the precise mechanism behind premature pregnancy and LBW remains unknown.

A systematic review and meta-analysis discovered that adolescent pregnancy is associated with low socioeconomic status, but that after controlling for high socioeconomic status (SES), adolescents were not at increased risk for LBW.^{8,23} This finding implies that SES had a significant influence on LBW.²⁴ Low SES is a complicated issue that necessitates the implementation of government legislation.²⁵ We recommend that the government provide access to high-quality prenatal care for teenage mothers from poor socioeconomic backgrounds to lower the frequency of LBW births.

The research has a limitation in that variable such as calorie consumption, weight gain during pregnancy, and the high socioeconomic level was not included since they may be associated with LBW. Additionally, the outcomes of this research were not compared to those of adolescents who did not receive antenatal or prenatal care or those of any external comparator. We obtained data primarily from single-center academic hospital, and hence cannot generalize to other hospitals.

Acknowledgment

The authors liked to express their gratitude to Budi Utomo and Atika for their support with statistical analysis. The research was developed and evaluated by Budi Prasetyo, Widati Fatmaningrum, and AH Modi. The article was written by Roy Prasojko.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

1. About Teen Pregnancy | CDC, <https://www.cdc.gov/teenpregnancy/about/index.htm> (2019, accessed 20 April 2022).
2. WHO. Adolescent Pregnancy. Geneva: World Health Organization, <http://www.who.int/mediacentre/factsheets/fs364/en/> (2014).
3. Giovanni M, Prabowo GI, Fatmaningrum W. Infant Birth Weight in Mothers with Maternal Anemia at Dupak Public Health Center Surabaya Working Area in 2017. *Biomol Heal Sci J* 2019; 2: 53.
4. Nair A, Devi S. Obstetric outcome of teenage pregnancy in comparison with pregnant women of 20-29 years: a retrospective study. *Int J Reprod Contraception, Obstet Gynecol* 2015; 4: 1319–1323.
5. Kamini S, Avvaru KV. Teenage Pregnancy: Maternal and Fetal Outcomes. *IOSR J Dent Med Sci e-ISSN* 2014; 13: 41–44.
6. Aprilia DN, Prasetyo B, Sulistiawati S. Correlation Between Nutritional Status of Pregnant Women Based on Upper Arm Circumference and Preeclampsia/Eclampsia Severity Degree at Jagir Public Health Center During January 2014 - March 2014. *Biomol Heal Sci J* 2018; 1: 120.
7. Veloso HJF, da Silva AAM, Bettiol H, et al. Low birth weight in São Luís, northeastern Brazil: Trends and associated factors. *BMC Pregnancy and Childbirth*; 14. Epub ahead of print 2014. DOI: 10.1186/1471-2393-14-155.
8. DeMarco N, Twynstra J, Ospina MB, et al. Prevalence of Low Birth Weight, Premature Birth, and Stillbirth Among Pregnant Adolescents in Canada: A Systematic Review and Meta-analysis. *Paediatr Perinat Epidemiol* 2021; 33: 530–537.
9. Hussein Eldessouki KM, Abd Elhakim Quodi M, Mahmoud Ahmed Hassane S. Assessment of The Impact of Teenage Pregnancy on Pregnancy Outcome among Pregnant Women in Minia University Hospitals, Minia Governorate. *Egypt J Community Med* 2020; 38: 87–95.
10. Silvestrin S, Hirakata VN, da Silva CH, et al. Inequalities in birth weight and maternal education: a time-series study from 1996 to 2013 in Brazil. *Sci Rep* 2020; 10: 8707.
11. Harville EW, Madkour AS, Xie Y. Predictors of birth weight and gestational age among adolescents. *American Journal of Epidemiology*; 176. Epub ahead of print 2012. DOI: 10.1093/aje/kws231.
12. Martin JA, Hamilton BE, Osterman MJK, et al. Births: Final data for 2017. *Natl Vital Stat Reports* 2018; 67: 1–49.
13. Canada S. Live births, by age group and marital status of mother, Canada, place of residence of mother, annual. CANSIM, <https://www150.statcan.gc.ca/t1/tb11/en/cv.action?pid=1310042001>.
14. Wong SPW, Twynstra J, Gilliland JA, et al. Risk Factors and Birth Outcomes Associated with Teenage Pregnancy: A Canadian Sample. *J Pediatr Adolesc Gynecol* 2020; 33: 153–159.
15. Kusparlina EP. Hubungan Antara Umur Dan Status Gizi Ibu Berdasarkan Ukuran Lingkar Lengan Atas Dengan Jenis Bblr Di Puskesmas Tawangrejo Kota Madiun. *J Delima Harapan* 2019; 6: 7–16.
16. Da Silva CH, Hernandez AR, Agranonik M, et al. Maternal age and low birth weight: A reinterpretation of their association under a demographic transition in southern Brazil. *Maternal and Child Health Journal* 2013; 17: 539–544.
17. Jain LH, Van Eyk N, Woolcott C, et al. Characteristics and Outcomes of Adolescent Births in Nova Scotia: A Retrospective Cohort Study. *J Obstet Gynaecol Canada* 2018; 40: 1459–1465.
18. N. K, H.B. S, H. K, et al. Premature birth, low birth weight and birth defects after assisted reproductive therapies. A 18-year comparative study. *Tunisie Medicale* 2017; 95: 103–108.
19. Fleming N, Ng N, Osborne C, et al. Adolescent Pregnancy Outcomes in the Province of Ontario: A Cohort Study. *J Obstet Gynaecol Canada* 2013; 35: 234–245.
20. Cutland CL, Lackritz EM, Mallett-Moore T, et al. Low birth weight: Case definition & guidelines for data collection, analysis, and presentation of maternal immunization safety data. *Vaccine* 2017; 35: 6492.
21. Luu TM, Katz SL, Leeson P, et al. Preterm birth: risk factor for early-onset chronic diseases. *C Can Med Assoc J* 2016; 188: 736.
22. Muhili A, Sudfeld CR, Smith ER, et al. Risk factors for small-for-gestational-age and preterm births among 19,269 Tanzanian newborns. *BMC Pregnancy Childbirth* 2016; 16: 1–12.
23. Marvin-Dowle K, Soltani H. A comparison of neonatal outcomes between adolescent and adult mothers in developed countries: A systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol X* 2020; 6: 6100109.
24. Sudhir A, Rasquinha VC, Rao SB., et al. Outcome of Pregnancy in Adolescent Age Group. *J Evol Med Dent Sci* 2015; 4: 7483–7488.
25. Amjad S, Adesunkanmi M, Twynstra J, et al. Social Determinants of Health and Adverse Outcomes in Adolescent Pregnancies. *Semin Reprod Med* 2021; 33: 88.