

Duration of SARS-CoV-2 RNA Shedding Is Significantly Influenced by Disease Severity, Bilateral Pulmonary Infiltrates, Antibiotic Treatment, and Diabetic Status: Consideration for Isolation Period

by Muhammad Vitanata Arfijanto

Submission date: 08-May-2023 09:50AM (UTC+0800)

Submission ID: 2086969270

File name: RNA_Shedding_Is_Significantly_Influenced_by_Disease_Severity.pdf (1.15M)

Word count: 6790

Character count: 35876

Article

Duration of SARS-CoV-2 RNA Shedding Is Significantly Influenced by Disease Severity, Bilateral Pulmonary Infiltrates, Antibiotic Treatment, and Diabetic Status: Consideration for Isolation Period

Muhammad Vitanata Arfijanto ¹, Tri Pudy Asmarawati ^{1,2,*}, Bramantono Bramantono ¹, Musofa Rusli ¹, Brian Eka Rachman ^{1,2}, Bagus Aulia Mahdi ¹, Nasronudin Nasronudin ^{1,2} and Usman Hadi ¹

¹ Department of Internal Medicine, Faculty of Medicine, Airlangga University, Surabaya 60132, Indonesia
² Universitas Airlangga Hospital, Airlangga University, Surabaya 60115, Indonesia
 * Correspondence: tpasmarawati@fk.unair.ac.id



Citation: Arfijanto, M.V.; Asmarawati, T.P.; Bramantono, B.; Rusli, M.; Rachman, B.E.; Mahdi, B.A.; Nasronudin, N.; Hadi, U. Duration of SARS-CoV-2 RNA Shedding Is Significantly Influenced by Disease Severity, Bilateral Pulmonary Infiltrates, Antibiotic Treatment, and Diabetic Status: Consideration for Isolation Period. *Pathophysiology* **2023**, *30*, 186–198. <https://doi.org/10.3390/pathophysiology30020016>

Academic Editor: Jonathan Steven Alexander

Received: 9 November 2022
Revised: 24 April 2023
Accepted: 28 April 2023
Published: 4 May 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) ribonucleic acid (RNA) shedding is an important parameter for determining the optimal length of isolation period required for coronavirus disease 2019 (COVID-19) patients. However, the clinical (i.e., patient and disease) characteristics that could influence this parameter have yet to be determined. In this study, we aim to explore the potential associations between several clinical features and the duration of SARS-CoV-2 RNA shedding in patients hospitalized with COVID-19. A retrospective cohort study involving 162 patients hospitalized for COVID-19 in a tertiary referral teaching hospital in Indonesia was performed from June to December 2021. Patients were grouped based on the mean duration of viral shedding and were compared based on several clinical characteristics (e.g., age, sex, comorbidities, COVID-19 symptoms, severity, and therapies). Subsequently, clinical factors potentially associated with the duration of SARS-CoV-2 RNA shedding were further assessed using multivariate logistic regression analysis. As a result, the mean duration of SARS-CoV-2 RNA shedding was found to be 13 ± 8.44 days. In patients with diabetes mellitus (without chronic complications) or hypertension, the duration of viral shedding was significantly prolonged (≥ 13 days; $p = 0.001$ and $p = 0.029$, respectively). Furthermore, patients with dyspnea displayed viral shedding for longer durations ($p = 0.011$). The multivariate logistic regression analysis reveals that independent risk factors associated with the duration of SARS-CoV-2 RNA shedding include disease severity (adjusted odds ratio [aOR] = 2.94; 95% CI = 1.36–6.44), bilateral lung infiltrates (aOR = 2.79; 95% CI = 1.14–6.84), diabetes mellitus (aOR = 2.17; 95% CI = 1.02–4.63), and antibiotic treatment (aOR = 3.66; 95% CI = 1.74–7.71). In summary, several clinical factors are linked with the duration of SARS-CoV-2 RNA shedding. Disease severity is positively associated with the duration of viral shedding, while bilateral lung infiltrates, diabetes mellitus, and antibiotic treatment are negatively linked with the duration of viral shedding. Overall, our findings suggest the need to consider different isolation period estimations for specific clinical characteristics of patients with COVID-19 that affect the duration of SARS-CoV-2 RNA shedding.

Keywords: COVID-19; viral shedding; SARS-CoV-2; risk factors; infectious disease

1. Introduction

Coronavirus disease 19 (COVID-19), an infectious disease caused by the novel severe acute respiratory syndrome coronavirus type-2 (SARS-CoV-2), encompasses varying manifestations ranging from asymptomatic and mild illness to life-threatening complications including acute respiratory distress syndrome (ARDS). The World Health Organization (WHO) announced COVID-19 as a pandemic on 11 March 2020; as of February 2023, over 758 million confirmed cases and over 6.8 million deaths had been reported [1]. In Indonesia,

COVID-19 has affected over 6.7 million people and more than 160,000 deaths (2.4%) have been reported [2].

The duration of SARS-CoV-2 RNA shedding is the period from the onset of symptoms to the virus not being detected upon polymerase chain reaction (PCR) examination [3]. Some studies have reported that the viral shedding time of SARS-CoV-2 from infected patients is affected by transmission source, age, disease severity, drugs, comorbidities, and immunity status [3–6]. The most prolonged duration is about 60 days, and the shortest is about 3 days [7]. However, the duration of SARS-CoV-2 viral shedding is variable, and the pattern and influencing factors are yet to be determined.

Recognizing the prolonged viral shedding of SARS-CoV-2 is mandatory in deciding the isolation period and treatment of patients with COVID-19. The risk of transmission due to viral shedding cannot be ignored, even if there is a resolution that is not accompanied by any symptoms. The risks vary depending on the infectiousness level of the virus and the patient population involved [7].

The prolonged infection and evolution of the SARS-CoV-2 virus in immunocompromised individuals may lead to the emergence of new variants, the increase in virulence levels, and potential immune escape, thus posing the risk of complicating pandemic control [8,9]. Thus, clinical judgments should be made based on clinical features, guidelines, the detectable viral load, transmission source, infectiousness level, and patient-related factors [7,10–14]. It is critical to further study the factors affecting the viral shedding time of SARS-CoV-2 to estimate the optimal duration of isolation required by patients based on certain conditions. As such, we intend to investigate related factors, particularly host factors (age, sex, and comorbidities) and symptoms, with respect to the duration of SARS-CoV-2 viral shedding.

2. Materials and Methods

2.1. Study Design and Sample Selection

This research is a single-center retrospective observational study on hospitalized COVID-19 patients in the isolation ward of a tertiary referral teaching hospital in Surabaya, Indonesia. This study was performed from June to December 2021. We aimed to analyze several factors, such as age, gender, comorbidity, disease severity, and COVID-19-related symptoms, and their association with the duration of SARS-CoV-2 RNA shedding in a retrospective design (Figure 1). The inclusion criteria were confirmed COVID-19 patients over 17 years old who had been hospitalized until being declared free from isolation after 2 consecutive negative RT-PCR tests. The exclusion criteria were confirmed COVID-19 patients who died or were discharged from the hospital before the RT-PCR test showed negative results. Diagnosis of COVID-19 was based on the confirmation of SARS-CoV-2 from RT-PCR using Roche COBAS Z480 analyzer (PT. Roche Indonesia, Jakarta). Severity classification was divided into 5 categories based on the Indonesia COVID-19 management guidelines, which are detailed as follows [15]:

- (1) Asymptomatic case.
- (2) Mild disease: symptomatic patients without evidence of viral pneumonia or hypoxia. Symptoms might appear as fever; cough; fatigue; anorexia; shortness of breath; myalgia; sore throat; nasal congestion; headache; diarrhea; nausea and vomiting; anosmia; or loss of appetite.
- (3) Moderate disease: patients with clinical signs of pneumonia (fever, cough, shortness of breath, rapid breathing) but no symptoms of severe pneumonia, including oxygen saturation >93% on room air.
- (4) Severe disease: patients with clinical signs of pneumonia (fever, cough, shortness of breath) plus respiratory rate >30 times/minute, severe respiratory distress, or SpO₂ < 93% on room air.
- (5) Critical disease: patients with acute respiratory distress syndrome (ARDS), sepsis, and septic shock.




Figure 1. Retrospective observational study design to analyze the association between clinical parameters and the duration of SARS-CoV-2 RNA shedding.

2.2. Definition

The clinical parameters studied, such as age, gender, comorbidities (diabetes mellitus [DM], malignancy, autoimmune disease, human immunodeficiency virus [HIV] infection, liver cirrhosis, obesity), and symptoms, were retrieved from medical records. RT-PCR examination was conducted every five days based on the local hospital standard procedure. The duration of viral shedding was calculated from the onset of the symptoms until the day of 2 consecutive negative detections of SARS-CoV-2 RNA within 24 h [16–18]. The cut-off value for prolonged viral shedding was determined from the average duration obtained from data analysis. The duration of SARS-CoV-2 RNA shedding was considered prolonged if it is equal to or more than 13 days (Figure 1). The primer and probe for RT-PCR for SARS-CoV-2 detection targeted the N, E, and ORF1ab genes, with a cycle threshold (CT) value <38 considered positive.

2.3. Statistical Analysis

We analyzed the data using SPSS version 24 (Chicago, IL, USA; RRID: SCR_002865). The two groups were divided based on the average duration of viral shedding. The differences between the two groups were analyzed using the chi-square and Fisher's exact tests because of the abnormally distributed data. The *p*-value that was considered significant was <0.05 with a confidence interval of 95%. Variables with significant differences in the bivariate analysis were included in the logistic regression multivariate analysis to determine factors that influenced the duration of SARS-CoV-2 viral shedding. The results were presented in the form of an odds ratio (OR) with the *p*-value considered significant being <0.05 and a confidence interval of 95%, and the adjusted OR (aOR) showed the results of the multivariate analysis.

2.4. Ethical Clearance

The ethical clearance (no. 0825/LOE/301.4.2/III/2022) for this study was granted by the Ethical Committee of Dr. Soetomo Hospital, Surabaya, Indonesia.

3. Results

3.1. Baseline Characteristics

The patient characteristics are shown in Table 1. This study included 162 patients as subjects with 83 (51.23%) men and 79 (48.76%) women. Most study participants (75.31%) were under 60 years old. The most frequent clinical symptoms were cough, fever, and dyspnea; some other symptoms were expectoration and hemoptysis. With respect to disease severity, mild cases were the most prevalent (43.83%) followed by severe cases (26.54%) and moderate cases (20.99%). The average length of hospitalization was 7.1 ± 8.82 days. Chest X-rays mostly showed bilateral pulmonary infiltrates (67.9%).

Table 1. COVID-19 patient (cohort) characteristics.

Characteristics	Total (n = 162)
Sex, n (%)	
Male	83 (51.23)
Female	79 (48.76)
Age, n (%)	
≥ 60 years	40 (24.69)
<60 years	122 (75.31)
Clinical symptoms, n (%)	
Fever	83 (51.23)
Cough	85 (52.47)
Expectorations	35 (21.60)
Hemoptysis	1 (0.62)
Dyspnea	80 (49.38)
Disease severity, n (%)	
Mild	71 (43.83)
Moderate	34 (20.99)
Severe	43 (26.54)
Critical	14 (8.64)
Length of hospitalization, mean ± SD	7.41 ± 8.82
Chest X-ray, n (%)	
Normal	32 (19.75)
Unilateral infiltrate	15 (9.26)
Bilateral infiltrates	110 (67.90)
Atypical	5 (3.09)
Comorbidities, n (%)	
Diabetes mellitus	55 (33.95)
DM with chronic complication	12 (7.41)
DM without chronic complication	43 (26.54)
Hypertension	46 (28.40)
HIV/AIDS	11 (6.79)
Malignancy	17 (10.49)
Liver cirrhosis	11 (6.79)
Autoimmune disease	7 (4.32)
Obesity	9 (5.55)
Treatment, n (%)	
Antiviral therapy:	
Lopinavir/ritonavir	34 (20.99)
Favipiravir	37 (22.84)
Remdesivir	44 (27.16)
Oseltamivir	25 (15.43)
Corticosteroid	63 (38.89)
Convalescent plasma	6 (3.70)

Table 1. Cont.

Characteristics	Total (n = 162)
Antibiotic	77 (47.53)
Oxygen therapy	89 (54.94)
Output, n (%)	
Death	11 (6.79)
Cured	151 (93.21)
Duration of viral shedding (days) (Mean \pm SD)	13 \pm 8.44

The most prevalent comorbidities were diabetes mellitus (33.95%) and hypertension (28.54%). Other comorbidities included malignancy, HIV/AIDS, liver cirrhosis, and autoimmune diseases. The antiviral treatments given were remdesivir (27.16%), favipiravir (22.84%), lopinavir/ritonavir (20.99%), and oseltamivir (15.43%). Oxygen therapy was delivered to 89 patients (54.94%).

Most patients were cured and discharged from the isolation ward (93.21%), while mortality was reported in 11 subjects (6.79%). The average length of stay was 7.41 ± 8.82 days. The median duration of viral shedding in this study was 10 days, with the shortest duration being 5 days and the most prolonged duration recorded being 42 days, with an interquartile range (IQR) of 7–17.25 days.

3.2. Bivariate Analysis

The bivariate analysis was based on the mean duration of viral shedding (Table 2). Most subjects aged <60, namely 79 subjects (79.8%), had viral shedding <13 days. Subjects aged 60 years and over had a more significant proportion who experienced prolonged viral shedding ≥ 13 days, although this difference was not statistically significant ($p = 0.097$). The duration of viral shedding was not significantly different for men and women ($p = 0.816$). However, there was a more significant proportion of more prolonged viral shedding for men (33 subjects; 52.4%) than for women. Patients with comorbid hypertension had a more considerable proportion who experienced a viral shedding duration of ≥ 13 days (38.1% vs. 22.2%), and the difference was statistically significant ($p = 0.029$). A total of 55 patients (33.95% of the total subjects) had DM, which was further categorized into DM with chronic complications in 12 patients (7.41% of the total subjects) and DM without chronic complications in 43 patients (26.54% of the total subjects). Most COVID-19 patients with DM had prolonged viral shedding ($p = 0.001$). The duration of viral shedding was not statistically different in COVID-19 patients with liver cirrhosis ($p = 0.27$). More COVID-19 patients with malignancy experienced viral shedding <13 days, although there was no statistically significant difference ($p = 0.748$). A total of 11 patients out of the total COVID-19 subjects (6.79%) experienced comorbid HIV/AIDS infection. Among these patients there was no significant difference in the duration of viral shedding with $p = 0.27$. Most of the COVID-19 patients with comorbid autoimmune diseases in this study had a course of viral shedding <13 days, but this difference was not statistically significant ($p = 0.172$). Obesity in this study was found in 9 patients (5.55%). There was no significant difference in the duration of viral shedding in obese patients with COVID-19 ($p = 0.725$). Patients who experienced dyspnea had a significantly longer duration of viral shedding ($p = 0.011$). Other symptoms, such as fever, cough, expectoration, and hemoptysis, did not differ substantially with respect to the duration of viral shedding.

Table 2. Characteristics based on the duration of viral shedding.

Characteristics	Total (n = 162)	Duration of Viral Shedding		p-Value
		<13 days (n = 99)	≥13 days (n = 63)	
Sex				
Male	83	50 (50.5%)	33 (52.4%)	0.816
Female	79	49 (49.5%)	30 (47.6%)	
Age				
≥60 years	40	20 (20.2%)	20 (31.7%)	0.097 *
<60 years	122	79 (79.8%)	43 (68.3%)	
Clinical symptoms				
Fever	83	45 (45.5%)	38 (60.3%)	0.065 *
Cough	85	48 (48.5%)	37 (58.7%)	0.203 *
Expectoration	35	18 (18.2%)	17 (27.0%)	0.184 *
Hemoptysis	1	1 (1.0%)	0 (0%)	0.484
Dyspnea	80	41 (41.4%)	39 (61.9%)	0.011 *
Disease severity				
Mild	71	50 (50.5%)	21 (33.3%)	0.012 *
Moderate	34	24 (24.2%)	10 (15.9%)	
Severe	43	19 (19.2%)	24 (38.1%)	
Critically ill	14	6 (6.1%)	8 (12.7%)	
Chest X-ray				
No infiltrate	32	26 (26.3%)	6 (9.5%)	0.009 *
Unilateral infiltrate	15	12 (12.1%)	3 (4.8%)	0.115 *
Bilateral infiltrate	110	56 (56.6%)	54 (85.7%)	0.00 *
Atypical	5	4 (4.0%)	1 (1.6%)	0.379
Comorbidities				
Diabetes Mellitus	55	24 (24.2%)	31 (49.2%)	0.001 *
- with chronic complications	12	5 (6.1%)	7 (9.5%)	0.151 *
- without chronic complications	43	18 (18.2%)	25 (39.7%)	0.003 *
Hypertension	46	22 (22.2%)	24 (38.1%)	0.029 *
HIV/AIDS	11	5 (5.1%)	6 (9.5%)	0.27
Malignancy	17	11 (11.1%)	6 (9.5%)	0.748
Liver cirrhosis	11	5 (5.1%)	6 (9.5%)	0.27
Autoimmune disease	7	6 (6.1%)	1 (1.6%)	0.172 *
Obesity	9	5 (5.1%)	4 (6.3%)	0.725
Treatment				
Lopinavir/ritonavir	34	10 (10.1%)	24 (38.1%)	0.00 *
Favipiravir	37	29 (29.3%)	8 (12.7%)	0.014 *
Remdesivir	44	27 (27.3%)	17 (27.0%)	0.968
Oseltamivir	25	17 (17.2%)	8 (12.7%)	0.442
Corticosteroid	63	32 (32.3%)	31 (49.2%)	0.032
Convalescent plasma	6	2 (2.0%)	4 (6.3%)	0.155 *

Table 2. Cont.

Characteristics	Total (n = 162)	Duration of Viral Shedding		p-Value
		<13 days (n = 99)	≥13 days (n = 63)	
Antibiotic	77	36 (36.4%)	41 (65.1%)	0.00 *
Oxygen therapy	89	44 (44.4%)	45 (71.4%)	0.001 *
Outcome				
Death	12	7 (7.1%)	4 (6.3%)	0.859
Cured	151	92 (92.9%)	59 (93.7%)	

* p-value chi-square and Fischer exact test <0.25.

3.3. Multivariate Analysis

The multivariate analysis of several variables with significant differences in the bivariate analysis included disease severity; dyspnea; bilateral pulmonary infiltrates on chest X-ray; hypertension; diabetes; and treatments including oxygen supplementation, antibiotics, oral antivirals, and corticosteroids. In the stepwise analysis, it was found that the factors related to the duration of viral shedding of SARS-CoV-2 were disease severity (aOR 2.94; CI 95%, 1.34–6.44), bilateral pulmonary infiltrates (aOR 2.79; CI 95%, 1.14–6.84), hypertension (OR 2.15; CI 95%, 1.08–4.32), and antibiotic therapy (aOR 3.66; CI 95%, 1.74–7.71). Age, gender, and symptoms were not associated with the duration of SARS-CoV-2 viral shedding (see Table 3).

Table 3. Multivariate analysis of parameters potentially associated with SARS-CoV-2 viral shedding.

Variable	Multivariate Analysis			Stepwise Analysis		
	OR	95% CI	p-Value	aOR	95% CI	p-Value
≥60 years	1.88	0.89–3.78	0.09			
Disease severity	3.05	1.56–5.97	0.001 *	2.94	1.34–6.44	0.007 *
Fever	1.82	0.96–3.46	0.065			
Dyspnea	2.29	1.20–4.39	0.011 *			
Cough	1.51	0.79–2.86	0.203			
Expectoration	1.66	0.78–3.54	0.184			
No infiltrate	0.299	0.11–0.76	0.009 *			
Unilateral infiltrate	0.363	0.09–1.34	0.115			
Bilateral pulmonary infiltrates	4.61	2.05–10.35	0.000 *	2.79	1.14–6.84	0.025 *
Hypertension	2.15	1.08–4.32	0.029 *			
Diabetes mellitus	3.027	1.54–5.94	0.001 *	2.17	1.02–4.63	0.046 *
Autoimmune disease	0.25	0.03–2.13	0.172			
Oxygen therapy	3.13	1.59–6.14	0.001 *			
Antibiotic	3.26	1.69–6.31	0.000 *	3.66	1.74–7.71	0.001 *
Oral antiviral	1.36	0.70–2.62	0.36	1.96	0.89–4.29	0.093
Corticosteroid	2.03	1.06–3.88	0.032 *			

OR, odds ratio; aOR, adjusted odds ratio. * p-value < 0.05.

4. Discussion

The characteristics of the research subjects revealed that gender had nearly equal proportions with 51.23% men and 48.76% women. Most research subjects (75.31%) were under 60 years old. This phenomenon might have been related to the disease severity pattern in this study, which was dominated by mild illness (43.83%). As reported in

52
previous studies, older age is associated with COVID-19 mortality [19,20]. The elderly mostly exhibit severe or critical disease that requires ICU treatment and is not included in this study. The clinical symptoms observed in this study were similar to those reported by Qi et al., including cough, fever, and shortness of breath. Some of the patients also reported having expectoration and hemoptysis [16]. The most prevalent comorbidities reported in this study were diabetes mellitus and hypertension, followed by malignancy, HIV/AIDS infection, liver cirrhosis, and autoimmune diseases. Several studies have reported hypertension as the most prevalent comorbidity (11–83%), while in this study, hypertension was found in 28.4% of subjects [3,16,17,21].

This study's reported mean duration of RNA viral shedding was 13 ± 8.44 days with a median of 10 days (interquartile range 7–17 days). Several similar studies have reported different mean and median viral shedding durations. A Chinese study in early 2020 involving 101 hospitalized patients reported a median viral shedding period of 11 days (interquartile range: 8–14.3 days) [3]. Another study of 147 patients at a COVID-19 treatment center outside Wuhan, China, reported a median viral shedding period of 17 days (interquartile range: 12–21 days) [16]. A multicenter retrospective cohort study during the early pandemic involving 191 inpatients reported a median viral shedding period of 20 days in survivors, but SARS-CoV-2 was detectable until death in non-survivors. The most prolonged observed duration of viral shedding in survivors was 37 days [19]. Differences in viral shedding times may have been due to a patient's condition and disease severity.

COVID-19 occurring in elderly patients is often associated with disease severity and the prolonged duration of SARS-CoV-2 viral shedding due to weakened immune systems and the presence of comorbidities. A study by Zhou et al. reported that older age was associated with the persistence of the SARS-CoV-2 virus; in these patients the virus could survive for 111 days in the respiratory tract [22]. This study found that more subjects over 60 experienced a longer duration of viral shedding ≥ 13 days, while subjects under 60 years experienced a shorter duration. However, the difference was not statistically significant. Several similar studies have also reported that there is no association between age and the duration of viral shedding [17,21]. This suggests that factors other than age contribute to the prolonged duration of viral shedding in SARS-CoV-2.

In this study, there was no difference in the viral shedding time of SARS-CoV-2 between men and women. This phenomenon has also been reported in several similar studies [16,18,21]. At the same time, a study by Xu et al. stated that the prolonged duration of SARS-CoV-2 viral shedding was found more commonly in men [17]. The widely reported influence of sex on COVID-19 is with respect to disease severity and mortality, where males tend to experience more severe cases with higher mortality. Immune responses and hormonal systems play a role in the pathogenesis of COVID-19, although this may still be multifactorial [23].

COVID-19 patients with hypertension had a significantly higher proportion who experienced a longer duration of viral shedding. A similar study by Xu et al. also reported prolonged viral shedding beyond 15 days in patients with comorbid hypertension [17]. Hypertension has been reported to be associated with COVID-19 disease progression and ARDS risk through oxidative stress mechanisms, endothelial dysfunction, and RAS modification [12,24].

Patients with type 2 diabetes mellitus are at risk of complications of COVID-19, and several researchers have reported an association between COVID-19 and diabetes. Type 2 diabetes mellitus is also a risk factor for prolonged viral shedding in SARS-CoV-2 [25]. In human monocytes, elevated blood sugar levels directly increase SARS-CoV-2 replication; glycolysis maintains SARS-CoV-2 replication through reactive oxygen species (ROS) production and the activation of hypoxia-inducible factor 1 α ; thus, hyperglycemia is a condition that supports viral proliferation [26]. In this study, DM patients experienced a significantly longer viral shedding time ($p = 0.001$). There was no significant difference in the duration of SARS-CoV-2 viral shedding in DM patients with chronic complications.

By contrast, there was a significant difference in the duration of viral shedding in uncomplicated DM. These results indicate that the influence of diabetes on the duration of viral shedding is determined by the diabetes status and not by the complications of diabetes. Some other studies have reported no significant difference in the course of viral shedding in COVID-19 patients with comorbid diabetes. All three of these studies mainly involved mild to moderate COVID-19 patients [3,16,17,21]. A study by Buetti reported that diabetes was a risk factor for the prolonged release of the SARS-CoV-2 virus, but the patients included in this study were COVID-19 patients in a critical condition [25].

The duration of viral shedding was not statistically different in COVID-19 patients with liver cirrhosis. Research on COVID-19 in the cirrhotic population is still very limited. Nonetheless, several other studies regarding the duration of viral shedding in COVID-19 patients with comorbid cirrhosis have reported no difference in the duration of viral shedding for SARS-CoV-2 [3,18,27].

The COVID-19 patients with malignancy in this study did not suffer differences in the duration of SARS-CoV-2 viral shedding. Another similar study also reported no difference in the period of viral shedding in COVID-19 patients with and without malignancy [3,9,18,21]. Some of these studies did not state the type of malignancy, such as hematological malignancies or solid tumors, as in this study. A case report of a COVID-19 patient with hematologic malignancy indicated a prolonged duration of viral shedding of up to 97 days [8].

This study showed no significant difference in the viral shedding time of COVID-19 patients with HIV/AIDS ($p = 0.27$) (Table 2). The depletion and impaired cellular function of CD4+ T cells might have been the cause behind the prolonged viral shedding time. The duration of SARS-CoV-2 viral shedding reported in HIV and COVID-19 coinfection was 28–85 days; it is therefore recommended that the isolation period of people living with HIV be longer than 14 days [28–30]. The different results of this study may have occurred due to the small sample of HIV/AIDS coinfection (6.79% of the total subjects).

A total of 7 patients (4.32% of the total study subjects) had an autoimmune disease. In this study, most COVID-19 patients with an autoimmune disease had a viral shedding duration of <13 days, but the difference was not statistically significant, with $p = 0.172$ (Table 2). A disturbance of the immune system in autoimmune disease will affect the body's ability to fight pathogens, including SARS-CoV-2. Treatment with immunosuppressants may also affect the prognosis of COVID-19 patients [31]. However, there have been no reports on the impact of autoimmune disease on the duration of SARS-CoV-2 viral shedding.

In some literature reviews, obesity has been said to be associated with increased transmission rates, aggravating disease severity, increased mortality, and prolonged SARS-CoV-2 viral shedding [32,33]. A study by Zhang et al. reported that obesity was an independent risk factor for protracted SARS-CoV-2 viral shedding in the respiratory tract with an OR of 3.31; 95% CI: 1.08–10.09 [21]. The difference in the results of this study may have been due to the small number of obese patients included (5.55% of the total sample). The proportion of obese patients with a SARS-CoV-2 viral shedding duration ≥ 13 days was slightly higher than <13 days, but it was not statistically significant.

COVID-19 patients who presented with dyspnea in this study had a significantly longer duration of viral shedding. Dyspnea as the primary signal of worsening respiratory failure supported by the presence of lung lesions is common in moderate to severe COVID-19 cases. The extent of the opacity on the CT of the lungs is related to the duration of viral shedding in airway samples [34].

In influenza A virus infection, there is a relationship between tympanic temperature and viral shedding [35]. As the most common symptom of COVID-19 in this study, fever showed a trend related to the prolonged duration of SARS-CoV-2 viral shedding. However, this was not statistically significant, which was possibly due to the limited sample size. Another study reported that in addition to protracted viral shedding, higher temperatures were significantly associated with the progression of COVID-19. Thus, patients with

high temperatures upon entry require more careful monitoring and the consideration of undergoing a more extended quarantine [16].

Based on this study, disease severity, bilateral pulmonary infiltrates on chest X-ray, and antibiotic therapy were independently associated with the duration of SARS-CoV-2 viral shedding. Figure 2 describes the proposed mechanism by which diabetes might prolong SARS-CoV-2 RNA shedding through viral entry and replication facilitation. Diabetes mellitus also affects the immune response to virus elimination. The pathogenic host response to SARS-CoV-2 could result in a hyperinflammatory state which determines disease severity. Disease severity related to a high viral load affects the length of viral clearance from the respiratory tract of COVID-19 patients [34]. The appearance of bilateral pulmonary infiltrates on a chest X-ray also reflects disease severity due to the excessive inflammatory response in COVID-19. Antibiotic therapy and its association with the duration of SARS-CoV-2 viral shedding may be related to bacterial coinfection commonly found in severe COVID-19 patients. Hence, it would affect the viral clearance of SARS-CoV-2. Previous studies have also reported that bacterial coinfection was more common in COVID-19 patients with poor outcomes [14,36].




Figure 2. Factors associated with prolonged viral shedding in COVID-19 patients.

This study's limitation was that the viral shedding of SARS-CoV-2 was determined based on PCR viral particle detection; thus, the presence of viral shedding in the respiratory tract could not be defined. This research was conducted in June–December 2021 when the dominating SARS-CoV-2 variant was the Delta variant, which may have affected the severity and duration of SARS-CoV-2 shedding. Viral RNA sequencing tests are still not being conducted routinely. However, differences in the SARS-CoV-2 variants had minimal or no impact on the virus properties in general. The SARS-CoV-2 PCR tests in this study were carried out every five days based on the protocol applied in the hospital, thus allowing for bias in determining the duration of SARS-CoV-2 viral shedding. This study's small sample size and observational method brought a high risk of confounding variables.

5. Conclusions

Several clinical parameters are related to the duration of SARS-CoV-2 RNA shedding. Disease severity is positively associated with the duration of viral shedding, while bilateral pulmonary infiltrates and antibiotic therapy are negatively related to the duration of viral shedding. Comorbidity diabetes mellitus is also associated with prolonged viral shedding. Overall, our findings suggest the need to consider different isolation period estimations for the specific clinical characteristics of patients with COVID-19 such as those with severe disease, bilateral pulmonary infiltrates, antibiotic treatment, and diabetes mellitus.

Author Contributions: Conceptualization; methodology; software; validation; formal analysis; investigation; data curation; and writing—original draft preparation, T.P.A., B.A.M. and M.V.A.; writing—review and editing, T.P.A., M.V.A., B.B., M.R., B.E.R., B.A.M., N.N. and U.H.; supervision, M.V.A. and U.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The ethical clearance (no. 0825/LOE/301.4.2/III/2022) for this study was granted by the Ethical Committee of Dr. Soetomo Hospital, Surabaya, Indonesia.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study. Written informed consent was obtained from the patient(s) to publish this paper.

Data Availability Statement: The raw data are available in DOI: <https://doi.org/10.6084/m9.figshare.22724183>.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Weekly Epidemiological Update on COVID-19—1 March 2023. Available online: <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19---1-march-2023> (accessed on 1 March 2023).
- Infeksi Emerging Kementerian Kesehatan RI. Available online: <https://infeksiemerging.kemkes.go.id/dashboard/covid-19> (accessed on 12 March 2023).
- Li, T.Z.; Cao, Z.H.; Chen, Y.; Cai, M.T.; Zhang, L.Y.; Xu, H.; Zhang, J.-Y.; Ma, C.-H.; Liu, Y.; Gao, L.-J.; et al. Duration of SARS-CoV-2 RNA shedding and factors associated with prolonged viral shedding in patients with COVID-19. *J. Med. Virol.* **2020**, *93*, 506–512. [CrossRef]
- Zhou, B.; She, J.; Wang, Y.; Ma, X. Duration of Viral Shedding of Discharged Patients With Severe COVID-19. *Clin. Infect. Dis.* **2020**, *71*, 2240–2242. [CrossRef]
- Wang, Y.; Zhang, L.; Sang, L.; Ye, F.; Ruan, S.; Zhong, B.; Song, T.; Alshukairi, A.N.; Chen, R.; Zhang, Z.; et al. Kinetics of viral load and antibody response in relation to COVID-19 severity. *J. Clin. Investig.* **2020**, *130*, 5235–5244. [CrossRef] [PubMed]
- Fu, Y.; Han, P.; Zhu, R.; Bai, T.; Yi, J.; Zhao, X.; Tao, M.; Quan, R.; Chen, C.; Zhang, Y.; et al. Risk factors for viral RNA shedding in COVID-19 patients. *Eur. Respir. J.* **2020**, *56*, 2001190. [CrossRef] [PubMed]
- Karia, R.; Nagraj, S. A Review of Viral Shedding in Resolved and Convalescent COVID-19 Patients. *SN Compr. Clin. Med.* **2020**, *2*, 2086–2095. [CrossRef] [PubMed]
- Leung, W.F.; Chorlton, S.; Tyson, J.; Al-Rawahi, G.N.; Jassem, A.N.; Prystajek, N.; Masud, S.; Deans, G.D.; Chapman, M.G.; Mirzanejad, Y.; et al. COVID-19 in an immunocompromised host: Persistent shedding of viable SARS-CoV-2 and emergence of multiple mutations: A case report. *Int. J. Infect. Dis.* **2021**, *114*, 178–182. [CrossRef]
- Chen, C.F.; Tsai, T.Y.; Yu, C.H.; Cheng, H.L.; Yeh, T.Y. Prolonged viral shedding and new mutations of COVID-19 could complicate the control of the pandemic. *Access Microbiol.* **2020**, *2*, acmi000133. [CrossRef]

10. Wibisono, E.; Hadi, U.; Bramantono Arfijanto, M.V.; Rusli, M.; Rahman, B.E.; Asmarawati, T.P.; Choirunnisa, M.L.; Rahayu, D.R.P. National early warning score (NEWS) 2 predicts hospital mortality from COVID-19 patients. *Ann. Med. Surg.* **2022**, *76*, 103462. [[CrossRef](#)] [[PubMed](#)]
11. Nugroho, C.W.; Suryantoro, S.D.; Yuliasih, Y.; Rosyid, A.N.; Asmarawati, T.P.; Andrianto, L.; Setiawan, H.W.; Mahdi, B.A.; Windradi, C.; Agustin, E.D.; et al. Optimal use of tocilizumab for severe and critical COVID-19: A systematic review and meta-analysis. *F1000Research* **2021**, *10*, 73. [[CrossRef](#)]
12. Soegiarto, G.; Wulandari, L.; Purnomosari, D.; Dhia Fahmita, K.; Ikhwan Gautama, H.; Tri Hadmoko, S.; Prasetyo, M.E.; Mahdi, B.A.; Arafah, N.; Prasetyaningtyas, D.; et al. Hypertension is associated with antibody response and breakthrough infection in health care workers following vaccination with inactivated SARS-CoV-2. *Vaccine* **2022**, *40*, 4046–4056. [[CrossRef](#)]
13. Asmarawati, T.P.; Suryantoro, S.D.; Rosyid, A.N.; Marfiani, E.; Windradi, C.; Mahdi, B.A.; Sutanto, H. Predictive Value of Sequential Organ Failure Assessment, Quick Sequential Organ Failure Assessment, Acute Physiology and Chronic Health Evaluation II, and New Early Warning Signs Scores Estimate Mortality of COVID-19 Patients Requiring Intensive Care Unit. *Indian J. Crit. Care Med.* **2022**, *26*, 466–473. [[CrossRef](#)] [[PubMed](#)]
14. Asmarawati, T.P.; Rosyid, A.N.; Suryantoro, S.D.; Mahdi, B.A.; Windradi, C.; Wulaningrum, P.A.; Arifianto, M.V.; Bramantono, B.; Triyono, E.A.; Rusli, M.; et al. The clinical impact of bacterial co-infection among moderate, severe and critically ill COVID-19 patients in the second referral hospital in Surabaya. *F1000Research* **2021**, *10*, 1–16. [[CrossRef](#)]
15. Burhan, E.; Susanto, A.D.; Isbaniah, F.; Sally, A.N.; Eka, G.; Ceva, W.P.; Susilo, A.; Firdaus, I.; Santoso, A.; Juzar, D.A.; et al. *Pedoman Tatalaksana COVID-19*, 3rd ed.; PDPI; PERKI; PAPDI; PERDATIN; IDAI: Jakarta, Indonesia, 2020; pp. 36–37.
16. Qi, L.; Yang, Y.; Jiang, D.; Tu, C.; Wan, L.; Chen, X.; Li, Z. Factors associated with the duration of viral shedding in adults with COVID-19 outside of Wuhan, China: A retrospective cohort study. *Int. J. Infect. Dis.* **2020**, *96*, 531–537. [[CrossRef](#)] [[PubMed](#)]
17. Xu, K.; Chen, Y.; Yuan, J.; Yi, P.; Ding, C.; Wu, W.; Li, Y.; Ni, Q.; Zou, R.; Li, X.; et al. Factors Associated With Prolonged Viral RNA Shedding in Patients with Coronavirus Disease 2019 (COVID-19). *Clin. Infect. Dis.* **2020**, *71*, 799–806. [[CrossRef](#)]
18. Feng, Z.; Li, J.; Yao, S.; Yu, Q.; Zhou, W.; Mao, X.; Li, H.; Kang, W.; Ouyang, X.; Mei, J.; et al. Clinical Factors Associated with Progression and Prolonged Viral Shedding in COVID-19 Patients: A Multicenter Study. *Aging Dis.* **2020**, *11*, 1069–1081. [[CrossRef](#)]
19. Zhou, F.; Yu, T.; Du, R.; Fan, G.; Liu, Y.; Liu, Z.; Xiang, J.; Wang, Y.; Song, B.; Gu, X.; et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: A retrospective cohort study. *Lancet* **2020**, *395*, 1054–1062. [[CrossRef](#)]
20. Aly, M.H.; Rahman, S.S.; Ahmed, W.A.; Alghamedi, M.H.; AlShehri, A.A.; Alkalkami, A.M.; Hassan, M.H. Indicators of Critical Illness and Predictors of Mortality in COVID-19 Patients. *Infect. Drug Resist.* **2020**, *13*, 1995–2000. [[CrossRef](#)]
21. Zhang, G.; Hu, C.; Luo, L.; Fang, F.; Chen, Y.; Li, J.; Peng, Z.; Pan, H. Clinical features and short-term outcomes of 221 patients with COVID-19 in Wuhan, China. *J. Clin. Virol.* **2020**, *127*, 104364. [[CrossRef](#)]
22. Zhou, C.; Zhang, T.; Ren, H.; Sun, S.; Yu, X.; Sheng, J.; Shi, Y.; Zhao, H. Impact of age on duration of viral RNA shedding in patients with COVID-19. *Aging* **2020**, *12*, 22399–22404. [[CrossRef](#)]
23. Pradhan, A.; Olsson, P.-E. Sex differences in severity and mortality from COVID-19: Are males more vulnerable? *Biol. Sex Differ.* **2020**, *11*, 53. [[CrossRef](#)]
24. Muhamad, S.A.; Ugusman, A.; Kumar, J.; Skiba, D.; Hamid, A.A.; Aminuddin, A. COVID-19 and Hypertension: The What, the Why, and the How. *Front. Physiol.* **2021**, *12*, 665064. [[CrossRef](#)] [[PubMed](#)]
25. Buetti, N.; Trimboli, P.; Mazzuchelli, T.; Priore, E.L.; Balmelli, C.; Trkola, A.; Conti, M.; Martinetti, G.; Elzi, L.; Ceschi, A.; et al. Diabetes mellitus is a risk factor for prolonged SARS-CoV-2 viral shedding in lower respiratory tract samples of critically ill patients. *Endocrine* **2020**, *70*, 454–460. [[CrossRef](#)]
26. Codo, A.C.; Davanzo, G.G.; de Brito Monteiro, L.; de Souza, G.F.; Muraro, S.P.; Virgilio-Da-Silva, J.V.; Prodonoff, J.S.; Carregari, V.C.; de Biagi Junior, C.A.O.; Crunfli, F.; et al. Elevated Glucose Levels Favor SARS-CoV-2 Infection and Monocyte Response through a HIF-1 α /Glycolysis-Dependent Axis. *Cell Metab.* **2020**, *32*, 437–446.e5. [[CrossRef](#)]
27. Zhang, S.; Zhu, H.; Ye, H.; Hu, Y.; Zheng, N.; Huang, Z.; Xiong, Z.; Fu, L.; Cai, T. Risk factors for prolonged virus shedding of respiratory tract and fecal in adults with severe acute respiratory syndrome coronavirus-2 infection. *J. Clin. Lab. Anal.* **2021**, *35*, e23923. [[CrossRef](#)]
28. Cunha, M.D.P.; Vilela, A.P.P.; Molina, C.V.; Acuna, S.M.; Muxel, S.M.; Barroso, V.D.M.; Baroni, S.; de Oliveira, L.G.; de Souza, A.Y.; Peron, J.P.S.; et al. Atypical Prolonged Viral Shedding With Intra-Host SARS-CoV-2 Evolution in a Mildly Affected Symptomatic Patient. *Front. Med.* **2021**, *8*, 1–11. [[CrossRef](#)]
29. Menghua, W.; Xin, Z.; Jianwei, L.; Yu, Z.; Qinwei, Y. Case report: One case of coronavirus disease 2019 (COVID-19) in a patient co-infected by HIV with a normal CD4+ T cell count. *AIDS Res. Ther.* **2020**, *17*, 46. [[CrossRef](#)] [[PubMed](#)]
30. Yousaf, M.; Hameed, M.; Alsoub, H.; Khatib, M.; Jamal, W.; Ahmad, M. COVID-19: Prolonged viral shedding in an HIV patient with literature review of risk factors for prolonged viral shedding and its implications for isolation strategies. *Clin. Case Rep.* **2021**, *9*, 1397–1401. [[CrossRef](#)] [[PubMed](#)]
31. Askanase, A.D.; Khalili, L.; Buyon, J.P. Thoughts on COVID-19 and autoimmune diseases. *Lupus Sci. Med.* **2020**, *7*, e000396. [[CrossRef](#)] [[PubMed](#)]
32. Aghili, S.M.M.; Ebrahimpur, M.; Arjmand, B.; Shadman, Z.; Pejman Sani, M.; Qorbani, M.; Larijani, B.; Payab, M. Obesity in COVID-19 era, implications for mechanisms, comorbidities, and prognosis: A review and meta-analysis. *Int. J. Obes.* **2021**, *45*, 998–1016. [[CrossRef](#)]

33. De Leeuw, A.J.M.; Oude Luttikhuis, M.A.M.; Wellen, A.C.; Müller, C.; Calkhoven, C.F. Obesity and its impact on COVID-19. *J. Mol. Med.* **2021**, *99*, 899–915. [[CrossRef](#)]
34. Bitker, L.; Dhelft, F.; Chauvelot, L.; Frobert, E.; Folliet, L.; Mezidi, M.; Trouillet-Assant, S.; Belot, A.; Lina, B.; Wallet, F.; et al. Protracted viral shedding and viral load are associated with ICU mortality in Covid-19 patients with acute respiratory failure. *Ann. Intensiv. Care* **2020**, *10*, 167. [[CrossRef](#)] [[PubMed](#)]
35. Ryoo, S.M.; Kim, W.Y.; Sohn, C.H.; Seo, D.W.; Oh, B.J.; Lee, J.H.; Lee, Y.S.; Lim, K.S. Factors promoting the prolonged shedding of the pandemic (H1N1) 2009 influenza virus in patients treated with oseltamivir for 5 days. *Influenza Other Respir. Viruses* **2012**, *7*, 833–837. [[CrossRef](#)] [[PubMed](#)]
36. Langford, B.J.; So, M.; Raybardhan, S.; Leung, V.; Westwood, D.; MacFadden, D.R.; Soucy, J.-P.R.; Daneman, N. Bacterial co-infection and secondary infection in patients with COVID-19: A living rapid review and meta-analysis. *Clin. Microbiol. Infect.* **2020**, *26*, 1622–1629. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

Duration of SARS-CoV-2 RNA Shedding Is Significantly Influenced by Disease Severity, Bilateral Pulmonary Infiltrates, Antibiotic Treatment, and Diabetic Status: Consideration for Isolation Period

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

14%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	www.hindawi.com Internet Source	1%
2	Rawan Nehme, Arthur Chervet, Caroline Decombat, Lucie Longechamp et al. "Aspalathus linearis (Rooibos) Targets Adipocytes and Obesity-Associated Inflammation", <i>Nutrients</i> , 2023 Publication	1%
3	Submitted to University of Bedfordshire Student Paper	1%
4	eurjmedres.biomedcentral.com Internet Source	1%
5	jiaphd.org Internet Source	1%
6	Snigdha Alur-Gupta, Mary Regina Boland, Anuja Dokras, Melissa A. Haendel et al. "Risk of severe acute respiratory syndrome	<1%

coronavirus 2 infection among women with polycystic ovary syndrome", Fertility and Sterility, 2023

Publication

7

Submitted to University of Witwatersrand

Student Paper

<1 %

8

Guo-Qing Qian, Xue-Qin Chen, Ding-Feng Lv, Ada Hoi Yan Ma, Li-Ping Wang, Nai-Bin Yang, Xiao-Min Chen. "Duration of SARS-CoV-2 viral shedding during COVID-19 infection", Infectious Diseases, 2020

Publication

<1 %

9

Submitted to Universidad de Sevilla

Student Paper

<1 %

10

Elicia Pettiroso, Michelle Giles, Stephen Cole, Megan Rees. "COVID - 19 and pregnancy: A review of clinical characteristics, obstetric outcomes and vertical transmission", Australian and New Zealand Journal of Obstetrics and Gynaecology, 2020

Publication

<1 %

11

Laksmi Wulandari, Berliana Hamidah, Cennikon Pakpahan, Nevy Shinta Damayanti et al. "Initial Study on TMPRSS2 p.Val160Met Genetic Variant in COVID-19 patients", Research Square, 2021

Publication

<1 %

12	www.lenus.ie Internet Source	<1 %
13	bsdwebstorage.blob.core.windows.net Internet Source	<1 %
14	www.thieme-connect.de Internet Source	<1 %
15	Umair Mallick. "Chapter 16 Diabetes Mellitus and COVID19", Springer Science and Business Media LLC, 2022 Publication	<1 %
16	assets.researchsquare.com Internet Source	<1 %
17	apicareonline.com Internet Source	<1 %
18	stacks.cdc.gov Internet Source	<1 %
19	www.cimm-icmm.org Internet Source	<1 %
20	Wen-Yu Su, Pin-Xian Du, Harvey M. Santos, Tzong-Shiann Ho et al. "Antibody Profiling in COVID-19 Patients with Different Severities by Using Spike Variant Protein Microarrays", Analytical Chemistry, 2022 Publication	<1 %
21	www.ncbi.nlm.nih.gov Internet Source	<1 %

<1 %

22

Adeline Penchenat, Zarinn Alavi, Christopher Payan, Jacques - Olivier Pers. "Prolonged 59 - day course of COVID - 19: The case of a SARS - CoV - 2 shedding persistency in a healthcare provider", Clinical Case Reports, 2020

Publication

<1 %

23

Submitted to Feng Chia University

Student Paper

<1 %

24

Submitted to Lindenwood University

Student Paper

<1 %

25

Manina Günter, Karin Anne Lydia Mueller, Mathew Salazar, Sarah Gekeler et al. "Immune signature of patients with cardiovascular disease – in-depth immunophenotyping predicts increased risk for a severe course of COVID-19", Cold Spring Harbor Laboratory, 2023

Publication

<1 %

26

www.ajtmh.org

Internet Source

<1 %

27

www.ijccm.org

Internet Source

<1 %

28

www.ingentaconnect.com

Internet Source

<1 %

29

Submitted to Çankırı Karatekin University

Student Paper

<1 %

30

Cristina Rodríguez-Grande, Javier Adán-Jiménez, Pilar Catalán, Luis Alcalá et al. "Inference of Active Viral Replication in Cases with Sustained Positive Reverse Transcription-PCR Results for SARS-CoV-2", Journal of Clinical Microbiology, 2021

Publication

<1 %

31

Tri Pudy Asmarawati, Okla Sekar Martani, Bramantono Bramantono, Muhammad Vitanata Arfijanto. "Prolonged fever and exaggerated hypercoagulopathy in malaria vivax relapse and COVID-19 co-infection: a case report", Malaria Journal, 2022

Publication

<1 %

32

biotech.chinaxiv.org

Internet Source

<1 %

33

journals.lww.com

Internet Source

<1 %

34

www.ethndis.org

Internet Source

<1 %

35

covid19-data.nist.gov

Internet Source

<1 %

36	warm.dovepress.com Internet Source	<1 %
37	1library.net Internet Source	<1 %
38	docksci.com Internet Source	<1 %
39	iosrjournals.org Internet Source	<1 %
40	jkms.org Internet Source	<1 %
41	translational-medicine.biomedcentral.com Internet Source	<1 %
42	Brand, Amy, Liz Allen, Micah Altman, Marjorie Hlava, and Jo Scott. "Beyond authorship: attribution, contribution, collaboration, and credit", Learned Publishing, 2015. Publication	<1 %
43	Rentha Monica Simamora, Muhammad Vitanata Arfijanto, Musofa Rusli, Budi Utomo, Cennikon Pakpahan, Garry Prasetyo Adi. "Clinical Signs and Laboratory Parameters as Predictors of Mortality among Hospitalized Human Immunodeficiency Virus-Infected Adult Patients at Tertiary Hospital in Surabaya", Open Access Macedonian Journal of Medical Sciences, 2021	<1 %

44	up.edu.ph Internet Source	<1 %
45	www.mdpi.com Internet Source	<1 %
46	www.science.gov Internet Source	<1 %
47	www.sciencegate.app Internet Source	<1 %
48	Shayne Mason, Mari van Reenen, Theresa Rossouw, Zander Lindeque, Roan Louw. "Phenylalanine metabolism and tetrahydrobiopterin bio-availability in COVID-19 and HIV", Heliyon, 2023 Publication	<1 %
49	lppm-unissula.com Internet Source	<1 %
50	www.cdc.gov Internet Source	<1 %
51	www.microbiologyresearch.org Internet Source	<1 %
52	Anthony Ravussin, Anna Hayman Robertson, Asia-Sophia Wolf, Kristine Blix et al. "Determinants of humoral and cellular immune responses to three doses of mRNA	<1 %

SARS-CoV-2 vaccines in older adults: a longitudinal cohort study", The Lancet Healthy Longevity, 2023

Publication

53

Hendra Gunawan, Eridani Eridani. "Fractional Integrals and Generalized Olsen Inequalities", Kyungpook mathematical journal, 2009

Publication

54

Muslim Suardi. "Clinical Outcomes of Antiviral Therapy on Comorbid and non-Comorbid T2-DM COVID-19 Patients at Rasidin Hospital", JOPS (Journal Of Pharmacy and Science), 2022

Publication

55

Nissar Shaikh, Mohammad Al Wraidat, Anood A. Al-Assaf, Salma K. Al-Kaabi et al. "COVID-19 in Patients with Cancer: Insights from a Tertiary Care Intensive Care Unit", Research Square Platform LLC, 2023

Publication

56

Pooja Lal, Dhyanesh A. Patel, Michael F. Vaezi. "Gastrointestinal Manifestations of COVID-19", Elsevier BV, 2024

Publication

57

Smaranda Gliga, Nadine Luebke, Alexander Killer, Henning Gruell et al. "Rapid selection of sotrovimab escape variants in SARS-CoV-2 Omicron infected immunocompromised patients", Clinical Infectious Diseases, 2022

<1 %

<1 %

<1 %

<1 %

<1 %

58 Tingxuan Yin, Yuanjun Li, Ying Ying, Zhijun Luo. "Prevalence of comorbidity in Chinese patients with COVID-19: systematic review and meta-analysis of risk factors", BMC Infectious Diseases, 2021
Publication

59 Zoya Morani, Saumil Patel, Sudeshna Ghosh, Falah Abu Hassan et al. "COVID-19 in HIV: a Review of Published Case Reports", SN Comprehensive Clinical Medicine, 2020
Publication

60 nrl.northumbria.ac.uk
Internet Source

61 repository.unair.ac.id
Internet Source

62 www.coursehero.com
Internet Source

63 www.e-trd.org
Internet Source

64 www.tandfonline.com
Internet Source

65 Karl Hagman, Magnus Hedenstierna, Johan Rudling, Patrik Gille-Johnson et al. "Duration of SARS-CoV-2 viremia and its correlation to mortality and inflammatory parameters in

patients hospitalized for COVID-19: a cohort study", Diagnostic Microbiology and Infectious Disease, 2022

Publication

66

Lingye Chen, Lyra B. Olson, Ibtehaj A. Naqvi, Bruce A. Sullenger, Smita K Nair, Loretta G. Que, Thomas N. Denny, Bryan D. Kraft. "SARS-CoV-2 viremia but not respiratory viral load is associated with ventilator complications", Research Square Platform LLC, 2022

Publication

<1 %

67

Marliana S. Rejeki, Nana Sarnadi, Retno Wihastuti, Vininta Fazharyasti et al. "Convalescent plasma therapy in patients with moderate-to-severe COVID-19: A study from Indonesia for clinical research in low- and middle-income countries", EClinicalMedicine, 2021

Publication

<1 %

68

Mohammad Mahdi Rezaei Tolzali, Maryam Noori, Pौरya Shokri, Shayan Rahmani et al. "Efficacy of tocilizumab in the treatment of COVID - 19: An umbrella review", Reviews in Medical Virology, 2022

Publication

<1 %

69

Siggeir F. Brynjolfsson, Hildur Sigurgrimsdottir, Olafur Gudlaugsson, Mar Kristjansson et al. "Determining SARS-CoV-2

<1 %

non-infectivity state–A brief overview", Frontiers in Public Health, 2022

Publication

70

www.uk-essen.de

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Duration of SARS-CoV-2 RNA Shedding Is Significantly Influenced by Disease Severity, Bilateral Pulmonary Infiltrates, Antibiotic Treatment, and Diabetic Status: Consideration for Isolation Period

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13