

Prolonged Fever And Exaggerated Hypercoagulopathy In Malaria Vivax Relapse And COVID-19 Co-Infection: A Case Report

by Muhammad Vitanata Arfijanto

Submission date: 08-May-2023 09:51AM (UTC+0800)

Submission ID: 2086970531

File name: alaria_Vivax_Relapse_And_COVID-19_Co-Infection_A_Case_Report.pdf (1.26M)

Word count: 4559

Character count: 25359

CASE REPORT

Open Access

Prolonged fever and exaggerated hypercoagulopathy in malaria vivax relapse and COVID-19 co-infection: a case report



Tri Pudy Asmarawati^{1,2,3*}, Okla Sekar Martani⁴, Bramantono Bramantono^{1,3} and Muhammad Vitanata Arfijanto^{1,3}

48

Abstract

Background: Coronavirus disease 2019 (COVID-19) often causes atypical clinical manifestations similar to other infectious diseases. In malaria-endemic areas, the pandemic situation will very likely result in co-infection of COVID-19 and malaria, although reports to date are still few. Meanwhile, this disease will be challenging to diagnose in areas with low malaria prevalence because the symptoms closely resemble COVID-19.

Case presentation: A 23-year-old male patient presented to the hospital with fever, anosmia, headache, and nausea 1 week before. He was diagnosed with COVID-19 and treated for approximately 10 days, then discharged to continue self-quarantine at home. 2 weeks later, he returned to the hospital with a fever raised intermittently every 2 days and marked by a chilling-fever-sweating cycle. A laboratory test for malaria and a nasopharyngeal swab for SARS CoV-2 PCR were conducted, confirming both diagnoses. The laboratory examination showed markedly elevated D-dimer. He was treated with dihydroartemisinin-piperazine (DHP) 4 tablets per day for 3 days and primaquine 2 tablets per day for 14 days according to Indonesian National Anti-malarial Treatment Guidelines. After 6 days of treatment, the patient had no complaints, and the results of laboratory tests had improved. This report describes the key points in considering the differential diagnosis and prompt treatment of malaria infection during the pandemic of COVID-19 in an endemic country to prevent the worse clinical outcomes. COVID-19 and malaria may also cause a hypercoagulable state, so a co-infection of those diseases may impact the prognosis of the disease.

Conclusion: This case report shows that considering the possibility of a co-infection in a COVID-19 patient who presents with fever can prevent delayed treatment that can worsen the disease outcome. Paying more attention to a history of travel to malaria-endemic areas, a history of previous malaria infection, and exploring anamnesis regarding the fever patterns in patients are important points in making a differential diagnosis of malaria infection during the COVID-19 pandemic.

Keywords: COVID-19, malaria, *Plasmodium vivax*, Infectious diseases

* Correspondence: tpasmarawati@fk.unair.ac.id

¹Tropical and Infectious Diseases Division, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java 60115, Indonesia

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

29

37

total number of 94,119 deaths or a case fatality rate of 2.76% [2]. The SARS-CoV-2 infection causes various clinical manifestations ranging from asymptomatic to a broad spectrum of symptoms, such as mild symptoms of upper respiratory tract disorders and life-threatening sepsis. Recent studies have also reported that COVID-19 patients frequently develop hypercoagulopathy and a high prevalence of thromboembolic events [3]. The most common feature in hospitalized COVID-19 patients is fever (70–90%), impaired sense of smell and taste (64–80%), dry cough (60–86%), shortness of breath (53–80%), fatigue (38%), nausea/vomiting or diarrhoea (15–39%), and myalgia (15–44%) [4]. Coagulopathy with abnormally elevated D-dimer levels has been reported in 260 of 560 cases of COVID-19 (46.4%), with the prevalence of 43% in non-severe patients compared with 60% in critically ill ICU patients.

Malaria is an infectious disease caused by protozoa of the genus *Plasmodium*, which is transmitted through the bite of an *Anopheles* mosquito. Malaria infections can cause severe clinical manifestations and death if not promptly diagnosed and treated. There are five species of *Plasmodium* that cause malaria in humans, with the most deaths caused by *Plasmodium falciparum*. *Plasmodium vivax* predominates as a cause of morbidity, while *Plasmodium ovale* and *Plasmodium malariae* rarely cause severe malaria. *Plasmodium knowlesi* primarily infects macaques in Southeast Asia, but humans living around these animals can become infected [5].

Malaria remains a serious global health problem, especially in endemic countries. In 2019, an estimated 229 million malaria cases occurred worldwide, with 3% of all cases caused by *Plasmodium vivax*, which is the predominant parasite responsible for 51.7% of malaria cases in Southeast Asia, including Indonesia [6]. Patients with malaria will usually experience paroxysmal fever, fatigue, malaise, and myalgia [7]. A co-infection between COVID-19 and malaria is rarely reported, and the mechanism is unclear. It is suggested that a co-infection leads to excess proinflammatory responses and pro-coagulant states that might result in more severe manifestations and poor prognosis. Heightened clinical suspicion is needed to diagnose other infections during the COVID-19 pandemic, especially for patients with a travelling history of malaria-endemic areas. In this case report, we found a confirmed COVID-19 patient diagnosed with malaria.

5 Case presentation

A 23-year-old male patient came to the emergency department at Universitas Airlangga Hospital presenting fever, anosmia, headache, and nausea 5 days before. He was diagnosed with COVID-19, confirmed by the RT-PCR amplification of SARS-CoV-2 virus nucleic



Fig. 1 Chest x-ray on first admission

acid on a nasopharyngeal swab. The patient had abnormal laboratory findings, such as lymphopenia, neutrophilia, thrombocytopenia, and high CRP, with a normal chest X-ray (Fig. 1). The patient was treated with Favipiravir at 1600 mg per 12 h on day 1, then 600 mg per 12 h on day 2–day 5, intravenous drip of paracetamol 1000 mg every 8 h, oral vitamin C 500 mg per 8 h, and oral vitamin D 1000 IU per day according to Indonesian national COVID-19 treatment guidelines. After 10 days of treatment, the patient was still PCR positive despite no complaints. He was then discharged to continue self-quarantine at home.

He returned to the hospital two weeks later with general weakness and fever. The fever did not resolve during his self-quarantine, was raised intermittently every 2 days, and was marked by a chilling-fever-sweating cycle. The fever temperature measured was around 39.7°–40 °C. The fever improved when the patient took paracetamol, but the fever returned 6 h later. The fever was accompanied by malaise and muscle pain. The next day the patient was fever-free and able to carry out his daily activities. The physical examination revealed a body temperature of 38 °C, a blood pressure of 112/72 mmHg, a heart rate of 93 bpm, a respiratory rate of 20 breaths/minute, and oxygen saturation of 98% under ambient air. The patient weighs 65 kg with a height of 170 cm (BMI 22.5 kg/m²). Other physical examinations were normal.

The laboratory results reflected a white blood cell count of 8750/μL with 76.3% neutrophils, 13.5% lymphocytes, and 0.6% eosinophils. Haemoglobin level and platelet counts were 12.2 g/dL and 231,000/μL, respectively. CRP and D-dimer levels were 41.51 mg/L and 9.52 mg/L, respectively. Serum electrolytes and renal function tests were normal, with urea and creatinine serum 6 mg/dL and 0.4 mg/dL, respectively. The liver enzymes test was normal, with ALT being 14 IU/L and AST being 25 IU/L. Electrocardiography showed normal sinus rhythm and

axis. The chest X-ray was unremarkable (Fig. 2). The RT-PCR amplification of the SARS-CoV-2 virus nucleic acid test from the nasopharyngeal swab was still positive.

From the previous history, he said that he traveled to Timika District, Papua, for office work for the last 2 years. In May 2021, the patient was infected with *P vivax*, with initial symptoms including fever, dizziness, nausea, and muscle pain. The patient was treated until declared fully cured. A laboratory test for malaria was performed. The rapid malaria test was positive for PAN antigen, and microscopic diagnosis on blood smear revealed *Plasmodium vivax* on ring form, trophozoite, and gametocyte stage (Figs. 3, 4, 5). Based on anamnesis, physical examination, and laboratory results, the patient was diagnosed with confirmed COVID-19 with hypercoagulopathy




Fig. 2 Chest x-ray on second admission

and malaria vivax relapse (Fig. 6). According to Indonesian National Guidelines for Antimalarial Treatment [8], he was treated with dihydroartemisinin-piperazine (DHP) 4 tablets per day for 3 days and primaquine 2 tablets per day for 14 days. The G6PD status was not tested. The patient was given an intravenous drip of paracetamol 1000 mg every 8 h and a subcutaneous injection of heparin 5000 IU every 12 h during treatment. After six days of treatment, the patient had no complaints, and the results of laboratory tests had improved. The patient was discharged from the hospital, continued self-isolation at home, and followed up with the internal medicine outpatient clinic two weeks later. Furthermore, the patient was called later and reported feeling healthy with no complaints.

Discussion

Indonesia is one of the tropical countries in the world still facing a high risk of malaria. Approximately 80% of cities or districts in Indonesia are endemic to malaria, one of




Fig. 3 Thick blood smear




Fig. 4 Gametocyte in thin blood smear




Fig. 5 Ring stage in thin blood smear

which is Papua. Based on the data from Annual Parasite Incidence (API) in 2019, the number of positive cases of malaria in Papua is 64.03% per 1000 population [9]. *Plasmodium vivax* malaria still causes significant morbidity in endemic areas. Reactivation of the dormant stage of hypnozoite can cause *P. vivax* relapse [10]. A single inoculation by a female *Anopheles* mosquito can be followed by multiple relapses [11].

Plasmodium vivax is capable of undergoing early and frequent relapses. In addition to blood-stage parasite replication, *P. vivax* has a dormant stage of hypnozoites that can persist in the liver and reactivate to cause relapses weeks or even years later [12]. This reactivation may be triggered by the host's inflammatory response to systemic illnesses or parasitic and bacterial infections, but not viral infections [13]. The estimated ratio of *P. vivax* relapses to total infections is 76–90% in the Papua New Guinea cohort study and 79% in the Thailand cohort study [12]. Based on the study conducted by Chu et al. *P. vivax* parasitaemia recurrences, mainly relapses, occurred in 377 from 644 (59%) patients treated with artesunate, chloroquine, or chloroquine-primaquine. A history of the previous malaria was more common in patients with recurrences (61%) than without recurrence (39%). About 90% of all *P. vivax* recurrences occurred by week 16 following treatment. Primaquine, as the radical cure with an estimated efficacy of 92%, would reduce the risk of




Fig. 6 COVID-19 and Malaria relapse disease course

relapses and lessen the significant burden of morbidity caused by *P. vivax* [14].

The pathophysiology behind the co-infection of COVID-19 with *P. vivax* malaria remains unclear. It is unknown whether SARS-CoV-2 infection reduces the immunity that leads to malaria reactivation or if complications from malaria increase the susceptibility to get COVID-19 [15]. Based on the disease course, this patient was suspected of having reactivation of *P. vivax* infection due to COVID-19 infection. The mechanism that caused the reactivation is still unclear and may be attributed to the cytokine response to COVID-19. There is still a possibility of natural reactivation or reinfection not related to COVID-19 infection that cannot be completely ruled out [16]. The excessive proinflammatory cytokine response generated in co-infection of COVID-19 and Malaria can worsen the prognosis [17].

The patient was also suspected of having prolonged viral shedding of SARS-CoV-2; the PCR was still positive on day 38 after symptom onset. This phenomenon may be attributable to several risk factors, including male sex, disease severity, delayed hospital admission, underlying comorbidities, and a higher pro-inflammatory cytokine level [18–20]. In a large Italian cohort study, Mondì et al. stated that pro-inflammatory marker changes, especially lymphopenia, and elevated D-dimer, were strongly associated with prolonged viral shedding [19]. A higher neutrophil count and CRP were also mentioned in a report by Gao et al. as independent risk factors of prolonged viral shedding found in COVID-19 patients with bacterial co-infection [20]. Rather than bacterial infection, the abnormal proinflammatory cytokine and infection markers related to prolonged SARS-CoV-2 RNA shedding in this case were exaggerated by malaria vivax relapse.

Early symptoms of SARS-CoV-2 infection, such as fever, fatigue, and myalgia, similar to those of malaria, can lead to delayed diagnosis, especially in malaria-endemic areas [16]. In contrast to severe malaria, which is often caused by *P. falciparum*, where neurological symptoms are predominant and more severe symptoms such as loss of consciousness, signs of focal neurological abnormalities, and severe anaemia can further narrow the diagnosis of malaria. The results of blood tests can also be confusing. Therefore, a diagnostic test for both malaria and COVID-19 is crucial. The principal diagnosis of malaria to date is the microscopic examination to detect the presence of *Plasmodium* at all stages. However, the sensitivity and specificity of this examination are highly dependent on the examiner's subjectivity. If a microscopy test is not available, malaria rapid diagnostic test (RDT) can be an alternative to confirm the diagnosis. Another method using a PCR is more sensitive than microscopy, but the results take too long, so it is rarely

used [21]. *Plasmodium* spp and COVID-19 have incubation periods that are not much different. For COVID-19, the incubation period reaches 14 days from exposure with a median value of 4–5 days, while for malaria, the incubation period varies from 7 to 30 days [21]. Hence, it is essential to gain a thorough anamnesis in differentiating these two diagnoses.

Hypercoagulopathy is common in COVID-19 patients, especially in those with severe disease [22]. SARS-CoV-2 induces tissue factor expression, a primary initiator of the coagulation cascade, by cytokines produced by inflammatory cells. Moreover, SARS-CoV-2 causes endothelial dysfunction through an angiotensin-converting enzyme-2 (ACE-2) receptor expressed on the surface of vascular endothelial cells and induces neutrophil extracellular traps (NETs) release, which activates the coagulation pathways and platelet [23]. Abnormal coagulation parameters such as elevated D-dimer and fibrin degradation product levels and prolonged prothrombin time are related to a poor outcome. The most common manifestations of COVID-19 hypercoagulopathy are venous thromboembolism and arterial thrombotic complications, including pulmonary embolism and stroke [3]. COVID-19 patients are at risk for developing disseminated intravascular coagulation (DIC), pulmonary haemorrhage, and thrombosis. Thrombocytopenia associated with a higher risk of severe COVID-19 is suspected to be caused by platelet consumption in the lungs and infected haematopoietic stem cells and megakaryocytes [24].

Malaria is also strongly associated with a hypercoagulopathy condition through activation of the coagulation cascade triggered by proinflammatory cytokines [e.g., tumour necrosis factor (TNF) and interleukin (IL)-6] [25]. The most common coagulopathy condition is microthrombotic complications, besides thrombosis of large vessels, including cerebral venous thrombosis and pulmonary embolism. Thrombocytopenia is a common finding (60–80%) that may be due to impaired coagulation, splenomegaly, bone marrow disorders, antibody-mediated platelet destruction, oxidative stress, and platelet aggregation. DIC and bleeding are related to high mortality, occurring only in severe malaria. Tissue factors released from damaged vascular endothelial cells and the lysis of activated platelets contribute to the development of a pro-coagulant state similar to the underlying mechanism in COVID-19 [26]. Therefore, *Plasmodium* spp. and SARS-CoV-2 co-infection could lead to severe coagulopathy and worse outcomes than with either infection alone. In this case, the patient had a high risk of mortality due to hypercoagulopathy, characterized by markedly increased D-dimer levels which was also aggravated by malaria vivax relapse. Regarding this condition,

Table 1 A Literature review of COVID-19 and *Plasmodium vivax* Co-infection

No	Author	Country	Year of Publication	Age	Clinical symptoms	History of <i>P. vivax</i> infection
1	Shahid et al. [11]	Qatar	2021	55	Dry cough, high-grade fever, chills, rigors, profuse sweating, lethargy	Documented (1 year before)
2	Ray et al. [27]	India	2020	67	Fever, shortness of breath	Not documented
3	Kishore et al. [16]	India	2020	10	High-grade fever, chills, rigors, headache, cold, cough, abdominal pain	Documented (6 months before)
4	Sardar et al. [17]	Qatar	2020	34	Fever, myalgia, vomiting, right upper quadrant abdominal pain	Not documented (but there was a travel history to Pakistan 3 months before)

thromboprophylaxis has a crucial role in reducing the risk of thrombotic events.

There was also some similar report regarding malaria and COVID-19 co-infection, as shown in Table 1. The first case report was described by Sardar et al. [17] in Qatar as a possible *P. vivax* reactivation secondary to COVID-19, similar to the case reports presented by Kishore et al. [16] in India and Shahid et al. [11] in Qatar. Ray et al. reported a concomitant infection between malaria vivax and COVID-19 in India without suggesting the possibility of *P. vivax* re-activation [27]. However, this report is the first case of malaria vivax and COVID-19 co-infection in Indonesia that discussed the possibility of *P. vivax* relapse related to COVID-19 with symptoms of prolonged fever and exaggerated hypercoagulopathy.

Based on World Health Organization's recommendations, in terms of facing challenges caused by the COVID-19 pandemic, such as the disruption of the malaria rapid test kits supply, the shortage of health workers and personal protective equipment, as well as the limited facilities of the Intensive Care Unit (ICU), the diagnosis of malaria must always be considered in all cases of fever in malaria-endemic countries [21]. Considering the differential diagnosis of malaria infection other than SARS-CoV-2 infection can reduce the risk of morbidity and mortality due to delayed and inappropriate treatment.

Various clinical trials to determine the appropriate COVID-19 therapeutic regimen are still ongoing. The massive use of hydroxychloroquine to treat COVID-19 in Malaria endemic areas will further increase the risk of anti-malarial drug resistance [17]. In this case report, the patient was treated with Favipiravir, a nucleoside analogue antiviral efficacious in various clinical trials for treating mild-to-moderate COVID-19 [28]. In a 2021 review study, Zenchenko et al. concluded that nucleoside analogue antivirals could treat a parasitic infection, such as malaria caused by *P. falciparum*. Meanwhile, studies on the nucleoside analogue

antiparasitic effect in vivax malaria are limited [29]. Artemisinin is a very potent anti-malarial drug and can overcome the problem of resistance to quinolones. Artemisinin is also able to inhibit endocytosis more strongly than chloroquine. Artesunate, a semisynthetic derivative of artemisinin, has lately attracted much attention to be tested as a COVID-19 therapy because of its anti-viral and anti-inflammatory effects through inhibition of Nuclear Factor kappa B (NF- κ B) downregulation and protein synthesis in the early stages of viral replication [30]. Further research is required to study the role of artemisinin in treating COVID-19 and the prognosis of COVID-19 co-infection with malaria.

Conclusion

This case report shows that considering the possibility of a secondary infection diagnosis other than COVID-19 in a patient who presents with fever can prevent delayed treatment that can worsen the disease outcome. Paying more attention to a history of travel to malaria-endemic areas, a history of previous malaria infection, and exploring anamnesis regarding the fever patterns in patients are important points in making a differential diagnosis of malaria infection during the COVID-19 pandemic. Hypercoagulopathy is another phenomenon that needs further analysis in Malaria and COVID-19 co-infection. Further research is required to study the role of artemisinin in treating COVID-19 and the prognosis of COVID-19 co-infection with malaria.

Abbreviations

ACE-2: Angiotensin-converting enzyme-2; ALT: Alanine transferase; APT: Annual parasite incidence; AST: Aspartate transaminase; COVID-19: Coronavirus disease 2019; CRP: C-reactive protein; DHP: Dihydroartemisinin-piperazine; DIC: Disseminated intravascular coagulation; ICU: Intensive care unit; IL-6: Interleukin-6; IU: International unit; LDH: Lactate dehydrogenase; NETs: Neutrophil extracellular traps; NF- κ B: Nuclear factor kappa B; RDT: Rapid diagnostic test; RT-PCR: Real-time polymerase chain reaction; SARS CoV-2: Severe acute respiratory syndrome coronavirus-2; TNF: Tumour necrosis factor.

Acknowledgements

Not applicable.

Author contributions

TP and OS contributed to the conception and manuscript preparation. BB and MV contributed to the literature search. All authors have read and approved the submitted version. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

Not applicable.

Declarations**Ethics approval and consent to participate**

Not applicable.

Consent for publication

Written informed consent for publication of their clinical details and/or clinical images was obtained from the patient. A copy of the consent form is available for review by the Editor of this journal.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Tropical and Infectious Diseases Division, Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java 60115, Indonesia; ²Universitas Airlangga Hospital, Surabaya, East Java 60115, Indonesia; ³Dr. Soetomo General Teaching Hospital, Surabaya, East Java 60286, Indonesia; ⁴Faculty of Medicine, Universitas Airlangga, Surabaya, East Java 60115, Indonesia.

⁴²

Received: 14 September 2021 Accepted: 9 June 2022

Published online: 23 June 2022

References

- Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): a review. *JAMA*. 2020;324:782–93.
- Gugus Tugas Percepatan Penanganan COVID-19 : Peta Sebaran COVID-19 Indonesia 2021 [cited 2021 2 August 2021]. Available from: URL.
- Iba T, Levy JH, Levi M, Connors JM, Thachil J. Coagulopathy of coronavirus disease 2019. *Crit Care Med*. 2020;48:1358–64.
- Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382:1708–20.
- Moxon CA, Gibbins MP, McGuinness D, Milner DA Jr, Marti M. New insights into malaria pathogenesis. *Annu Rev Pathol*. 2020;15:315–43.
- WHO. World Malaria Report. 20 years of global progress and challenges. Geneva: World Health Organization; 2020. p. 2020.
- Cowman AF, Healer J, Marapana D, Marsh K. Malaria: biology and disease. *Cell*. 2016;167:610–24.
- Kemenkes. Kementerian Kesehatan RI. Buku Saku Tatalaksana Kasus Malaria. Jakarta: Kementerian Kesehatan RI; 2019. <https://www.malaria.id/en/report>.
- Kemenkes. Kementerian Kesehatan RI. Profil Kesehatan Indonesia Tahun 2019. Jakarta: Kementerian Kesehatan RI; 2020. <https://pusdatin.kemkes.go.id/resources/download/pusdatin/profil-kesehatan-indonesia/Profil-Kesehatan-indonesia-2019.pdf>.
- Dayananda KK, Achur RN, Gowda DC. Epidemiology, drug resistance, and pathophysiology of *Plasmodium vivax* malaria. *J Vector Borne Dis*. 2018;55:1–8.
- Shahid Z, Karim N, Shahid F, Yousaf Z. COVID-19 associated imported *Plasmodium vivax* malaria relapse: first reported case and literature review. *Res Rep Trop Med*. 2021;12:77–80.
- White MT, Karl S, Koepfli C, Longley RJ, Hofmann NE, Wampfler R, et al. *Plasmodium vivax* and *Plasmodium falciparum* infection dynamics: re-infections, recrudescences and relapses. *Malar J*. 2018;17:170.
- Shanks GD, White NJ. The activation of vivax malaria hypnozoites by infectious diseases. *Lancet Infect Dis*. 2013;13:900–6.
- Chu CS, Phyo AP, Lwin KM, Win HH, San T, Aung AA, et al. Comparison of the cumulative efficacy and safety of chloroquine, artesunate, and chloroquine-primaquine in *Plasmodium vivax* malaria. *Clin Infect Dis*. 2018;67:1543–9.
- Eid MM. Co-Infection with COVID-19 and malaria in a young man. *Dubai Med J*. 2021;4:1–3.
- Kishore R, Dhakad S, Arif N, Dar L, Mirdha BR, Aggarwal R, et al. COVID-19: possible cause of induction of relapse of *Plasmodium vivax* infection. *Indian J Pediatr*. 2020;87:751–2.
- Sardar S, Sharma R, Alyamani TYM, Aboukamar M. COVID-19 and *Plasmodium vivax* malaria co-infection. *IDCases*. 2020;21: e00879.
- Xu K, Chen Y, Yuan J, Yi P, Ding C, Wu W, et al. Factors associated with prolonged viral RNA shedding in patients with coronavirus disease 2019 (COVID-19). *Clin Infect Dis*. 2020;71:799–806.
- Mondi A, Lorenzini P, Castilletti C, Gagliardini R, Lalle E, Corpolongo A, et al. Risk and predictive factors of prolonged viral RNA shedding in upper respiratory specimens in a large cohort of COVID-19 patients admitted to an Italian reference hospital. *Int J Infect Dis*. 2021;105:532–9.
- Gao C, Zhu L, Jin CC, Tong YX, Xiao AT, Zhang S. Proinflammatory cytokines are associated with prolonged viral RNA shedding in COVID-19 patients. *Clin Immunol (Orlando, Fla)*. 2020;221: 108611. <https://doi.org/10.1016/j.clim.2020.108611>.
- Gao C, Zhu L, Jin CC, Tong YX, Xiao AT, Zhang S. Proinflammatory cytokines are associated with prolonged viral RNA shedding in COVID-19 patients. *Clin Immunol*. 2020;221: 108611.
- Terpos E, Ntanasis-Stathopoulos I, Elalamy I, Kastritis E, Sergentanis TN, Politou M, et al. Hematological findings and complications of COVID-19. *Am J Hematol*. 2020;95:834–47.
- Mori H, Ohkawara H, Togawa R, Rikimaru M, Shibata Y, Ikezoe T. Diagnosis and treatment of disseminated intravascular coagulation in COVID-19 patients: a scoping review. *Int J Hematol*. 2021;113:320–9.
- Lin J, Yan H, Chen H, He C, Lin C, He H, et al. COVID-19 and coagulation dysfunction in adults: a systematic review and meta-analysis. *J Med Virol*. 2021;93:934–44.
- Francischetti IM, Seydel KB, Monteiro RQ. Blood coagulation, inflammation, and malaria. *Microcirculation*. 2008;15:81–107.
- Autino B, Corbett Y, Castelli F, Taramelli D. Pathogenesis of malaria in tissues and blood. *Mediterr J Hematol Infect Dis*. 2012;4: e2012061.
- Ray M, Vazifdar A, Shivaprakash S. Co-infection with malaria and coronavirus disease-2019. *J Glob Infect Dis*. 2020;12:162–3.
- Manabe T, Kambayashi D, Akatsu H, Kudo K. Favipiravir for the treatment of patients with COVID-19: a systematic review and meta-analysis. *BMC Infect Dis*. 2021;21:489.
- Zenchenko AA, Drenichev MS, Il'icheva IA, Mikhailov SN. Antiviral and antimicrobial nucleoside derivatives: structural features and mechanisms of action. *Mol Biol*. 2021;55:786–812.
- Uzun T, Toptas O. Artesunate: could be an alternative drug to chloroquine in COVID-19 treatment? *Chin Med*. 2020;15:54.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Prolonged Fever And Exaggerated Hypercoagulopathy In Malaria Vivax Relapse And COVID-19 Co-Infection: A Case Report

ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

15%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	tel.archives-ouvertes.fr Internet Source	1%
2	aging-us.com Internet Source	1%
3	sostelemedicina.ucv.ve Internet Source	1%
4	daten-quadrat.de Internet Source	1%
5	www.dovepress.com Internet Source	1%
6	Cindy S Chu, Aung Pyae Phyo, Khin Maung Lwin, Htun Htun Win et al. "Comparison of the Cumulative Efficacy and Safety of Chloroquine, Artesunate, and Chloroquine-Primaquine in Plasmodium vivax Malaria", <i>Clinical Infectious Diseases</i> , 2018 Publication	1%

7	Internet Source	1 %
8	bmjopen.bmj.com Internet Source	1 %
9	docksci.com Internet Source	1 %
10	Parat Boonyarangka, Kittijarankon Phontham, Sabaithip Sriwichai, Kamonporn Poramathikul et al. "Co-Infection with Plasmodium vivax and COVID-19 in Thailand", Tropical Medicine and Infectious Disease, 2022 Publication	1 %
11	Dragan Ljutić, Goran Radovniković, Tina Tičinović Kurir. "Acute Renal Failure Associated with Rhabdomyolysis in Acute Q Fever", Renal Failure, 2009 Publication	<1 %
12	fdokumen.id Internet Source	<1 %
13	media.neliti.com Internet Source	<1 %
14	Khabab Abbasher Hussien Mohamed Ahmed, Mazin Salah-Eldien Hassan Haroun, Abbasher Hussien, Mohammed Eltahier Abdalla Omer. "COVID-19 and Malaria Co-infection; Same	<1 %

coin but different faces?", Research Square Platform LLC, 2021

Publication

15 Jing Lin, Han Yan, Hanchuan Chen, Chen He, Chunjin Lin, Haoming He, Sicheng Zhang, Songjing Shi, Kaiyang Lin. "COVID - 19 and coagulation dysfunction in adults: A systematic review and meta - analysis", Journal of Medical Virology, 2020

Publication

16 Saleh Al-Muhsen, Nouf S. Al-Numair, Narjes Saheb Sharif-Askari, Roaa Basamh et al. "Favipiravir Effectiveness and Safety in Hospitalized Moderate-Severe COVID-19 Patients: Observational Prospective Multicenter Investigation in Saudi Arabia", Frontiers in Medicine, 2022

Publication

17 Submitted to Padjadjaran University

Student Paper

18 jamanetwork.com

Internet Source

19 www.molvis.org

Internet Source

20 Dan Huang, Jing Qi, ChuanZheng Sun, JingChao Lei, Xi Chen, Qiong Cui. "Clinical prognostic value of elevated CK-MB in severe

multiple trauma patients with nonmyocardial contusion", Research Square Platform LLC, 2023

Publication

21

Qian Huang, Wen-Jie Xu, Xiao-Xiao Wang, Xuan Zhang, Ke-Nu Pan, Jia-Qi Zhang, Hua-Liang Chen, Wei Ruan, Li-Nong Yao. "SARS-CoV-2 and Plasmodium falciparum Co-Infection in a Returning Traveler", Frontiers in Public Health, 2022

Publication

<1 %

22

pubs.sciepub.com

Internet Source

<1 %

23

Niels Hvidt, Alex Kappel Kørup, Farr Curlin, Klaus Baumann et al. "The NERSH International Collaboration on Values, Spirituality and Religion in Medicine: Development of Questionnaire, Description of Data Pool, and Overview of Pool Publications", Religions, 2016

Publication

<1 %

24

Lu Cai, Jiankang Wang, Yongjie Li, Min Qin, Xueming Yin, Zhangjiang He, Jichuan Kang. "Global regulatory factor FsveA up-regulate the production of antitumor substance in endophytic Fusarium solani", Research Square Platform LLC, 2022

Publication

<1 %

25	iopscience.iop.org Internet Source	<1 %
26	www.tandfonline.com Internet Source	<1 %
27	"Malarial Drug Delivery Systems", Springer Science and Business Media LLC, 2023 Publication	<1 %
28	Submitted to Thomas University Student Paper	<1 %
29	journal.einstein.br Internet Source	<1 %
30	journal.fk.unpad.ac.id Internet Source	<1 %
31	Submitted to Monash University Student Paper	<1 %
32	Tomas Palecek. "Papillary Fibroelastoma Arising from the Left Ventricular Apex Associated with Nonspecific Systemic Symptoms", Echocardiography, 5/2008 Publication	<1 %
33	Zubair Shahid, Nadia Karim, Fakhar Shahid, Zohaib Yousaf. "COVID-19 Associated Imported Plasmodium vivax Malaria Relapse: First Reported Case and Literature Review", Research and Reports in Tropical Medicine, 2021	<1 %

34

estudogeral.uc.pt

Internet Source

<1 %

35

Christian Bleilevens, Josefin Soppert, Adrian Hoffmann, Thomas Breuer et al. "Macrophage Migration Inhibitory Factor (MIF) Plasma Concentration in Critically Ill COVID-19 Patients: A Prospective Observational Study", *Diagnostics*, 2021

Publication

<1 %

36

Rashmi Kishore, Shivram Dhakad, Nazneen Arif, Lalit Dar, B. R. Mirdha, Richa Aggarwal, S. K. Kabra. "COVID-19: Possible Cause of Induction of Relapse of Plasmodium vivax Infection", *The Indian Journal of Pediatrics*, 2020

Publication

<1 %

37

sajid.co.za

Internet Source

<1 %

38

weekly.chinacdc.cn

Internet Source

<1 %

39

www.degruyter.com

Internet Source

<1 %

40

Grigoris T. Gerotziafas, Mariella Catalano, Mary-Paula Colgan, Zsolt Pecsvarady et al. "Guidance for the Management of Patients with Vascular Disease or Cardiovascular Risk

<1 %

Factors and COVID-19: Position Paper from VAS-European Independent Foundation in Angiology/Vascular Medicine", Thrombosis and Haemostasis, 2020

Publication

41

Toshiaki Iba, Jerrold H. Levy, Marcel Levi, Jean Marie Connors, Jecko Thachil. "Coagulopathy of Coronavirus Disease 2019", Critical Care Medicine, 2020

Publication

<1 %

42

crossmark.crossref.org

Internet Source

<1 %

43

esmed.org

Internet Source

<1 %

44

f6publishing.blob.core.windows.net

Internet Source

<1 %

45

files.covid19treatmentguidelines.nih.gov

Internet Source

<1 %

46

journals.lww.com

Internet Source

<1 %

47

referencecitationanalysis.com

Internet Source

<1 %

48

www.cdc.gov

Internet Source

<1 %

49

Indrani Sarkar, Saurabh Singh Rathore, Gyan Dev Singh, Ram Pratap Singh. "Whole-genome

<1 %

and mitogenome based in silico analysis of select Plasmodium and identification of a novel drug molecule against the malaria parasite", Cold Spring Harbor Laboratory, 2022

Publication

50

Khabab Abbasher Hussien Mohamed Ahmed, Elfatih A. Hasabo, Mazin S. Haroun, Moh. Mah. Fadelallah Eljack et al. "Clinical characteristics, complications, and predictors of outcome of hospitalized adult Sudanese patients with COVID - 19 and malaria coinfection in Sudan: A multicenter retrospective cross - sectional study", Journal of Medical Virology, 2022

<1 %

Publication

51

Yong-Jian Geng, Zhi-Yao Wei, Hai-Yan Qian, Ji Huang, Robert Lodato, Richard J. Castriotta. "Pathophysiological Characteristics and Therapeutic Approaches for Pulmonary Injury and Cardiovascular Complications of Coronavirus Disease 2019", Cardiovascular Pathology, 2020

<1 %

Publication

52

Fatemeh Nemati Zargaran, Mosayeb Rostamian, Sara Kooti, Hamid Madanchi, Keyghobad Ghadiri. "Co-infection of COVID-19

<1 %

and parasitic diseases: A systematic review",
Parasite Epidemiology and Control, 2023

Publication

53

George D Liatsos. "SARS-CoV-2 induced liver injury: Incidence, risk factors, impact on COVID-19 severity and prognosis in different population groups", World Journal of Gastroenterology, 2023

Publication

<1 %

54

Junjie Zou, Jian Yin, Lei Fang, Mingjun Yang, Tianyuan Wang, Weikun Wu, Michael A. Bellucci, Peiyu Zhang. "Computational Prediction of Mutational Effects on SARS-CoV-2 Binding by Relative Free Energy Calculations", Journal of Chemical Information and Modeling, 2020

Publication

<1 %

55

Kieran A. Walsh, Karen Jordan, Barbara Clyne, Daniela Rohde et al. "SARS-CoV-2 detection, viral load and infectivity over the course of an infection", Journal of Infection, 2020

Publication

<1 %

56

Sreekanth Kokkonda, Farah El Mazouni, Karen L. White, John White et al. " Isoxazolopyrimidine-Based Inhibitors of Dihydroorotate Dehydrogenase with Antimalarial Activity ", ACS Omega, 2018

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Prolonged Fever And Exaggerated Hypercoagulopathy In Malaria Vivax Relapse And COVID-19 Co-Infection: A Case Report

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
