


Editor-in-Chief:
Dr. Syed A. A. Rizvi

Journal of Medicinal and Chemical Sciences



www.jmchemsci.com



Editorial Board



Editor-in-Chief

Professor Dr. Syed A. A. Rizvi Department of Pharmaceutical Sciences, College of Pharmacy and Health Sciences, Mercer University, Atlanta, GA, USA.

Professor of Pharmaceutical Sciences

- pharmaps.nova.edu/profile.cfm?BioID=srizvi
- srizvinova.edu
- +1 954-262-8311
- [0000-0002-2385-5672](https://orcid.org/0000-0002-2385-5672)
- h-index: 30
- [More](#)




Editor-in-Chief

Professor Dr. Rassoul Dinarvand Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran PO Box 14155-6451, IRAN

Professor of Pharmaceutics, drug delivery, nanomedicine

- isid.research.ac.ir/Rassoul_Dinarvand
- dinarvandtums.ac.ir
- +98 21 66959095
- [0000-0003-0694-7556](https://orcid.org/0000-0003-0694-7556)
- h-index: 66
- [More](#)



Co-Editor-in-Chief

Professor Dr. Ali Nokhodchi Pharmaceutics Research Laboratory, School of Life Sciences, University of Sussex, Brighton BN1 9QJ, UK

Professor of Pharmaceutics and Drug Delivery

- www.sussex.ac.uk/lifesci/nokhodchilab/index
- a.nokhodchisussex.ac.uk
- +44 1273872811
- [0000-0002-3244-2482](https://orcid.org/0000-0002-3244-2482)
- h-index: 65
- [More](#)



Editorial Board

Professor Dr. Khosro Khajeh Dept. of Biochemistry, Faculty of Biological Sciences, Tarbiat Modares University, P.O.Box: 14115, Tehran, Iran

Professor of Biochemistry

- ibj.pasteur.ac.ir/files/site1/files/CV_Khajeh.pdf
- khajehmodares.ac.ir
- (+98-21) 82884718
- [0000-0002-5916-0338](https://orcid.org/0000-0002-5916-0338)
- h-index: 42
- [More](#)



Associate Editor

Dr. Zeinab Arzhegar Department of Chemistry, Payame Noor University, PO BOX 19395-4697 Tehran, Iran.

Assistant Professor in Organic chemistry

- arzhgaryahoo.com
- +98 84 32226101
- 0000-0903-3774-4348
- h-index: 13



Director-in-Charge

Dr. Sami Sajjadifar Department of Chemistry, Payame Noor University, PO BOX 19395-4697 Tehran, Iran.

Assistant Professor in Organic Chemistry

- chemsajjadifar.blogspot.com
- ss.sajjadifargmail.com
- +98 84 32226101
- 0000-0001-8661-1264
- h-index: 24



Senior Editor

Professor Dr. Mohammad Mansoob Khan Chemical Sciences, Faculty of Science, Universiti Brunei Darussalam, Jalan Tungku Link, Gadong, BE 1410, Brunei Darussalam, Tanzania.

Professor in Inorganic Chemistry

- expert.ubd.edu.bn/mansoob.khan.php
- mansoob.khanubd.edu.bn
- +673 246 0922 / 246 0923
- 0000-0002-8633-7493
- h-index: 55
- [More](#)



Senior Editor

Professor Dr. Ali Delpisheh Department of Child and Reproductive Health, Liverpool School of Tropical Medicine, Liverpool, UK.

Professor of Clinical Epidemiology

- www.feedage.com/feeds/2454625/most-cited-full-text-articles
- alidelpishe@yahoo.com
- +98-841-3334060
- h-index: 41
- [More](#)



Editorial Board

Dr. Ahmad Reza Moosavi-Zare Hamedan University of Technology, Hamedan, 65155, Iran.

Associate Professor of Organic Chemistry

- che.sjau.ac.ir/%D8%A%D8%A9%D8%AA%D8%B1-%D8%A7%D8%AD%D9%85%D8%AF...
- moosavizare@yahoo.com
- 08133117804
- [0000-0003-0321-9326](https://orcid.org/0000-0003-0321-9326)
- h-index: 46
- [More](#)



Editorial Board

Dr. Ali Maleki Department of Chemistry, Iran University of Science and Technology (IUST), Tehran 16846-13114, IRAN

Associate Professor of Organic Chemistry

- www.iust.ac.ir/find.php?item=20_10920_19490_fa
- malekiust.ac.ir
- 0098 (21) 77240540
- [0000-0001-5490-3350](https://orcid.org/0000-0001-5490-3350)
- h-index: 69



Editorial Board

Dr. Majid Darroudi Department of Modern Sciences and Technologies & Nuclear Medicine Research Center, School of Medicine, Mashhad University of Medical Sciences (MUMS), Mashhad, Iran.

Assistant Professor of Nanomedicine

- isid.research.ac.ir/Majid_Darroudi
- majiddarroudigmail.com
- 05138002286
- [0000-0002-2624-7242](https://orcid.org/0000-0002-2624-7242)
- h-index: 51
- [More](#)



International Editorial Board

Professor Dr. Nenad L. Ignjatovic Centre for Fine Particles Processing and Nanotechnologies, Institute of Technical Sciences of the Serbian Academy of Sciences and Arts, Knez Mihailova 35/4, 11000 Belgrade, Serbia.

Professor of Biomaterials

- www.ftn.sanu.ac.rs/nenadignjatovicen.htm
- nenad.ignjatovicitn.sanu.ac.rs
- +381 63 8115494
- [0000-0002-5749-094X](https://orcid.org/0000-0002-5749-094X)
- h-index: 35
- [More](#)



Editorial Board

Professor Dr. Behrooz Maleki Department of Organic Chemistry, University of Mazandaran, Babolsar, Iran

Professor of Organic Chemistry

- b.malekiumz.ac.ir
- h-index: 46
- [More](#)



Editorial Board

Professor Dr. Mehrdad Hamidi Zanzan University of Medical Sciences (ZUMS)disabled, Zanzan, Iran

Professor of Pharmaceutics

- isid_research.ac.ir/Mehrdad_Hamidi
- hamidimzums.ac.ir
- [0000-0001-7977-5252](tel:+984179775252)
- h-index: 40
- [More](#)



Editorial Board

Professor Dr. Ghodsi Mohammadi Ziarani Department of Chemistry, Alzahra University, Tehran, Iran.

Professor of Organic Chemistry

- gmohammadialzahra.ac.ir
- [0000-0001-5177-7889](tel:+9821000151777889)
- h-index: 45
- [More](#)



International Editorial Board

Dr. Yasser Fakri Mustafa Hussein Department of Pharmaceutical Chemistry, College of Pharmacy, University of Mosul, Mosul, Iraq.

Professor in Medicinal Chemistry

- www.researchgate.net/profile/Yasser_Mustafa3
- dr.yassermustafauomosul.edu.iq
- +9647701615864
- [0000-0002-0926-7428](tel:+9649267428)
- h-index: 31
- [More](#)



Editorial Board

Dr. Gholamabbas Chehardoli Department of Medicinal Chemistry, School of Pharmacy, Hamadan University of Medical Science, Hamadan, Iran

Associate Professor of Organic Chemistry

- cheh1002@gmail.com
- +98 811 8381594
- [0000-0002-8760-3837](https://orcid.org/0000-0002-8760-3837)
- h-index: 21



International Editorial Board

Dr. Ali H. Jawad Al-Taie Faculty of Applied Sciences, University of Technology MARA (UiTM) Shah Alam, Selangor, Malaysia

Associate Professor in Biochemical and Environment

fsg.uitm.edu.my/v1/research/194-center-of-coals-a-biomass-energy-research.html

- ali288salam.uitm.edu.my
- (+603)55211721
- [0000-0002-4827-9093](https://orcid.org/0000-0002-4827-9093)
- h-index: 53
- [More](#)



Editorial Board

Professor Dr. Asghar Mesbahi Medical Physics Department, Medical school, Tabriz University of Medical Sciences, Tabriz, Iran

Professor at School of Medicine

isid.research.ac.ir/Asghar_Mesbahi

- amesbahi2010@gmail.com
- [0000-0001-9159-2168](https://orcid.org/0000-0001-9159-2168)
- h-index: 26
- [More](#)




Editorial Board

Dr. Masoud Mohammadi Department of Chemistry, Faculty of Science, Ilam University, P.O. Box, 69315516, Ilam, Iran.

Assistant Professor of Organic Chemistry

chemistrypmu2.blogfa.com/

- tbr.masoudgmail.com
- [0000-0002-1043-3470](https://orcid.org/0000-0002-1043-3470)
- h-index: 23
- [More](#)



• [More](#)


International Editorial Board

Dr. Azahar Ali Affiliation: Electrical and computer Engineering, Coover Hall, Iowa State University, Ames, IA-50011, USA.

Associate Professor of Nanoscience and Nanotechnology

www.info.iastate.edu/individuals/info/259711/Ali-Azahar

- azahariastate.edu
- +1-515-598-6440
- 0000-0001-5752-8808
- h-index: 37



Editorial Board

Professor Dr. Ali Reza Modarresi-Alam Department of Chemistry, Faculty of Science, University of Sistan and Baluchestan, Zahedan, Iran, Postcode: 9816745785.

Professor in Organic Chemistry

- modaresichem.usb.ac.ir
- +98-54-33431146
- 0000-0003-4055-4633
- h-index: 20
- [More](#)



International Editorial Board

Professor Dr. Yogesh Chandra Tripathi Chemistry and Bioprospecting Division, Forest Research Institute, P.O. New Forest, Dehradun-248006, India. RG=82.77

Professor of Chemistry and Medicinal Chemistry

www.researchgate.net/profile/YOGESH_TRIPATHI/

- tripathicycfre.org
- +91-135-224207
- 0000-0003-1367-5122
- h-index: 19
- [More](#)



Editorial Board

Dr. Majid Hajifaraji Dept. of Nutrition and Food Policy & Planning Research, National Nutrition & Food Technology, Research Institute (NNFTRI), Shahid Beheshti University of Medical Sciences (SBUMS), Iran.

Associate Professor in Nutrition

www.linkedin.com/in/majid-hajifaraji-06bb5138/

- m.hajifarajinnftri.ac.ir
- +98(21) 22357486
- [0000-0002-4353-7866](https://orcid.org/0000-0002-4353-7866)
- h-index: 21
- [More](#)



International Editorial Board

Professor Dr. Ali Sabea Hammood Head of Biomedical Materials Engineering Track-Faculty of Engineering-University of Kufa-Iraq.

Professor in Biomedical Materials Engineering

www.researchgate.net/profile/Ali_Hammood

- alis.altameemiukufa.edu.iq
- 00964(0)7801035379
- [0000-0002-0047-2900](https://orcid.org/0000-0002-0047-2900)
- h-index: 10
- [More](#)



International Editorial Board

Professor Dr. Roberto Acevedo University of Virginia, Santiago Province, Chile.

Professor Dr. in Chemistry

www.roberto-acevedo.cl/

- roberto.acevedo.llanos@gmail.com
- +569 4209-5982
- h-index: 15



International Editorial Board

Professor Dr. Ehab AlShamaileh Department of Chemistry, School of Science, The University of Jordan, Amman 11942, Jordan

Professor of Physical Chemistry

- ehabju.edu.jo
- +9626 5355000 ext 22133
- h-index: 15



Language Editor

Dr. Behroz Jamalvandi Ilam Farhangyan University, Ilam, Iran.

University lecturer in applied linguistics

- behrouzjamlvaandigmail.com
- 0000-0003-4162-2052
- h-index: 5



Language Editor

Dr. Fatemeh Ramezani Ph.D. Student, EFL Teaching, Tehran university, Tehran, Iran.

- ramezani.tv@yahoo.com




Language Editor

Dr. Nadereh Shirvani Ph.D. Student at Ilam University, Ilam, Iran.

- n.shirvani865gmail.com

Number of Articles: 30



Intensity Modulated Radiation and Volumetric Modulated Arc Therapies in Breast Cancer


May Z. Saddik; Fatiheea F. Hassan

Volume 6, Issue 9, September 2023, Pages 1925-1934

<https://doi.org/10.26655/JMCHMSCI.2023.9.1>

Abstract Background: Breast cancer is the most common malignant neoplasm in women, accounting for 25% of all cancers. Radiotherapy is one of the safest methods compared with other treatments. Methods: ... [Read More](#)

[View Article](#) [PDF 974.79 K](#)




Clinical and Laboratory Characteristics of COVID-19 Patients with Fecal SARS-CoV-2 Detection at a Tertiary Hospital in Indonesia

isoforms of prostaglandin G-H synthetase (I ... [Read More](#)

[View Article](#) [PDF 1.06 M](#)

Review Article



Perception of Patient Safety Incident Reporting System among Healthcare Workers in Palestinian Government Hospital

Salam Al Ratrout

Volume 6, Issue 9, September 2023, Pages 1953-1969

<https://doi.org/10.26655/JMCHMSCI.2023.9.4>

Abstract The aim of the current study was to determine the system perception for reporting patient safety incidents among workers in the field of health care, evaluating the factors affecting ... [Read More](#)

[View Article](#) [PDF 1.01 M](#)

Original Article


Andrew Adhytia Lieputra; Rebrarina Hapsari; Cecilia Oktaria Permatadewi; Irfan Kesumayadi; Iva Puspitasari; Purnomo Hadi; Nur Farhanah; Hendro Wahyono; Hery Djagat Purnomo

Volume 6, Issue 9, September 2023, Pages 1935-1942

<https://doi.org/10.26655/JMCHMSCI.2023.9.2>

Abstract Transmission prevention is important to prevent the spread of COVID-19. Although most cases are transmitted through droplets and aerosols, several studies have shown the possibility ... [Read More](#)

[View Article](#) [PDF 852.69 K](#)




Density Functional Theory, ADME, and Molecular Docking of Some Anthranilic Acid Derivatives as Cyclooxygenase Inhibitors

Karam S. Atrushi; Dana M. Ameen; Shaymaa H. Abdulrahman; Faris T. Abachi

Volume 6, Issue 9, September 2023, Pages 1943-1952

<https://doi.org/10.26655/JMCHMSCI.2023.9.3>

Abstract Nonsteroidal anti-inflammatories (NSAIDs), are very effective agents in relieving mild to moderate pain and inflammation by inhibiting two



Communicable Diseases (CDs) Among Paramedics in Private Healthcare in India


Soumya Ranjan Ray Choudhury; B. Chandra Mohan Patnaik; Ipseeta Satpathy

Volume 6, Issue 9, September 2023, Pages 1967-1974

<https://doi.org/10.26655/JMCHMSCI.2023.9.5>

Abstract Humanitarian emergencies including disasters associated with natural hazards, conflict, complex emergencies, and famines can pose significant risks to public health, especially when ... [Read More](#)

[View Article](#) [PDF 844.49 K](#)



Synthesis of Benzimidazole and Phthaloylamino Acid derivatives and Antibacterial Activity


Hussniya A. AlDifar; Basma S. Baaiu; Khalid M. A. Darwish; Mohamad F. Ali; Othman O. Dakhil; Mansour Abd-Alsalam; Haneen Al Difar

Volume 6, Issue 9, September 2023, Pages 1975-1984

<https://doi.org/10.26655/JMCIEMSCI.2023.9.6>

Abstract Seven novel benzimidazole and phthaloylamino acid derivatives were synthesized. In the presence of acetic acid, the phthalic anhydride reacted with benzidine in a cyclization reaction ... [Read More](#)

[View Article](#) [PDF 1.01 M](#)



Competitive Adsorption of Li+, Na+, and K+ Ions on Phillip Site/Chabazite Zeolitic Tuff from Jordan

Reem Naser Dabaibeh; Hiba Hamed Amayrah

Volume 6, Issue 9, September 2023, Pages 1985-1997

<https://doi.org/10.26655/JMCIEMSCI.2023.9.7>

Abstract The adsorption of alkali metal ions on zeolite has no such possibility of precipitation. In the current work, we have studied the adsorption isotherms, and the effect of pH and temperature ... [Read More](#)

[View Article](#) [PDF 1.23 M](#)

Original Article

Disorder


Ameer Ali Althabhaawee; Taghreed Fadhil Zaidan Zaidan

Volume 6, Issue 9, September 2023, Pages 2010-2017

<https://doi.org/10.26655/JMCIEMSCI.2023.9.9>

Abstract There has been a general uptick in the number of people diagnosed with major depressive disorder (MDD) worldwide, which has led to a commensurate rise in the need for oral medicine ... [Read More](#)

[View Article](#) [PDF 743.15 K](#)



Predictors of Mortality in Sepsis Patients Resulting from Severe and Critical COVID-19


Nur Farhanah; Supriadi Supriadi; Hendro Wahjono; Suharyo Hadisaputro; Muhammad Hussein Gasem

Volume 6, Issue 9, September 2023, Pages 2018-2027

<https://doi.org/10.26655/JMCIEMSCI.2023.9.10>

Abstract Patients with severe and critical COVID-19 may exhibit sepsis and mortality resulting from multi-organ failure. Neutrophil-lymphocyte-ratio (NLR) values, C-reactive protein (CRP) levels, ... [Read More](#)

[View Article](#) [PDF 488.04 K](#)



Critical Role of Monocyte Chemotactic Protein-1 and Macrophage Migration Inhibitory Factor in Children who Have Pneumonia


Gehad Nabil Abd El-Aal; Shimaa Moustafa Kamel; Mona Sayed Mohammad Attaya; Dalia Mahmoud Eldewi; Fatma Elzahraa AE. Diab; Marwa Mohsen Ramadan Hassan; Suzan Mohamed Abdel Kareem Mohamed

Volume 6, Issue 9, September 2023, Pages 1998-2009

<https://doi.org/10.26655/JMCIEMSCI.2023.9.8>

Abstract Background: The leading cause of death for children under the age of five years old worldwide is pneumonia. The assessment of illness progression and clinical stage is necessary for ... [Read More](#)

[View Article](#) [PDF 1.05 M](#)



Assessment of Salivary Alpha-Amylase, Cortisol, and Prevalence of Temporomandibular Disorder in Patients with Major Depressive Disorder



Impact of Pharmacological and Non-Pharmacological Intervention among Pregnant Women Complaining of Nausea and Vomiting


Shaymaa S. Maqsood; Ibrahim H. Mustafa

Volume 6, Issue 9, September 2023, Pages 2028-2037

<https://doi.org/10.26655/JMCIEMSCI.2023.9.11>

Abstract This study was conducted with the aim of investigating the effect of pharmaceutical and non-pharmacological interventions in pregnant women complaining of nausea and vomiting because ... [Read More](#)

[View Article](#) [PDF 862.02 K](#)




Survival Analysis of Delta versus Omicron Variants in COVID-19 Intensive Care Unit (ICU)

Somayeh Mehrpour; Mojdeh Bahadorzadeh; Hassan Safdari; Mansour Shakeri; Reihane Tabarai

<https://doi.org/10.26655/JMCHMSCI.2023.9.12>

Abstract Objective: COVID-19 has presented numerous epidemiological and clinical pictures from its beginning and much effort has been paid to detect the behavior of disease and its new types. ... [Read More](#)

[View Article](#) [PDF 794.21 K](#)



Ghrelin and Serotonin Expression on Restricted Feed with Curcuma xanthorrhiza Roxb. and Iron Supplementation

Amallia Nuggetsiana Setyawati; Diky Syahrul Ramadhan; Kusmiyati Tjahjono; Ryan Halleyantoro; Puspita Kusuma Dewi

Volume 6, Issue 9, September 2023, Pages 2046-2055

<https://doi.org/10.26655/JMCHMSCI.2023.9.13>

Abstract Background: Nutritional shortages and acute malnutrition are caused by insufficient protein and calorie consumption. With severe short- and long-term morbidity and death, malnutrition ... [Read More](#)

[View Article](#) [PDF 797.25 K](#)

Original Article

Abstract Background: Malaria resistance to artemisinin combination therapy has prompted researchers to explore anti-plasmodium materials. One of the marine biotas thought to have the potential ... [Read More](#)

[View Article](#) [PDF 1.02 M](#)



Effectiveness Comparison of Decontamination Sputum Specimen between Kudoh and Petroff Modification Technique in Mycobacterium Culture

Andritta Febriana; Ni Made Mertaniasih; Pepy Dwi Endraswari; Lindawati Alimsardjono; Eko Budi Koendhori

Volume 6, Issue 9, September 2023, Pages 2096-2103


<https://doi.org/10.26655/JMCHMSCI.2023.9.16>

Abstract Objectives: Tuberculosis (TB) is the common cause of death due to a single infectious agent. MTB culture is still the gold standard for TB diagnosis. For optimal care of pulmonary TB, ... [Read More](#)

[View Article](#) [PDF 853.77 K](#)



Modified Lung Ultrasound Score to Determine the Degree of Perioperative Atelectasis in Adults Undergoing Elective



Rigidity and Flexibility of Pyrazole, s-Triazole, and v-Triazole Derivative of Chloroquine as Potential Therapeutic against COVID-19


Adel M. Najar; Asma Eswayah; Moayed Ben Moftah; Ruwida Omar M.K.; Eman Bobtaina; Mohamed Najwa; Tawfeg A. Elhisadi; Aeyad Tahani; Salha M.Tawati; Aliaa M. M.Khalifa; Aly Abdou; Ahmed Emhamed DowAltome

Volume 6, Issue 9, September 2023, Pages 2056-2084

<https://doi.org/10.26655/JMCHMSCI.2023.9.14>

Abstract Based on the core unit of chloroquine, new types of N-heterocyclic compounds that are fused together have been made. The compounds were put into two groups. In series A, the five-member ... [Read More](#)

[View Article](#) [PDF 2.85 M](#)



Comparing Black Trepan and Curryfish Extract's Antimalarial Activity Using In Vitro Screening

Prawesty Diah Utami; M. Fathi Ilmawan; Herin Setianingsih

Volume 6, Issue 9, September 2023, Pages 2085-2095

<https://doi.org/10.26655/JMCHMSCI.2023.9.15>

Laparoscopic Surgery Under General Anaesthesia- A Prospective Observational Study


Dhivya R; Arun Prasath D; Yachendra V.S.G; Surya R; Lakshmi Ramakrishnan

Volume 6, Issue 9, September 2023, Pages 2104-2110

<https://doi.org/10.26655/JMCHMSCI.2023.9.17>

Abstract Background: Lung ultrasound score (LUS) is a semiquantitative assessment tool that has been proposed to assess lung aeration changes in mechanically ventilated patients after therapeutic ... [Read More](#)

[View Article](#) [PDF 851.34 K](#)



Correlation Between the Bacterial and Fungal Profiles from the Clinical Specimens with the CD4 Counts and the NLR Values of HIV/AIDS Patients at Tertiary Referral Hospital in Indonesia


Nafdzu Makhmudatul Muna; Agung Dwi Wahyu Widodo; Pepy Dwi Endraswari; M. Vitanata Arfijanto

Volume 6, Issue 9, September 2023, Pages 2111-2127

<https://doi.org/10.26655/JMCHMSCI.2023.9.18>

Abstract Human immunodeficiency virus (HIV) poses a significant problem for the global public health with high morbidity and mortality rates. HIV/AIDS cases in Indonesia, the country with the ... [Read More](#)

[View Article](#) [PDF 968.99 K](#)



Synthesis, Antibacterial Evaluation, and Docking Studies of Some Azo Compounds and Schiff Bases Derived from Sulfonamide

Munther Abduljaleel Muhammad-Ali; Ekhlas Qanber Jasim; Abdullah H. Al-Saadon

Volume 6, Issue 9, September 2023, Pages 2128-2139

<https://doi.org/10.26655/JMCEMMSCI.2023.9.19>


Abstract Objective: The aim of study was to synthesize new products of sulfonamide compounds containing azo and Schiff base fragments and confirm the structures by 1H-NMR and FT-IR spectroscopy, ... [Read More](#)

[View Article](#) [PDF 1.51 M](#)

Original Article



Nigella sativa + black pepper with Letrozole + Tamoxifen as treatments for infertile women with PCOS



Removal of Nickel(II) by Silica Aerogel-Activated Carbon Nanocomposite from Wastewater

Lubna Salah Muhammed

Volume 6, Issue 9, September 2023, Pages 2140-2153

<https://doi.org/10.26655/JMCEMMSCI.2023.9.20>

Abstract HMs (HMs) are important pollutants in the environment. The main purpose of this study was to evaluate the efficiency of nickel(II) adsorption process by Silica Aerogel-Activated Carbon ... [Read More](#)

[View Article](#) [PDF 1.38 M](#)

Original Article

Comparative Effect of Nigella sativa + black pepper with Letrozole + Tamoxifen as treatments for infertile women with PCOS

The pregnancy rate was significantly higher in the intervention group (48.9%) compared to the control group (11.1%). The intervention group also showed significant differences in endometrial thickness, dominant follicle size, and number of follicles on day 12 of the menstrual cycle

Comparative Effect of Nigella Sativa+ Black Pepper and Letrozole + Tamoxifen on Female Infertility in Women with Polycystic Ovarian Syndrome: A Randomized Clinical Trial

Athar Rasekh Jahromi; Asra Ansari; Zahra Zareibabaraabi; Vahid Rahmanian; Nader Sharifi

Volume 6, Issue 9, September 2023, Pages 2154-2163

<https://doi.org/10.26655/JMCEMMSCI.2023.9.21>

Abstract Background: Polycystic ovary syndrome (PCOS), one of the most common reasons for infertility in terms of anovulation, influences 4–7% of reproductive women. Objectives: This study ... [Read More](#)

[View Article](#) [PDF 1020.71 K](#)

Prevalence of Respiratory Disorders among Petrol Pump Workers: A Cross-Sectional Descriptive Study


Mustafa Mohammad Ameen; Salih Ahmed Abdulla

Volume 6, Issue 9, September 2023, Pages 2164-2176

<https://doi.org/10.26655/JMCEMMSCI.2023.9.22>

Abstract Objectives: Gas station workers who are continuously exposed to gasoline/diesel smoke, are at high risk of causing respiratory problems. This study was conducted to determine the prevalence ... [Read More](#)

[View Article](#) [PDF 1.12 M](#)



The Role of Dynamic (99m) Tc Pertechnetate Scintigraphy in Early Detection and Differentiation of Hyperthyroidism


Ramë Miftari; Hana Maloku; Afërdita Bajqinca; Bujar Rexhaj; Sanije Miftari; Ylli Kaqiu

Volume 6, Issue 9, September 2023, Pages 2177-2185

<https://doi.org/10.26655/JMCEMMSCI.2023.9.23>

Abstract Thyrotoxicosis is hypermetabolic states that is not always associated with hyperfunction of thyroid gland and can present a problem in diagnostic differentiation. We have tried to prove ... [Read More](#)

[View Article](#) [PDF 893.03 K](#)



Survey on Enhancement of Nuclear Cardiology Images Based on Image Processing Techniques for Diagnosis of Ischemic Patients

Loai Abudaqa; Nabil Al-Swari; Shady Hegazi; Osama Alyasseen; Maro Gharbi; Farah Alaila; Hadeel Lozon; Arif Al Nooryani


Volume 6, Issue 9, September 2023, Pages 2186-2197

<https://doi.org/10.26655/JMCEMMSCI.2023.9.24>

Abstract A blockage of the blood vessels feeding the area causes ischemia, which is defined as a localized absence of blood flow. If an organ is not getting enough oxygen and blood flow, such ... [Read More](#)

[View Article](#) [PDF 976.48 K](#)

Original Article



Upregulation of INSR Gene Expression and Restoration of Melatonin Levels by Tinospora Cordifolia Stem Extract and Alkaloid Rich Fraction against Continuous Light Induced Polycystic Ovary Syndrome in Rodents


Murali Krishna Moka; Sumithra M

Volume 6, Issue 9, September 2023, Pages 2198-2217

<https://doi.org/10.26655/JMCEMMSCI.20239.25>

Abstract Females of childbearing age often suffer from polycystic ovary syndrome (PCOS), a hormonal condition. Symptoms including insulin resistance, high androgen levels, irregular menstrual ... [Read More](#)

[View Article](#) [PDF 1.87 M](#)



Tofacitinib in Pharmaceutical Solid Dosage from Dissolution Study: Development and Validation of RP-HPLC Method

Krishna Prasad Narapereddy; Devi Sravanthi Alladi

Volume 6, Issue 9, September 2023, Pages 2218-2227


Alireza Ghanbari; Majid Nouri; Mohammad Darvishi

Volume 6, Issue 9, September 2023, Pages 2234-2241

<https://doi.org/10.26655/JMCEMMSCI.2023.9.28>

Abstract Introduction: One of the important risk factors for diabetic patients, including foot ulcers, is lack of blood sugar control, the standard criterion for which in the long run is the ... [Read More](#)

[View Article](#) [PDF 859.73 K](#)



Effects of Dietary Supplements and Anabolic Hormones on Hematological and Biochemical Parameters among Young Bodybuilders in Kurdistan Region of Iraq

Saman Sulaiman; Safin Hussein; Karzan Qurbani; Kale Husen; Shara Rasul; Avesta Mustafa

Volume 6, Issue 9, September 2023, Pages 2242-2248

<https://doi.org/10.26655/JMCEMMSCI.2023.9.29>


Abstract Athletes and bodybuilders frequently utilize vitamins and anabolic hormones. Utilizing these substances, gymgoers can achieve their goals faster by consuming a combination of supplements ... [Read More](#)

[View Article](#) [PDF 758.46 K](#)

<https://doi.org/10.26655/JMCEMMSCI.2023.9.26>

Abstract A simple, sensitive, precise, and accurate reversed phase high performance liquid chromatographic (RP-HPLC) method has been developed for the dissolution study of Tofacitinib in Tofacitinib ... [Read More](#)

[View Article](#) [PDF 826.64 K](#)



Microcapsules of Protein-Polysaccharide Complexes Produced on a Variety of Matrices


Ali L Alfalluji; Fadhel Omran Essa

Volume 6, Issue 9, September 2023, Pages 2228-2233


<https://doi.org/10.26655/JMCEMMSCI.2023.9.27>

Abstract In this work, the chance of the microcapsules development by layer-by-layer adsorption of biopolyelectrolytes was investigated on the cores of inorganic (particles CaCO₃) and polymer ... [Read More](#)

[View Article](#) [PDF 863.62 K](#)



Evaluation of Relationship between Serum Hemoglobin A1C Level and Severity of Diabetic Foot Ulcers Based on Wagner Criteria



Type-2 Diabetes Among Investors Community in India

Srutiva Mishra; B. Chandra Mohan Mohan Patnaik; Ipseeta Satpathy

Volume 6, Issue 9, September 2023, Pages 2249-2258

<https://doi.org/10.26655/JMCEMMSCI.2023.9.30>

Abstract Comorbidity and mortality are high among people with type-2 diabetes. Globally, diabetes is a major health issue, with a growing number of patients at risk of complications. A change ... [Read More](#)



Original Article

Correlation between the Bacterial and Fungal Profiles from the Clinical Specimens with the CD4 Counts and the NLR Values of HIV/AIDS Patients at Tertiary Referral Hospital in Indonesia

Nafdzu Makhmudatul Muna¹ , Agung Dwi Wahyu Widodo^{2,3*} , Pepy Dwi Endraswari^{2,3,4} , M. Vitanata Arfijanto^{5,6}

¹Study Program of Clinical Microbiology Specialist, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

²Department of Medical Microbiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

³Department of Clinical Microbiology, Dr. Soetomo Academic Hospital, Surabaya, Indonesia

⁴Unit of Clinical Microbiology, Airlangga University Hospital, Surabaya, Indonesia

⁵Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

⁶Department of Internal Medicine, Dr. Soetomo Academic Hospital, Surabaya, Indonesia

ARTICLE INFO

Article history

Receive: 2023-01-07

Received in revised: 2023-03-25

Accepted: 2023-04-03

Manuscript ID: JMCS-2302-1951

Checked for Plagiarism: Yes

Language Editor:

Dr. Fatimah Ramezani

Editor who approved publication:

Dr. Gholamabbas Chehardoli

DOI:10.26655/JMCHMSCI.2023.9.18

KEYWORDS

HIV

Bacterial profiles

Fungal profiles

CD4


NLR

Opportunistic infections

ABSTRACT

Human immunodeficiency virus (HIV) poses a significant problem for the global public health with high morbidity and mortality rates. HIV/AIDS cases in Indonesia, the country with the fifth highest risk of HIV/AIDS in Asia, continue to increase. A decrease in CD4 levels indicates an increase in the opportunistic infection rates of HIV patients. In addition, Neutrophil to Lymphocyte Ratio (NLR) can serve as the basis for bacterial infection, infection rate, and antibiotic therapy. This observational analytic study applies a retrospective cross-sectional design and aims to observe the correlation between bacterial and fungal profiles of clinical specimens and the CD4 cell counts and the NLR values of HIV/AIDS inpatients at Dr. Soetomo Tertiary Referral Hospital from September 2021 to August 2022. A total of 192 bacterial and fungal identification test results from 76 patients were collected, consisting of 83.3% bacterial culture tests, consisting of 52.5% negative cultures, 12% gene expert TB tests, and 4.7% fungal culture tests. *Pseudomonas aeruginosa* and *Escherichia coli* ESBL were the highest numbers of Gram-negative bacteria found in this study, amounting to 51.33%, followed by *Streptococcus viridans*, the highest number of Gram-positive bacteria found in this study, amounting to 48.67%. A total of 34.2% commensal bacteria and 21.05% Multi-Drug Resistant Organism (MDRO) bacteria were also found in this study. The gene expert TB tests resulted in 73.9% undetected *M. tuberculosis*, while the fungal culture tests resulted in 88.9% *Candida albicans*. Furthermore, there was no correlation between the bacterial and fungal profiles from clinical specimens with CD4 counts or NLR values in HIV/AIDS-infected patients.

GRAPHICAL ABSTRACT



* Corresponding author: Agung Dwi Wahyu Widodo

✉ E-mail: agungimunologi@gmail.com

© 2023 by SPC (Sami Publishing Company)

Introduction

Human immunodeficiency virus (HIV) poses a significant problem for global public health [1] and is one of the most life-threatening infectious diseases [2] with high morbidity and mortality rates [3]. Data shows that HIV/AIDS cases in Indonesia, the country with the fifth highest risk of HIV and AIDS in Asia, continue to increase annually, although it tends to fluctuate [4, 5]. In addition, this disease has a high mortality rate in Indonesia [3].

Although HIV is the initial cause of AIDS, most deaths occur in HIV-infected patients with complications of Opportunistic Infections (OI). OI quickly attacks patients with low cellular and humoral immune defense conditions [6]. The World Health Organization (WHO) states that patients with stage III and IV HIV have a 9.4 and 22.6 times higher risk of developing OI than those with stage I of HIV. HIV-infected patients with a T-CD4 cell count of fewer than 200 cells per mm³ were found to have a 4.9 times higher risk of developing OI than HIV-infected patients with a CD4-T cell count of more than 350 cells per mm³ [6].

Opportunistic infections arise due to decreased immunity and can attack several organs, such as the respiratory tract, digestive tract, neurological organs, skin, and so forth [7, 8]. Such infections are a significant cause of hospitalization, morbidity, and mortality (accounting for 94.1% of all deaths) in patients with HIV/AIDS [9], especially those in low- and middle-income countries [10-12].

Severe bacterial infections other than tuberculosis should be recognized as HIV-associated opportunistic diseases [13] and the most common complications [1], as well as a primary cause of morbidity and death in patients with HIV/AIDS [14]. Meanwhile, fungi are the main contributors to OI that affect patients with HIV/AIDS and contribute significantly to OI in patients with late-stage HIV infection [15].

CD4 cell count is the most appropriate choice of the early indicators of the severity of HIV infection. A decrease in the CD4 cell count indicates an increase in the OI rate in HIV-infected patients [16, 17]. The neutrophil-to-

lymphocyte ratio (NLR) can serve as a basis for bacterial infection, infection rate, and antibiotic therapy [18], as well as an inflammation marker for diagnosing HIV and a warning for HIV severity [19, 20]. NLR is obtained from the absolute neutrophil count divided by the absolute lymphocyte count through a laboratory test, which aims to determine the occurrence of an inflammatory process [21]. The higher the NLR value, the lower the CD4 cell count, indicating that the relevant patients with HIV/AIDS are susceptible to other infections. Therefore, NLR is used as a warning of the infection severity [22]. This study is the first to observe the correlation between the bacterial and fungal profiles with the CD4 counts and the NLR values of HIV/AIDS-infected patients.

Materials and Methods

This analytic observational study applied a retrospective cross-sectional approach. The study population included medical record data of HIV/AIDS-infected patients aged ≥ 18 years old treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital from September 2021 to August 2022.

The inclusion criteria in this study included the HIV/AIDS inpatients treated in the special inpatient ward for HIV/AIDS isolation who sent their results of bacterial or fungal identification tests of all specimens to the clinical microbiology laboratory with complete CD4 and NLR data. Ethical clearance for this study has been issued by the Health Research Ethics Committee of Dr. Soetomo Tertiary Referral Hospital, Surabaya, based on a certificate of ethical clearance No. 1060/LOE/301.4.2/X/2022.

Descriptive analysis is displayed as tables containing frequency and percentage data. Likewise, the analysis was performed using the SPSS software with a 95% confidence interval. Data were analyzed univariately to obtain descriptive results regarding the patients' characteristics, including age, sex, types of OI, bacterial and fungal profiles, and CD4 profiles. The correlation test was utilized to determine the correlation between the CD4 cell counts and the

bacterial and fungal profiles of HIV/AIDS patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Regional Public Hospital and the correlation between NLR value and the bacterial and fungal profiles of HIV/AIDS patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital.

Results and Discussion

Resistance rate of bacteremia isolates due to CRAB

The total number of HIV-infected patients treated in the special inpatient ward for HIV/AIDS isolation who sent their results of bacterial and

fungal identification tests in the Microbiology Laboratory at Dr. Soetomo Regional Public Hospital during the study period from September 2021 to August 2022 amounted to 206 patients, and those who had complete CD4 and NLR data amounted to 76 patients. A total of 192 microbiological examination results for the identification of bacteria and fungi were obtained, which were associated with the patients' CD4 counts and NLR values.

Characteristics of the HIV/AIDS-infected patients treated in the special inpatient ward for HIV/AIDS isolation

Table 1: Characteristics of the HIV/AIDS-infected patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital

Characteristics	Number	Percentage
Sex		
Male	59	77.6
Female	17	22.4
Age categories		
15-19 years old	0	0
20-24 years old	10	13.6
25-49 years old	55	76.3
≥ 50 years old	11	10.2
HIV infection stages		
Stage 1 (CD4 counts of > 500)	3	3.26
Stage 2 (CD4 counts of 200-499)	10	10.87
Stage 3 (CD4 counts < 200)	79	85.87
ARV History		
Treatment-naive HIV/AIDS	34	44.7
HIV/AIDS on ARV	37	48.7
Drug-withdrawal HIV/AIDS	5	6.6
NLR Values		
CD4 counts of > 500	7.6 + 0.73	
CD4 counts of 200-499	11 + 5.68	
CD4 counts of 100-199	7 + 4.54	
CD4 counts of 50-99	13.6 + 11.07	
CD4 counts of < 50	11.02 + 9.60	
Number of opportunistic infections		
No opportunistic infection	22	28.9
One type of opportunistic infection	30	39.5
Two types of opportunistic infection	21	27.6
Three types of opportunistic infection	2	2.6
Four types of opportunistic infection	1	1.3
Types of opportunistic infection		
Pulmonary TB	18	23.7
Extrapulmonary TB	1	1.3
Encephalitis	3	3.9

	PCP	10	13.2
	Toxoplasmosis	14	18.4
	Diarrhea	12	15.8
	Hepatitis A	1	1.3
	Hepatitis B	5	6.6
	Hepatitis C	2	2.6
	Oral candidiasis	16	21.1
	ISK	11	14.5
	Pneumonia	22	28.9
Administration of antibiotic therapy			
	Receive antibiotic therapy	159	82.8
	Do not receive antibiotic therapy	331	17.2
Albumin, hematological, & electrolyte abnormalities			
	Hypoalbuminemia	43	56.6
	Anemia	37	48.7
	Leukopenia	3	3.9
	Pancytopenia	3	3.9
	Thrombocytopenia	1	1.3
	Hyponatremia	34	44.7
	Hypokalemia	20	26.3
	Hyperkalemia	4	5.3
	Hypernatremia	3	3.9

Table 2 presents that the lower the CD4 counts, the more OI the patient has. A study at Bandar Lampung Hospital, Indonesia, indicated a strong correlation between the CD4 cell counts of patients with HIV/AIDS and the incidence of OI. CD4 counts can serve as a good marker for assessing the HIV development and the OI possibility [23-25]. Meanwhile, a study on HIV/AIDS patients who had been taking ARVs for six months at the VCT outpatient department of Surakarta Hospital, Indonesia, stated that the OI incidence was affected by CD4 cell counts, not by viral load and NLR values [26, 27].

A study conducted by Subhas B. *et al.*, in 2015-2018 in India, found that there were hematological abnormalities at all stages in HIV/AIDS patients. These disorders may include anemia, leukopenia, lymphocytopenia, and thrombocytopenia [28] as presented in Table 3. The hematological profile of HIV-infected patients appears to reflect the degree of viral replication, with severe abnormalities observed in patients with reduced CD4 cell counts and high viral loads [29]. It is in accordance with Table 3, which shows that the lower the CD4 count, the

higher the incidence of hematological abnormalities.

Cytopenia develops due to various mechanisms. HIV infection causes increased expression of proinflammatory cytokines, such as tumor necrosis factor-alpha (TNF- α), transforming growth factor-beta (TGF- β), and interleukin-1 (IL-1), which result in myelosuppression and changes in the bone marrow microenvironment. Immune-mediated destruction of blood cells may also occur. Opportunistic infections, malignancies, and HIV treatment can further contribute to developing cytopenias [30, 31].

Several studies have revealed anemia as the most common cytopenia in HIV-infected individuals, as seen in Table 3. The Brazilian study further determined that anemia in HIV patients was independently associated with death, making hemoglobin a valuable prognostic biomarker [32, 29].

Several factors contribute to anemia in HIV-infected patients. Several studies have shown that the pathophysiology of anemia in HIV is influential. Another factor that contributes to anemia in HIV-infected individuals is CD4 count. A study conducted by Sang Bagus *et al.* (2018), in

Bali, found a significant correlation between Hb levels and CD4 counts. Anemia can serve as a marker for decreased CD4 counts, so a simple complete blood count can support the diagnosis

of a decreased CD4 count [17]. It is in accordance with the results of this study presented in Table 3, which demonstrates that incidence of anemia increases with lower CD4 count.

Table 2: Distribution of opportunistic infections by CD4 counts

Opportunistic infections	CD4 counts of > 500 $\Sigma = 2$ n (%)	CD4 counts of 200-499 $\Sigma = 5$ n (%)	CD4 counts of 100-199 $\Sigma = 10$ n (%)	CD4 counts of 50-199 $\Sigma = 17$ n (%)	CD4 counts of < 50 $\Sigma = 42$ n (%)
Pulmonary TB					
Yes ($\Sigma = 18$)	0 (0)	0 (0)	4 (40)	6 (35.3)	8 (19)
No ($\Sigma = 58$)	2 (100)	5 (100)	6 (60)	11 (64.7)	34 (81)
Extrapulmonary TB					
Yes ($\Sigma = 1$)	0 (0)	0 (0)	0 (0)	0 (0)	1 (2.4)
No ($\Sigma = 75$)	2 (100)	5 (100)	10 (100)	17 (99.4)	41 (97.6)
Encephalitis					
Yes ($\Sigma = 3$)	0 (0)	0 (0)	0 (0)	1 (5.9)	2 (4.8)
No ($\Sigma = 73$)	2 (100)	5 (100)	10 (100)	16 (94.1)	40 (95.2)
PCP					
Yes ($\Sigma = 10$)	0 (0)	0 (0)	3 (30)	1 (5.9)	6 (14.3)
No ($\Sigma = 66$)	2 (100)	5 (100)	7 (70)	16 (94.1)	36 (85.7)
Toxoplasmosis					
Yes ($\Sigma = 14$)	0 (0)	0 (0)	2 (20)	4 (23.5)	8 (19)
No ($\Sigma = 62$)	2 (100)	5 (100)	8 (80)	13 (76.5)	34 (81)
Diarrhea					
Yes ($\Sigma = 12$)	0 (0)	0 (0)	5 (50)	2 (11.8)	5 (11.9)
No ($\Sigma = 64$)	2 (100)	5 (100)	5 (50)	15 (88.2)	37 (88.1)
Hepatitis A					
Yes ($\Sigma = 1$)	0 (0)	0 (0)	1 (10)	0 (0)	0 (0)
No ($\Sigma = 75$)	2 (100)	5 (100)	9 (90)	17 (100)	42 (100)
Hepatitis B					
Yes ($\Sigma = 5$)	0 (0)	0 (0)	0 (0)	3 (17.6)	2 (4.8)
No ($\Sigma = 71$)	2 (100)	5 (100)	10 (100)	14 (82.4)	40 (95.2)
Hepatitis C					
Yes ($\Sigma = 2$)	0 (0)	0 (0)	0 (0)	0 (0)	2 (4.8)
No ($\Sigma = 74$)	2 (100)	5 (100)	10 (100)	17 (100)	40 (95.2)
Oral candidiasis					
Yes ($\Sigma = 16$)	0 (0)	0 (0)	3 (30)	1 (5.9)	12 (28.6)
No ($\Sigma = 60$)	2 (100)	5 (100)	7 (70)	16 (94.1)	30 (71.4)
ISK					
Yes ($\Sigma = 11$)	1 (50)	1 (20)	1(10)	2 (11.8)	6 (14.3)
No ($\Sigma = 65$)	1 (50)	4 (80)	9 (90)	15 (88.2)	36 (85.7)
Pneumonia					
Yes ($\Sigma = 22$)	0 (0)	1 (20)	1 (10)	3 (12.6)	17 (40.5)
No ($\Sigma = 58$)	2 (100)	4 (80)	9 (90)	14 (82.4)	25 (59.5)

Table 3: Albumin, hematological, and electrolyte abnormalities in HIV/AIDS-infected patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital based on CD4 counts

Albumin, hematological, and electrolyte abnormalities	CD4 counts of >500 $\Sigma = 2$ n (%)	CD4 counts of 200-499 $\Sigma = 5$ n (%)	CD4 counts of 100-199 $\Sigma = 10$ n (%)	CD4 counts of 50-99 $\Sigma = 17$ n (%)	CD4 counts of <50 $\Sigma = 42$ n (%)
Hypoalbuminemia					
Yes ($\Sigma = 43$)	2 (100)	2 (40)	4 (40)	10 (58.8)	25 (59.5)
No ($\Sigma = 33$)	0 (0)	3 (60)	6 (60)	7 (41.2)	17 (40.5)
Anemia					
Yes ($\Sigma = 37$)	1 (50)	3 (60)	5 (50)	12 (70.6)	16 (38.1)
No ($\Sigma = 39$)	1 (50)	2 (40)	5 (50)	5 (29.4)	26 (61.9)
Leukopenia					
Yes ($\Sigma = 3$)	0 (0)	0 (0)	0 (0)	0 (0)	3 (7.1)
No ($\Sigma = 73$)	2 (100)	5 (100)	10 (100)	17 (100)	39 (92.9)
Thrombocytopenia					
Yes ($\Sigma = 1$)	0 (0)	0 (0)	0 (0)	1 (5.9)	0 (0)
No ($\Sigma = 75$)	2 (100)	5 (100)	10 (100)	16 (94.1)	42 (100)
Pancytopenia					
Yes ($\Sigma = 3$)	0 (0)	0 (0)	0 (0)	0 (0)	3 (7.1)
No ($\Sigma = 73$)	2 (100)	5 (100)	10 (100)	17 (100)	39 (92.9)
Hyponatremia			-		-
Yes ($\Sigma = 34$)	0 (0)	0 (0)	4 (40)	8 (47.1)	22 (52.4)
No ($\Sigma = 42$)	2 (100)	5 (100)	6 (60)	9 (52.9)	20 (47.6)
Hypokalemia					
Yes ($\Sigma = 20$)	1 (50)	1 (20)	6 (60)	2 (11.8)	10 (23.8)
No ($\Sigma = 56$)	1 (50)	4 (80)	4 (40)	15 (88.2)	32 (76.2)
Hypernatremia					
Yes ($\Sigma = 5$)	1 (50)	0 (0)	0 (0)	0 (0)	2 (4.8)
No ($\Sigma = 71$)	1 (50)	5 (100)	10 (100)	17 (100)	40 (95.2)
Hyperkalemia					
Yes ($\Sigma = 4$)	0 (0)	0 (0)	0 (0)	1 (5.9)	3 (7.1)
No ($\Sigma = 72$)	2 (100)	5 (100)	10 (100)	16 (94.1)	39 (92.9)

Characteristics of bacterial and fungal identification results of HIV/AIDS-infected patients treated in the inpatient ward at Dr. Soetomo Regional Public Hospital

The gram-negative bacteria with the highest prevalence in a study by Tan *et al.* in the same hospital in the previous period were *Klebsiella pneumoniae* (35.15%), followed by *Escherichia coli* (10.8%), *Pseudomonas aeruginosa* (8.1%), and *Acinetobacter baumannii* (8.1%) (Table 4) [33]. Meanwhile, in this study, as depicted in Table 5, the gram-negative bacteria with the highest prevalence were *Escherichia coli* ESBL and *Pseudomonas aeruginosa*, followed by *Klebsiella pneumoniae* and *Escherichia coli*. The

gram-positive bacteria with the highest prevalence in the study by Tan *et al.* were *Streptococcus mitis/oralis* (30.7%), followed by *Staphylococcus aureus* (23.1%) and *Staphylococcus epidermidis* (15.4%) [33].

Table 5 indicates that gram-negative bacteria have a higher percentage than gram-positive bacteria. It is in line with a study conducted on inpatients with HIV/AIDS at the same hospital from August 2019 to February 2020, which stated that the most common types of bacteria were gram-negative bacteria at 44.6%, followed by gram-positive bacteria at 15.7%, while negative culture results reached 37.3% [33]. This study obtained more negative culture results at 52.5%.

Meanwhile, in this study, as displayed in Table 5, the gram-positive bacteria with the highest prevalence were *Viridans streptococci* (24.32%), followed by coagulase-negative *Staphylococci* (16.22%), *Staphylococcus haemolyticus* (13.51%), *Staphylococcus aureus* (10.81%), *Staphylococcus haemolyticus* (13.51%), *Staphylococcus hominis* (8.11%), and methicillin-resistant *Staphylococcus aureus* (5.41%). In comparison, those with the lowest prevalence of 2.70% were vancomycin-resistant *Enterococcus faecalis*, *Mycobacterium tuberculosis*, *Staphylococcus sciuri*, *Staphylococcus carnosus*, *Pediococcus pantosaceus*, *Enterococcus faecium*, *Enterococcus faecalis*, and *Corynebacterium tuberculosis*.

In this study, the prevalence of bacteriuria/candiduria was lower at 51.16%, most of which were found in female patients (53.8%), and most were aged 25-49 years old (59.1%). The isolates with the highest prevalence

were *E. coli* ESBL at 13.95%, followed by *Pseudomonas aeruginosa* and *E. coli* at 4.65%. The prevalence of ESBL bacteria in this study was lower than that in a study conducted at Tertiary Referral Hospital in Central Europe, which stated that 16.8% of HIV/AIDS patients with positive UTIs were infected with bacteria that produce Extended-spectrum beta-lactamases (ESBLs) [34].

This study also shows that the lower the CD4 count, the higher the prevalence of bacteriuria/candiduria, which is in line with a study conducted by Tan *et al.*, which stated that HIV-infected individuals with CD4 counts of >200 cells/mm³ are less likely to experience a UTI than HIV-infected individuals with a CD4 count of <200 cells/mm³. They also stated that there was no significant correlation between other risk factors in their study [16].

Table 4: Percentage of bacteria and fungi based on specimens in HIV/AIDS-infected patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital

Bacterial and fungal profiles by specimens	N (%/SPECIMENS)	% Total specimens
Sputum specimens	69	35.94
Bacterial cultures	39 (56.5)	20.3
There was no growth of M. TB bacteria	2 (2.9)	1
There was no growth of bacteria	2 (2.9)	1
Gram-negative bacteria	18 (26.1)	9.37
<i>Acinetobacter baumannii</i>	1 (1.4)	0.5
<i>Citrobacter koseri</i>	1 (1.4)	0.5
<i>Enterobacter cloacae</i>	2 (2.9)	1
<i>Eschericia coli</i>	1 (1.4)	0.5
<i>Eschericia coli</i> ESBL	2 (2.9)	1
<i>Klebsiella pneumonia</i>	3 (4.3)	1.6
<i>Klebsiella pneumonia</i> ESBL	1 (1.4)	0.5
<i>Carbapenem-resistant Klebsiella pneumonia</i>	1 (1.4)	0.5
<i>Moraxella spp</i>	1 (1.4)	0.5
<i>Pseudomonas aeruginosa</i>	5	2.6
Gram-positive bacteria	17 (24.6)	8.8
<i>Staphylococcus aureus</i>	2 (2.9)	1
<i>Coagulase-negative Staphylococci</i>	6 (8.7)	3.1
<i>Viridans streptococci</i>	9 (13)	4.7
Fungal cultures	8 (11.6)	4.2
<i>Candida albicans</i>	7 (10.1)	3.6
<i>Candida tropicalis</i>	1 (1.4)	0.5
TB expert gene tests	22 (31.9)	11.5
<i>Not detected M. tuberculosis</i>	16 (23.2)	8.3
<i>Rifampicin-sensitive M. tuberculosis</i>	6 (8.7)	3.1
BLOOD SPECIMENS	67	34.9

There was no growth of bacteria	57 (85.1)	29.7
Gram-positive bacteria	9 (13.4)	4.7
<i>Pediococcus pentosaceus</i>	1 (1.5)	0.5
<i>Staphylococcus aureus</i>	1 (1.5)	0.5
<i>Staphylococcus haemolyticus</i>	3 (4.5)	1.6
<i>Staphylococcus hominis</i>	3 (4.5)	1.6
<i>Corynebacterium urealyticum</i>	1 (1.5)	0.5
Gram-negative bacteria	1 (1.5)	0.5
<i>Kluyvera intermedia</i>	1 (1.5)	0.5
Urine specimens	43 (100)	22.4
Bacterial cultures	42 (97.7)	21.9
There was no growth of bacteria	21 (48.9)	10.9
Gram-negative bacteria	16 (37.2)	8.3
<i>Acinetobacter baumannii</i>	1 (2.3)	0.5
<i>Citrobacter koseri</i>	1 (2.3)	0.5
<i>Eschericia coli</i>	2 (4.7)	1
Carbapenem-resistant <i>Eschericia coli</i>	1 (2.3)	0.5
<i>Eschericia coli</i> ESBL	6 (14)	3.1
<i>Klebsiella pneumoniae</i>	1 (2.3)	0.5
<i>Klebsiella pneumoniae</i> ESBL	1 (2.3)	0.5
<i>Pseudomonas aeruginosa</i>	2 (4.7)	1
<i>Proteus vulgaris</i>	1 (2.3)	0.5
Gram-positive bacteria	5 (11.6)	2.5
<i>Enterococcus faecalis</i>	1 (2.3)	0.5
Vancomycin-resistant <i>Enterococcus faecalis</i>	1 (2.3)	0.5
<i>Enterococcus faecium</i>	1 (2.3)	0.5
<i>Staphylococcus carnosus</i>	1 (2.3)	0.5
<i>Staphylococcus haemolyticus</i>	1 (2.3)	0.5
Fungal cultures	1 (2.3)	0.5
<i>Candida albicans</i>	1 (2.3)	0.5
PUS specimens	8 (100)	4.2
Gram-negative bacteria	4 (50)	2.1
<i>Eschericia coli</i>	1 (12.5)	0.5
<i>Eschericia coli</i> ESBL	1 (12.5)	0.5
<i>Pseudomonas aeruginosa</i>	2 (25)	1
Gram-positive bacteria	4 (50)	2.1
<i>Staphylococcus aureus</i>	1 (12.5)	0.5
Methicillin-resistant <i>Staphylococcus aureus</i>	2 (25)	1
<i>Staphylococcus sciuri</i>	1 (12.5)	0.5
Cerebrospinal fluid specimens	4 (100)	2.1
There was no growth of bacteria	1 (25)	0.5
<i>Staphylococcus haemolyticus</i>	1 (25)	0.5
<i>M. tuberculosis</i> (<i>M.tb</i> culture)	1 (25)	0.5
Not detected <i>M. tuberculosis</i> (gene expert)	1 (25)	0.5
Stool specimens	1 (100)	0.52
There was no growth of bacteria	1 (100)	0.5

Table 5: Bacterial and fungal identification results in all specimens in HIV/AIDS-infected patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital

Bacterial and fungal profiles by specimens	N (%/TESTS)	% Total specimens
Bacterial culture tests	160 (100)	83.3
Non-MDRO bacteria	60 (37.5)	31.25
Gram-negative bacteria	26 (16.25)	13.54
<i>Acinetobacter baumannii</i>	2 (1.25)	1.04
<i>Citrobacter koseri</i>	2 (1.25)	1.04
<i>Enterobacter cloacae</i>	2 (1.25)	1.04
<i>Eschericia coli</i>	4 (2.5)	2.08
<i>Klebsiella pneumonia</i>	4 (2.5)	2.08
<i>Kluyvera intermedia</i>	1 (0.62)	0.52
<i>Moraxella spp</i>	1 (0.62)	0.52
<i>Pseudomonas aeruginosa</i>	9 (5.63)	4.69
<i>Proteus vulgaris</i>	1 (0.62)	0.52
Gram-positive bacteria	34 (21.25)	17.89
<i>Corynebacterium urealyticum</i>	1 (0.62)	0.52
<i>Enterococcus faecalis</i>	1 (0.62)	0.52
<i>Enterococcus faecium</i>	1 (0.62)	0.52
<i>Pediococcus pantosaceus</i>	1 (0.62)	0.52
<i>Staphylococcus aureus</i>	4 (2.5)	2.08
<i>Staphylococcus carnosus</i>	1 (0.62)	0.52
<i>Coagulase-negative Staphylococci</i>	6 (3.75)	3.12
<i>Staphylococcus haemolyticus</i>	5 (3.13)	2.60
<i>Staphylococcus hominis</i>	3 (1.88)	1.57
<i>Staphylococcus sciuri</i>	1 (0.62)	0.52
<i>Streptococcus viridans</i>	9 (5.63)	4.69
<i>Mycobacterium tuberculosis</i>	1 (0.62)	0.52
MDRO bacteria	16 (0.1)	8.3
Gram-negative bacteria	13 (8.13)	6.77
<i>Eschericia coli</i> ESBL	9 (5.63)	4.69
<i>Carbapenem-resistant Eschericia coli</i>	1 (0.62)	0.52
<i>Klebsiella pneumonia</i> ESBL	2 (1.25)	1.04
<i>Carbapenem-resistant Klebsiella pneumonia</i>	1 (0.62)	0.52
Gram-positive bacteria	3 (1.88)	1.56
<i>Vancomycin-resistant Enterococcus faecalis</i>	1 (0.6)	0.52
<i>Methicillin-resistant Staphylococcus aureus</i>	2 (1.25)	1.04
There was no growth of bacteria	84 (52.5)	43.75
<i>M. tuberculosis</i> Gene Expert Tests	23 (100)	12
Not detected <i>M. tuberculosis</i>	17 (73.9)	8.88
Rifampicin-sensitive <i>M. tuberculosis</i>	6 (26.1)	3.12
FUNGAL culture tests	9 (100)	4.70
<i>Candida albicans</i>	8 (88.9)	4.18
<i>Candida tropicalis</i>	1 (11.1)	0.52

Analysis of the correlation between the CD4 counts and the bacterial and fungal profiles of HIV/AIDS-infected patients Treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital

A study on urine samples of HIV/AIDS patients in Nigeria showed that CD4 counts might have little or no effect on bacteriuria. Still, with CD4 counts of ≤ 200 cells/mm³, the study subjects were more likely to show UTI symptoms even at low bacterial counts [35]. In accordance with this

study, as listed in Table 6, the UTI incidence is more common in patients with CD4 counts of <200, especially in patients with CD4 cell counts of <50.

CD4 is the target cell for HIV infection. Until now, CD4 cell counts remain an essential parameter in HIV/AIDS patients. Decreased CD4 counts in HIV infection make the patient susceptible to infection. HIV causes a cytopathic effect on CD4 by activating the immune system, resulting in PMN cell apoptosis and damage to stem cells and

lymphoid tissue, including the thymus, so that no new cells are formed. It interferes with the function and number of neutrophils. Neutrophil dysfunction makes HIV patients susceptible to bacterial infections [36]. The correlation between CD4 counts and bacterial and fungal profiles, as demonstrated in Table 6, indicated insignificant results in bivariate analysis using Spearman's rank correlation coefficient, with a p-value of 0.350.

Table 6: Bacterial and fungal profiles of HIV/AIDS-infected patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital based on CD4 count categories

Species name	CD4 counts of > 500	CD4 counts of 200-499	CD4 counts of 100-199	CD4 counts of 50-99	CD4 counts of < 50	Total
<i>Acinetobacter baumannii</i>	0 (0)	0 (0)	0 (0)	1(2.6)	1 (0.9)	2 (1)
<i>Citrobacter koseri</i>	0 (0)	0 (0)	1 (3)	1(2.6)	0 (0)	2 (1)
<i>Enterobacter cloacae</i>	0 (0)	0 (0)	0 (0)	0 (0)	2 (1.9)	2 (1)
<i>Eschericia coli</i>	0 (0)	0 (0)	0 (0)	0 (0)	4 (3.7)	4 (2.1)
<i>Klebsiella pneumonia</i>	0 (0)	0 (0)	3 (9.1)	0 (0)	1 (0.9)	4 (2.1)
<i>Kluyvera intermedia</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
<i>Moraxella spp</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
<i>Pseudomonas aeruginosa</i>	1 (20)	0 (0)	0 (0)	2 (5.1)	6 (5.6)	9 (4.7)
<i>Proteus vulgaris</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
<i>Eschericia coli</i> ESBL	0 (0)	0 (0)	0 (0)	1(2.6)	8 (7.5)	9 (4.7)
<i>Carbapenem-resistant Eschericia coli</i>	1 (20)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.5)
<i>Klebsiella pneumonia</i> ESBL	0 (0)	0 (0)	0 (0)	1(2.6)	1 (0.9)	2 (1)
<i>Carbapenem-resistant Klebsiella pneumonia</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
<i>Corynebacterium urealyticum</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
<i>Enterococcus faecalis</i>	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	1 (0.5)
<i>Enterococcus faecium</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (100)	1 (0.5)
<i>Pediococcus pantosaceus</i>	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	1 (0.5)
<i>Staphylococcus aureus</i>	0 (0)	0 (0)	0 (0)	1(2.6)	3 (2.8)	
<i>Staphylococcus carnosus</i>	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	1 (0.5)
<i>Coagulase-negative Staphylococci</i>	0 (0)	0 (0)	1 (3)	2 (5.1)	3 (2.8)	6 (3.1)
<i>Staphylococcus haemolyticus</i>	0 (0)	1 (12.5)	0 (0)	2 (5.1)	2 (1.9)	5 (2.6)
<i>Staphylococcus hominis</i>	0 (0)	1 (12.5)	0 (0)	1(2.6)	1 (0.9)	3 (1.6)
<i>Staphylococcus sciuri</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
<i>Streptococcus viridans</i>	0 (0)	0 (0)	2 (6.1)	1(2.6)	6 (5.6)	9 (4.7)
<i>Mycobacterium tuberculosis</i>	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.9)	1 (0.5)
<i>Vancomycin-resistant Enterococcus faecalis</i>	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	1 (0.5)

<i>Methicillin-resistant Staphylococcus aureus</i>	0 (0)	0 (0)	0 (0)	0 (0)	2 (1.9)	2 (1)
<i>Sterile</i>	3 (60)	4 (50)	16 (48.5)	19 (48.7)	42 (39.3)	84 (43.8)
<i>Rifampicin-sensitive M. tuberculosis</i>	0 (0)	0 (0)	1 (3)	3 (7.7)	2 (1.9)	6 (3.1)
<i>Not detected M. tuberculosis</i>	0 (0)	2 (25)	3 (9.1)	1(2.6)	11 (10.3)	17 (8.9)
<i>Candida albicans</i>	0 (0)	0 (0)	1 (3)	3 (7.7)	4 (3.7)	8 (4.2)
<i>Candida tropicalis</i>	0 (0)	0 (0)	1 (3)	0 (0)	0 (0)	1 (0.5)
Total	5 (100)	8 (100)	33 (100)	39 (100)	107 (100)	192 (100)

The bivariate analysis showed an insignificant correlation between the CD4 cell counts and the bacterial and fungal profiles, with a p-value of 0.350.

Analysis of the correlation between the CD4 counts and the positive values of bacterial and fungal identification of HIV/AIDS-infected patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital

The correlation between CD4 counts and the positive values of bacterial or fungal identification, as provided in [Table 7](#), also revealed insignificant results in bivariate analysis using Spearman's rank correlation coefficient, with a p-value of 0.238.

Some studies show a lack of association between CD4 and bacteriuria. For instance, a study in Nigeria noticed that bacteriuria in HIV-infected women was not affected by CD4 counts [37], bacteriuria was not affected by CD4 counts of less than 200 cells/mm³ [38], and CD4 cell counts of <200 cells/mm³ does not affect the occurrence of asymptomatic bacteriuria [39]. Other studies yielded different results in that bacteriuria was found to correlate to CD4 cell counts and/or viral loads [40-42].

Bacteriuria may not be affected by CD4 counts because research shows that immunity to UTIs results from complex interactions between host factors and the invading organism. Defenses against UTIs include urinary pH, urinary flushing

effect, high salt concentration, chemical inhibition secreted by the epithelial cells lining the urinary tract, secretory IgA, and lactoferrin activity (an iron-binding protein that absorbs iron, invasion of starved bacteria that causes decreased growth rate). The prostate gland in men secretes an infection-fighting substance. The epithelial cells lining the urinary tract also secrete antimicrobial peptides that directly kill bacteria. The mucous lining of the bladder contains antimicrobial substances that help eliminate invading pathogens. This defense mechanism is not influenced by the host's CD4 counts. This partly explains why individuals with CD4 cell counts of less than 100 cells/mm³ can have sterile cultures, despite the lack of exposure to UTI causative agents can be responsible for that. It suggests that the host's CD4 counts may have little or no effect on bacteriuria [35].

The neutrophil-to-lymphocyte ratio (NLR) is a simple parameter utilized to assess the inflammatory status and results from the division of total neutrophils by total lymphocytes [43]. NLR is a marker of inflammation and HIV severity [19]. [Table 1](#) presents the highest average NLR values in the 50-99 CD4 count range and the lowest in the 100-199 count CD4 range. The average NLR value varies in different CD4 count ranges, which can be caused by the administration of antibiotics received by the patients.

Table 7: Positive values of bacterial and fungal identification based on the CD4 count categories

Bacterial and fungal profiles	CD4 counts of > 500 n (%)	CD4 counts of 200-499 n (%)	CD4 counts of 100-199 n (%)	CD4 counts of 50-99 n (%)	CD4 counts of < 50 n (%)	Total n (%)
Results (+)	3 (60)	6 (75)	19 (57.6)	21 (53.8)	53 (49.5)	102 (53.1)
Results (-)	2 (40)	2 (25)	14 (42.4)	18 (46.2)	86 (48)	90 (46.9)
Total	5 (100)	8 (100)	33 (100)	39 (100)	107 (100)	192 (100)

The bivariate analysis revealed an insignificant correlation between the CD4 cell counts and the positive bacterial or fungal identification values, with a p-value of 0.238.

Analysis of the correlation between the NLR values and the bacterial and fungal profiles of HIV/AIDS-infected patients treated in the special inpatient

ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospita

The correlation between NLR value and bacterial and fungal profiles, as shown in [Table 8](#), indicated insignificant results in bivariate analysis using Spearman's rank correlation coefficient, with a p-value of 0.956.

Table 8: Bacterial and fungal profiles based on NLR values

Bacterial and fungal profiles	NLR
There was no growth of bacteria	11.5 + 9.80
Gram-positive bacteria	10.14 + 7.67
Gram-negative bacteria	13.09 + 13.56
MDRO	9.67 + 5.11
<i>Rifampicin-sensitive M. tuberculosis</i>	10.07 + 5.15
<i>Not detected M. tuberculosis</i>	8.23 + 5.86
Fungus	7.13 + 4.63

The bivariate analysis showed an insignificant correlation between the NLR values and the bacterial and fungal profiles, with a p-value of 0.557.

Analysis of the correlation between the NLR values and the positive values of bacterial and fungal identification of HIV/AIDS patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital

The correlation between NLR value and the positive values of bacterial or fungal identification, as illustrated in [Table 9](#), also revealed insignificant results in bivariate analysis using Spearman's rank correlation coefficient, with a p-value of 0.557. This study shows that the bacterial and fungal profiles have no significance to the NLR values because, as depicted in [Table 1](#), 82.8% of the isolates were exposed to antibiotics. Only 17.2% of isolates were not exposed to

antibiotics, even though antibiotics could not quickly turn the results of the gene expert TB tests into negative. In addition, the gene expert tests in this study only account for 12%, so it can be concluded that the antibiotics administration can affect the NLR values. Commensal bacteria at 34.2% in the bacterial culture results indicated that the bacteria obtained were normal flora, even though they can serve as causative agents in immunocompromised patients.

NLR is affected by many conditions, including age, race, medications, and chronic diseases, such as coronary heart disease, stroke, diabetes, obesity, psychiatric diagnosis, solid organ cancer, anemia, and stress [43]. In Asian races, NLR values will usually be higher in women than those in men because estrogen can inhibit apoptosis. Older people, especially women, have lower NLR values due to decreased estrogen due to menopause [44, 27].

Table 9: Positive values of bacterial and fungal identification based on the average NLR value

Bacterial and fungal profiles	NLR
Bacterial and fungal identification (+)	9.71 ± 7.19
Bacterial and fungal identification (-)	10.80 ± 9.85

The bivariate analysis indicated an insignificant correlation between the NLR values and the positive values of bacterial and fungal identification of HIV/AIDS-infected patients treated in the inpatient ward at Dr. Soetomo Regional Public Hospital, with a p-value of 0.956. The distribution of HIV/AIDS-infected patients in the samples of this study showed that male patients had a dominant percentage of 77.6%. It differs from the data in global HIV statistics issued by UNAIDS, which stated that 54% of HIV/AIDS-infected people were women and girls [23]. Still, the percentage of male patients in this study was slightly higher than that in the 2020 HIV data from the Data and Information Center of the Ministry of Health of the Republic of Indonesia, which stated that the percentage of male HIV-infected patients reached 64.5% and that of male AIDS-infected patients reached 68.60% [24]. The percentage of male patients is higher than that of female patients because men are more likely to have risky sexual intercourse than women and more likely to use illegal drugs (injecting needles) [25].

Most HIV/AIDS patients in this study were aged 25-49 years old, amounting to 72.4%. It is higher than that in the data and information center of the Ministry of Health of the Republic of Indonesia in 2021, which states that most HIV patients come from the age range of 25-49 years old, amounting to 69.7% [24]. UNAIDS also states that men aged 25-49 occupy the largest population at 42% [26]. The percentage of HIV/AIDS patients aged 25-49 years old in this study was higher than that in data from the Ministry of Health and UNAIDS because this study only included adult inpatients. In comparison, data from the Ministry of Health and UNAIDS included all patients from 0 year old. The age range of 25-49 years old is a productive age, and it is very likely for those in this age range to get infected with HIV due to high sexual activity.

Conclusion

According to the profiles of HIV/AIDS patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Tertiary Referral Hospital from September 2021 to August 2022, most of them were male patients at 77.6% aged 20-24 years old at 13.2%, with the highest HIV stage being the AIDS stage at 90.8%. The highest OIs included pneumonia at 28.9%, followed by pulmonary TB at 23.7% and oral candidiasis at 21.1%. The bacterial culture tests of 83.3% showed 52.5% negative cultures, 47.5% positive cultures, and 51.33% gram-negative bacteria, with *Pseudomonas aeruginosa* and *Escherichia coli* ESBL as the highest number of isolates, 48.67% gram-positive bacteria with *Streptococcus viridans* as the highest number of isolates, 34.2% commensal bacteria, 65.8% pathogenic bacteria, and 21.05% MDRO bacteria. The TB expert gene tests of 12% resulted in 73.9% undetected *M. tuberculosis* and 26.1% positive, rifampicin-sensitive *M. tuberculosis*, while the fungal culture tests of 4.7% resulted in 88.9% *Candida albicans* and 11.1% *Candida tropicalis*. There is no correlation between the profile of bacteria and fungi from the clinical specimens with CD4 counts and NLR values of HIV/AIDS patients treated in the special inpatient ward for HIV/AIDS isolation at Dr. Soetomo Regional Public Hospital from September 2021 to August 2022. The profiles of HIV/AIDS patients in this study did not describe all the patients as a whole because the data were collected only from patients with complete medical records of CD4 counts and NLR values.

Acknowledgments

The authors present their gratitude to Dr. Soetomo Public Academic Hospital and the Department of Microbiology, Faculty of Medicine, Universitas Airlangga, for providing all necessary support in this study.

Disclosure Statement

No potential conflict of interest was reported by the authors.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' Contributions

All authors contributed to data analysis, drafting, and revising of the paper and agreed to be responsible for all the aspects of this work.

Orcid

Nafduz Makhmudatul Muna

<https://orcid.org/0000-0003-3871-3455>

Agung Dwi Wahyu Widodo

<https://orcid.org/0000-0002-3449-768X>

Pepy Dwi Endraswari

<https://orcid.org/0000-0002-0271-8505>

M. Vitanata Arfijanto

<https://orcid.org/0000-0003-4510-755X>

References

- [1]. Khademi F., Bacterial Co-infections in HIV/AIDS-positive Subjects: A Systematic Review and Meta-analysis', *Folia medica*, 2018, **60**:339 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [2]. Sutini S., Rahayu S.R., Saefurrohman M.Z., Al Ayubi M.T.A., Wijayanti H., Wandastuti A.D., Miarso D., Susilastuti M.S., Prevalence and Determinants of Opportunistic Infections in HIV Patients: A Cross-Sectional Study in the City of Semarang, *Ethiopian Journal of Health Sciences*, 2022, **32**:809 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [3]. Witaningrum A.M., Khairunisa S.Q., Yunifiar M.Q., Bramanathi R., Rachman B.E., Opportunistic infection of HIV / AIDS patients in West Papua', In *IOP Conference Series: Earth and Environmental Science*, 2018, **125**:012062 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [4]. Infodatin Kemenkes R.I., Infodatin HIV', *Kementerian Kesehatan RI*, 2020, 1–12 [[Google Scholar](#)],
- [5]. Prabowo K.A, Gayatri A.A.A.Y., Sukmawati N.M.D.D., Utama M.S., Somia K.A., Merati K.T.P., HIV-related toxoplasma encephalitis: a tertiary care experience, *Indonesia Journal of Biomedical Science*, 2021, **15**:198 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [6]. Damtie D., Yismaw G., Woldeyohannes D., Anagaw B., Common Opportunistic Infections and Their CD4 Cell Correlates among HIV-Infected Patients Attending at Antiretroviral Therapy Clinic of Gondar University Hospital, Northwest Ethiopia, *BMC research notes*, 2013, **6**:1 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [7]. Jayani I., Sulistyawati W., The Correlation between CD4 Count Cell and Opportunistic Infection among HIV/AIDS Patients', *Journal of Physics: Conference Series*, 2020, **1569**:032066 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [8]. Fazelipour S., Moghadam F.A., Davudi P., Tootian Z., Assadi F., Histometrical study of ovarian follicles of immature mice treated with methylphenidate, *Journal of Veterinary Research*, 2015, **70**:3 [[Google Scholar](#)], [[Publisher](#)]
- [9]. Alebel A., Demant D., Petručka P., Sibbritt D., Effects of undernutrition on opportunistic infections among adults living with HIV on ART in Northwest Ethiopia: Using inverse-probability weighting', *PLoS ONE*, 2022, **17**:e0264843 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [10]. Dagnaw Tegegne K., Cherie N., Tadesse F., Tilahun L., Kassaw M.W., Biset G., Incidence and Predictors of Opportunistic Infections Among Adult HIV Infected Patients on Anti-Retroviral Therapy at Dessie Comprehensive Specialized Hospital, Ethiopia: A Retrospective Follow-Up Study', *HIV/AIDS - Research and Palliative Care*, 2022, **14**:195 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [11]. Mazlominezhad A., Moghadam F.A. Evaluation of quality of life and self-efficacy in adolescents with amblyopia, *Journal of Medicine and Life*, 2022, **15**:499 [[Crossref](#)], [[Google Scholar](#)]
- [12]. Daryabari S.H., Asadollah A., Moghadam F.A., Dorostkar R., Bahramifar A., Aghamollaei H., Correction to: Detection of COVID-19 in tears of ICU-admitted patients with SARS-CoV-2 infection, *International Ophthalmology*, 2022, **42**:723 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

- [13]. Anglaret X., Eholie S.P., Severe bacterial infections: overlooked opportunistic diseases', *The Lancet HIV*, 2017, **4**:e94 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [14]. Rameshkumar R.M., Arunagirinathan N., Drug-Resistant Bacterial Infections in HIV Patients', *Advances in HIV and AIDS Control*, 2018, **83** [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [15]. Limper A.H., Adenis A., Le T., Harrison T.S., Fungal infections in HIV/AIDS', *The Lancet Infectious Diseases*, 2017, **17**:e334 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [16]. Harningtyas C.D., Damayanti, Maylita S., Muhammad Y.L., Diah M.I., Linda A., Budi U., Dwi M., Setyana W., Astindari, Afif N.H., Relationship between CD4 levels and mucocutaneous manifestations in HIV-AIDS patients at Dr. Soetomo General Academic Teaching Hospital, Surabaya, Indonesia, *Bali Medical Journal*, 2022, **11**:216 [[Google Scholar](#)]
- [17]. Muhamad R.U.F., Solichul H., The Correlation between total lymphocyte count, hemoglobin levels, lymphocyte/leukocyte ratio (LLR), and lymphocyte/neutrophil ratio (LNR) to CD4 levels in patients with Human Immunodeficiency Virus infection at Sanglah Hospital, *Bali Medical Journal*, 2019, **8**:429 [[Google Scholar](#)], [[Publisher](#)]
- [18]. Kusnadi D., Liwang M.N.I., Katu S., Mubin A.H., Halim R., Correlation between the neutrophil-lymphocyte count ratio and bacterial infection in patient with human immunodeficiency virus', In *IOP Conference Series: Earth and Environmental Science*, 2018, **125**:012029 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [19]. Emokpae M.A., Aruomaren A.I., Mrakpor B.A., Association of Neutrophil-to-lymphocyte ratio with Respiratory burst enzymes in Human Immunodeficiency virus type 1 infected Africans, *Journal of Medical Discovery*, 2017, **2**:1 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [20]. Kumbara C.I.Y.K., Wibawa I.D.N., Somayana I.G., Suryadarma I.G.A., & Mariadi I.K. Diagnostic performance of APRI, RPR, FIB-4, GPR, AAR, and NLR in assessing the degree of liver fibrosis in patients with chronic hepatitis B, *Indonesia Journal of Biomedical Science*, 2020, **14**:122 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [21]. Wahyuni R.D., Sallatu N.A.P.R.S, Harun H., Rasjid M., Perbandingan Rasio Neutrofil Limfosit (RNL) pada Pasien TB Paru BTA Positif dan BTA Negatif di RSUD Anutapura Palu Tahun 2017, *Continuing Medical Education*, 2019, **1** [[Google Scholar](#)], [[Publisher](#)]
- [22]. Handayani K., Katu S., Bakri S., Halim R., Aman A.M., Rasyid H., Kasim H., Seweng A., Correlation of CD4 Count And Neutrophil-Lymphocyte Ratio in Human Immunodeficiency Virus-Acquired Immunodeficiency Syndrome (HIV- AIDS) Patients, *European Journal of Molecular & Clinical Medicine*, 2020, **7**:2020 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [23]. UNAIDS. Global HIV Statistics', Fact Sheet 2021, 1 [[Publisher](#)]
- [24]. Kemenkes RI. Pusat Data dan Informasi, 2022 [[Publisher](#)]
- [25]. Ladyani F., Kiristianingsih A., Hubungan Antara Jumlah CD4 Pada Pasien Yang terinfeksi HIV / AIDS Dengan Infeksi Oportunistik di Rumah Sakit Umum Abdul Moeloek Bandar Lampung Tahun 2016', *Jurnal Kedokteran Universitas Lampung*, 2019, **3**:34 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [26]. UNAIDS. UNAIDS Global AIDS Update 2022, 2022, Geneva [[Publisher](#)]
- [27]. Kurniawati V.V., Harioputro D.R., Susanto A.J., 'Evaluasi Kadar Sel Cd4, Viral Load, Dan Neutrophil Lymphocyte Ratio (Nlr) Terhadap Infeksi Oportunistik Pada Pasien Hiv/Aids', *Biomedika*, 2022, **14**:99 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [28]. Bhardwaj S., Almaeen A., Wani F.A., Thirunavukkarasu A., Hematologic derangements in HIV/AIDS patients and their relationship with the CD4 counts: a cross-sectional study', *International journal of clinical and experimental pathology*, 2020, **13**:756 [[Google Scholar](#)], [[Publisher](#)]
- [29]. Vaswani P.P.M., Senga-Tang I.R., Catapia J.R.M., Abad C.L.R., Dumagay T.E., The hematologic profile of Filipino HIV-infected individuals and its association with CD4 counts', *Hematology, Transfusion and Cell Therapy*, 2022, **44**:307 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [30]. Gunda D.W., Godfrey K.G., Kilonzo S.B., Mpondo B.C., Cytopenias among ART-naive patients with advanced HIV disease on enrolment

- to care and treatment services at a tertiary hospital in Tanzania: A cross-sectional study', *Malawi Medical Journal*, 2017, **29**:43 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [31]. Solati H., Sahebalzamani M., Moghadam F.A., 'Effect of Family-Based Care Training by Tele-nursing on Emotional Reactions in Mothers of Children with Bone Marrow Transplantation', *Journal of Mazandaran University of Medical Sciences*, 2021, **30**:156 [[Google Scholar](#)], [[Publisher](#)]
- [32]. Nardo M., Brunetta D.M., Vilar F.C., Machado A.A., De Santis G.C., Hemoglobin concentration increment is associated with a better prognosis in HIV patients with anemia', *International Journal of Infectious Diseases*, 2012, **16**:e703 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [33]. Tan F., Triyono E.A., Wahyunitisari M.R., Microbial Patterns of Hospitalized Hiv Positive Patients Admitted in Dr. Soetomo General Hospital, Surabaya Indonesia', *Journal of Vocational Health Studies*, 2021 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [34]. Klasinc R., Rieger A., Presterl E., Wrba T., Diab-Elschahawi M., Epidemiology of Urinary Tract Infections in HIV Positive Patients at a Tertiary Care University Hospital in Central Europe (2011 - 2016)', *Infectious Disorders-Drug Targets (Formerly Current Drug Targets-Infectious Disorders)*, 2018, **18**:199 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [35]. Danjuma Z.A., Asuquo A.E., Pokam T., Ibeneme E., Bacteriuric Profiles and CD4 Counts of Human Immunodeficient Virus (HIV) Seropositive Individuals Seeking Care in a Nigerian Health Facility Sample Collection and Processing Methods Study Subjects Analysis of Whole Blood Cells Ethical Clearance Statistic', *International Journal of Biomedical Laboratory Science*, 2019, **8**:1 [[Google Scholar](#)], [[Publisher](#)]
- [36]. Handayani K., Correlation of CD4 Count and Neutrophil- Lymphocyte Ratio in Human Immunodeficiency Virus-Acquired Immunodeficiency Syndrome (HIV- AIDS) Patients', *European Journal of Molecular & Clinical Medicine*, 2020, **7**:2020 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [37]. Akinbami A., Bode-Shojobi I., Ajibola S., Oshinaike O., Adediran A., Ojelabi O., Ismail K., Osikomaiya B., Prevalence of Asymptomatic Bacteriuria in HIV Infected Patients in a Tertiary Hospital in Lagos, Nigeria', *World Journal of AIDS*, 2013, **03**:105 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [38]. Omoregie R., Eghafona N.O., Urinary tract infection among asymptomatic HIV patients in Benin City, Nigeria', *British Journal of Biomedical Science*, 2009, **66**:190 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [39]. Widmer T.A., Theron G., Grove D., Prevalence and risks of asymptomatic bacteriuria among HIV-positive pregnant women', *Southern African Journal of Epidemiology and Infection*, 2010, **25**:28 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [40]. Ezechi O.C., Gab-Okafor C.V., Oladele D.A., Kalejaiye O.O., Oke B.O., Ekama S.O., Audu R.A., Okoye R.N., Ujah I.A., Prevalence and risk factors of asymptomatic bacteriuria among pregnant Nigerians infected with HIV', *The Journal of Maternal-Fetal and Neonatal Medicine*, 2013, **26**:402 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [41]. Abderrazzack A.F., Salou M., Patassi A., Yehadji D., Ameyapoh Y., Correlation between Asymptomatic Bacteriuria and HIV-1 Viral Load Level and CD4 Count in Pregnant Women on Antiretroviral Therapy in N'djamena (Chad)', *World Journal of AIDS*, 2015. **05**:308 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [42]. Olowe O., Ojo-Johnson B., Makanjuola O., Olowe R., Mabayoje V., Detection of bacteriuria among human immunodeficiency virus seropositive individuals in Osogbo, southwestern Nigeria', *European Journal of Microbiology and Immunology*, 2015, **5**:126 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]
- [43]. Záhorec R., Ratio of neutrophil to lymphocyte counts-rapid and simple parameter of systemic inflammation and stress in critically ill, *Bratislavske lekarske listy*, 2001, **102**:5 [[Google Scholar](#)], [[Publisher](#)]
- [44]. Wu L., Zou S., Wang C., Tan X., Yu M., 'Neutrophil-to-lymphocyte and platelet-to-lymphocyte ratio in Chinese Han population from Chaoshan region in South China', *BMC Cardiovascular Disorders*, 2019, **19**:1 [[Crossref](#)], [[Google Scholar](#)], [[Publisher](#)]

HOW TO CITE THIS ARTICLE

Nafdzu Makhmudatul Muna, Agung Dwi Wahyu Widodo, Pepy Dwi Endraswari, M. Vitanata Arfijanto. Correlation between the Bacterial and Fungal Profiles from the Clinical Specimens with the CD4 Counts and the NLR Values of HIV/AIDS Patients at Tertiary Referral Hospital in Indonesia. *J. Med. Chem. Sci.*, 2023, 6(9) 2111-2127

DOI: <https://doi.org/10.26655/JMCHEMSCI.2023.9.18>

URL: http://www.jmchemsci.com/article_169455.html