

# Prevalence And Susceptibility Profile Of Carbapenem- Resistant Pseudomonas Aeruginosa (CRPA) At Dr. Soetomo Public Hospital, Surabaya, From January To December 2021

*by* Muhammad Vitanata Arfijanto

---

**Submission date:** 08-May-2023 10:15AM (UTC+0800)

**Submission ID:** 2086997276

**File name:** tant\_Pseudomonas\_Aeruginosa\_CRPA\_At\_Dr.\_Soetomo\_Public\_Hospi.pdf (407.06K)

**Word count:** 5016

**Character count:** 27958

# Prevalence and susceptibility profile of carbapenem-resistant pseudomonas aeruginosa (CRPA) at Dr. Soetomo Public Hospital, Surabaya, from January to December 2021



Andritta Febriana<sup>1</sup>, Agung Dwi Widodo<sup>2\*</sup>, Muhammad Vitanata Arfijanto<sup>3</sup>

## ABSTRACT

**Introduction:** *Pseudomonas aeruginosa* is a pathogen that frequently causes healthcare-associated infections (HAI), which has also been associated with high mortality and morbidity rates. Carbapenems have been widely utilized as empiric therapy for *P. aeruginosa* infections since these organisms have intrinsic resistance to various antibiotics. Therefore, the high rate of CRPA infection became the reason for conducting this study to determine the prevalence and susceptibility pattern of CRPA in Dr. Soetomo Public Hospital, Surabaya, from January to December 2021.

**Method:** The researcher employed descriptive observational study from secondary data with the first isolate sample of *P. aeruginosa* per specimen per patient that has been identified by the BD Phoenix™ automated identification and susceptibility testing system from specimens of urine, blood, sterile fluid, pus, tissue, and sputum that are phenotypically resistant to meropenem or imipenem antibiotics, examined at the Clinical Microbiology Unit of Dr. Soetomo Public Hospital, Surabaya, from January to December 2021.

**Result:** Of the total *P. aeruginosa* isolates, 149 CRPA isolates were obtained. The researcher conveyed that the majority of the samples were male (59.1%); the most comorbid cases were diabetes mellitus with complications (22.4%) found in the intensive care unit (40.3%); the majority of specimens were from the respiratory tract (43%); the highest antibiotic susceptibility was amikacin (62.4%); the prevalence of CRPA in Dr. Soetomo Public Hospital, Surabaya, from January to December 2021 reached 21.25%.

**Conclusion:** In this study, CRPA isolates showed the highest sensitivity to amikacin, and the highest distribution of CRPA events was found in the intensive care unit.

**Keywords:** CRPA, incidence rate, susceptibility profile.

**Cite This Article:** Febriana, A., Widodo, A.D., Arfijanto, M.V. 2023. Prevalence and susceptibility profile of carbapenem-resistant pseudomonas aeruginosa (CRPA) at Dr. Soetomo Public Hospital, Surabaya, from January to December 2021. *Bali Medical Journal* 12(1): 571-576. DOI: 10.15562/bmj.v12i1.4098

<sup>1</sup>Clinical Microbiology Specialist program-1, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia;

<sup>2</sup>Department of Clinical Microbiology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Surabaya, Indonesia;

<sup>3</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Surabaya, Indonesia;

\*Corresponding author:

Agung Dwi Widodo;  
Department of Clinical Microbiology,  
Faculty of Medicine, Universitas  
Airlangga. Dr. Soetomo General  
Academic Hospital, Surabaya, Indonesia  
andritta.micro77@gmail.com

Received: 2022-11-08

Accepted: 2023-01-11

Published: 2023-02-04

## INTRODUCTION

The worldwide increase in infection with multidrug-resistant gram-negative (MDR) pathogens is a crucial global health problem. Southeast Asia has a high prevalence of MDR gram-negative bacteria, including multidrug-resistant non-fermenters such as *P. aeruginosa*.<sup>1,2</sup> *P. aeruginosa* is an opportunistic pathogen that requires minimal nutrition for its survival, enabling it to easily adapt to and survive in an environment with extreme conditions, such as a hospital environment.<sup>3,4</sup> These pathogens are commonly found on abiotic surfaces such as medical equipment, rooms, and showers and even can contaminate distilled water, and are widely isolated, especially in

immunocompromised patients, patients treated in the ICU, or patients with severe comorbidities.<sup>5,6</sup> Thus, *P. aeruginosa* is commonly linked to healthcare-associated infections.<sup>7,8</sup>

*P. aeruginosa* has an innate resistance to several antibiotics, such as ampicillin, amoxicillin, ampicillin-sulbactam, amoxicillin-clavulanic acid, cefotaxime, ceftriaxone, ertapenem, tetracycline, and tigecycline.<sup>9,10</sup> Carbapenem is a broad-spectrum  $\beta$ -lactam antibiotic with bacterial cell wall synthesis inhibitory activity. Carbapenem has extremely high potency against Gram-negative and Gram-positive bacteria among other  $\beta$ -lactam drugs. In addition, carbapenem is a last-line drug for treating cases of patients with MDR pathogenic infections, including *P.*

*aeruginosa*.<sup>7,11,12</sup> The cause of carbapenem-resistant *P. aeruginosa* (CRPA) is generally acquired due to a chromosomal mutation that causes loss or inactivation of the OprD (Outer membrane protein D) porin and/or overexpression of the efflux pump. Resistance to carbapenems can also be obtained through the acquisition of horizontal gene transfer (HGT) of the gene encoding carbapenemase. This B-lactamase is capable of hydrolyzing carbapenems and provides resistance to all  $\beta$ -lactam antibiotics.<sup>8,13,14</sup>

The increase in CRPA rates can be caused by the emergence of resistance in these organisms after exposure to carbapenems and can occur due to the spread of CRPA from patient to patient.<sup>15</sup> Resistance to these carbapenems,

especially when mediated by transferable carbapenemase-coding genes, can spread rapidly and cause outbreaks.<sup>11</sup> The increase in CRPA cases has become significant health and economic issue that deserves more attention given the limited options for an effective and safe therapy.<sup>1,16</sup> In addition, carbapenemase-producing CRPA strains cannot be treated with classical  $\beta$ -lactamase inhibitors, thus, it can lead to poor clinical conditions, such as an increased risk of mortality, morbidity, length of hospitalization, and increased treatment costs.<sup>5</sup>

HAI-related CRPA cases have increased globally in recent years.<sup>5,17</sup> In 2017, WHO classified CRPA as a priority pathogen of the 'critical pathogen' category for research and development of the latest antibiotics.<sup>18</sup> In Indonesia, the incidence of CRPA from the ICU of Dr Cipto Mangunkusumo Hospital reached 21.9% in 2011.<sup>19</sup> Based on the surveillance of antibiotic resistance in class A and B hospitals in Indonesia in 2020, the prevalence of CRPA amounted to 25% of all specimens and rooms from all hospitals.<sup>20</sup> Therefore, the high rate of CRPA infection became the reason for conducting this study to determine the prevalence and susceptibility pattern of CRPA in Dr. Soetomo Public Hospital, Surabaya, from January to December 2021.

## MATERIAL AND METHODS

The research sample included all data on *Pseudomonas aeruginosa* isolates identified by the BD Phoenix™ automated identification and susceptibility testing system from urine, blood, sterile fluid, sputum, pus, and tissue specimens that are phenotypically resistant to carbapenem antibiotics examined at the Clinical Microbiology Unit of Dr. Soetomo Public Hospital, Surabaya. The isolates taken were the first isolates from each type of

specimen in the same patient from January 1 to December 31, 2021. This research was approved by the Health Research Ethics Committee of Dr. Soetomo Public Hospital, Surabaya, based on Ethical Clearance No. 1411/122/4/IV/2022.

The data analyzed in this study included CRPA distribution by age and sex, CRPA distribution by comorbidities, CRPA distribution by the patient clinical outcome, CRPA distribution by care unit, CRPA distribution by the specimen, and CRPA antibiotic susceptibility pattern.

The data on the results of the antibiotic susceptibility test of carbapenem resistant *P. aeruginosa* (CRPA) isolates were stated in the Susceptible, Intermediate, and Resistant categories. The methods employed were disk diffusion for testing the antibiotic, such as fosfomycin, and the semi-automatic BD Phoenix™ automated identification and susceptibility testing system for testing antibiotics, such as gentamicin, amikacin, imipenem, meropenem, ceftazidime, cefepime, ciprofloxacin, levofloxacin, aztreonam, piperacillin-tazobactam, according to the group *P. aeruginosa* test at the Clinical and Laboratory Standards Institute (CLSI). The data obtained were presented in the form of tables containing frequency and percentage data for analysis.

## RESULTS

This research collected 701 data from culture and susceptibility test of *P. aeruginosa*, which was divided into 149 (21.25%) carbapenem-resistant *P. aeruginosa* (CRPA) and 552 carbapenem susceptible *P. aeruginosa* (CSPA), which derived from all specimens of blood, sterile fluid, urine, sputum, pus, and tissue recorded at the Clinical Microbiology unit of Dr. Soetomo Public Hospital, Surabaya. Based on the distribution data of the study sample, CRPA was most commonly found

in the male sex, reaching 88 (59.06%) people, while in the female sex, it reached 61 (40.93%) people. Meanwhile, based on the distribution of samples by age, the most CPRA was found in patients aged > 45 years, as many as 78/149 (52.34%), which were divided into middle age adults (54 patients or 36.2%) and seniors adult (24 patients or 16.1%), as presented in table 1.

The distribution of CRPA patients with comorbid factors based on the Charlson comorbidity index indicated that 107/149 (71.81%) patients had comorbidities and one patient could have more than one comorbid. The highest frequency of comorbidity was diabetes, which reached 42 (28.18%) cases, which was divided into 18 (12.1%) cases of diabetes without complications and 24 (16.1%) cases of diabetes with complications. The second most common comorbidities were Cerebrovascular disease and chronic lung disease, each with 12 (8.1%) cases (Table 2).

This study revealed that more than 70% of patients were estimated to have comorbidities, but no specific comorbidities were associated with an increased risk of CRPA incidence. The most comorbid in this study was diabetes, which reached 28.18% of cases, divided into diabetes without complications (12.1%) and diabetes with complications (16.1%). The clinical outcomes of CRPA patients in this study demonstrated that 88 (59.06%) patients were alive and 61 (40.93%) patients were deceased (Table 3). From all care units, the highest number of CRPA was found in intensive care units, including ICU, NICU, ROI, and HCU rooms, which reached 60 (40.3%) cases, followed by surgical care unit, amounting to 32 (21.5%) cases (Table 4).

The highest incidence of CRPA derived from respiratory tract specimens which

**Table 1. CRPA distribution by age and sex.**

Age (years)	Toddler (0-5)	Child (6-11)	Teenager (12-25)	Adult (26-45)	Middle Age Adult (46-65)	Senior Adult (> 65)	Total
Male	15 (17%)	5 (5.7%)	10 (11.4%)	14 (15.9%)	35 (39.8%)	9 (10.2%)	88 (59.06%)
Female	5 (8.2%)	0 (0.0%)	7 (11.5%)	15 (24.6%)	19 (31.1%)	15 (24.6%)	61 (40.93%)
<b>Total</b>	20 (13.4%)	5 (3.4%)	17 (11.4%)	29 (19.5%)	54 (36.2%)	24 (16.1%)	149 (100%)

reached 64 (43%) specimens, which consisted of 23 spontaneous sputum specimens, 40 endotracheal tube aspirates, and 1 bronchial lavage. CRPA isolates from pus specimens were 41 (27.5%) specimens, from urine specimens were 30 (20.8%) specimens, from blood specimens were 8 (5.4%) specimens, and from sterile fluid specimens were 5 (3.4%) specimens (Table 5).

The results of the antibiotic susceptibility test of all CRPA isolates demonstrated the highest susceptibility to *amikacin* as much as 93/149 (62.4%), followed by *piperacillin-tazobactam* at 57/149 (38.3%), *aztreonam* at 40/149 (26.8%), *gentamicin* at 38/148 (25.7%), *ceftazidime* at 31/148 (20.9%), *cefepime* at 28/148 (18.9%), *ciprofloxacin* (17.4%), *levofloxacin* (16.2%), *meropenem* (13.5%), *fosfomycin* (10.7%),

and *imipenem* (2.7%). The highest number of CRPA isolates resistant to *imipenem* antibiotics was 140/149 (94%) isolates. *Fosfomycin* revealed resistance of 122/140 (87.1%), *cefepime* and *levofloxacin* each of 120/148 (80.5%), *ciprofloxacin* of 119/149 (79.9%), *meropenem* of 117/148 (79.1%), and *gentamicin* of 106/148 (71.6%) (Figure 1).

## DISCUSSION

In this study, the prevalence of CRPA reached 21.25% which is in line with previous study which revealed the prevalence of CRPA from the ICU of Dr. Cipto Mangunkusumo Hospital in 2011 was 21.9%.<sup>21</sup> However, the prevalence of CRPA in this study was not restricted to the ICU. The results of this study were higher than the study conducted by Tartof et al. (2018) which demonstrated a prevalence of CRPA of 20.2% in hospitalized patients in 14 medical centers from 2011 to 2015.<sup>22</sup> Meanwhile, other study revealed the prevalence of HAIs caused by CRPA was 40%, however, this study did not distinguish between HAIs patients and non-HAI patients.<sup>19</sup>

Based on the aforementioned data, CRPA was mostly found in male patients (59.06%). These data are in accordance with a study conducted by Kang et al. (2003), which demonstrated that most patients with CRPA were male (66.2%).<sup>21</sup> Similarly, in their study, Tartof et al. (2018) also revealed that most patients with CRPA were male (59.4%).<sup>22</sup> In this study, patients over 45 were identified to have the highest prevalence of CRPA. Male patients from middle age adults to senior adults included in the category with higher rates of infection acquisition. This result is in line with other research findings that age has become a significant risk factor for *P. aeruginosa* infection and that nosocomial infection generally occurs in males.<sup>23,24</sup> Additionally, elderly patients are more susceptible to infection with multiresistant microorganisms. This is brought on by a decline in immune system activity, which also affects the body's structure and function, leading to other conditions such as the necessity for a catheter, malnutrition, and prolonged treatment.<sup>5,26,32</sup>

The most comorbid in this study was diabetes, in a study conducted by

**Table 2. CRPA distribution by comorbidity.**

Comorbidity	Total (n= 107)
Diabetes with complications	24 (22.4%)
Diabetes without complications	18 (16.8%)
Cerebrovascular disease	12 (11.2%)
Chronic lung disease	12 (11.2%)
Mild liver disease	11 (10.3%)
Malignant solid tumor	11 (10.3%)
Kidney illness	10 (9.3%)
Peripheral vascular disease	8 (7.5%)
Tumors with metastases	5 (4.7%)
Congestive heart failure	3 (2.8%)
Acute myocardial infarction	2 (1.9%)
HIV/AIDS	2 (1.9%)
Hemiplegia/paraplegia	2 (1.9%)
Dementia	1 (0.9%)
Leukemia	1 (0.9%)
Lymphoma	1 (0.9%)
Rheumatism	0 (0%)
Moderate to serious liver disease	0 (0%)
Peptic ulcer	0 (0%)

**Table 3. CRPA distribution by patient clinical outcome.**


Living Patient n (%)	Deceased patient n (%)	Total n (%)
88 (59.06%)	61 (40.93%)	149 (100%)

**Table 4. CRPA distribution by care unit.**

Care Units	CRPA n (%)
ICU, NICU, ROI, HCU	60 (40.3)
Surgical Care Unit	32 (21.5)
Medical Care Unit	27 (18.1)
Pediatric Care Unit	15 (10.1)
Special Isolation Room	5 (3.4)
Obstetrics and Gynaecology	4 (2.7)
Emergency Room	6 (4)
Total	149 (100)

**Table 5. CRPA distribution by specimen.**

Blood n (%)	Sterile fluid n (%)	Sputum n (%)	Urine n (%)	Pus and tissue n (%)	Total n (%)
8 (5.4)	5 (3.4)	64 (43)	31 (20.8)	41 (27.5)	149 (100)



**Figure 1.** CRPA antibiotic susceptibility pattern.

Palavutitoti (2018), diabetes was the third most common comorbid after hypertension and chronic kidney disease (CKD).<sup>7</sup> Patients with CRPA infection have a higher risk of receiving appropriate antimicrobial therapy late, resulting in prolonged hospitalization, increased risk of subsequent antibiotic-resistant infections, as well as increased morbidity and mortality.<sup>22,27</sup> The presence of virulence factors such as flagella, pili, and lipopolysaccharides as well as biologically active processes such as toxin secretion, biofilm formation, and quorum sensing has an impact on the outcome of *P. aeruginosa* infection.<sup>25</sup> In this study, the mortality rate reached 40.93%. Suarez et al., (2009) stated that CRPA was not associated with a higher mortality rate in hospitalized patients with bacteremia.<sup>28</sup>

The global incidence of nosocomial infections caused by *P. aeruginosa* was estimated at 10-15% and around 10-20% was obtained from intensive care units.<sup>29,30</sup> These research findings are in line with the research conducted by Tartof et al. (2018).<sup>22</sup> The high rate of cross-transmission in intensive care units (Zavascki et al., 2006) and the use of a variety of broad-spectrum antibiotics in the ICU (Alhussain, 2021) contributed significantly to the incidence of CRPA infection.<sup>23,31</sup> In addition, the use of invasive medical devices, such as the installation of ventilators and urinary catheters, are also at risk for infection

with CRPA. Patients in the intensive care unit are patients with severe diseases who frequently require invasive medical devices, which increases their risk of recurrent infections and complications.<sup>7</sup> Several studies have discovered that using invasive medical devices is a significant determinant of CRPA.<sup>22</sup> Patients on a ventilator have eleven times higher risk of infection with *P. aeruginosa* than those without a ventilator.<sup>23</sup>

In this study, the most CRPA specimens were obtained from respiratory tract samples, i.e., sputum by 43%, where this result is in line with research conducted by Zavascki et al. (2006) with 50.3% specimens<sup>31</sup>; Karuniawati (2011)<sup>19</sup> with 61/81 (75%) specimens; Tartof et al. (2018) with 61.45% specimens<sup>22</sup>; and Alhussain (2021) with 60% specimens<sup>14</sup>. In this regard, Hospital Acquired Pneumonia (HAP) and Ventilator-Associated Pneumonia (VAP) were the most common nosocomial infections.<sup>32</sup> *P. aeruginosa* was the most common cause of hospital pneumonia, the third most common cause of urinary tract infection, the fourth most common cause of surgical site infection, and the seventh most common pathogen isolated from blood.

*P. aeruginosa* has a strong tendency to become a pathogen that is resistant to various intrinsic and acquired antibiotics.<sup>8</sup> *P. aeruginosa* susceptibility decreased significantly to carbapenem, ceftazidime,

and ciprofloxacin (Alhussain, 2021).<sup>15</sup> Additionally, *P. aeruginosa* is able to survive for a long time even under unfavorable conditions. These bacteria have various virulence factors, one of which produces biofilms that play a major role in the occurrence of nosocomial infections and resistance to antibiotics.<sup>33,34</sup>

CRPA is often resistant to various drugs, causing difficulties in treating and controlling the infection.<sup>35</sup> In this study, CRPA resistance ranged from 22.8 - 80.5% with the highest susceptibility to amikacin (62.4%), while susceptibility to other antipseudomonal ranged from 2.7 - 38.3%. This study is in line with the research conducted by Alhussain et al. (2021) which indicated the highest susceptibility to amikacin at 86.7%.<sup>23</sup> Meanwhile, research conducted by Anggraini et al. (2018) showed that MDR-PA had the highest susceptibility to amikacin at 50.9%, lower than the susceptibility of amikacin in non-MDR-PA strains of 76.9%.<sup>36</sup> CRPA has more resistance than CSPA.<sup>22,35</sup> Amikacin is a semisynthetic aminoglycoside that is extremely resistant to enzyme modification, so many bacteria are sensitive to this antibiotic. A fairly good susceptibility to amikacin indicates that the use of amikacin is not too frequent considering that amikacin may cause the side effects of ototoxicity and nephrotoxicity.<sup>36</sup>

Colistin is the last line of therapy for CRPA multidrug treatment. The analytical study conducted by Goli et al. (2016) and Sheikh et al., (2019) revealed that colistin still has a good susceptibility value to MDR CRPA infection. However, these antibiotics are rarely used since they are nephrotoxic and neurotoxic.<sup>37,38</sup> Although the prevalence of resistance to colistin is still low, the incidence of resistance to colistin has been reported in various countries.<sup>37,38</sup> In this study, only three CRPA isolates were tested against colistin with 100% of the isolates being resistant. Due to the limited number, this statement cannot generally be concluded.

The increased incidence of imipenem resistance in *P. aeruginosa* is often associated with resistance to other antipseudomonal drugs. This statement is in line with the research conducted by Palavutitoti et al. (2018), which

demonstrated that among the CRPA strains, 53.5% were XDR-PA, and 24.8% were MDR-PA. Patients with the XDR-PA strain tended to receive inappropriate empiric therapy and as many as 78.6% of them received carbapenem monotherapy.<sup>7</sup>

CRPA is generally acquired as a result of chromosomal mutations leading to loss or inactivation of the OprD porin and/or overexpression of the efflux pump. Resistance to carbapenems can also be obtained through the acquisition of horizontal gene transfer (HGT) of the gene encoding carbapenemase. This  $\beta$ -lactamase is able to hydrolyze carbapenems and provides resistance to all classes of  $\beta$ -lactam antibiotics (Botelho et al., 2019).<sup>5</sup> Additionally, increased CRPA can be caused by selective pressure due to exposure to antibiotics and transmission through health workers or between patients. Thus, it requires efforts to control and prevent nosocomial infections as well as strict regulation and monitoring in the wise and rational use of antibiotics.<sup>15</sup> This study had several limitations, including (1) The sample size of this study was limited to 149 inpatients at one hospital. (2) this research was conducted in Surabaya, so the situation may be different in other countries. (3) This study did not routinely test susceptibility to colistin. (4) Clinical data, laboratory and duration of use of antibiotics were not included in this study.

## CONCLUSION

In this study, CRPA isolates showed the highest sensitivity to amikacin, and the highest distribution of CRPA events was found in the intensive care unit. The findings of this study are of paramount relevance for case identification, control measures, and optimal treatment strategies of *P. aeruginosa* infection in ICUs. The increasing prevalence of antimicrobial resistance among hospitalized patients continues to pose a challenge for practitioners.

## ACKNOWLEDGMENTS

We appreciate Dr. Soetomo General Academic Hospital and the Department of Microbiology, Faculty of Medicine, Airlangga University for providing all necessary support in this research.

16

## CONFLICT OF INTEREST

The authors declare that they have no financial or personal conflicts of interest in this work.

7

## AUTHOR CONTRIBUTION

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

## FUNDING

None.

## REFERENCES

1. Treatment Options for Carbapenem-resistant Gram-negative Bacterial Infections. Clin Infect Dis. 2019;69(7):565-575. doi: <https://doi.org/10.1093/cid/ciz830>.
2. Ansori ANM, Fadholy A, Hayaza S, Susilo RJK, Inayatillah B, Winarni D, Husen SA. A Review on Medicinal Properties of Mangosteen (*Garcinia mangostana* L.). Res J Pharm Technol. 2020; 13(2):974-982. doi: <https://doi.org/10.5958/0974-360X.2020.00182.1>.
3. Pachori P, Goyalwal R, Gandhi P. Emergence of antibiotic resistance *Pseudomonas aeruginosa* in intensive care unit; a critical review. Genes Dis. 2019;6(2):109-119. Published 2019 Apr 17. doi: <https://doi.org/10.1016/j.gendis.2019.04.001>.
4. Hisham A, Abbas, Fathy M, Serry, Eman M, EL-Masry. Synergic interaction between antibiotics and the artificial sweeteners xylitol and sorbitol against *Pseudomonas aeruginosa* biofilms. Asian J. Pharm. Res. 2012;2(4):129-131.
5. Tsao LH, Hsin CY, Liu HY, Chuang HC, Chen LY, Lee YJ. Risk factors for healthcare-associated infection caused by carbapenem-resistant *Pseudomonas aeruginosa*. J Microbiol Immunol Infect. 2018;51(3):359-366. doi: <https://doi.org/10.1016/j.jmii.2017.08.015>.
6. Kharisma VD, Ansori ANM, Jakhmola V, Rizky WC, Widyandana MH, Probojati RT, Murtadlo AAA, Rebezov M, Scherbakov P, Burkov P, Matrosova Y, Romanov A, Sihombing MAEM, Antonius Y, Zainul R. Multi-strain human papillomavirus (HPV) vaccine innovation via computational study: A mini review. Res J Pharm Technol. 2022; 15(8). doi: <https://doi.org/10.52711/0974-360X.2022.00638>.
7. Palavutitotai N, Jitmuang A, Tongchai S, Kiratisin P, Angkasekwinai N. Epidemiology and risk factors of extensively drug-resistant *Pseudomonas aeruginosa* infections. PLoS One. 2018;13(2):e0193431. doi: <https://doi.org/10.1371/journal.pone.0193431>.
8. Botelho J, Grosso F, Peixe L. Antibiotic resistance in *Pseudomonas aeruginosa* - Mechanisms, epidemiology and evolution. Drug Resist Updat. 2019;44:100640. doi: <https://doi.org/10.1016/j.drug.2019.07.002>.
9. Clinical and Laboratory Standards Institute. 'M100 Performance standards for antimicrobial susceptibility testing (32th Edition)'. Wayne,

PA: Clinical and laboratory standards institute. 2021.

10. Gupta A. Synthesis of Novel Nitro Substituted Benzothiazole Derivatives and Antibacterial activity against *Pseudomonas aeruginosa*. Research J. Pharm. and Tech. 2019;12(10):4663-4668. doi: <https://doi.org/10.5958/0974-360X.2019.00803.5>.
11. Meletis G. Carbapenem resistance: overview of the problem and future perspectives. Ther Adv Infect Dis. 2016;3(1):15-21. doi: <https://doi.org/10.1177/2049936115621709>.
12. Raheem HQ, Hussein EF, Rasheed AH. Class 1,2 Integron Genes Distribution in *Pseudomonas aeruginosa* Isolated from Clinical Specimens. Research Journal of Pharmacy and Technology. 2022;15(7):3165-8. doi: <https://doi.org/10.52711/0974-360X.2022.00529>.
13. AL-Shimmary SMH, Mohamed NS, Safaa A. S. Al-Qaysi, Almohaidi AMS. Phylogeny Analysis of gyrB Gene and 16S rRNA Genes of *Pseudomonas aeruginosa* Isolated from Iraqi Patients. Research Journal of Pharmacy and Technology. 2021; 14(5):2517-1. doi: <https://doi.org/10.52711/0974-360X.2021.00443>.
14. Al-Obaidi RD, Hussein O.M. Al-Dahmishi. Molecular Study of Secretion Systems Virulence Protein among *Pseudomonas aeruginosa* Isolated from different Clinical samples. Research Journal of Pharmacy and Technology 2022; 15(10):4577-3. doi: <https://doi.org/10.52711/0974-360X.2022.00768>.
15. Aloush V, Navon-Venezia S, Seigman-Igra Y, Cabili S, Carmeli Y. Multidrug-resistant *Pseudomonas aeruginosa*: risk factors and clinical impact. Antimicrob Agents Chemother. 2006;50(1):43-48. doi: <https://doi.org/10.1128/AAC.50.1.43-48.2006>.
16. Dantas RC, Ferreira ML, Gontijo-Filho PP, Ribas RM. *Pseudomonas aeruginosa* bacteraemia: independent risk factors for mortality and impact of resistance on outcome. J Med Microbiol. 2014;63(Pt 12):1679-1687. doi: <https://doi.org/10.1099/jmm.0.073262-0>.
17. Raman G, Avendano EE, Chan J, Merchant S, Puzniak L. Risk factors for hospitalized patients with resistant or multidrug-resistant *Pseudomonas aeruginosa* infections: a systematic review and meta-analysis. Antimicrob Resist Infect Control. 2018;7:79. Published 2018 Jul 4. doi: <https://doi.org/10.1186/s13756-018-0370-9>.
18. Horcajada JP, Montero M, Oliver A, et al. Epidemiology and Treatment of Multidrug-Resistant and Extensively Drug-Resistant *Pseudomonas aeruginosa* Infections. Clin Microbiol Rev. 2019;32(4):e00031-19. doi: <https://doi.org/10.1128/CMR.00031-19>.
19. Karuniawati A, Saharman YR, Lestari DC. Detection of carbapenemase encoding genes in Enterobacteriaceae, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii* isolated from patients at Intensive Care Unit Cipto Mangunkusumo Hospital in 2011. Acta Med Indones. 2013;45(2):101-106.
20. Anggraini D, Kuntaman K, Karuniawati A. Surveilans resistensi antibiotik rumah sakit kelas A dan B di Indonesia tahun 2020. Deepublish. 2021.

21. Kang CI, Kim SH, Kim HB, et al. Pseudomonas aeruginosa bacteremia: risk factors for mortality and influence of delayed receipt of effective antimicrobial therapy on clinical outcome. Clin Infect Dis. 2003;37(6):745-751. doi: <https://doi.org/10.1086/377200>.
22. Tartof SY, Kuntz JL, Chen LH, et al. Development and Assessment of Risk Scores for Carbapenem and Extensive  $\beta$ -Lactam Resistance Among Adult Hospitalized Patients With Pseudomonas aeruginosa Infection. JAMA Netw Open. 2018;1(6):e183927. doi: <https://doi.org/10.1001/jamanetworkopen.2018.3927>.
23. Alhussain FA, Yenugadhati N, Al Eidan FA, Al Johani S, Badri M. Risk factors, antimicrobial susceptibility pattern and patient outcomes of Pseudomonas aeruginosa infection: A matched case-control study. J Infect Public Health. 2021;14(1):152-157. doi: <https://doi.org/10.1016/j.jiph.2020.11.010>.
24. Salih NS, Yahya WI, Yousif Al-Labban HMY, Aljanaby AAJ. Schiff bases compounds prepared from Phenyl hydrazine as a starting material were Synthesized, Characterized, and their Biological activity was Investigated. Research Journal of Pharmacy and Technology. 2022;15(8):3595-8. doi: <https://doi.org/10.52711/0974-360X.2022.00602>.
25. Jeong SJ, Yoon SS, Bae IK, Jeong SH, Kim JM, Lee K. Risk factors for mortality in patients with bloodstream infections caused by carbapenem-resistant Pseudomonas aeruginosa: clinical impact of bacterial virulence and strains on outcome. Diagn Microbiol Infect Dis. 2014;80(2):130-135. doi: [10.1016/j.diagmicrobio.2014.07.003](https://doi.org/10.1016/j.diagmicrobio.2014.07.003).
26. Al-Hamamy HR, Salih WH, Al-Mudallal NH. Bacterial Isolates and Antibiotic Susceptibility of Ear Infections in Al-Kindy Teaching Hospital, Baghdad, Iraq. Research Journal of Pharmacy and Technology. 2021; 14(12):6503-6. doi: <https://doi.org/10.52711/0974-360X.2021.01124>.
27. Murugan T, Murugan M, Albino Wins J. Rhamnolipid Biosurfactants produced by Pseudomonas sp. MK5 and its efficacy on Pharmaceutical Application. Research J. Pharm. and Tech. 2017; 10(8): 2645-2649. doi: <https://doi.org/10.5958/0974-360X.2017.00470.X>.
28. Suárez C, Peña C, Gavalda L, et al. Influence of carbapenem resistance on mortality and the dynamics of mortality in Pseudomonas aeruginosa bloodstream infection. Int J Infect Dis. 2010;14 Suppl 3:e73-e78. doi: <https://doi.org/10.1016/j.ijid.2009.11.019>.
29. Strateva T, Yordanov D. Pseudomonas aeruginosa - a phenomenon of bacterial resistance. J Med Microbiol. 2009;58(Pt 9):1133-1148. doi: <https://doi.org/10.1099/jmm.0.009142-0>.
30. Biswal I, Arora BS, Kasana D, Neetushree. Incidence of multidrug resistant pseudomonas aeruginosa isolated from burn patients and environment of teaching institution. J Clin Diagn Res. 2014;8(5):DC26-DC29. doi: <https://doi.org/10.7860/JCDR/2014/7483.438>.
31. Zavascki AP, Barth AL, Gaspareto PB, et al. Risk factors for nosocomial infections due to Pseudomonas aeruginosa producing metallo-beta-lactamase in two tertiary-care teaching hospitals. J Antimicrob Chemother. 2006;58(4):882-885. doi: <https://doi.org/10.1093/jac/dkl327>.
32. Kumar ST, Yassin A, Bhowmick T, Dixit D. Recommendations From the 2016 Guidelines for the Management of Adults With Hospital-Acquired or Ventilator-Associated Pneumonia. P T. 2017;42(12):767-772.
33. Mahon CR, Lechman DC. Nonfermenting and miscellaneous gram-negative bacilli. Dalam Textbook of diagnostic microbiology 3th Edition. St. Louis, Missouri: Elsevier Health Sciences, Edisi 3. Elsevier. 2019:480-83.
34. Thi MTT, Wibowo D, Rehm BHA. Pseudomonas aeruginosa Biofilms. Int J Mol Sci. 2020 Nov 17;21(22):8671. doi: <https://doi.org/10.3390/ijms21228671>.
35. Onguru P, Erbay A, Bodur H, et al. Imipenem-resistant Pseudomonas aeruginosa: risk factors for nosocomial infections. J Korean Med Sci. 2008;23(6):982-987. doi: <https://doi.org/10.3346/jkms.2008.23.6.982>.
36. Anggraini D, Yulindra UG, Savira M, Djojogugito FA, Hidayat N. Prevalensi dan pola sensitivitas antimikroba multidrug resistant Pseudomonas aeruginosa di RSUD Arifin Achmad. Majalah kedokteran Bandung. 2018;50(1). doi: <https://doi.org/10.15395/mkb.v50n1.1150>.
37. Goli HR, Nahaei MR, Ahangarzadeh Rezaee M, Hasani A, Samadi Kafil H, Aghazadeh M. Emergence of colistin resistant Pseudomonas aeruginosa at Tabriz hospitals, Iran. Iran J Microbiol. 2016;8(1):62-69.
38. Farajzadeh Sheikh A, Shahin M, Shokoohzadeh L, Halaji M, Shahcheraghi F, Ghanbari F. Molecular epidemiology of colistin-resistant Pseudomonas aeruginosa producing NDM-1 from hospitalized patients in Iran. Iran J Basic Med Sci. 2019;22(1):38-42. doi: <https://doi.org/10.22038/ijbms.2018.29264.7096>.



This work is licensed under a Creative Commons Attribution

# Prevalence And Susceptibility Profile Of Carbapenem-Resistant Pseudomonas Aeruginosa (CRPA) At Dr. Soetomo Public Hospital, Surabaya, From January To December 2021

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

13%

PUBLICATIONS

5%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Submitted to Universitas Airlangga 1 %  
Student Paper
- 2 [journal.unhas.ac.id](http://journal.unhas.ac.id) 1 %  
Internet Source
- 3 Su Jin Jeong, Sang Sun Yoon, Il Kwon Bae, Seok Hoon Jeong, June Myung Kim, Kyungwon Lee. "Risk factors for mortality in patients with bloodstream infections caused by carbapenem-resistant Pseudomonas aeruginosa: clinical impact of bacterial virulence and strains on outcome", Diagnostic Microbiology and Infectious Disease, 2014 1 %  
Publication
- 4 Submitted to Monash University 1 %  
Student Paper
- 5 Yanyan Hu, Yan Qing, Jiawei Chen, Congcong Liu, Jiayue Lu, Qi Wang, Shufang Zhen, Hongwei Zhou, Ling Huang, Rong Zhang. "Prevalence, Risk Factors, and Molecular



# Epidemiology of Intestinal Carbapenem-Resistant *Pseudomonas aeruginosa*", Microbiology Spectrum, 2021

Publication

---

6	<a href="http://repub.eur.nl">repub.eur.nl</a> Internet Source	1 %
7	<a href="http://microbiologyjournal.org">microbiologyjournal.org</a> Internet Source	1 %
8	<a href="http://sciencescholar.us">sciencescholar.us</a> Internet Source	1 %
9	<a href="http://www.scilit.net">www.scilit.net</a> Internet Source	1 %
10	Submitted to Universitas Sebelas Maret Student Paper	1 %
11	<a href="http://e-journal.unair.ac.id">e-journal.unair.ac.id</a> Internet Source	1 %
12	<a href="http://journals.plos.org">journals.plos.org</a> Internet Source	1 %
13	<a href="http://www.spandidos-publications.com">www.spandidos-publications.com</a> Internet Source	<1 %
14	"13th European Congress of Clinical Microbiology and Infectious Diseases", Clinical Microbiology and Infection, 2003 Publication	<1 %
15	<a href="http://aricjournal.biomedcentral.com">aricjournal.biomedcentral.com</a> Internet Source	

---

<1 %

16

[www.ijsrp.org](http://www.ijsrp.org)

Internet Source

<1 %

17

[downloads.hindawi.com](http://downloads.hindawi.com)

Internet Source

<1 %

18

[journals.lww.com](http://journals.lww.com)

Internet Source

<1 %

19

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

20

Submitted to Syiah Kuala University

Student Paper

<1 %

21

Submitted to Badan PPSDM Kesehatan  
Kementerian Kesehatan

Student Paper

<1 %

22

E Al Otaibi Fawzia, Fuad Al Hulaily Fatma.  
"Imipenem-resistant Pseudomonas aeruginosa:  
Epidemiology and susceptibility patterns at a  
Teaching Hospital in Riyadh, Saudi Arabia",  
African Journal of Microbiology Research,  
2012

Publication

<1 %

23

João Botelho, Filipa Grosso, Luísa Peixe.  
"Antibiotic resistance in Pseudomonas  
aeruginosa – Mechanisms, epidemiology and  
evolution", Drug Resistance Updates, 2019

<1 %

24

[mail.phcogj.com](mailto:mail.phcogj.com)

Internet Source

<1 %

---

25

[link.springer.com](http://link.springer.com)

Internet Source

<1 %

---

26

A. Arhin. "The Outer Membrane Protein OprQ and Adherence of *Pseudomonas aeruginosa* to Human Fibronectin", *Microbiology*, 01/21/2010

Publication

<1 %

---

27

Laksmi Wulandari, Gatot Soegiarto, Anna Febriani, Farah Fatmawati, Sahrin.

"Comparison of Detection of Epidermal Growth Factor Receptor (EGFR) Gene Mutation in Peripheral Blood Plasma (Liquid Biopsy) with Cytological Specimens in Lung Adenocarcinoma Patients", *Indian Journal of Surgical Oncology*, 2020

Publication

<1 %

---

28

Li-Hsin Tsao, Chih-Yi Hsin, Hsin-Yi Liu, Han-Chuan Chuang, Li-Yuan Chen, Yuarn-Jang Lee.

"Risk factors for Healthcare-associated Infection caused by carbapenem-resistant *Pseudomonas aeruginosa*", *Journal of Microbiology, Immunology and Infection*, 2017

Publication

<1 %

---

29

Zavascki, A.P.. "Risk factors for imipenem-resistant *Pseudomonas aeruginosa*: a comparative analysis of two case-control studies in hospitalized patients", *Journal of Hospital Infection*, 200502

Publication

&lt;1 %

30

[www.jidmr.com](http://www.jidmr.com)

Internet Source

&lt;1 %

31

Dewi Anggraini, Dewi Santosaningsih, Pepy Dwi Endraswari, Novira Jasmin, Fajri Marindra Siregar, Usman Hadi, Kuntaman Kuntaman. "Multicenter Study of the Risk Factors and Outcomes of Bloodstream Infections Caused by Carbapenem-Non-Susceptible *Acinetobacter baumannii* in Indonesia", *Tropical Medicine and Infectious Disease*, 2022

Publication

&lt;1 %

32

Luma I Al-Allaf, Rana KA Attarbashee, Jawnaa K Mammdoh. "THE EFFECT OF CYCLOPHOSPHAMIDE ON HIPPOCAMPAL STRUCTURE OF ADULT MALE RATS (ROLE OF ROSUVASTATIN)", *Military Medical Science Letters*, 2022

Publication

&lt;1 %

33

Raziyeh Ramazani, Rabeeh Izadi Amoli, Mojtaba Taghizadeh Armaki, Abazar Pournajaf, Hami Kaboosi. "A molecular New

&lt;1 %

# Update on the Biofilm Production and Carbapenem Resistance Mechanisms in Clinical *Pseudomonas aeruginosa* Isolates", Iranian Journal of Medical Microbiology, 2022

Publication

---

34	<a href="https://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
35	<a href="https://hdl.handle.net">hdl.handle.net</a> Internet Source	<1 %
36	<a href="https://inabj.org">inabj.org</a> Internet Source	<1 %
37	<a href="https://lume.ufrgs.br">lume.ufrgs.br</a> Internet Source	<1 %
38	<a href="https://lup.lub.lu.se">lup.lub.lu.se</a> Internet Source	<1 %
39	<a href="https://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a> Internet Source	<1 %
40	<a href="https://www.frontiersin.org">www.frontiersin.org</a> Internet Source	<1 %
41	<a href="https://www.scielo.br">www.scielo.br</a> Internet Source	<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Prevalence And Susceptibility Profile Of Carbapenem-Resistant Pseudomonas Aeruginosa (CRPA) At Dr. Soetomo Public Hospital, Surabaya, From January To December 2021

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---