

Height-for-age in Children under 5 Years Old with Down Syndrome and Hypothyroidism

by Nur Rochmah

Submission date: 14-Mar-2023 11:58AM (UTC+0800)

Submission ID: 2036745139

File name: dren_under_5_Years_Old_with_Down_Syndrome_and_Hypothyroidism.pdf (386.25K)

Word count: 3283

Character count: 17241

2 Height-for-age in Children under 5 Years Old with Down Syndrome and Hypothyroidism

Muhammad Faizi^{1,2}, Nur Rochmah^{1,2}, Yuni Hisbiyah^{1,2}, Anang Endaryanto^{1,2}, Soetjipto^{1,3}

¹Doctoral Program of Medical Science, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia; ²Researcher, Department of Child Health, Dr. Soetomo General Hospital, Faculty of Medicine Universitas Airlangga, Surabaya, East Java, Indonesia; ³Professor, Department of Medical Biochemistry, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

Abstract

Background: Children with Down syndrome tend to have accompanying comorbidities, such as hypothyroidism, and late detection of this ailment leads to reduced growth of the child. This study aimed to assess the growth patterns in children with Down Syndrome and hypothyroidism at diagnosis.

²⁰**Methods:** This cross-sectional study was conducted from September to December 2020 with 56 subjects at the Pediatric Endocrine Outpatient Clinic of Dr. Soetomo Hospital. Diagnosis of Down Syndrome was confirmed by karyotyping, while the weights and heights were assessed using a standardized curve for children with Down Syndrome and then calculated using Peditools. Fifty-six children with Down Syndrome and hypothyroidism were included, comprising 32 boys and 24 girls (mean age, 37.75 ± 34.26 months). Majority of the subjects had normal weight, height, and Body Mass Index (36/56 [mean z-score, -1.62 ± 2.36], 33/56 [mean height-z-score, -0.43 ± 2.74], and 30/56 [mean z-score, -2.00 ± 2.06], respectively). Furthermore, the mean onset of diagnosis was 17.07 ± 32.23 months, where 23 out of the 56 children had short stature and had gotten diagnosed at over 12 months of age.

²**Conclusion:** From the results obtained, hypothyroidism can be said to be associated with reduced growth in children with DS.

Key Words: down syndrome, growth, short stature, hypothyroidism

Introduction

⁵Short stature is one of the most recognized features in children with Down syndrome (DS), and in a healthy population of children with this condition, 35.7% were found to have short stature¹⁻³.

⁵Consequently, several adapted growth charts are used to assess the growth of children with DS⁴⁻⁷. The prevalence of short stature is increased by 46.2% in children with DS and hypothyroidism compared to those without hypothyroidism⁸. Children with DS experience late growth until 12–15 years, when they are approximately 20 cm shorter than children at the same age but without the condition⁹. Short stature is one of many complications caused by hypothyroidism, which is, as a result, often suspected if it occurs in adolescence. It should be suspected in children with short stature, as a study conducted in India showed

Corresponding author:

Muhammad Faizi

Mayjend Prof. Dr. Moestopo No. 6-8, Surabaya, East Java, Indonesia, 60286

Tel +628123402094,

mail : muhammad.faizi@fk.unair.ac.id

that almost 50% of these had hypothyroidism ¹⁰.

Hypothyroidism is a common comorbidity accompanying DS, and previous studies show that the incidence in people with this condition is 12 to 22 times that of individuals without ¹¹. Neonates suspected of having DS often demonstrate an increase ⁵ in plasma thyroid-stimulating hormone (TSH), and hypothyroidism is found in 4%–8% of this population ¹². Thyroid hormone is associated with cognitive development in the central nervous system during childhood, and any abnormality in its function may interrupt brain development, and consequently, general growth and development ^{10,11}. Consequently, hypothyroidism is believed to impair the growth of children with DS and increase their chances of having short stature. Although the detection of congenital hypothyroidism should be performed for every baby born ¹³, it is different in babies with DS. A study stated that autoimmune hypothyroidism occurs at five (5) months, and many were diagnosed at eight (8) years ¹⁴. Hence, detecting congenital hypothyroidism in babies with DS can happen after a few years ¹⁵.

Research on DS with hypothyroidism has not been widely studied in Indonesia, especially regarding the growth of children, although growth studies have been performed on subjects with Turner syndrome ¹⁶. Therefore, this study aims to assess how hypothyroidism impacts the growth pattern of children with DS.

A greater understanding of the growth pattern of children with DS and hypothyroidism could lead to earlier diagnosis and intervention. Early intervention has proven to offer good results in preventing and treating growth disorders in patients with these conditions ¹⁷. Indeed, current guidelines suggest evaluating thyroid function in DS just after birth and every six (6) months afterward to detect problems earlier ¹⁸. Moreover, the high incidence of short stature in patients with hypothyroidism has been shown to improve with earlier diagnosis and appropriate therapy ¹⁷.

Materials and Methods

¹ This study used a cross-sectional design and was performed from September to December 2020. The data were obtained from the patients' medical records, and the subjects were interviewed to collect hypothyroid history and nutritional status. This process was performed during their visit to the Pediatric Endocrine Outpatient Clinic of Dr. Soetomo Hospital.

Sample

This study involved children diagnosed with both DS and hypothyroidism and registered at the Pediatric Endocrine Outpatient Clinic of Dr. Soetomo Hospital. The inclusion criteria were patients aged 1–5 years, diagnosed with both conditions, while critically ill patients were excluded. Fifty-six (56) patients between 1 and 5 years of age (32 boys and 24 girls) were the subjects in this study. The weight-for-age, height-for-age, body mass index (BMI)-for-age, and BMI were presented.

Down syndrome

The DS diagnosis was confirmed physically by referring to the Fried *et al.* ¹⁹ criteria, which evaluated the abundant neck skin, downturned mouth corners, general hypotonia, and flat face. Other indications are dysplastic ear, epicanthic eye-fold, the gap between the first and second toes, and protruding tongue. This diagnosis was confirmed by a karyotyping test.

Instrument / Nutritional status

⁹ The nutritional status of children with DS was classified by Zemel *et al.* ² after recalculation of the standard growth curve of children with the condition. Using the Peditools.org website, the z-score of each growth parameter was calculated. The instrument, peditools.org/downinfant/, was used for children aged 0–2 years, while peditools.org/downpedi/ was used for those older than 2 years. Then, the height-for-age was classified either as short stature or normal based on the DS curve and z-score classification by Peditools.

According to WHO child growth standards, 2006, a z-score of -3.00 is severely stunted, -3.00 to -2.01 is moderately stunted, while -2.00 to 1.01 is mild.

Evaluation of data

A descriptive test was used to analyze the data using SPSS 17.0 software (IBM SPSS), while the mean and standard deviation of each element was used for the baseline and clinical characteristics. The Kolmogorov-Smirnov test was also used to evaluate the normality of each dataset.

Ethical permission

This study received ethical approval from the Ethics Committee overseeing health research at Dr. Soetomo Hospital (Ref. No.: 1960/KEKP/IV/2020).

Results and Discussion

This cross-sectional study included 56 children with DS and hypothyroidism, comprising 32 boys and 24 girls, at 57.1% and 42.9%, respectively, with an average age of 37.75 ± 34.26 months old. The baseline characteristics and history of these patients are presented in Tables 1 and 2. Most of the subjects had normal weights, heights, and BMI (36/56 [mean z-score, -1.62 ± 2.36], 33/56 [mean height-z-score, -0.43 ± 2.74], and 30/56 [mean z-score, -2.00 ± 2.06], respectively). Based on the nutritional status, 20 of the 56 subjects, at 35.7%, were revealed to be underweight, while 23, at 41.1%, had short statures. The mean duration of hypothyroid illness was 22.97 ± 23.24 months, while the mean onset age was 21.65 ± 39.76 months. Finally, the mean z-score of all criteria was within the normal limit of -2 < z < 2.

Table I. Frequency of subject characteristics

Characteristic	N (%)
Sex	
- Boy	32 (57.1)
- Girl	24 (42.9)
Weight-for-age	
- Severely underweight	12 (21.4)
- Underweight	8 (14.3)
- Normal	36 (64.3)
Height-for-age	
- Short stature	23 (41.1)
- Normal	33 (58.9)
BMI-for-age	
- Severely wasted	11 (19.6)
- Wasted	15 (26.8)
- Normal	30 (53.6)

Table II. Baseline characteristics of subjects

Characteristic	Mean (SD)	Minimum	Maximum	N
Age (months)	47.75 (34.23)	2	129	56
Illness duration (months)	23.25 (23.63)	0	90	56
Onset age (months)	17.07 (32.23)	0	163	56
¹² Weight-for-age z-score	-1.62 (2.36)	-10.13	1.66	56
Height-for-age z-score	-0.43 (2.74)	-5.66	4.88	56
BMI-for-age z-score	-2.00 (2.06)	-9.68	0.80	56

Based on the results, 23, i.e., 41.1% out of the 56 children with DS and hypothyroidism, had short statures, while the remaining 33, at 58.9%, were classified as normal. This finding is consistent with the report by Nam *et al.*, which concluded that infants with DS are likely to have short statures, especially when accompanied by comorbidities, such as a history of infectious or non-infectious diseases²⁰. Adequate and complete nutrient intake is also a strong predictor of nutritional status, including height-for-age in infants²¹. Furthermore, a study by the Leung brothers had similar findings to this investigation that hypothyroidism can stunt growth and cause short stature in the absence of immediate therapy to improve the growth gap²². Growth failure in the thyroid may be related to the need for cartilage maturation to T3²³. Histologically, individuals with chronic hypothyroidism will demonstrate a delay in bone maturation and resting on the epiphysial plate²⁴.

¹ As seen in Table 2, the mean age at diagnosis of hypothyroidism was 17.07 months, while some subjects were diagnosed as late as 20–23 months old. This is a cause for concern, as delayed diagnosis causes physical and intellectual disorders^{13,25}, and substitution therapy can improve hypothyroidism symptoms⁸. A study conducted in Turkey reported the average age of diagnosis to be 49 months²⁶. Early diagnosis is crucial to prevent sequelae, as although a

delay in diagnosis can lead to poor growth, it can be reversed⁸, unlike intellectual delays²⁷.

Zemel *et al.* conducted an epidemiological study on nutritional status and DS in the United States². It showed that the mean z-scores of height-for-age (-1.7 ± 1.2), weight-for-age (-0.8 ± 1.2), Head Circumference (HC)-for-age (-1.6 ± 1.0), and BMI-for-age (0.9 ± 1.0) were accompanied by a BMI value of 21.1 ± 5.8 . These values are similar to our study, although only individuals with DS and hypothyroidism were included. However, several studies reported no decrease in the z-score of the nutritional status as a result of hypothyroidism²⁸. The hypothyroid condition is a hypometabolic state in which the basal metabolic rate (BMR) decreases^{29,30}, hence, some individuals with hypothyroidism experience excess nutrients rather than a lack³¹. Also, the BMR in healthy individuals and those with DS devoid of other comorbidities are similar, meaning that DS does not affect BMR³².

¹¹ Concerning the nutritional status of the patients in this study, 35.7% were underweight and severely underweight, compared with the research by Al-Fahham *et al.*, where the prevalence in the DS group was only 14.3%³. However, this previous research also found congenital heart disease (CHD) in DS patients, a common comorbidity in this condition, similar to

this study, where 25% of the sample also had CHD¹⁵. Conversely, Al-Fahham found that the incidence of underweight subjects in the CHD group at 34.2%, which was greater than that observed in this study. Hypothyroidism is a chronic disease that can affect an individual's nutritional status, especially during growth and development⁸. A study from Al-Aaraj *et al.* showed an improvement in the nutritional status of individuals with DS undergoing hypothyroidism therapy at height-for-age and BMI-for-age⁸.

The main limitations of the current study were that it was single-centered and included only a small proportion of children with DS and hypothyroidism that had short stature, meaning more research of similar samples is required to confirm this relationship. Also, it did not compare a population of children with DS without hypothyroidism. This should be considered in prospective studies, as it would assist in drawing conclusions about the association between both conditions. Also, it will be important in future studies to perform early screening of hypothyroidism when examining the incidence of short stature in a similar population of children. If possible, the Indonesian multi-center epidemiological research will be performed to obtain more representative results on these conditions.

Conclusion and Acknowledgement

Hypothyroidism, as a comorbidity, has been shown to be associated with reduced growth of subjects with DS. Although late diagnosis is common, earlier detection could prevent these issues before they worsen, making the evaluation of the thyroid hormone status of patients born with suspected DS important. Finally, adolescents with short stature should undergo testing for hypothyroidism, and education should be provided to the community to create awareness of these associations and their consequences if left undiagnosed.

We thank the Director of Dr. Soetomo General Hospital who has given the permission to conduct this research.

3
Conflict of Interest: The author reports no conflicts of interest in this work

Source of Funding: This research has no funding sources

References

1. Aburawi EH, Nagelkerke N, Deeb A, Abdulla S, Abdulrazzaq YM. National growth charts for United Arab Emirates children with Down syndrome from birth to 15 years of age. *J Epidemiol.* 2015;25(1):20–9.
2. Zemel BS, Papan M, Stallings VA, Hall W, Schadt K, Freedman DS, et al. Growth charts for children with down syndrome in the United States. *Pediatrics.* 2015;136(5):e1204–11.
3. Al-Fahham M, Kotby A, Elabd H, Tawfik A, Ali H. Nutritional assessment of children with down syndrome with and without associated congenital heart defects. *J Pediatr Gastroenterol Nutr.* 2018;66:980.
4. Cronk C, Crocker AC, Pueschel SM, Shea AM, Zackai E, Pickens G, et al. Growth charts for children with Down syndrome: 1 month to 18 years of age. *Pediatrics.* 1988;81(1):102–10.
5. van Gameren-Oosterom HBM, van Dommelen P, Oudesluis-Murphy AM, Buitendijk SE, van Buuren S, van Wouwe JP. Healthy growth in children with down syndrome. *PLoS One.* 2012;7(2):1–8.
6. Piro E, Pennino C, Cammarata M, Corsello G, Greci A, Giudice CL, et al. Growth charts of Down syndrome in Sicily: Evaluation of 382 children 0-14 years of age. *Am J Med Genet.* 1990;70(SUPPL. 7):66–70.
7. Fernandes A, Mourato AP, Xavier MJ, Andrade D, Fernandes C, Palha M. Characterisation of the somatic evolution of Portuguese children with Trisomy 21--preliminary results. *Downs Syndr Res Pract.* 2001;6(3):134–8.
8. AlAaraj N, Soliman AT, Itani M, Khalil A, De Sanctis V. Prevalence of thyroid dysfunctions in

- infants and children with Down Syndrome (DS) and the effect of thyroxine treatment on linear growth and weight gain in treated subjects versus DS subjects with normal thyroid function: a controlled study. *Acta Biomed.* 2019;90(8S):36–42.
9. Bertapelli F, Agiovlasis S, Machado MR, Roso R do V, Guerra-Junior G. Growth charts for Brazilian children with Down syndrome: Birth to 20 years of age. *J Epidemiol [Internet]*. 2017;27(6):265–73. Available from: <http://dx.doi.org/10.1016/j.je.2016.06.009>
 10. Virmani A, Menon PSN, Kamarkar MG, Kochupillai N, Seth V, Ghai OP, et al. Evaluation of thyroid function in children with undiagnosed short stature in north India. *Ann Trop Paediatr.* 1987;7(3):205–9.
 11. Uppal H, Chandran S, Potluri R. Risk factors for mortality in Down syndrome. *J Intellect Disabil Res.* 2015;59(9):873–81.
 12. Bull MJ, Saal HM, Braddock SR, Enns GM, Gruen JR, Perrin JM, et al. Clinical report - Health supervision for children with Down syndrome. *Pediatrics.* 2011;128(2):393–406.
 13. Seth A, Aggarwal V, Maheshwari A. Hypothyroidism in children beyond 5 y of age: Delayed diagnosis of congenital hypothyroidism. *Indian J Pediatr.* 2012;79(7):891–5.
 14. Shalitin S, Phillip M. Autoimmune thyroiditis in infants with Down's syndrome. *J Pediatr Endocrinol Metab.* 2002;15(5):649–52.
 15. Graber E, Chacko E, Regelman MO, Costin G, Rapaport R. Down Syndrome and Thyroid Function. *Endocrinol Metab Clin North Am [Internet]*. 2012;41(4):735–45. Available from: <http://dx.doi.org/10.1016/j.ecl.2012.08.008>
 16. Novina N, Gunardi H, Pulungan AB. Height and Upper/Lower Body Ratio in Turner Syndrome Adolescents in Indonesia; Is There any Significant Difference Based on Karyotype? In: 58th Annual ESPE meeting. Vienna, Austria: European Society for Paediatric Endocrinology; 2019.
 17. Ranke MB, Schwarze CP, Mohnike K, Von Mühlendahl KE, Keller E, Willgerodt H, et al. Catch-up growth after childhood-onset substitution in primary hypothyroidism: Is it a guide towards optimal growth hormone treatment in idiopathic growth hormone deficiency? *Horm Res.* 1998;50(5):264–70.
 18. Hardy O, Worley G, Lee MM, Chaing S, Mackey J, Crissman B, et al. Hypothyroidism in Down syndrome: Screening guidelines and testing methodology. *Am J Med Genet.* 2004;124A(4):436–7.
 19. FRIED K. a Score Based on Eight Signs in the Diagnosis of Down Syndrome in the Newborn. *J Intellect Disabil Res.* 1980;24(3):181–5.
 20. Nam JY, Choi Y, Jung MK, Shin J, Cho KH, Kim W, et al. The effect of neonatal hypothyroidism and low family income on intellectual disability: A population-based cohort study. *PLoS One.* 2018;13(11):1–13.
 21. Lutter CK, Rivera JA. Nutritional status of infants and young children and characteristics of their diets. In: *Journal of Nutrition.* 2003.
 22. Leung AKC, Leung AAC. Evaluation and management of the child with hypothyroidism. *World J Pediatr [Internet]*. 2019;15(2):124–34. Available from: <https://doi.org/10.1007/s12519-019-00230-w>
 23. Williams GR. Thyroid Hormone Actions in Cartilage and Bone. *Eur Thyroid J.* 2012;3–13.
 24. Boersma B, Otten BJ, Stoelinga GBA, Wit JM. Catch-up growth after prolonged hypothyroidism. *Eur J Pediatr.* 1996;155(5):362–7.
 25. Pulungan AB, Oldenkamp ME, van Trotsenburg ASP, Windarti W, Gunardi H. Effect of delayed diagnosis and treatment of congenital hypothyroidism on intelligence and quality of life: An observational study. *Med J Indones.* 2019;28(4):396–401.

26. Tarim OF, Yordam N. Congenital hypothyroidism in Turkey: a retrospective evaluation of 1000 cases. *Turk J Pediatr.* 1992;34(4):197–202.
27. Seo MK, Yoon JS, So CH, Lee HS, Hwang JS. Intellectual development in preschool children with early treated congenital hypothyroidism. *Ann Pediatr Endocrinol Metab.* 2017;22(2):102.
28. Zafar Y, Suddaby SR, Shafiq M. Hypothyroidism Manifesting as a Combination of Ascites and Malnutrition Requiring Total Parenteral Nutrition: A Unique Presentation. *Cureus.* 2019;11(8):5338.
29. Mullur R, Liu YY, Brent GA. Thyroid hormone regulation of metabolism. *Physiol Rev.* 2014;94(2):355–82.
30. Johansen K, Hansen JM, Skovsted L. The Preferential Role of Triiodothyronine in the Regulation of Basal Metabolic Rate in Hyper- and Hypothyroidism. *Acta Med Scand.* 1978;204(1–6):357–9.
31. Ghizlane EM, Bouznad N, El AN. Effects of hypothyroidism on nutritional status: which impact? *Endocr Abstr.* 2017;
32. SCHAPIRO MB, RAPOPORT SI. Basal metabolic rate in healthy Down's syndrome adults. *J Intellect Disabil Res.* 1989;33(3):211–9.

Height-for-age in Children under 5 Years Old with Down Syndrome and Hypothyroidism

ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

15%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.researchgate.net Internet Source	2%
2	www.semanticscholar.org Internet Source	2%
3	www.ijphrd.com Internet Source	1%
4	ejournal.poltekkes-smg.ac.id Internet Source	1%
5	www.frontiersin.org Internet Source	1%
6	scite.ai Internet Source	1%
7	pr.hec.gov.pk Internet Source	1%
8	"Handbook of Growth and Growth Monitoring in Health and Disease", Springer Nature, 2012 Publication	1%

9	"6th World Congress of PGHAN: Abstracts", Journal of Pediatric Gastroenterology & Nutrition, 2021 Publication	1 %
10	Fandy Wicaksono, Yudhistira Pradnyan Kloping, Fikri Rizaldi, Doddy Moesbadianto Soebadi. "Ejaculatory duct cyst causing bilateral obstruction and subsequent infertility: A rare case report from Indonesia", Radiology Case Reports, 2022 Publication	1 %
11	worldwidescience.org Internet Source	1 %
12	dro.deakin.edu.au Internet Source	1 %
13	jehp.net Internet Source	1 %
14	"Abstracts", Hormone Research in Paediatrics, 2019 Publication	1 %
15	synapse.koreamed.org Internet Source	1 %
16	www.ijthyroid.org Internet Source	1 %
17	www.ijpediatrics.com Internet Source	<1 %

18

d197for5662m48.cloudfront.net

Internet Source

<1 %

19

van Dommelen, P., and S. van Buuren.
"Methods to obtain referral criteria in growth
monitoring", *Statistical Methods in Medical
Research*, 2013.

Publication

<1 %

20

www.e-mjm.org

Internet Source

<1 %

21

www.repositorio.unicamp.br

Internet Source

<1 %

22

Harvinder Kaur, Anil Kumar Bhalla, Inusha
Panigrahi. "Percentile Charts for Body Mass
Index of Indian Down Syndrome Children",
Journal of Pediatric Genetics, 2021

Publication

<1 %

23

Adi Wasis Prakosa, Muhammad
Miftahussurur, Juniastuti Juniastuti, Langgeng
Agung Waskito et al. "Characterization of
Helicobacter pylori tlyA and Its Association
with Bacterial Density", *Digestive Diseases*,
2021

Publication

<1 %

24

Eleni Magdalini Kyritsi, Christina Kanaka-
Gantenbein. "Autoimmune Thyroid Disease in
Specific Genetic Syndromes in Childhood and

<1 %

Adolescence", Frontiers in Endocrinology, 2020

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Height-for-age in Children under 5 Years Old with Down Syndrome and Hypothyroidism

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
