

Effectiveness of Health Education toward Healthcare Knowledge Improvement about Congenital Hypothyroidism Newborn Screening

by Nur Rochmah

Submission date: 11-Apr-2023 04:27PM (UTC+0800)

Submission ID: 2061404940

File name: mprovement_about_Congenital_Hypothyroidism_Newborn_Screening.pdf (303.63K)

Word count: 2589

Character count: 13552



Effectiveness of Health Education toward Healthcare Knowledge Improvement about Congenital Hypothyroidism Newborn Screening

Muhammad Faizi¹, Nur Rochmah¹, Yuni Hisbiyah¹, RayiKurnia Perwitasari¹, WikaYuli Deakandi² and Feelin Fatwa Titiharja³

Department of Child Health, Faculty of Medicine, Dr. Soetomo General Hospital, Universitas Airlangga¹, Surabaya, East Java, Faculty of Medicine, Universitas Islam Malang², Malang, East Java, Faculty of Medicine, Muhammadiyah University of Malang³, Malang, East Java, Indonesia

Background: The most frequent congenital endocrine disorder in children is congenital hypothyroidism (CH). In Indonesia, CH's prevalence is greater than the data currently available. However, CH newborn screening has only been carried out in a few locations across Indonesia. Therefore, there is a need to raise primary healthcare practitioners' understanding of the CH Screening Program. **Materials and Methods:** This study used a pre-posttest design to gauge the healthcare professionals' knowledge of newborn congenital hypothyroid screening. After intervention, a post-test was given. The categorical measures' descriptive statistical analysis results were presented as a mean, median, percentage (%), and frequency (n) distribution table. The improvement in healthcare professionals' understanding of newborn CH screening was evaluated using the *Paired Samples test*. The cutoff for statistical significance was $p < 0.05$ using SPSS version 20.0. **Results:** In our study, most of the participants were midwives with a total of 21/53 (39.62%) followed by nurses with 19/53 (35.83%). There was also an increase in the score where previously in the pretest only 1 person got a score above 80, then it increased to half of the participants, namely 49.06% (26/53) who got better scores. months and 37/54 (68.5%). With the Paired Sample test, we prove that there is a significant increase in the participants' scores, and the data obtained is $p < 0.05$, which means that there is an increase in the average data from the pretest to the post-test. **Conclusion:** There was an improvement in pre-posttest knowledge of healthcare who had been given education about CH newborn screening.

Key Words: Congenital hypothyroidism, Newborn screening, Knowledge

Introduction

Congenital hypothyroidism (CH) is the most prevalent congenital endocrine condition in children.^{1,2)} Thyroid hormones are essential in growth and neurological development, especially during the first years of life. Thus hypothyroid screening during childbirth can prevent mental retardation and disability.³⁾

The global prevalence of CH is 1 per 3000 to 4000 births.^{1,2)} In Indonesia, the incidence of CH is estimated to be higher than the existing data.⁴⁾ The prevalence in Indonesia is 1 per 2513 birth, based on CH screening in 14 provinces. CH rarely manifests during childbirth and becomes misdiagnosed if not screened.⁵⁾ However, neonatal CH screening in Indonesia has not been implemented nationally, only in selected regions and hospitals.⁴⁾ Hence, CH often

Received October 21, 2022 / Accepted November 7, 2022

Correspondence: Nur Rochmah, MD, PhD, Department of Child Health, Faculty of Medicine, Dr. Soetomo General Hospital, Universitas Airlangga, Mayjend Prof. Dr. Moestopo No. 6-8, Surabaya, East Java 60286, Indonesia
Tel: 62-81703501118, Fax: 6231-5022068, E-mail: nur-r@fk.unair.ac.id

Copyright © the Korean Thyroid Association. All rights reserved.

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

goes undetected and is found too late.⁶

The newborn screening program for CH has been mandatory by the Minister of Health of the Republic of Indonesia since 2014. However, the implementation is still challenging due to the screening not being covered by universal health insurance.⁷

The first level of health facility (Public Health Center or Puskesmas) plays a vital role in the Congenital Hypothyroid Screening program. Puskesmas not only does the specimen collection and examination, but also provides communication, information, education, follow-up on screening results, diagnosis, management, case monitoring, organization, and program monitoring-evaluation.⁸ This study aims to increase the awareness and knowledge of CH and Congenital Hypothyroid Screening programs among primary healthcare professionals.

Materials and Methods

In this study, a pre-posttest design was used in order to assess the healthcare in their awareness of congenital hypothyroid screening for newborns. Congenital hypothyroid screening for newborn in-

struction was delivered through online meetings with experts as part of the strategy. Indonesian Bahasa was used to teach health education through seminars and group discussions. Participants took a pre-test evaluating their general knowledge of congenital hypothyroid screening for newborns before the intervention. A post-test was taken once the intervention was over. Volunteers from Indonesian healthcare facilities in East Java (Driyorejo Public Health Center, Gresik) were made the study's sample. The following criteria were used to determine inclusion: age above 18, communication skills, and willingness to participate in a survey. Consent from participants was gained in advance using consecutive sampling. This project received ethical clearance from the connected university under number 138/EC/KEPK/FKUA/2022. In this study, samples' age, gender, employment status, and educational background were recorded via an online self-reported questionnaire that participants completed. Using a self-created questionnaire based on our previous study, participants' knowledge of the fundamentals of congenital hypothyroid screening in the newborn was assessed.

Statistical Analyses

Descriptive statistical analysis of the categorical measures was reported as mean, median, percentage (%), and frequency (n) distribution table. The improvement of knowledge of healthcare about CH screening for newborns was analyzed by *Paired Samples Test*. Statistical significance was set at $p < 0.05$. The statistical analyses were performed using SPSS version

Table 1. Demographic characteristics of samples

Variables	n (%) (n=53)
Gender	
Male	7 (13.21)
Female	46 (86.79)
Age	38.87 ± 8.63
Educational background	
Diploma degree	35 (66.04)
Bachelor degree	17 (32.08)
Doctoral degree	1 (1.89)
Job	
Nurse	19 (35.85)
Midwife	21 (39.62)
Medical doctor	9 (16.98)
Pharmacist	1 (1.89)
Nutritionist	3 (5.66)

Table 3. Interpretation of sample's knowledge

Interpretation	Range	Pretest n (%)	Posttest n (%)
Good	80-100	1 (1.89)	26 (49.06)
Average	60-79	29 (54.72)	18 (33.96)
Poor	<60	23 (43.40)	9 (16.98)

Table 2. Comparison of pretest and posttest scores

Variable	N	Mean	Std. Deviation	Min	Max	Sig. (2-tailed)
Pre-test	53	59.24	7.69	46.67	80	0.000
Post-test	53	73.20	13.23	46.67	86.67	0.000

20.0. The Kolmogorov-Smirnov test was used to determine whether the data from the pretest and post-test scores were normal.

Results

This study had 53 participants and of them there were doctors totaling 9/53 (16.98%). Among 53 participants, most of them were Diploma degree and also 1 doctoral degree. The least participant is from the

pharmacist (Table 1).

There was an improvement in the posttest score compared to the pretest where previously there were 23/53 (43.40%) participants who scored poorly in the pretest and increased in the posttest to 26 (49.06%) participants who scored well (Tables 2, 3). The data acquired is $p < 0.05$, which indicates that there has been an increase in the average data from the pretest to the post-test. Using the Paired Sample test, we demonstrate that the participant scores have sig-

Table 4. Questionnaire of congenital hypothyroidism (CH) newborn screening

No.	Questionnaire	Pretest		Post-test	
		Correct n (%)	Wrong n (%)	Correct n (%)	Wrong n (%)
Guidelines for CH newborn screening					
1	Examination of congenital hypothyroidism in newborns was done by checking the TSH	49 (92.45)	4 (7.55)	52 (98.11)	1 (1.89)
2	The most ideal blood specimen collection was immediately after the newborn until the age of 72 hours.	7 (13.21)	46 (86.79)	11 (20.75)	42 (79.25)
3	CH screening in newborns was positive if the TSH level was ≥ 20 mU/L.	50 (94.34)	3 (5.66)	48 (90.57)	5 (9.43)
4	Babies with positive screening results could already establish the diagnosis of CH, so there was no need to confirm with the re-examination of serum TSH and FT4.	24 (45.28)	29 (54.72)	38 (71.70)	15 (28.30)
5	The diagnosis of CH was made when the TSH level was low and the FT4 was high	12 (22.64)	41 (77.36)	31 (58.49)	22 (41.51)
6	More than 95% of newborns with CH have no clinical signs at birth.	50 (94.34)	3 (5.66)	47 (88.68)	6 (11.32)
7	CH is one of the causes of mental retardation that could be prevented by early therapy.	51 (96.23)	2 (3.77)	52 (98.11)	1 (1.89)
8	Signs and symptoms that can appear in children with CH were lethargy (decreased activity), jaundice (yellow), macroglossia (large tongue), umbilical hernia (bulk), flat nose, constipation, and dry skin.	44 (83.02)	9 (16.98)	51 (96.23)	2 (3.77)
9	The sample was used in the form of blood taken by pricking the baby's heel (heelprick).	53 (100)	0 (0)	53 (100)	0 (0)
10	The puncture site was the medial part of the left or right heel.	0 (0)	53 (100)	20 (37.74)	33 (62.26)
11	Puncture the heel with a disposable sterile lancet to a depth of 2 mm.	51 (96.23)	2 (3.77)	53 (100)	0 (0)
12	The blood that comes out is dripped on a special filter paper until the paper circle was full of blood, then after it was dry, it was sent to the CH screening laboratory.	50 (94.34)	3 (5.66)	53 (100)	0 (0)
13	After the heel was pierced, the first drop of blood was directly dripped onto the filter paper.	12 (22.64)	41 (77.36)	31 (58.49)	22 (41.51)
14	In order to be examined, at least two full circles of filter paper blood specimens are required.	3 (5.66)	50 (94.34)	1 (1.89)	52 (98.11)
15	If the blood didn't come out, the examiner can make a squeezing motion on the baby's heel.	11 (20.75)	41 (77.36)	32 (60.38)	21 (39.62)

FT4: free thyroxine, TSH: thyroid stimulating hormone

nificantly increased.

The detail of the questionnaire is described in Table 4.

Discussion

This study showed that education should be viewed as an important factor in empowerment for all healthcare. This approach should enable all healthcare to use knowledge and practical skills in a newborn screening program of CH for better care. The clinical manifestation of CH, which is undetected at or around birth, occurs too late for treatment to reverse the resulting mental and growth retardation. CH may not be detected at all in many neonates, which places an unwarranted financial, social, and medical burden on the family. Infants who receive early diagnosis and treatment typically grow up without mental disabilities and can contribute positively to society. With effective screening, the newborn with CH can be diagnosed early and treated early about their disease.⁹⁾

From the result, we can also see that the subject mostly were female at the age around 38, and mostly of their job were nurses where it was also reported in a previous journal in Brazil that it was known that many of their subjects were female caregivers who were more concerned with pediatric patients because many of them also had children.¹⁰⁾

Our study showed significant gains in knowledge and attitude following counseling in a healthcare unit. In previous study, there was a comparable improvement in attitude was shown with CH screening for newborn tests. The results also led to a comparable modification in practice.¹¹⁾

This result also shows that there was a difference in the pretest and post-test scores in subjects after being given education about the importance of newborn screening for CH, it was intended that pediatric patients suffering from CH can be treated early. This was also stated in another journal where there was a significant difference in the quality of life of a child with CH in which one of them was given early treatment after newborn screening was done compared to children who already have symptoms of hypothyroidism and had just been treated.¹²⁾

In conclusion, the findings of several studies highlight the difficulties healthcare faces in screening for CH in newborns. Although this course of screening is simple, it may not provide adequate education. According to the data presented, that well-educated healthcare with essential information about CH in newborns and its screening processes present higher chances of early screening of CH in newborns.

Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

Orcid

Muhammad Faizi: <https://orcid.org/0000-0002-7009-4896>

Nur Rochmah: <https://orcid.org/0000-0002-9626-9615>

Yuni Hisbiyah: <https://orcid.org/0000-0002-1362-108X>

RayiKurnia Perwitasari: <https://orcid.org/0000-0002-8699-4063>

WikaYuli Deakandi: <https://orcid.org/0000-0002-5656-8725>

Feelin Fatwa Titiharja: <https://orcid.org/0000-0002-6296-7191>

References

- 1) LaFranchi SH, Austin J. *How should we be treating children with congenital hypothyroidism?* *J Pediatr Endocrinol Metab* 2007;20(5):559-78.
- 2) Ghasemi M, Hashemipour M, Hovsepian S, Heiydari K, Sajadi A, Hadian R, et al. *Prevalence of transient congenital hypothyroidism in central part of Iran.* *J Res Med Sci* 2013; 18(8):699-703.
- 3) Cherella CE, Wassner AJ. *Congenital hypothyroidism: insights into pathogenesis and treatment.* *Int J Pediatr Endocrinol* 2017;2017:11.
- 4) Endokrinologi UKK, Indonesia IDA. *Diagnosis dan tata laksana hipotiroid kongenital. Panduan Praktik Klinis Ikatan Dokter Anak Indonesia Jakarta* 2017;1-21.
- 5) Tariq B, Ahmed A, Habib A, Turab A, Ali N, Soofi SB, et al. *Assesment of knowledge, attitudes and practices towards newborn screening for congenital hypothyroidism before and after a health education intervention in pregnant women in a hospital setting in Pakistan.* *Int Health* 2018;10(2):100-7.
- 6) Rustama DS, Fadil MR, Harahap ER, Primadi A. *Newborn screening in Indonesia.* *Southeast Asian J Trop Med Public Health* 2003;34 Suppl 3:76-9.
- 7) Pulungan AB, Oldenkamp ME, van Trotsenburg ASP, Windarti W, Gunardi H. *Effect of delayed diagnosis and treatment of congenital hypothyroidism on intelligence and quality of life: an observational study.* *Medical J Indones*

- 2019;28(4):396-401.
- 8) Soepardi J, Anggraeni ND, Rustama DS, Soenggoro EP, Tridjaja B, Roeslani RD, et al. *Pedoman skrining hipotiroid. Jakarta, Indonesia: Kementerian Kesehatan RI; 2014. p. 1-71.*
 - 9) Brito LNS, Andrade CLO, Alves CAD. *Adhesion to treatment by children with congenital hypothyroidism: knowledge of caregivers in Bahia State, Brazil. Rev Paul Pediatr 2021;39:e2020074.*
 - 10) Buyukgebiz A. *Newborn screening for congenital hypothyroidism. J Clin Res Pediatr Endocrinol 2013;5 Suppl 1:8-12.*
 - 11) Neethu J, Anwar F, Alex A, Reshma R, Rajan R, Prasobh GR. *Assessment of risk factors of congenital hypothyroidism and impact of patient counselling on improving knowledge, attitude and practice of the disease in post natal mothers-a pilot study. Am J Pharm Tech Res 2020;8(5):2321-3647.*
 - 12) Rochmah N, Faizi M, Dewanti C, Suryawan A. *Pediatric quality of life in congenital hypothyroidism: an Indonesian study. Int J Thyroidol 2020;13(2):150-4.*

Effectiveness of Health Education toward Healthcare Knowledge Improvement about Congenital Hypothyroidism Newborn Screening

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Valentina Drozd, Daniel I. Branovan, Christoph Reiners. "Increasing Incidence of Thyroid Carcinoma: Risk Factors and Seeking Approaches for Primary Prevention", *International Journal of Thyroidology*, 2020
Publication 2%
- 2 www.koreamed.org
Internet Source 2%
- 3 Wahyu Setyaningsih, Ratna Dwi Wulandari. "The Evaluation of Congenital Hypothyroidism Screening Program in Indonesia: A Literature Review", *Jurnal Aisyah : Jurnal Ilmu Kesehatan*, 2022
Publication 1%
- 4 brieflands.com
Internet Source 1%
- 5 Dong Woo Lee, Hyoung Shin Lee. "A Case of Papillary Thyroid Carcinoma Accompanied by Zenker's Diverticulum: Considerations for

Surgery", International Journal of Thyroidology, 2022

Publication

6	medicopublication.com Internet Source	1 %
7	Myung-Chul Chang. "Multiple Endocrine Neoplasia 2A Detected by Peptic Ulcer Perforation with Hypercalcemia", International Journal of Thyroidology, 2022 Publication	1 %
8	doc.rero.ch Internet Source	1 %
9	www.scielo.br Internet Source	1 %
10	Sang Hwon Cho, Jin Seok Lee, Hyeok Jun Yun, Yong Sang Lee, Hang-Seok Chang. "Effect of Afatinib for Lung Cancer on Papillary Thyroid Carcinoma", International Journal of Thyroidology, 2022 Publication	<1 %
11	Vivian Soetikno. "How important is early detection and prompt treatment in congenital hypothyroidism patients?", Medical Journal of Indonesia, 2019 Publication	<1 %
12	poj.org.pk Internet Source	<1 %

13

Hamenudin Hamzah, Rosnah Sutan, Azmi Mohd Tamil, Aminah Bee Mohd Kassim, Adliah Mohamed Soid, Amar Singh HSS. "Assessment of healthcare workers' behavior in implementing a newborn screening program for congenital hypothyroidism in Perak, Malaysia", Journal of Health Research, 2020

Publication

<1 %

14

Minjing Zou, Yufei Shi. "Chapter 2 Congenital Hypothyroidism and Thyroid Cancer", IntechOpen, 2012

Publication

<1 %

15

Peng Xue, Yuqi Yang, Qi Yun, Yue Cui, Bin Yu, Wei Long. "Variant of TSHR is Not a Frequent Cause of Congenital Hypothyroidism in Chinese Han Patients", International Journal of General Medicine, 2021

Publication

<1 %

16

www.researchgate.net

Internet Source

<1 %

17

Catia Mio, Giorgio Grani, Cosimo Durante, Giuseppe Damante. "Molecular defects in thyroid dysgenesis", Clinical Genetics, 2019

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Effectiveness of Health Education toward Healthcare Knowledge Improvement about Congenital Hypothyroidism Newborn Screening

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
