

SOUTH AFRICAN JOURNAL OF CHILD HEALTH

March 2020 | Volume 14 | No. 1

The South African Journal of Child Health is the official journal of the South African Paediatric Association

EDITORIAL

- 2 South African undergraduate paediatric clinical training: A call to action
A George, Z Dangor, S G Lala

RESEARCH

- 4 A descriptive study of children with cerebral palsy at Chris Hani Baragwanath Academic Hospital
N Mahlaba, F L Nakwa, J R Rodda
- 10 Factors associated with preterm very low birthweight infant mortality at a tertiary hospital in Limpopo Province, South Africa
T S Ntuli, M P A Mashego, N Shipalana, C Sutton, M H K Hamese
- 15 Impact of the child support grant on the diet and nutritional status of children under 5 years old in Mogalakwena Municipality, Limpopo Province, South Africa
M J Kekana, N S Mabapa, X G Mbhenyane
- 20 The quality of life, health needs and knowledge of children living with congenital heart disease in KwaZulu-Natal, South Africa
P Pillay, N Pefile, S Cobbing, T Gurayah
- 25 Children's ability to consent to medical management in South Africa
J van Heerden, R Delpont, M Kruger
- 30 Correlation between metabolic control, serum electrolyte levels, duration of illness and QT interval length parameters in children with type 1 diabetes mellitus
N Rochmah, D Fiasry, M Faizi, T Ontoseno
- 34 Patterns of healthcare utilisation and barriers affecting access to child healthcare services in low-income urban South African settings
S A Adedini, M Sello, D Thaele, S A Madhi
- 40 Congenital anomalies of the kidney and the urinary tract in a South African paediatric nephrology setting
N C Okoronkwo, A Mudi, C Levy, T Khumalo, G Moonsamy
- 45 The South African Sodium Regulation (R214): Does it make provision for processed foods frequently consumed by young children?
M Korff, M Wicks, T van Zyl, B van der Westhuizen

REVIEW

- 50 Children with oculocutaneous albinism in Africa: Characteristics, challenges and medical care
J C Kromberg

CASE REPORT

- 55 *Bacillus* species cultured repeatedly from the breastmilk of a donor milk mother
C J Opperman, E J Kruger, J Wright

SAPA NEWS

CPD

EDITOR

Prof. J M Pettifor

FOUNDING EDITOR

Prof. N P Khumalo

EDITORIAL BOARD

Prof. M Adhikari (University of KwaZulu-Natal, Durban)
Prof. M Kruger (Stellenbosch University)
Prof. H Rode (Red Cross War Memorial Children's Hospital, Cape Town)
Prof. L Spitz (Emeritus Nuffield Professor of Paediatric Surgery, London)
Prof. A Venter (University of the Free State, Bloemfontein)
Prof. A Westwood (Red Cross War Memorial Children's Hospital, Cape Town)
Prof. D F Wittenberg (University of Pretoria)

SOUTH AFRICAN MEDICAL ASSOCIATION

GENERAL MANAGER

Dr Vusumuzi Nhlapho

EXECUTIVE EDITOR

Bridget Farham

MANAGING EDITORS

Claudia Naidu
Naadia van der Bergh

TECHNICAL EDITORS

Naadia van der Bergh
Kirsten Morreira

PRODUCTION & DISTRIBUTION OFFICER

Emma Jane Couzens

SENIOR DESIGNER

Clinton Griffin

HEAD OF PUBLISHING

Diane Smith | Tel. 012 481 2069
Email: dianes@samedical.org

ONLINE SUPPORT

Gertrude Fani | Tel. 021 532 1281
Email: publishing@samedical.org

HEAD OFFICE

Block F, Castle Walk Corporate Park,
Nossob Street, Erasmuskloof Ext. 3,
Pretoria, 0181

EDITORIAL OFFICE

Suite 11, Lonsdale Building,
Lonsdale Way, Pinelands, 7405
Tel. 021 532 1281
Email: publishing@samedical.org

ISSN 1999-7671

Use of editorial material is subject to the Creative Commons Attribution – Non-commercial Works Licence. <http://creativecommons.org/licenses/by-nc/4.0>



Published by the South African Medical Association |
Suite 11, Lonsdale Building | Lonsdale Way | Pinelands | 7405
Papers for publication should be addressed to the Editor, via the
website: www.sajch.org.za

©Copyright: South African Medical Association (Pty) Ltd

Correlation between metabolic control, serum electrolyte levels, duration of illness and QT interval length parameters in children with type 1 diabetes mellitus

N Rochmah, MD (Paed); D Fiasry, MD (Paed); M Faizi, MD (Paed); T Ontoseno, MD (Paed) (Cardio) FIHA

Department of Child Health, Faculty of Medicine, Airlangga University-Dr. Soetomo Hospital, Surabaya, Indonesia

Corresponding author: N Rochmah (drnurochmah@gmail.com)

Background. Cardiac autonomic neuropathy (CAN) is a common complication of type 1 diabetes mellitus (T1DM) that can cause arrhythmia and increase mortality risk. Prolonged corrected-QT (QTc) and dispersion-QT (QTd) intervals can be used for early detection. Few studies have investigated relationships between metabolic control, serum electrolyte levels, duration of illness and CAN in T1DM children.

Objective. To investigate serum haemoglobin A1c (HbA1c) level, serum electrolyte levels, duration of illness and QTc and QTd interval parameters in T1DM children.

Methods. This cross-sectional study included T1DM patients up to 18 years of age at the Paediatric Outpatients Clinic, Dr. Soetomo Hospital, Surabaya, Indonesia, between January 2016 and July 2017. Serum HbA1c and sodium and potassium levels were analysed. QTc and QTd intervals were recorded from 12-lead electrocardiograms.

Results. Of the 40 patients, 70% were found to have poor metabolic control. Among patients with poor metabolic control, prolonged QTc and QTd intervals were observed in 21 (75.0%) and 16 (57.1%), respectively. Serum HbA1c levels were significantly correlated with QTc ($p=0.001$; $r=0.541$) and QTd interval lengths ($p=0.018$; $r=0.373$). Hypokalaemia was observed in 28 patients, with prolonged QTc and QTd intervals seen in 85% ($n=17$) and 55% ($n=11$), respectively. We found significant correlations between potassium level and QTc interval length ($p\leq 0.05$; $r=0.556$), but not between potassium level and QTd interval length.

Conclusion. We suggest that prolonged QTc and QTd intervals could be predictive of a cardiovascular complication in T1DM in children.

S Afr J Child Health 2020;14(1):30-33. <https://doi.org/10.7196/SAJCH.2020.v14.i1.1624>

The global prevalence of type 1 diabetes mellitus (T1DM) is increasing.^[1,2] In 2017, an estimated 1 106 500 people younger than 20 years were affected and the number of newly diagnosed cases each year was estimated at 132 600.^[2]

Cardiac autonomic neuropathy (CAN) is a serious complication in diabetes,^[3] which is estimated to account for 5 per 1 000 cardiovascular-related T1DM deaths each year.^[2] Arrhythmia, which forms part of CAN, can be detected as prolonged corrected QT (QTc) and dispersion QT (QTd) intervals.^[4,5] Prolonged QTc and QTd intervals can trigger ventricular arrhythmia and it may lead to sudden death.^[6,7] Poor metabolic control and electrolyte imbalance in diabetic ketoacidosis (DKA) are known to cause prolonged QTc and QTd intervals.^[8-10]

Duration of illness also appears to contribute to the process of CAN. In 2018, the International Society for Pediatric and Adolescent Diabetes (ISPAD) recommended screening T1DM patients from the age of 11 for CAN complications within 2 - 5 years of disease onset, followed up annually by physical examination and clinical testing.^[11] The American Diabetes Association further recommends screening hyperglycaemic patients for cardiovascular disease by measuring serum haemoglobin A1c (HbA1c) every 3 months as a means of monitoring blood glucose control, with the target set at $<7.5\%$.^[12]

Our study aimed to investigate the correlation between serum HbA1c levels, serum electrolyte levels, duration of illness and parameters of QT intervals among Indonesian T1DM patients 18 years old and younger. Few studies have considered the correlations between these parameters and none so far in an Indonesian population.

Methods

We conducted a cross-sectional study at the paediatric endocrinology outpatient clinic and the paediatric ward of the Dr. Soetomo Hospital in Surabaya, Indonesia, from January 2016 to July 2017. All T1DM patients up to 18 years were included. Patients with a history of congenital disorders, including congenital heart disease, were not included in the study. According to the 2014 ISPAD guidelines,^[1] diagnosis criteria of T1DM was (i) classic symptoms of diabetes or hyperglycaemic crisis, with a plasma glucose concentration ≥ 11.1 mmol/L (200 mg/dL); (ii) a fasting plasma glucose level ≥ 7.0 mmol/L (≥ 126 mg/dL); (iii) a glucose level ≥ 11.1 mmol/L (≥ 200 mg dL) two hours post load as determined by an oral glucose tolerance test; or (iv) HbA1c $>6.5\%$. All patients in our study used basal-bolus insulin injection (detemir and short-acting insulin). As per ISPAD criteria, DKA was defined as a condition of hyperglycaemia (blood glucose >11 mmol/L (200 mg/dL)), acidosis (pH <7.3 or bicarbonate levels <15 mmol/L), ketonaemia and ketonuria.^[13]

Nutritional status was classified according to the World Health Organization BMI-for-age values defined for children and adolescents (5 - 18 years of age): severe malnutrition ($z < -3$); moderate malnutrition ($-3 < z < -2$); normal ($-2 < z \leq 1$); overweight ($1 < z < 2$); and obesity ($z > 2$).^[14] We grouped the subjects into three groups: normal; malnourished; and overweight/obese.

Blood samples were analysed for HbA1c and sodium and potassium levels at the clinical pathology laboratory of the Dr.

Soetomo Hospital, Surabaya. HbA1c results were classified as poor (>9%), suboptimal (7.5 - 9%) or good (<7.5%).^[15] Results were subsequently classified as indicative of poor or good metabolic control (HbA1c >9% and HbA1c <9%, respectively). Potassium levels <3.5 mmol/L and >5.5 mmol/L were considered indicative of hypokalaemia and hyperkalaemia, respectively. Potassium levels of 3.5 - 5.5 mmol/L were considered normal.^[16] Similarly, sodium levels <136 mmol/L and >145 mmol/L were classified as indicative of hyponatraemia and hypernatraemia, respectively. Sodium levels of 136 - 145 mmol/L were considered normal.^[16]

The QTc and QTd values were obtained from 12-lead electrocardiography (Bionet CardioTouch 3000, Bionet, USA). The QTc interval was calculated according to Bazett's formula ($QTc = \frac{QT}{\sqrt{RR}}$), with the RR interval measured relative to the previous QRS complex. A prolonged QTc was taken as >440 ms in boys and >460 ms in girls. QTd was calculated as the difference between QTmax and QTmin; the interval was considered prolonged if the difference was >60 ms.^[6]

Statistical analysis

Means and associated standard deviations were calculated for normally distributed data. Data that were not normally distributed were described as medians and the associated range. Correlations between serum levels for HbA1c, sodium and potassium, duration of illness and QT interval parameters were assessed according to a phi association test with the correlation coefficient *r*. All data analyses were performed using SPSS software version 17.0 (SPSS Inc., USA). A significance level of *p*≤0.05 and a 95% confidence interval (CI) were used in all analyses.

Ethical approval

Approval was granted by the ethics committee overseeing health research at the Dr. Soetomo Hospital, Surabaya, Indonesia (ref. no. 555/Panke.KKE/IX/2016).

Results

The study included 40 patients with T1DM, of whom 18 were male. Close to two-thirds had a normal nutritional status (60%). DKA was observed in 57.5% of cases. The characteristics of the study population are shown in Table 1. Prolonged QTc and QTd intervals were observed in 23 and 18 patients, respectively. Approximately 90% of these patients showed poor metabolic control. We did not observe any cases of hyperkalaemia, but found that all of the subjects with hypernatraemia (*n*=3) presented with a prolonged QTc interval, as shown in Table 2.

Discussion

Our results showed significant correlations between serum HbA1c and both QTc (*p*=0.001; *r*=0.541) and QTd intervals (*p*=0.018; *r*=0.373). Prolonged QTc and QTd intervals were observed in 21 (75.0%) and 16 (57.1%) patients with poor metabolic control. The observation is similar to that of Stern *et al.*,^[8] who reported an association between serum HbA1c and QTc interval length (*p*<0.001). Tesfaye *et al.*^[9] suggested that high serum HbA1c levels are caused by neuropathy in T1DM, with a reported odds ratio of 2.48 (*p*<0.001, 95% CI). Findings from the Diabetes Control and Complications Trial (DCCT)^[13] suggested that complications of T1DM may be prevented by optimal metabolic control. The DCCT further reported that intensive treatment reduced the risk of cardiovascular disease by 42% (*p*=0.02).^[17] Circulating endothelial cells, identified by fluorescence microscopy, were

Table 1. Baseline characteristics of the study population (N=40)

Characteristics	Total
Gender, <i>n</i> (%)	
Male	18 (45.0)
Female	22 (55.0)
Age (months), mean (SD)	144 (38.7)
Weight (kg), mean (SD)	34.5 (13.5)
Height (cm), mean (SD)	135.6 (14.2)
DKA during the study, <i>n</i> (%)	
Yes	23 (57.5)
No	17 (42.5)
Nutritional status, <i>n</i> (%)	
Normal	24 (60.0)
Malnutrition	14 (34.5)
Overweight/obesity	2 (5.0)
Serum HbA1c (%), mean (SD)	10.27 (2.2)
Potassium level (mmol/L), median (range)	3.8 (1.7 - 5.9)
Sodium level (mmol/L), mean (SD)	136.1 (6.6)
Duration of illness (months), median (range)	16.5 (0 - 132)
QTc interval (ms), mean (SD)	460 (43.0)
QTd interval (ms), median (range)	50 (40 - 120)

SD = standard deviation; HbA1c = haemoglobin A1c.

suggested as a cause of endothelial vascular damage and have been reported to be elevated in T1DM patients with raised HbA1c levels.^[18] This contradicts the results of a study by Lu *et al.*,^[19] which found that serum HbA1c was not significantly correlated with the length of the QTc interval (*p*=0.77).

CAN is considered a cardiovascular complication of diabetic neuropathy. Earlier studies reported that T1DM patients may suffer cardiovascular events by the age of 30,^[20] which often appear after 20 years of the disease.^[21] Cardiovascular disease appears not only to occur earlier in T1DM than in type 2 diabetes mellitus, but also to affect men and women at a similar rate by the age of 40.^[22] Poor glycaemic control is a risk factor for cardiovascular events in both types of diabetes mellitus.^[12] Hypertension, proteinuria, obesity, high serum HbA1c and lipid levels, smoking and a long disease duration may be important risk factors for cardiovascular events.^[12] Therefore, screening for prolonged QTc and QTd intervals should be performed to predict a possible cardiovascular event.

Our results showed a significant correlation between serum potassium level and QTc interval length, but not between potassium level and QTd interval length. Serum sodium level was not significantly correlated with a prolonged QTc or QTd interval. No significant correlation was found between the length of the QTc interval and serum levels of potassium (*p*=0.139) or sodium (*p*=0.962) in DKA patients in a study by Aygün *et al.*,^[23] similar to findings regarding the correlation between potassium level and QTc interval length in DKA patients reported by Kuppermann *et al.*^[10] Hypokalaemia, hyperkalaemia, hypocalcaemia, hypophosphataemia and hypomagnesaemia may develop in DKA.^[24]

Our results showed prolonged QTc and QTd intervals in 18 (66.7%) and 14 (51.9%) patients, respectively, within 2 years of disease onset. However, the correlation between duration of illness and QT interval lengths was not significant. Uysal *et al.*^[25] reported significant differences in QTd interval length in newly diagnosed T1DM patients at follow-up after 1 year, within 5 years and later than 5 years compared

Table 2. Correlation between serum haemoglobin A1c levels, serum electrolytes and QTc and QTd intervals (N=40)

Variable	QTc interval				QTd interval			
	Normal, n (%)	Elongated, n (%)	p-value	r	Normal, n (%)	Elongated, n (%)	p-value	r
Metabolic control*								
Good (N=12)	10 (83.3)	2 (16.7)	0.001 [†]	0.541	10 (83.3)	2 (16.7)	0.018 [†]	0.373
Poor (N=28)	7 (25.0)	21 (75.0)			12 (42.9)	16 (57.1)		
Potassium								
Normal (N=20)	14 (70.0)	6 (30.0)	0.000 ^{††}	0.556	13 (65.0)	7 (35.0)	0.204	0.201
Hypokalaemia (N=20)	3 (15.0)	17 (85.0)			9 (45.0)	11 (55.0)		
Sodium								
Normal (N=23)	11 (47.8)	12 (57.2)	0.769	0.048	14 (60.9)	9 (39.1)	0.175	0.286
Hyponatraemia (N=14)	6 (42.9)	8 (57.1)			6 (42.9)	8 (57.1)		
Hypernatraemia [‡] (N=3)	0	3 (100)	–	–	2 (66.7)	1 (33.3)	0.846	–0.038
Duration of illness, months								
<24 (N=27)	9 (33.3)	18 (66.7)	0.091	–0.267	13 (48.1)	14 (51.9)	0.209	–0.198
≥24 (N=13)	8 (61.5)	5 (38.5)			9 (69.2)	4 (30.8)		

HbA1c = haemoglobin A1c.

*HbA1c <6.5% considered indicative of good metabolic control; HbA1c ≥6.5% considered indicative of poor metabolic control.

[†]Statistically significant, $p \leq 0.05$.

[‡]Not analysed.

with controls ($p=0.005$, $p<0.001$, $p<0.001$, respectively). However, the correlation between QTd interval length and disease duration was not significant.^[25]

Study limitations

Owing to demographic and racial differences in our study population, our findings cannot be generalised.

Conclusion

The results of this study showed a significant correlation between serum HbA1c levels and both QTc and QTd interval length in children diagnosed with T1DM. We suggest that screening for prolonged QTc and QTd intervals may be used to predict cardiovascular complications in this patient group.

Declaration. None.

Acknowledgements. The authors thank the paediatric endocrinology team at the Dr Soetomo General Hospital, Surabaya, for their support.

Author contributions. All authors contributed to conceptualisation and design of the study and analysis, and interpretation of the data. NR, MF and TO were responsible for drafting the manuscript and critical revision of the scientific content. All authors approved the final manuscript for publication.

Funding. None.

Conflicts of interest. None.

- Craig ME, Jefferies C, Dabelea D, Balde N, Seth A, Donaghue K. Definition, epidemiology, and classification of diabetes in children and adolescents. *Pediatr Diabetes* 2014;15(Suppl.20):4-17. <https://doi.org/10.1111/pedi.12186>
- International Diabetes Federation. IDF Diabetes Atlas 8th edition 2017. Brussels: IDF, 2017. <http://diabetesatlas.org/resources/2017-atlas.html> (accessed 16 September 2018).
- Tang ZH, Zeng F, Li Z, Zhou L. Association and predictive value analysis for resting heart rate and diabetes mellitus on cardiovascular autonomic neuropathy in general population. *J Diabetes Res* 2014;2014:215473. <https://doi.org/10.1155/2014/215473>
- Zdárská D, Pelíšková P, Charvát J, et al. ECG body surface mapping (BSM) in type 1 diabetic patients. *Physiol Res* 2007;56(4):403-410.
- Voulgari C, Tentolouris N, Stefanadis C. The ECG vertigo in diabetes and cardiac autonomic neuropathy. *Exp Diabetes Res* 2011;2011:687624. <https://doi.org/10.1155/2011/687624>
- Park M. Cardiac arrhythmias. In: Park MK. *Park's Pediatric Cardiology for Practitioners*. 6th ed. Philadelphia: Elsevier, 2014:407-435.
- Clemente D, Pereira T, Ribeiro S. Ventricular repolarization in diabetic patients: Characterization and clinical implications. *Arq Bras Cardiol* 2012;99(5):1015-1022. <https://doi.org/10.1590/s0066-782x2012005000095>
- Stern K, Cho YH, Benitez-Aguirre P, et al. QT interval, corrected for heart rate, is associated with HbA1c concentration and autonomic function in diabetes. *Diabet Med* 2016;33(10):1415-1421. <https://doi.org/10.1111/dme.13085>
- Tesfaye S, Chaturvedi N, Eaton SE, et al. Vascular risk factors and diabetic neuropathy. *N Engl J Med* 2005;352(4):341-350. <https://doi.org/10.1056/NEJMoa032782>
- Kuppermann N, Park J, Glatter K, Marcin JP, Glaser NS. Prolonged QT interval corrected for heart rate during diabetic ketoacidosis in children. *Arch Pediatr Adolesc Med* 2008;162(6):544-549. <https://doi.org/10.1001/archpedi.162.6.544>
- Donaghue KC, Marcovecchio ML, Wadwa RP, et al. ISPAD Clinical Practice Consensus Guidelines 2018: Microvascular and macrovascular complications in children and adolescents. *Pediatr Diabetes* 2018;19(Suppl 27):262-274. <https://doi.org/10.1111/pedi.12742>
- De Ferranti SD, De Boer IH, Fonseca V, et al. Type 1 diabetes mellitus and cardiovascular disease: A scientific statement from the American Heart Association and American Diabetes Association. *Diabetes Care* 2014;37(10):2843-2863. <https://doi.org/10.2337/dc14-1720>
- Wolfsdorf JI, Allgrove J, Craig ME, et al. Diabetic ketoacidosis and hyperglycemic hyperosmolar state. *Pediatr Diabetes* 2014;15(Suppl 20):154-179. <https://doi.org/10.1111/pedi.12165>
- World Health Organization. BMI for age 5 - 19 years. Geneva: WHO, 2007. http://www.who.int/growthref/who2007_bmi_for_age/en/ (accessed 16 September 2018).
- Rewers MJ, Pillay K, De Beaufort C, et al. Assessment and monitoring of glycemic control in children and adolescents with diabetes. *Pediatr Diabetes* 2014;15(Suppl 20):102-114. <https://doi.org/10.1111/pedi.12190>
- Andropoulos DB. Pediatric normal laboratory values. In: Gregory GA, Andropoulos DB, eds. *Gregory's Pediatric Anesthesia*, 5th ed. Texas: Blackwell, 2012:1300-1314.
- Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Study Research Group. Intensive diabetes treatment and cardiovascular disease in patients with Type 1 diabetes. *N Engl J Med* 2005;353:2643-2653. <https://doi.org/10.1056/nejmoa052187>
- Ascioglu E, Yavuz DG, Koc M, et al. Circulating endothelial cells are elevated in patients with type 1 diabetes mellitus. *Eur J Endocrinol* 2010;162(4):711-717. <https://doi.org/10.1530/eje-09-0795>
- Lu Z, Lense L, Sharma M, et al. Prevalence of QT prolongation and associated LVEF changes in diabetic patients over a four-year retrospective time period. *J Community Hosp Intern Med Perspect* 2017;7(2):87-94. <https://doi.org/10.1080/2009666.2017.1320203>
- Skrivarhaug T, Bangstad HJ, Stene LC, Sandvik L, Hanssen KF, Joner G. Long-term mortality in a nationwide cohort of childhood-onset type 1 diabetic patients in Norway. *Diabetologia* 2006;49(2):298-305. <https://doi.org/10.1007/s00125-005-0082-6>
- Secrest AM, Becker DJ, Kelsey SE, Laporte RE, Orchard TJ. Cause-specific mortality trends in a large population-based cohort with longstanding childhood-onset type 1 diabetes. *Diabetes* 2010;59(12):3216-3222. <https://doi.org/10.2337/db10-0862>

RESEARCH

22. Laing SP, Swerdlow AJ, Slater SD, et al. Mortality from heart disease in a cohort of 23,000 patients with insulin-treated diabetes. *Diabetologia* 2003;46(6):760-765. <https://doi.org/10.1007/s00125-003-1116-6>
23. Aygün D, Aygün F, Nişli K, Çıtak A. [Electrocardiographic changes in children with diabetic ketoacidosis and ketosis]. *Turk Pediatri Ars* 2017;52(4):194-201. <https://doi.org/10.5152/turkpediatriars.2017.4917>
24. Savage MW, Dhatariya KK, Kilvert A, et al. Joint British Diabetes Societies guideline for the management of diabetic ketoacidosis. *Diabet Med* 2011;28(5):508-515. <https://doi.org/10.1111/j.1464-5491.2011.03246.x>
25. Uysal F, Ozboyaci E, Bostan O, Sağlam H, Semizel E, Cil E. Evaluation of electrocardiographic parameters for early diagnosis of autonomic dysfunction in children and adolescents with type-1 diabetes mellitus. *Pediatr Int* 2014;56(5):675-680. <https://doi.org/10.1111/ped.12329>

Accepted 31 July 2019.



Source details

SAJCH South African Journal of Child Health

Open Access ⓘ

Scopus coverage years: from 2008 to 2022

Publisher: Health and Medical Publishing Group

ISSN: 1994-3032 E-ISSN: 1999-7671

Subject area: Medicine: Pediatrics, Perinatology and Child Health

Source type: Journal

CiteScore 2021

1.1 ⓘ

SJR 2021

0.277 ⓘ

SNIP 2021

0.531 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 ▼

$$1.1 = \frac{175 \text{ Citations 2018 - 2021}}{164 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$0.6 = \frac{101 \text{ Citations to date}}{159 \text{ Documents to date}}$$

Last updated on 05 April, 2023 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
— Pediatrics, Perinatology and Child Health	#194/298	35th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

