

Diabetic ketoacidosis in children an 11-year retrospective in Surabaya, Indonesia

by Nur Rochmah

Submission date: 15-Mar-2023 03:28PM (UTC+0800)

Submission ID: 2037639457

File name: in_children_an_11-year_retrospective_in_Surabaya,_Indonesia.pdf (81.32K)

Word count: 2069

Character count: 11533

Original Article

Diabetic ketoacidosis in children: an 11-year retrospective in Surabaya, Indonesia

Nur Rochmah, Muhammad Faizi, Netty Harjantien

10

Abstract

Background Diabetic ketoacidosis (DKA) is a complication of type I diabetic mellitus (T1DM). Early recognition and prompt treatment can reduce mortality.

Objective To evaluate the profiles of patients with diabetic ketoacidosis in Dr. Soetomo Hospital, Surabaya, Indonesia.

Methods Retrospective medical records of patients admitted to the pediatric intensive care unit with DKA in Dr. Soetomo Hospital, Surabaya, Indonesia from January 2002 to June 2013 were reviewed. The data collected included clinical parameters, laboratory and imaging results, predisposing factors, complications and outcomes.

Results During an 11 years period, there were 58 cases of DKA in children with T1DM. Eighteen subjects (31%) were boys. The severity of DKA was classified as follows: mild 13 (22.4%), moderate 23 (39.7%), and severe DKA 22 (37.9%). Recurrent DKA was diagnosed in 24 (41.4%) patients. Common clinical profiles recorded were dehydration 46 (79.3%), malaise 37 (63.8%), decreased consciousness 35 (60.3%), dyspnea 27 (46.6%), vomiting 26 (44.8%), fever 25 (43.1%), seizure 13 (22.4%), and decreased body weight 9 (15.5%). Laboratory results observed were as follows: hyponatremia 19 (32.8%), hyperkalemia 12 (20.7%) and acute renal failure 3 (5.2%). Head CT scans showed that 2 (3.4%) patients suffered from cerebral edema. Infections, as triggers of DKA, were found in 12 (20.6%) patients: 4 caries and periodontitis, 3 urinary tract infections, 2 acute diarrhea, 2 acute pharyngitis, and 1 otitis externa. Four out of 24 patients with recurrent DKA failed to take their insulin dose prior to DKA. The average of length of patient stay in the PICU was 3.26 (SD 3.50) days. No patients died during the study.

Conclusion Dehydration is the most common clinical profile of DKA in our study. More than half of the patients suffer from moderate to severe DKA. [Paediatr Indones. 2015;55:40-3].

Keywords: children, diabetic ketoacidosis, clinical manifestations

9

Diabetic ketoacidosis (DKA) is one of the acute complications in type 1 diabetes mellitus (T1DM). It may be the first clinical manifestation observed. Diabetic ketoacidosis is the predominant cause of mortality in children with diabetes. An estimated 65,000 children under 15 year old develop the disease each year worldwide, and the global incidence in children continues to increase at a rate of 3% per year.^{1,2} Between 10 to 70% of these diagnosed children present as diabetic ketoacidosis cases.³

Early recognition and prompt treatment can substantially reduce mortality in children with T1DM. To the best of our knowledge, this is the first study reporting on clinical manifestations of DKA in Indonesia. We aimed to study the profiles of pediatric patients with DKA between year 2002-2013 in Dr. Soetomo Hospital, Surabaya, Indonesia.

Methods

Retrospective medical record reviews of patients admitted to the pediatric intensive care unit with

From the Division of Endocrinology, Department of Child Health, Airlangga University Medical School/ Dr. Soetomo Hospital, Surabaya, Indonesia.

Reprint requests to: Nur Rochmah, Division of Endocrinology, Department of Child Health, Airlangga University Medical School/ Dr. Soetomo Hospital, Jl. Prof. Dr. Moestopo 6-8 Surabaya 60286. Tel. +6281703501118. E-mail: dnmurrochmah@gmail.com.

DKA in Dr. Soetomo Hospital, Surabaya, Indonesia from January 2002-June 2013 were performed. We collected data on clinical parameters, laboratory and imaging results, predisposing factors, complications and outcomes. The current criteria for DKA diagnosis, published by the *International Society for Paediatric and Adolescent Diabetes*, are blood glucose > 11 mmol/L, venous pH < 7.3 or bicarbonate < 15 mmol/L, and ketonemia and ketonuria. The severity of DKA was categorized by the degree of acidosis as follows: mild: venous pH < 7.3 or bicarbonate < 15 mmol/L, moderate: pH < 7.2, bicarbonate < 10 mmol/L, or severe: pH < 7.1, bicarbonate < 5 mmol/L.³

consciousness 35 (60.3%), dyspnea 27 (46.6%), vomiting 26 (44.8%), fever 25 (43.1%), seizure 13 (22.4%), and decreased body weight 9 (15.5%).

We observed the following laboratory results: hyponatremia 19 (32.8%), hyperkalemia 12 (20.7%) and acute renal failure 3 (5.2%). Two (3.4%) patients had cerebral edema on head CT scan. There were 12 (20.6%) patients with infections during the study, confirmed by clinical manifestations, complete blood counts and blood cultures (**Figure 2**).

Of the subjects with recurrent DKA, 4/24 patients failed to take their insulin days before their DKA episodes. Mean length of patient stay in the

Results

There were 58 DKA cases in children with T1DM during the study period. Characteristics of patients are shown in **Table 1**. Clinical manifestations of DKA are presented in **Figure 1**.

Common clinical profiles recorded were dehydration 46 (79.3%), malaise 37 (63.8%), decreased con-

Table 1. Characteristics of the subjects

| Characteristics | n (%) |
|-------------------------|-----------|
| Male gender | 18 (31) |
| DKA classification | |
| Mild | 13 (22.4) |
| Moderate | 23 (39.7) |
| Severe | 22 (37.9) |
| Recurrent DKA | 24 (41.4) |
| First DKA manifestation | 34 (58.6) |




Figure 1. Percentage of clinical manifestations of DKA in Dr. Soetomo Hospital

PICU was 3.26 (SD 3.50) days. No patients died during the study.




Figure 2. Types of infections as DKA triggers

Discussion

More than half of the subjects experienced DKA as their first clinical manifestation of T1DM. In contrast, a study stated that approximately one-third of newly-diagnosed T1DM patients presented with DKA.⁴ In Europe and North America, this figure ranged from approximately 15 to 70%.⁵⁻⁹ It remains unknown why some children present in DKA and whether the development of diabetic ketoacidosis is a consequence of delayed diagnosis and treatment or whether it reflects a particularly aggressive form of diabetes.¹⁰

In our study, 39.7% and 37.9% of patients suffered from moderate and severe DKA, respectively. Other studies reported that mild DKA was more frequent than moderate and severe. Klingensmith *et al.* reported 17% subjects with moderate or severe DKA, while Oyarzabal *et al.* found that moderate and severe forms were 34.4% and 17.8%, respectively.^{4,11}

Boys comprised 31% of our subjects. Usher-Smith JA *et al.* reported that of 21 studies on the effect of gender on DKA frequency, 20 found no effect, while 1 study by Neu *et al.* (n=2,121) reported a small but statistically significant increase in DKA frequency in girls in 2003.^{12,13}

The most common clinical manifestation

observed was dehydration, consistent with finding in earlier study.¹⁴ Other clinical manifestations included weakness, decreased consciousness, dyspnea, vomiting, fever, seizures, and decreased body weight. Similarly, another study stated that DKA often presented abdominal pain,^{15,16} weakness,¹⁶ changes in mental status,^{14,16,18} dyspnea,¹⁷ vomiting,¹⁵⁻¹⁷ and anorexia.¹⁶ Two studies also showed that children with DKA had significantly greater weight loss than those without.¹⁴⁻¹⁶ A study reported that classic triad (polyuria, polydipsia, and weight loss) had the highest percentage in DKA,¹⁸ meanwhile another study reported polydipsia, polyuria, weakness, and abdominal pain were the commonest symptoms observed.¹⁹

Hyponatremia, hyperkalemia and acute renal failure were improved after rehydration, insulin treatment and serum electrolyte correction. Poovazaghi *et al.* found that 11.5% of DKA patients had renal failure. Sepsis and shock were the common etiological factors. Mortality due to acute renal failure (ARF) in DKA was 40%.²⁰

Twelve (20.6%) patients suffered from infection during the study. Poovazaghi reported that infections were often the precipitating factors in DKA, as observed in 47% of his cases.²¹ History of infection or febrile illness was associated with an increased risk of DKA [odds ratios 6.50 (2.06 to 20.53) and 1.87 (1.05 to 3.33)].¹³ In our subjects, 6.9% failed to take their insulin among the ones with established T1DM, compared to 28% with poor compliance in the Poovazaghi study in India.²¹ Length of patient PICU stay was 3.26 (SD 3.50) days, longer than that in India which was about 1.94 days.²¹ No patients died during our study. However, Poovazaghi reported the mortality in established DKA and in new cases to be 7.1% and 12.7%, respectively.²¹

In conclusion, physicians must be aware that diabetic ketoacidosis may be the first clinical manifestation observed in T1DM. Increased awareness of T1DM in the medical and lay communities is needed to decrease the incidence of life-threatening complications.

References

1. DIAMOND Project Group. Incidence and trends of childhood type 1 diabetes worldwide 1990-1999. *Diabet Med.*

- 2006;23:857-66.
- EURODIAB ACE Study Group. Variation and trends in incidence of childhood diabetes in Europe. EURODIAB ACE Study Group. *Lancet*. 2000;355:873-6.
 - Wolfsdorf J, Craig M, Daneman D, Dunger D, Edge J, Lee W, et al. ISPAD clinical practice consensus guidelines 2009 compendium: diabetic ketoacidosis. *Pediatr Diabetes*. 2009;10:118-33.
 - Klingensmith GJ, Tamborlane WV, Wood J, Haller MJ, Silverstein J, Cengiz E. Diabetic ketoacidosis at diabetes onset: still an all too common threat in youth. *J Pediatr*. 2013;162:330-4.
 - Levy-Marchal C, Papoz L, de Beaufort C, Doutreix J, Froment V, Voirin J, et al. Clinical and laboratory features of type 1 diabetic children at the time of diagnosis. *Diabet Med*. 1992;9:279-84.
 - Komulainen J, Lounamaa R, Knip M, Kaprio EA, Akerblom HK. Ketoacidosis at the diagnosis of type 1 (insulin dependent) diabetes mellitus is related to poor residual beta cell function. Childhood Diabetes in Finland Study Group. *Arch Dis Child*. 1996;75:410-5.
 - Levy-Marchal C, Patterson CC, Green A. Geographical variation of presentation at diagnosis of type 1 diabetes in children: the EURODIAB study. *European and Diabetologia*. 2001;44:B75-80.
 - Hanas R, Lindgren F, Lindblad B. Diabetic ketoacidosis and cerebral oedema in Sweden—a 2-year paediatric population study. *Diabet Med*. 2007;24:1080-5.
 - Rodacki M, Pereira J, Nabuco De Oliveira Am, Barone B, Mac Dowell R, Perricelli P, et al. Ethnicity and young age influence the frequency of diabetic ketoacidosis at the onset of type 1 diabetes. *Diabetes Res Clin Pract*. 2007;78:259-62.
 - Neu A, Ehehalt S, Willasch A, Kehr M, Hub R, Ranke MB. Varying clinical presentations at onset of type 1 diabetes mellitus in children—epidemiological evidence for different subtypes of the disease? *Pediatr Diabetes*. 2001;2:147-53.
 - Oyarzabal-Irigoyen M, García Cuartero B, Barrio Castellanos R, Torres Lacruz M, Gómez Gila AL, González Casado I, et al. Ketoacidosis at onset of type 1 diabetes mellitus in pediatric age in Spain and review of the literature. *Pediatr Endocrinol Rev*. 2012;9:669-71.
 - Neu A, Willasch A, Ehehalt S, Hub R, Ranke MB, Becker SA, et al. Ketoacidosis at onset of type 1 diabetes mellitus in children—frequency and clinical presentation. *Pediatr Diabetes*. 2003;4:77-81.
 - Usher-Smith JA, Thompson MJ, Sharp SJ, Walter FM. Factors associated with the presence of diabetic ketoacidosis at diagnosis of diabetes in children and young adults: a systematic review. *BMJ*. 2011;343:d4092.
 - Onyriuka AN, Ifebi E. Ketoacidosis at diagnosis of type 1 diabetes in children and adolescents: frequency and clinical characteristics. *J Diabetes Metab Disord*. 2013;12:47:1-5.
 - Blanc N, Lucidarme N, Tubiana-Rufi N. Factors associated to ketoacidosis at diagnosis of type 1 diabetes in children. *Archives de Pédiatrie*. 2003;10:320-5.
 - Ting WH, Huang CY, Lo FS, Hung CM, Chan CJ, Li HJ, et al. Clinical and laboratory characteristics of type 1 diabetes in children and adolescents: experience from a medical center. *Acta Paediatr Taiwan*. 2007;48:119-24.
 - Xin Y, Yang M, Chen XJ, Tong YJ, Zhang LH. Clinical features at the onset of childhood type 1 diabetes mellitus in Shenyang, China. *J Paediatr Child Health*. 2010;46:171-5.
 - Alanani NMK, Alsulaimani AA. Epidemiological pattern of newly diagnosed children with type 1 diabetes mellitus, Taif, Saudi Arabia. *Scientific World J*. 2013;1-9.
 - Razavi Z. Frequency of ketoacidosis in newly diagnosed type 1 diabetic children. *Oman Med J*. 2010; 25:114-7.
 - Poovazhagi V, Senguttuvan P, Padmaraj R. Outcome of acute renal failure in children with diabetic keto acidosis (DKA). *Pediatr Oncall J*. 2011;8:63-70.
 - Poovazhagi V. Diabetic ketoacidosis among children with established diabetes from a pediatric tertiary care institute. *Int J Diabetes Dev Ctries*. 2014;34:95-9.

Diabetic ketoacidosis in children an 11-year retrospective in Surabaya, Indonesia

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Annunziata Lapolla, Flavia Amaro, Daniela Bruttomesso, Paolo Di Bartolo et al. "Diabetic Ketoacidosis: A Consensus Statement of the Associazione Medici Diabetologi (AMD), Società Italiana di Diabetologia (SID), Società Italiana di Endocrinologia e Diabetologia Pediatrica (SIEDP)", *Nutrition, Metabolism and Cardiovascular Diseases*, 2020
Publication 2%
- 2 www.smj.org.sa
Internet Source 2%
- 3 "Acute kidney injury and diabetic ketoacidosis in pediatric patients: risk factors", *Archivos Argentinos de Pediatría*, 2020
Publication 1%
- 4 pubmed.ncbi.nlm.nih.gov
Internet Source 1%
- 5 Mona M. Hassan, Mona H. Hafez, Hala M. Koura, Nagwa A. Mohamed, Mostafa A. Barakat, Abeer M. Osman. "Etiology, risk

factors, clinical features, and pitfalls in the management of newly diagnosed diabetic children and adolescents", Medical Research Journal, 2015

Publication

6

library.gunadarma.ac.id

Internet Source

1 %

7

Alphonsus N Onyiriuka, Emeka Ifebi. "Ketoacidosis at diagnosis of type 1 diabetes in children and adolescents: frequency and clinical characteristics", Journal of Diabetes & Metabolic Disorders, 2013

Publication

1 %

8

Helen Siyoum Atkilt, Muluken Gizaw Turago, Balewgizie Sileshi Tegegne. "Clinical Characteristics of Diabetic Ketoacidosis in Children with Newly Diagnosed Type 1 Diabetes in Addis Ababa, Ethiopia: A Cross-Sectional Study", PLOS ONE, 2017

Publication

1 %

9

www.karger.com

Internet Source

1 %

10

Mawaddah Batwa, Lujain Alharthi, Reem Ghazal, Meaad Alsulami, Rahaf Slaghour, Renad Aljuhani, Abdullah Bakhsh. "Diabetic Ketoacidosis at the Onset of Type 1 Diabetes Mellitus Among Children and Adolescents in Jeddah, Saudi Arabia: A Study From the

1 %

Emergency Department", Cureus, 2022

Publication

11

diabetestalk.net

Internet Source

1 %

12

D. B. Dunger, M. A. Sperling, C. L. Acerini, D. J. Bohn et al. "European Society for Paediatric Endocrinology/Lawson Wilkins Pediatric Endocrine Society Consensus Statement on Diabetic Ketoacidosis in Children and Adolescents", PEDIATRICS, 2004

Publication

1 %

13

Marcel Simon, Stephan Braune, Azien Laqmani, Maria Metschke, Christoph Berliner, Maria Kalsow, Hans Klose, Stefan Kluge. "Value of Computed Tomography of the Chest in Subjects With ARDS: A Retrospective Observational Study", Respiratory Care, 2016

Publication

1 %

14

orthorsds-fkua.org

Internet Source

1 %

15

Yun Jeong Lee, Young Ah Lee, Jae Hyun Kim, Hye Rim Chung, Min Jeong Gu, Ji Young Kim, Choong Ho Shin. "The durability and effectiveness of sensor-augmented insulin pump therapy in pediatric and young adult patients with type 1 diabetes", Annals of Pediatric Endocrinology & Metabolism, 2020

Publication

1 %

16 Naglaa Mohamed Kamal Alanani, Adnan Amin Alsulaimani. "Epidemiological Pattern of Newly Diagnosed Children with Type 1 Diabetes Mellitus, Taif, Saudi Arabia", The Scientific World Journal, 2013 1 %
Publication

17 Puruhito. "Availability and Limits of Intermediate Cardiovascular Technology", Artificial Organs, 2008 1 %
Publication

18 Ayse Nurcan Cebeci, Ayla Guven, Heves Kirmizibekmez, Metin Yildiz, Fatma Dursun. "Clinical features and management of diabetic ketoacidosis in different age groups of children: children less than 5 years of age are at higher risk of metabolic decompensation", Journal of Pediatric Endocrinology and Metabolism, 2012 1 %
Publication

19 V. Poovazhagi. "Factors associated with mortality in children with diabetic ketoacidosis (DKA) in South India", International Journal of Diabetes in Developing Countries, 2015 <1 %
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Diabetic ketoacidosis in children an 11-year retrospective in Surabaya, Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
