

---

## Fwd: [Kemas] Article Review Request

1 message

---

**trias mahmudiono** <triasmahmudiono@gmail.com>  
To: trias-m <trias-m@fkm.unair.ac.id>, nur.sahila-2018@fkm.unair.ac.id

Tue, Jun 6, 2023 at 6:06 PM

----- Forwarded message -----

From: **trias mahmudiono** <triasmahmudiono@gmail.com>  
Date: Tue, Jun 6, 2023 at 6:06 PM  
Subject: Re: [Kemas] Article Review Request  
To: Jurnal Kesmas <jurnalkesmas2.ui@gmail.com>

ini bisa di print termasuk bukti kerja mereview di Kesmas UI yang sudah Scopus

On Mon, Feb 13, 2023 at 2:33 PM Jurnal Kesmas <jurnalkesmas2.ui@gmail.com> wrote:  
Terima kasih banyak Pak. Sehat selalu

On Mon, Feb 13, 2023 at 12:45 PM trias mahmudiono <triasmahmudiono@gmail.com> wrote:  
Silahkan

On Tue, Feb 7, 2023 at 2:44 PM Jurnal Kesmas <jurnalkesmas2.ui@gmail.com> wrote:  
Dear Trias Mahmudiono, [S.KM](#), [M.PH](#) (Nutr.), GCAS, Ph.D

I believe that you would serve as an excellent reviewer of the manuscript titled "The Association of Obesity with Asthma Risk in Indonesian Adults" which has been submitted to Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal). The submission's abstract is inserted below, and I hope that you will consider undertaking this important task for us.

The review itself is due 2023-02-10.

Thank you for considering this request.

Best Regards,  
Dewi Susanna

### **The Association of Obesity with Asthma Risk in Indonesian Adults**

#### **Abstract**

Obesity and asthma are both global public health challenges. Mounting evidence has suggested that obesity may increase asthma risk in adults; however, the association by sex remains uncertain. This study examined the association of obesity with asthma risk in adults' Indonesian men and women. Data were obtained from the 2018 Indonesia Basic Health Research. The analysis included 299,837 men and 333,218 women aged  $\geq 18$  years. Asthma was identified by the self-report of a doctor's diagnosis. Obesity was defined as a Body Mass Index (BMI)  $\geq 30$  kg/m<sup>2</sup>. A logistic regression was used for data analysis. Asthma prevalence was 2.5% in men and 2.8% in women. The adjusted odds ratio (aOR) of having asthma for obese women and obese men were 1.25, 95% CI = 1.21-1.34 and 1.14, 95% CI = 1.05-1.25, respectively. In conclusion, asthma prevalence was slightly higher in women compared with men. A small increase in the odds of having asthma-related obesity was found in both sexes. The adjusted odds of having asthma-related obesity were higher in women than in men. The causality between asthma and obesity warrants further studies with a longitudinal design.

**Keywords:** Asthma, Obesity, Adults, Indonesia

--

**Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal)** is published by the Faculty of Public Health Universitas Indonesia since August 2006 and has been accredited by the Director General of Higher Education in July 2009 and re-accredited in 2012 (No.56/DIKTI/Kep/2012) and 2017 (No.32a/E/KPT/2017). Kesmas is indexed in Scopus and SINTA 1 and published quarterly in February, May, August and November.

<http://journal.fkm.ui.ac.id/kesmas>

**Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal)**

Faculty of Public Health  
Universitas Indonesia  
G301 Building G 3rd Floor  
Kampus Baru UI Depok 16424  
Phone: +62815 1141 6600

--

**Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal)** is published by the Faculty of Public Health Universitas Indonesia since August 2006 and has been accredited by the Director General of Higher Education in July 2009 and re-accredited in 2012 (No.56/DIKTI/Kep/2012) and 2017 (No.32a/E/KPT/2017). Kesmas is indexed in Scopus and SINTA 1 and published quarterly in February, May, August and November.

<http://journal.fkm.ui.ac.id/kesmas>

**Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal)**

Faculty of Public Health  
Universitas Indonesia  
G301 Building G 3rd Floor  
Kampus Baru UI Depok 16424  
Phone: +62815 1141 6600



UNIVERSITAS  
INDONESIA

Veritas, Probatum, Justitia

FAKULTAS

KESEHATAN  
MASYARAKAT

**Kesmas**

Jurnal Kesehatan Masyarakat Nasional  
(National Public Health Journal)

SERT-03/UN2.F10.D.13/PPM.00.04/II/2023

# CERTIFICATE

awarded to

**Trias Mahmudiono, S.KM, M.PH (Nutr.), GCAS, Ph.D**

as a reviewer in

**Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal)**

with the title of

**“The Association of Obesity and Asthma Risk in Indonesian Adults”**

period of **February 2023**, published in **Volume 18, Issue 1, February 2023**

Editor-in-Chief of

Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal)



Prof. Dr. Dewi Susanna, dra., MS.

NIP. 196312021988032002

# Obesity and Asthma Risk in Indonesian Adults: Findings from the 2018 Indonesia Basic Health Research

Hoirun Nisa

Department of Public Health, Faculty of Health Sciences, Universitas Islam Negeri (UIN) Syarif Hidayatullah Jakarta, South Tangerang, Indonesia

## Abstract

Obesity and asthma are both global public health challenges. Mounting evidence suggests that obesity may increase asthma risk in adults; however, the association by sex remains uncertain. This study examined the association of obesity with asthma risk in Indonesian adult men and women. Data were obtained from the 2018 Indonesia Basic Health Research. The analysis included 299,837 men and 333,218 women aged  $\geq 18$  years. Asthma was identified by the self-report of a doctor's diagnosis. Obesity was defined as a body mass index  $\geq 30$  kg/m<sup>2</sup>. A logistic regression was used for data analysis. Asthma prevalence was 2.7% (2.5% in men and 2.8% in women) and was higher in obese than non-obese adults. The adjusted odds ratio (aOR) of having asthma-related obesity was 1.22, 95% Confidence Interval (CI) = 1.17–1.28 (aOR = 1.25, 95% CI = 1.21–1.34 in women and aOR = 1.14, 95% CI = 1.05–1.25 in men). In conclusion, asthma prevalence is relatively low in Indonesian adults and slightly higher in women than men. Both men and women have a slight increase in the odds of having asthma-related obesity. Longitudinal studies are needed to understand better the causality association of obesity with asthma in adults.

**Keywords:** adults, asthma, Indonesia, obesity

## Introduction

Asthma affects about 262 million people globally,<sup>1</sup> and is among adults' most frequent chronic respiratory diseases.<sup>2</sup> The recent findings from the Global Asthma Network Phase I study, a multi-country population-based survey, reported that asthma prevalence among adults differed significantly within and between countries.<sup>3</sup> Although its global prevalence has decreased,<sup>4</sup> the burden of asthma remains substantial on healthcare systems and the community.<sup>5,6</sup> Findings from a systematic analysis of the 2017 global burden of disease study indicated that asthma was the second most common cause of death among chronic respiratory diseases.<sup>4</sup> In fact, the majority of deaths from asthma occurred in low and lower-middle-income countries.<sup>7,8</sup> In Indonesia, the prevalence of self-reported, doctor-diagnosed asthma had significantly increased from 1.9% in 2007,<sup>9</sup> to 2.4% in 2018.<sup>10</sup> In 2018, asthma prevalence among Indonesian adults ranged from 2.2% to 5.1%.<sup>10</sup> In addition, asthma prevalence in 2018 differed by sex, with a prevalence of 2.3% in men and 2.5% in women.<sup>10</sup>

Obesity has become another global public health concern in adults.<sup>11</sup> In Indonesia, obesity prevalence is high-

er in females than males, and obesity prevalence in adults has increased over the past two decades.<sup>9,10</sup> Obesity has been associated with many chronic diseases, including asthma.<sup>12</sup> The increase in obesity prevalence has implicated obesity-related diseases and health economics, subsequently burdening the healthcare system.<sup>13</sup> A review of obesity and asthma in adults reported a higher asthma prevalence in obese women and a predominantly positive association in women.<sup>14</sup> In addition, increased odds of having asthma were shown among obese adults in some cross-sectional studies.<sup>15-17</sup> A meta-analysis of Mendelian randomization analysis suggested an increased risk of asthma associated with a higher genetically predicted body mass index (BMI) in European adults; however, individual studies showed mixed results.<sup>18</sup> The role of sex in the obesity-asthma association remains controversial. For example, several studies reported a significant association only in females,<sup>19,20</sup> but another found a similar association between sexes.<sup>21</sup> Other studies found a stronger association in females than males.<sup>22,23</sup> Two population-based Indonesian studies showed positive associations of obesity with asthma in men and working women,<sup>24</sup> and non-working

**Correspondence\*:** Hoirun Nisa, Department of Public Health, Faculty of Health Sciences, Universitas Islam Negeri (UIN) Syarif Hidayatullah Jakarta, Kertamukti Street No. 5, Ciputat, South Tangerang, Banten 15419, Indonesia, E-mail: hoirun.nisa@uinjkt.ac.id, Phone: +62 21-74716718

Received : January 30, 2023  
Accepted : February 21, 2023  
Published : February 27, 2023



women.<sup>25</sup>

Given the increasing prevalence of both obesity and asthma in Indonesia and the prevalence differed by sex,<sup>9,10</sup> it is essential to assess the association between these conditions for preventive measures. Many studies in other countries reported inconclusive results regarding the potential role of sex on the association of obesity with adult asthma<sup>20,22,23</sup>; however, evidence from Indonesia is limited. A better understanding of the sex differences in asthma prevalence in adult populations and its association with obesity can benefit its management. Thus, this study aimed to examine the association of obesity with asthma in Indonesian adults, using national data from the 2018 Indonesia Basic Health Research.

## Method

Data were derived from the 2018 Indonesia Basic Health Research,<sup>10</sup> a national survey with multi-purposes to evaluate Indonesian health status. The 2018 Indonesia Basic Health Research was a cross-sectional survey conducted in 34 provinces, 416 districts, and 98 cities in Indonesia. Data were collected in March 2018 by the Indonesian Ministry of Health. To assess the asthma situation, the 2018 Indonesia Basic Health Research interviewed a representative sample of 1,017,290 individuals of all ages consisting of 510,714 males and 506,576 females.

In this study, the inclusion criteria included individuals aged  $\geq 18$  years with information on doctor-diagnosed asthma. A data set request proposal was submitted to the Data Management Laboratory, the Indonesian Ministry of Health, to obtain data set according to the inclusion criteria and proposed variables. Of the 641,951 individuals who met the inclusion criteria, 8,896 were excluded due to missing data on BMI and other covariates. The final analytic sample included 633,055 adults, of which 299,837 were men and 333,218 were women.

The National Institute of Health Research and Development Ethics Committee, the Indonesian Ministry of Health, approved the 2018 Indonesia Basic Health Research with the letter LB.02.01/3/KE.024/2018. Written informed consent was obtained from participants by interviewers of the National Institute of Health Research and Development of the Indonesian Ministry of Health.

Asthma was determined through a question, "Have you ever been diagnosed with asthma by a doctor?". Asthma prevalence was calculated by dividing the number of adults with asthma by the total number of corresponding individuals. Anthropometrics data were obtained by measuring height and weight. The BMI was estimated by dividing body weight (kg) by height (m<sup>2</sup>). Obesity was determined as a BMI  $\geq 30$  kg/m<sup>2</sup> according to World Health Organization's (WHO) classification.<sup>26</sup>

Information on covariates (age, residential area, marital status, education level, smoking status, physical activity, and history of hypertension and diabetes) was obtained from the 2018 Indonesia Basic Health Research questionnaire responses. Age was grouped into 18–44 years, 45–64 years, or  $\geq 65$  years. The residential area was classified as rural or urban areas. Marital status was categorized into single, married, or divorced. Education level was grouped into primary if the highest education was middle school, secondary if the highest education was high school, or higher education if the highest education was university. Information on smoking status was ascertained by asking individuals whether they had smoked daily in the past month. Smoking status was then categorized as never, former, or current smoking. Physical activity was categorized into active if individuals performed moderate physical activities for  $\geq 150$  minutes/week or vigorous physical activities for  $\geq 75$  minutes/week, or less active if individuals did not meet the WHO recommendation.<sup>27</sup> Self-report of a doctor's diagnosis was used to define hypertension and diabetes.

Data analyses were independently done for both sexes. Data were presented in mean  $\pm$  standard deviation for continuous variables and proportion for categorical variables. Microsoft Office Excel 365 for the web was used to draw a figure on the distribution of obesity according to asthma status. Variables significantly associated with asthma at a p-value of less than 0.05 in bivariate analysis were selected as covariates in multivariate analysis. Multivariate analyses were performed using a logistic regression test to examine the association of obesity with asthma in adults. In multivariate analysis, adjustment was made for age, residential area, marital status, education level, smoking status, physical activity, and records of hypertension and diabetes mellitus. The odd ratios (OR) and 95% confidence intervals (CIs) were presented. Statistical significance was declared if a two-sided p-value was less than 0.05.

## Results

Asthma prevalence in adults was 2.7%, with a mean age of 42.9 years. Over half of the adults were women, living in rural areas, married, less active, never smokers, and attained primary education. The adults' mean BMI was 23.9 kg/m<sup>2</sup>, and 12.6% of adults were obese. Adults with a records of hypertension and diabetes were 48.8% and 2.3%, respectively (Table 1). Asthma prevalence was higher in women (2.8%) than in men (2.5%). The mean age of men and women was 43.0 and 42.8 years, respectively. More than half of the men and women lived in rural areas, were married, and had primary education. In men, the proportions of current smokers, less active, and records of hypertension were relatively high, but the proportions of obesity and diabetes were relatively low. Most

**Table 1. Participants' Characteristics by Sex**

| Variable                 | Category         | All (n = 653,055) | Male (n = 299,837) | Female (n = 353,218) |
|--------------------------|------------------|-------------------|--------------------|----------------------|
|                          |                  | n (%)             | n (%)              | n (%)                |
| Asthma                   | No               | 616,023 (97.3)    | 292,276 (97.5)     | 323,747 (97.2)       |
|                          | Yes              | 17,032 (2.7)      | 7,561 (2.5)        | 9,471 (2.8)          |
| Sex                      | Male             | 299,837 (47.4)    | -                  | -                    |
|                          | Female           | 353,218 (52.6)    | -                  | -                    |
| Age, year*               |                  | 42.9±14.9         | 43.0±14.9          | 42.8±14.9            |
| Age group, year          | 18-44            | 360,820 (57.0)    | 168,784 (56.3)     | 192,036 (57.6)       |
|                          | 45-64            | 216,183 (34.1)    | 104,756 (34.9)     | 111,427 (33.4)       |
|                          | ≥65              | 56,052 (8.9)      | 26,297 (8.8)       | 29,755 (8.9)         |
| Residential area         | Rural            | 359,217 (56.7)    | 171,051 (57.0)     | 188,166 (56.5)       |
|                          | Urban            | 273,838 (43.3)    | 128,786 (43.0)     | 145,052 (43.5)       |
| Marital status           | Single           | 85,501 (13.5)     | 55,046 (18.4)      | 30,455 (9.1)         |
|                          | Married          | 484,568 (76.5)    | 250,831 (77.0)     | 253,737 (76.1)       |
|                          | Divorced         | 62,986 (9.9)      | 15,960 (4.7)       | 49,026 (14.7)        |
| Education level          | Primary          | 395,244 (62.4)    | 178,973 (59.7)     | 216,271 (64.9)       |
|                          | Secondary        | 174,492 (27.6)    | 92,350 (30.8)      | 82,142 (24.7)        |
|                          | Higher education | 63,319 (10.0)     | 28,514 (9.5)       | 34,805 (10.4)        |
| Physical activity        | Less active      | 452,499 (71.5)    | 168,585 (56.2)     | 283,914 (85.2)       |
|                          | Active           | 180,556 (28.5)    | 131,252 (43.8)     | 49,304 (14.8)        |
| Smoking status           | Never            | 397,677 (62.8)    | 76,680 (25.6)      | 320,997 (96.3)       |
|                          | Former           | 33,057 (5.2)      | 28,717 (9.6)       | 4,340 (1.3)          |
|                          | Current          | 202,321 (32.0)    | 194,440 (64.8)     | 7,881 (2.4)          |
| BMI, kg/m <sup>2</sup> * |                  | 23.9±4.6          | 23.0±4.0           | 24.7±4.9             |
| Obesity status           | No               | 553,057 (87.4)    | 278,626 (92.9)     | 274,431 (82.4)       |
|                          | Yes              | 79,998 (12.6)     | 21,211 (7.1)       | 58,787 (17.6)        |
| Hypertension             | No               | 324,371 (51.2)    | 132,469 (44.2)     | 191,902 (57.6)       |
|                          | Yes              | 308,684 (48.8)    | 167,368 (55.8)     | 141,316 (42.4)       |
| Diabetes                 | No               | 618,807 (97.7)    | 294,287 (98.1)     | 324,520 (97.4)       |
|                          | Yes              | 14,248 (2.3)      | 5,550 (1.9)        | 8,698 (2.6)          |

Notes: BMI = Body Mass Index, \*Data are presented as means±standard deviation.

women were never smokers and less active. In women, the proportions of obesity and hypertension were high, but the proportion of diabetes was low (Table 1).

Table 2 shows the characteristics of the participants based on their asthma status. Asthma prevalence was higher in rural, married, lower educated, and less active adults. Adults with asthma were likely to be older, had lower rates of hypertension and diabetes, and 23.1% of adults were current smokers. In men, the highest asthma prevalence was among current smokers. In contrast, the highest asthma prevalence in women was among never smokers.

Figure 1 presents the distribution of obesity according to asthma status. The obesity proportion was higher in adults with asthma than those without asthma. In addition, obesity was specifically higher in women with asthma than in men with asthma. Obesity was significantly associated with asthma regardless of sex.

Table 3 presents the OR associated with asthma according to obesity status. Obese adults had 1.23 times higher odds of having asthma than non-obese adults. The odds of having asthma associated with obesity differed between sexes. The odds of having asthma were 1.15 for obese men and 1.26 for obese women. Adjustment for

the covariates did not noticeably change the results. The multivariate-adjusted odds of having asthma were higher in obese women than in obese men (aOR (95% CI) = 1.25 (1.21–1.34) vs. 1.14 (1.05–1.25), p-value<0.05).

### Discussion

This study showed that asthma prevalence was relatively low in Indonesian adults and slightly higher in women than men. The prevalence of doctor-diagnosed asthma in this study was 2.5% in men and 2.8% in women. These findings were lower than the respective prevalence of 6.7% and 7.0% among men and women workers in Indonesia, which defined asthma based on symptoms.<sup>24</sup> In addition, several studies have reported that the prevalence of adult asthma differs between men and women.<sup>4,17,22,23,28,29</sup> Results from a national survey of Iranian adults in 2018 reported a variation in asthma prevalence between sexes and asthma definitions. In that study, asthma prevalence was higher in men using the European Community Respiratory Health Survey questionnaire but higher in women when using self-report of doctor-diagnosis.<sup>29</sup> Consistent with the current findings, many studies have shown a higher prevalence among women than men.<sup>4,17,22,23,28</sup> It has been suggested that

Table 2. Participants' Characteristics by Asthma Status


| Variable                  | Category         | All (n = 633,055) |                |          | Male (n = 299,837) |                |          | Female (n = 333,218) |                |          |
|---------------------------|------------------|-------------------|----------------|----------|--------------------|----------------|----------|----------------------|----------------|----------|
|                           |                  | Asthma            | Without Asthma | p-value* | Asthma             | Without Asthma | p-value* | Asthma               | Without Asthma | p-value* |
|                           |                  | n (%)             | n (%)          |          | n (%)              | n (%)          |          | n (%)                | n (%)          |          |
| Sex                       | Male             | 7,561 (44.4)      | 292,276 (47.4) | <0.001   | -                  | -              | -        | -                    | -              | -        |
|                           | Female           | 9,471 (55.6)      | 323,747 (52.6) |          |                    |                |          |                      |                |          |
| Age, year**               |                  | 47.0±16.2         | 42.8±14.8      | <0.001   | 49.8±16.7          | 42.8±14.8      | <0.001   | 44.7±15.4            | 42.8±14.9      | <0.001   |
| Age group, year           | 18-44            | 7,914 (46.5)      | 352,906 (57.3) |          | 2,911 (38.5)       | 165,873 (56.8) |          | 5,003 (52.9)         | 187,033 (57.8) |          |
|                           | 45-64            | 6,433 (37.8)      | 209,750 (34.0) | <0.001   | 3,077 (40.7)       | 101,679 (34.8) | <0.001   | 3,356 (35.4)         | 108,071 (33.4) | <0.001   |
|                           | 65               | 2,685 (15.8)      | 53,367 (8.7)   |          | 1,573 (20.8)       | 24,724 (8.5)   |          | 1,112 (11.7)         | 28,643 (8.8)   |          |
| Residential area          | Rural            | 9,287 (54.5)      | 349,930 (56.8) | <0.001   | 4,305 (56.9)       | 166,746 (57.1) | 0.845    | 4,982 (52.6)         | 183,184 (56.6) | <0.001   |
|                           | Urban            | 7,745 (45.5)      | 266,093 (43.2) |          | 3,256 (43.1)       | 125,530 (42.9) |          | 4,489 (47.4)         | 140,563 (43.4) |          |
| Marital status            | Single           | 1,738 (10.2)      | 83,765 (13.6)  |          | 976 (12.9)         | 54,070 (18.5)  |          | 760 (8.0)            | 29,695 (9.2)   |          |
|                           | Married          | 13,088 (76.8)     | 471,480 (76.5) | <0.001   | 5,982 (79.1)       | 224,849 (76.9) | <0.001   | 7,106 (75.0)         | 246,531 (76.2) | <0.001   |
|                           | Divorced         | 2,208 (13.0)      | 60,778 (9.9)   |          | 603 (8.0)          | 13,357 (4.6)   |          | 1,605 (16.9)         | 47,421 (14.6)  |          |
| Education level           | Primary          | 11,284 (66.3)     | 383,960 (62.3) |          | 5,106 (67.5)       | 173,867 (59.5) |          | 6,178 (65.2)         | 210,093 (64.9) |          |
|                           | Secondary        | 4,010 (23.5)      | 170,482 (27.7) | <0.001   | 1,783 (23.6)       | 90,567 (31.0)  | <0.001   | 2,227 (23.5)         | 79,915 (24.7)  | 0.003    |
|                           | Higher education | 1,738 (10.2)      | 61,581 (10.0)  |          | 672 (8.9)          | 27,842 (9.5)   |          | 1,066 (11.3)         | 33,739 (10.4)  |          |
| Physical activity         | Less active      | 13,254 (77.8)     | 439,245 (71.3) | <0.001   | 4,965 (65.7)       | 163,620 (56.0) | <0.001   | 8,289 (87.5)         | 275,625 (85.1) | <0.001   |
|                           | Active           | 3,778 (22.2)      | 176,778 (28.7) |          | 2,596 (34.3)       | 128,656 (44.0) |          | 1,182 (12.5)         | 48,122 (14.9)  |          |
| Smoking status            | Never            | 11,242 (66.0)     | 386,435 (62.7) |          | 2,278 (30.1)       | 74,402 (25.5)  |          | 8,964 (94.6)         | 312,033 (96.4) |          |
|                           | Former           | 1,848 (10.9)      | 31,209 (5.1)   | <0.001   | 1,606 (21.2)       | 27,111 (9.3)   | <0.001   | 242 (2.6)            | 4,098 (1.3)    | <0.001   |
|                           | Current          | 3,942 (23.1)      | 198,397 (32.2) |          | 3,677 (48.6)       | 190,763 (65.3) |          | 265 (2.8)            | 7,616 (2.4)    |          |
| BMI, kg/m <sup>2</sup> ** |                  | 23.8±5.2          | 23.9±4.6       | 0.019    | 22.5±4.4           | 23.0±4.0       | <0.001   | 24.9±5.5             | 24.7±4.9       | 0.004    |
| Hypertension              | No               | 9,683 (56.9)      | 314,688 (51.1) | <0.001   | 4,002 (52.9)       | 128,467 (44.0) | <0.001   | 5,681 (60.0)         | 186,221 (57.5) | <0.001   |
|                           | Yes              | 7,349 (43.1)      | 301,335 (48.9) |          | 3,559 (47.1)       | 163,809 (56.0) |          | 3,790 (40.0)         | 137,526 (42.5) |          |
| Diabetes mellitus         | No               | 16,347 (96.0)     | 602,460 (97.8) | <0.001   | 7,289 (96.4)       | 286,998 (98.2) | <0.001   | 9,058 (95.6)         | 315,462 (97.4) | <0.001   |
|                           | Yes              | 685 (4.0)         | 13,563 (2.2)   |          | 272 (3.6)          | 5,278 (1.8)    |          | 413 (4.4)            | 8,285 (2.6)    |          |

Notes: BMI = Body Mass Index, \*p-values were calculated by t-test for continuous variables and the Chi-square test for categorical variables, \*\*Data are presented as means±standard deviation.

the mixed results in asthma prevalence may be partially due to differences in diagnostic criteria.<sup>30</sup> A review article on the prevalence of adult asthma with a wide variation of asthma definitions in Asian countries, including Indonesia, showed that the asthma prevalence in adults varied from 0.7% to 11.9%.<sup>31</sup>

This study showed a slight increase in the odds of having asthma in obese adults regardless of sex, and the associations were statistically significant. The increase in the odds of having asthma was higher in obese women compared to obese men. However, the association for men should cautiously be interpreted because the 95% CI of the OR for asthma in obese men reached a null association. Obesity is a significant risk factor for developing asthma, worsening asthma symptoms, and controlling poor asthma.<sup>5,6</sup> Although this study found a slight increase in the odds of having asthma in obese adults, the prevalence of both asthma and obesity is rising.<sup>9,10</sup> Thus, asthma-related obesity may cause an immense burden of chronic disease prevention in the future public healthcare systems.

Obesity has many negative impacts on health,<sup>12</sup> and the risk of asthma associated with obesity may differ by sex.<sup>32</sup> The association of obesity with asthma in men remains unclear.<sup>23,32</sup> In addition, previous studies found



Notes: The association between obesity and asthma was evaluated using Chi-square tests, \*p-value<0.05.

Figure 1. Distribution of Obesity according to Asthma Status

an increase in the odds of having asthma associated with obesity in adults after adjusting for potential confounders; however, results for separate analyses were not shown for men and women.<sup>15,17</sup> Several other studies reported an increased risk of adult asthma associated with obesity only in women and not men.<sup>19,20</sup> Similar to the current findings, a few national-level cross-sectional stud-

**Table 3. Odds Ratio (95% Confidence Interval) Associated with Asthma According to Obesity Status**

| Variable | Category  | Unadjusted OR (95% CI) | p-value | Adjusted OR (95% CI) | p-value |
|----------|-----------|------------------------|---------|----------------------|---------|
| All      | Non-obese | 1.00 (reference)       | -       | 1.00 (reference)     | -       |
|          | Obese     | 1.25 (1.21–1.31)       | <0.001  | 1.22 (1.17–1.28)*    | <0.001  |
| Male     | Non-obese | 1.00 (reference)       | -       | 1.00 (reference)     | -       |
|          | Obese     | 1.15 (1.06–1.25)       | 0.001   | 1.14 (1.05–1.25)**   | 0.002   |
| Female   | Non-obese | 1.00 (reference)       | -       | 1.00 (reference)     | -       |
|          | Obese     | 1.26 (1.20–1.33)       | <0.001  | 1.25 (1.21–1.34)**   | <0.001  |

**Notes:** OR = Odds Ratio, CI = Confidence Interval

\*Adjusted with sex, age, residence area, marital status, educational attainment, physical activity, smoking status, and history of hypertension and diabetes mellitus.

\*\*Adjusted with age, residence area, marital status, educational attainment, physical activity, smoking status, and history of hypertension and diabetes mellitus.

ies reported a higher increase in the odds of having asthma associated with obesity in women than men.<sup>16,22,23</sup> A meta-analysis of 174,556 adults from 51 nationally representative surveys in which asthma prevalence remains common showed that obese men and women had a similarly high risk for doctor-diagnosed asthma.<sup>33</sup> Results from a prospective study including 523,245 adults found a higher risk of asthma in obese women compared to obese men after an 11-year follow-up.<sup>21</sup> In addition, a retrospective cohort study on more than 5 million adults with a BMI  $\geq 23.0$  kg/m<sup>2</sup> showed that the risk of adult asthma gradually increased with increasing BMI, and the increased risk of adult asthma associated with obesity was higher in women than men.<sup>34</sup>

The mechanisms for sex differences in asthma prevalence in adults remain unclear. It has been suggested that sex hormones may cause women a higher asthma risk after puberty.<sup>32</sup> In addition, weight gain may reduce the airway, resulting in airway obstruction in women with smaller airways than men.<sup>35</sup> Another mechanism is probably due to sex differences in the inflammatory response. A higher level of inflammation associated with obesity in obese women than in obese men may explain a higher risk of having asthma for women.<sup>36</sup>

This study had some limitations. First, the causality between obesity and adult asthma could not be determined due to the cross-sectional design. Second, recall bias might occur because the definition of asthma relies on self-reporting a doctor’s diagnosis. However, many epidemiologic studies have used this self-report of doctor-diagnosed asthma; thus, it is a standard definition used in asthma studies. Third, the large sample size of this study tended to detect significant associations, and a low asthma prevalence in both sexes precluded further stratification analysis from assessing the association. Nonetheless, a slight increase in the odds of having asthma in obese adults might be considered to prevent comorbidity of the two conditions and alleviate the burden

of chronic disease in future public healthcare systems.

### Conclusion

This study of Indonesian adults shows that asthma prevalence is relatively low and slightly higher in women. Obesity is associated with asthma in both sexes, and the increase in the odds of having asthma is higher in women than men. Longitudinal studies are needed to understand better the causality association of obesity with asthma in adults.

### Abbreviations

BMI: Body Mass Index; WHO: World Health Organization; OR: Odds Ratio; CI: Confidence Interval.

### Ethics Approval and Consent to Participate

This study was approved by the Ethics Committee of the National Institute of Health Research and Development, the Ministry of Health of the Republic of Indonesia.

### Competing Interest

The author declares that there is no significant competing financial, professional, or personal interest that might have affected the performance or presentation of the work described in this manuscript.

### Availability of Data and Materials

The data used in this study are not publicly available. A reasonable request for the dataset can be sent to the corresponding author.

### Authors’ Contribution

HN was responsible for the entire process, including conceptualization, data analysis, writing, and revising the manuscript.

### Acknowledgment

This study was supported by the Indonesia National Institute of Health and Research Development, the Ministry of Health of the Republic of Indonesia.



## References

1. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2020; 396 (10258): 1204–22.
2. Global Asthma Network. The Global Asthma Report 2022. *Int J Tuberc Lung Dis*. 2022; 26 (1): 1-104.
3. Mortimer K, Lesosky M, García-Marcos L, Asher MI, Pearce N, Ellwood E, et al. The burden of asthma, hay fever and eczema in adults in 17 countries: GAN Phase I study. *Eur Respir J*. 2022; 60 (3).
4. GBD Chronic Respiratory Disease Collaborators. Prevalence and attributable health burden of chronic respiratory diseases, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Respir Med*. 2020; 8 (6): 585-96.
5. Nunes C, Pereira AM, Morais-Almeida M. Asthma costs and social impact. *Asthma Res Pract*. 2017; 3: 1.
6. Czira A, Turner M, Martin A, Hinds D, Birch H, Gardiner F, et al. A systematic literature review of burden of illness in adults with uncontrolled moderate/severe asthma. *Respir Med*. 2022; 191: 106670.
7. Meghji J, Mortimer K, Agusti A, Allwood BW, Asher I, Bateman ED, et al. Improving lung health in low-income and middle-income countries: from challenges to solutions. *Lancet*. 2021; 397 (10277): 928-40.
8. Mortimer K, Reddel HK, Pitrez PM, Bateman ED. Asthma management in low and middle income countries: case for change. *Eur Respir J*. 2022; 60 (5).
9. Kementerian Kesehatan Republik Indonesia. Laporan Nasional Riskesdas 2007. Jakarta: Lembaga Penerbit Badan Penelitian dan Pengembangan Kesehatan (LPB).
10. Kementerian Kesehatan Republik Indonesia. Laporan Nasional Riskesdas 2018. Jakarta: Lembaga Penerbit Badan Penelitian dan Pengembangan Kesehatan (LPB).
11. Jaacks LM, Vandevijvere S, Pan A, McGowan CJ, Wallace C, Imamura F, et al. The obesity transition: stages of the global epidemic. *Lancet Diabetes Endocrinol*. 2019; 7 (3): 231-40.
12. Endalifer ML, Diress G. Epidemiology, Predisposing Factors, Biomarkers, and Prevention Mechanism of Obesity: A Systematic Review. *J Obes*. 2020; 2020: 6134362.
13. Wang Y, Zhao L, Gao L, Pan A, Xue H. Health policy and public health implications of obesity in China. *Lancet Diabetes Endocrinol*. 2021; 9 (7): 446-61.
14. Khalid F, Holguin F. A review of obesity and asthma across the life span. *J Asthma*. 2018; 55 (12): 1286-300.
15. Souza ECC, Pizzichini MMM, Dias M, Cunha MJ, Matte DL, Karloh M, et al. Body mass index, asthma, and respiratory symptoms: a population-based study. *J Bras Pneumol*. 2020; 46 (1): e20190006.
16. Forno E, Han YY, Libman IM, Muzumdar RH, Celedón JC. Adiposity and Asthma in a Nationwide Study of Children and Adults in the United States. *Ann Am Thorac Soc*. 2018; 15 (3): 322-30.
17. Tarraf H, Aydin O, Mungan D, Albader M, Mahboub B, Doble A, et al. Prevalence of asthma among the adult general population of five Middle Eastern countries: results of the SNAPSHOT program. *BMC Pulm Med*. 2018; 18 (1): 68.
18. Larsson SC, Burgess S. Causal role of high body mass index in multiple chronic diseases: a systematic review and meta-analysis of Mendelian randomization studies. *BMC Med*. 2021; 19 (1): 320.
19. Greenblatt R, Mansour O, Zhao E, Ross M, Himes BE. Gender-specific determinants of asthma among U.S. adults. *Asthma Res Pract*. 2017; 3: 2.
20. Lampalo M, Majer M, Ferara N, Milošević M, Barišić Kutija M, Jukić I. Gender Differences in Relationship between Body Mass Index and Asthma. *Psychiatr Danub*. 2019; 31 (Suppl 5): 786-91.
21. Brumpton B, Langhammer A, Romundstad P, Chen Y, Mai XM. General and abdominal obesity and incident asthma in adults: the HUNT study. *Eur Respir J*. 2013; 41 (2): 323-9.
22. Kang M, Sohn SJ, Shin MH. Association between Body Mass Index and Prevalence of Asthma in Korean Adults. *Chonnam Med J*. 2020; 56 (1): 62-7.
23. Wang L, Wang K, Gao X, Paul TK, Cai J, Wang Y. Sex difference in the association between obesity and asthma in U.S. adults: Findings from a national study. *Respir Med*. 2015; 109 (8): 955-62.
24. Tana L. Determinants of asthma disease in workers of productive age in Indonesia Basic Health Research 2015. *Bulletin Penelitian Kesehatan*. 2018; 46 (1): 11-22.
25. Maulana AP, N.A.; Yovsyah. Effects of Obesity with the Risk of Asthma on Productive Age Female in Indonesia. *Jurnal Epidemiologi Kesehatan Indonesia*. 2020; 4 (1).
26. World Health Organization (WHO) Consultation on Obesity. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. Geneva: World Health Organization; 2019.
27. World Health Organization. WHO Guidelines on physical activity and sedentary behaviour. Geneva: World Health Organization; 2020.
28. Akiki Z, Saadeh D, Farah R, Hallit S, Sacre H, Hosseini H, et al. Asthma prevalence and associated factors among Lebanese adults: the first national survey. *BMC Pulm Med*. 2021; 21 (1): 162.
29. Fazlollahi MR, Najmi M, Fallahnezhad M, Sabetkish N, Kazemnejad A, Bidad K, et al. The prevalence of asthma in Iranian adults: The first national survey and the most recent updates. *Clin Respir J*. 2018; 12 (5): 1872-81.
30. Global Initiative for Asthma. 2022 GINA Main Report, Global Strategy for Asthma Management and Prevention. Fontana: Global Initiative for Asthma; 2022.
31. Song WJ, Kang MG, Chang YS, Cho SH. Epidemiology of adult asthma in Asia: toward a better understanding. *Asia Pac Allergy*. 2014; 4 (2): 75-85.
32. Chowdhury NU, Guntur VP, Newcomb DC, Wechsler ME. Sex and gender in asthma. *Eur Respir Rev*. 2021; 30 (162).
33. Patra J, Maher YI, Mishra S, Bhatia M, Alam D, Malini DS, et al. Effects of body mass index, tobacco smoking, alcohol drinking and solid fuel use on the risk of asthma: Individual Participant Data (IPD) meta-analysis of 175 000 individuals from 51 nationally representative surveys. *BMJ Open Respir Res*. 2016; 3 (1): e000121.
34. Koebnick C, Fischer H, Daley MF, Ferrara A, Horberg MA, Waitzfelder B, et al. Interacting effects of obesity, race, ethnicity and sex on the incidence and control of adult-onset asthma. *Allergy Asthma Clin Immunol*. 2016; 12: 50.
35. LoMauro A, Aliverti A. Sex and gender in respiratory physiology. *Eur Respir Rev*. 2021; 30 (162).

36. Ekpruke CD, Silveyra P. Sex Differences in Airway Remodeling and Inflammation: Clinical and Biological Factors. *Front Allergy.* 2022; 3: 875295.



## Trias Mahmudiono

<https://www.webofscience.com/wos/author/rid/I-4020-2019>

Web of Science ResearcherID: I-4020-2019

ORCID: 0000-0002-3128-2173


### Current affiliation:

- Universitas Airlangga from 2003 until present

 Academy graduate & mentor

## Verified reviews

### Review Summary



### Reviewer Summary

For manuscripts reviewed from date range December 2018 - June 2023

(6) Open Access Macedonian Journal of Med...

(5) BMC Public Health

(4) Nutrition Journal

(3) Asia Pacific Journal of Public Health

(3) Public Health Nutrition

(2) Journal of Health, Population and Nutriti...

(2) The American Journal of Clinical Nutrition

(1) Revista de Saude Publica

---

(1) Malawi Medical Journal

(1) Frontiers in Public Health

---

(1) Malaysian Journal of Nutrition

(1) Journal of Nutrition and Metabolism

---

(1) Nutrition and Metabolic Insights

(1) Journal of Nutritional Science

---

(1) Media Gizi Indonesia

---

33 REVIEWS OF 28 MANUSCRIPTS

For manuscripts published from date range December 2018 - June 2023

---

Naturally Acquired Lactic Acid Bacteria from Fermented Cassava Improves Nutrient and Anti-dysbiosis Activity of Soy Tempeh

Reviewed: Nov 2021 for Open Access Macedonian Journal of Medical Sciences

---

STUNTING IS THE DOMINANT FACTOR ASSOCIATED WITH IRON DEFICIENCY IN ELEMENTARY SCHOOL CHILDREN

Reviewed: Nov 2021 for Open Access Macedonian Journal of Medical Sciences

---

Double Burden of Malnutrition as a Risk Factor for Overweight and Obesity

Reviewed: Nov 2021 for Revista de Saude Publica

---

Household food security during the COVID-19 pandemic in urban and semi-urban areas in Indonesia

2 rounds from Oct 2021 to Oct 2021 for Journal of Health, Population and Nutrition

---

Socio-demographic characteristics of children and young adults with varied asthma control- does it make a difference?

Reviewed: Oct 2021 for Malawi Medical Journal

---

The Primary Prevention Intervention Of Stunting In Pasaman Barat Regency West Sumatera Indonesia 2019

Reviewed: Oct 2021 for Open Access Macedonian Journal of Medical Sciences

---

Body Fat Composition Related to Fatty Snack Habit Consumption of Office Workers in Urban Area

Reviewed: Sep 2021 for Open Access Macedonian Journal of Medical Sciences

---

An Analysis Of Baby Stunting Prevalence Causing Factors In The Pandemy Era Covid-19

Reviewed: Sep 2021 for Open Access Macedonian Journal of Medical Sciences

---

The Improving the performance of children meal using meal report at Kindergarten School in Indonesia

Reviewed: Sep 2021 for Open Access Macedonian Journal of Medical Sciences

---

Association Between Food Choices Motivators, Physical Activity in Body Image (Dis)satisfaction in Portuguese Adolescents

Reviewed: Feb 2021 for Frontiers in Public Health

---

Zikabra: A Networking Initiative for International Cooperation in Health

Reviewed: Dec 2020 for BMC Public Health

---

An initiative of cooperation in Zika virus research: the experience of the ZIKABRA study in Brazil

Reviewed: Dec 2020 for BMC Public Health

---

Energy, Protein, and Vitamin C Intakes Are Associated to Stunting Among Preschool Children in Central Jakarta, Indonesia: a Case-control Study

Reviewed: Aug 2020 for Malaysian Journal of Nutrition

---

Prevalence Stunting Among Preschool Children (1-5Years) in Bushenyi District Southwestern Uganda- a Health Center Based Study

Reviewed: Aug 2020 for Journal of Nutrition and Metabolism

---



---

Geographical, Socioeconomical and Behavioral Factors Related to Obesity Among Adults in the Republic of Palau Based on the WHO STEPwise Approach to Surveillance 2011–2013

Reviewed: Jul 2020 for Asia Pacific Journal of Public Health

---

Social circumstances and cultural beliefs influence maternal nutrition, breastfeeding and child feeding practices in South Africa

2 rounds from Mar 2020 to May 2020 for Nutrition Journal

---

Intra-individual Double Burden of Overweight Or Obesity and Micronutrient Deficiencies Or Anemia Among Women of Reproductive Age in 17 Population-based Surveys

2 rounds from Jan 2020 to Apr 2020 for The American Journal of Clinical Nutrition

---

Scaled-Up Nutrition Education on Pulse-Cereal Complementary Food in Ethiopia: A Cluster-Randomized Trial

Reviewed: Mar 2020 for BMC Public Health

---

Overfed-Undernourished: The Nutritional Dual Burden of Malnutrition

Reviewed: Feb 2020 for Nutrition and Metabolic Insights

---

The Effectiveness of Nutritional Educational Comics on Increasing Haemoglobin Levels in Anaemic Adolescent Girls

Reviewed: Feb 2020 for Public Health Nutrition

---

Nutrient Adequacy of Japanese Schoolchildren By Household Income: Comparison Between Days with and Without School Lunch

Reviewed: Oct 2019 for Asia Pacific Journal of Public Health

---

Video Animasi Dan Pengetahuan Gizi Remaja

Reviewed: Sep 2019 for Media Gizi Indonesia

---

The association of dietary patterns and adherence to WHO healthy diet with metabolic syndrome in children and adolescents: Tehran lipid and glucose study

2 rounds from Jul 2019 to Sep 2019 for BMC Public Health

---

Nutrient adequacy of Japanese schoolchildren on days with and without a school lunch by household income

Reviewed: Aug 2019 for Asia Pacific Journal of Public Health

---

Coping through a drought: the association between child nutritional status and household food insecurity in the district of iLembe, South Africa

Reviewed: Jul 2019 for Public Health Nutrition

---

Parent–child cooking meal together may alleviate parental concerns about the diets of their toddlers and preschoolers: a cross-sectional analysis in Japan

2 rounds from May 2019 to Jul 2019 for Nutrition Journal

---

The double burden of overnutrition and undernutrition in mother-child dyads in Kenya: demographic and health survey data, 2014

Reviewed: Jul 2019 for Journal of Nutritional Science

---

Coping Through a Drought: The Association Between Child Nutrition and Household Food Insecurity in the District of iLembe, South Africa

Reviewed: Apr 2019 for Public Health Nutrition

---