

01. Matrix Metalloproteinase-3 Down

by Ferdian Ramadhan

Submission date: 16-May-2023 02:02PM (UTC+0800)

Submission ID: 2094407118

File name: 01._Matrix_Metalloproteinase-3_Down.pdf (619.23K)

Word count: 4762

Character count: 24940

RESEARCH ARTICLE

Matrix Metalloproteinase-3 Down Regulation and Cell Migration Inhibition in Human Pterygium Fibroblasts by Mitomycin-C, Curcumin and Fibrin GlueFerdian Ramadhan¹, Jamaluddin², Ismi Zuhria³, Luki Indriaswati³, Evelyn Komaratih^{3,*}¹Ophthalmology Resident, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Hospital, Jl. Mayjen. Prof. Dr. Moestopo No. 47, Surabaya, Indonesia²Rumah Sakit Mata Masyarakat Jawa Timur, Jl. Ketintang Baru Sel. I No. 1, Surabaya 60232, Indonesia³Department of Ophthalmology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Hospital, Jl. Mayjen. Prof. Dr. Moestopo No. 47, Surabaya, Indonesia

*Corresponding author. E-mail: risetpublikasi@gmail.com

Received date: Feb 8, 2022; Revised date: Mar 25, 2022; Accepted date: Apr 12, 2022

Abstract

BACKGROUND: Pterygium is an ocular surface disease that often occurs in tropical countries with a high recurrence rate. Matrix metalloproteinase-3 (MMP-3) play a key role in the inflammatory process of pterygium. This study aims to investigate the ability of curcumin and fibrin glue (FG) in suppressing the expression of MMP-3, and whether can be expected as adjuvant therapy to reduce pterygium recurrence.

METHODS: Human pterygium fibroblasts (HPF) obtained from primary cultured of pterygium were treated with no treatment, curcumin, mitomycin-C (MMC), and FG. MMP-3 expression was analyzed using immunocytochemistry and the intensity measurement was done using ImageJ software. Cell migration was measured by scratching and stratification of fibroblast culture after cell confluence, and assessed for 48 hours.

RESULTS: The expression of MMP-3 were lower in the HPF treated with 100 mol/mL curcumin, 200 mol/mL, and FG (2205.84±86.1 pg/mL, 1002.51±25.22 pg/mL, 1131.55±17.71 pg/mL, respectively) in comparison with untreated HPF (4703.49±108.9 pg/mL). The expression of MMP-3 were significantly different between groups ($p < 0.001$). Cell migration of HPF after scratching with curcumin intervention at 200 mol/mL decrease from 178.67±2.85 (24 hours) to 88.83±1.48 (48 hours). Meanwhile the migration in FG group also decrease from 180.4±2.56 (24 hours) to 72.45±1.25 (48 hours).

CONCLUSION: Curcumin and FG able to reduce the expression of MMP-3 and inhibit the migration of HPF cells.

KEYWORDS: curcumin, mitomycin C, fibrin glue, human pterygium fibroblast, MMP-3

Indones Biomed J. 2022; 14(2): 172-9

Introduction

Pterygium is a degenerative and inflammatory disease of the conjunctiva with the clinical appearance of triangular fibrovascular tissue on the nasal side of the conjunctiva leading to the limbus. Risk factors associated with the development of pterygium include UV exposure, viral infections such as human papillomavirus, herpes simplex virus, and cytomegalovirus.(1-3)

The prevalence of pterygium around the world is quite alarming (ranging from 7% to 15%) and the term band pterygium is known for the region with the highest prevalence of pterygium across the equator. According to the 2010 Indonesian Basic Health Study, among the prevalence of pterygium among all detected eye disease cases in Indonesia reached 3.2% in both eyes and 1.9% in one eye.(1,3)

The relatively high recurrence rate after pterygium resection is another issue that clinicians must face. The



incidence of postoperative recurrence varied from 3.8% to 89%. The pathophysiology of recurrence is not fully understood, but the most likely causes are genetic factors, repeated exposure to risk factors, and surgical technique. Bare sclera technique was the first to be adopted and is still quite high in practice in Indonesia as a modality for the treatment of pterygium, but it is positively associated with a high recurrence rate (up to 24-89%). Treatment modification by using conjunctiva autografts or by adding adjuvant therapy, such as the antimetabolites 5-fluorouracil (5-FU) or mitomycin-C (MMC), as antimitotic and antiapoptotic are often done by clinical practitioners. The use of antimetabolites was successful in reducing the recurrence rate of pterygium, but some serious complications including corneal edema, corneal perforation, or scleral calcification have become new obstacles.(1,3,4)

Recent studies have shown that herbs and spices have a significant attention as natural antioxidants. (5,6) Curcumin, which is a polyphenol derivative from turmeric, has pharmacological properties as antioxidants, antiapoptotic and antiproliferative properties. Curcumin has been found to have anti-inflammatory action through inhibition of nuclear factor kappa-B (NF- κ B), antioxidant action through inhibition of free radicals, suppression of lipid peroxidation, increase of antioxidant molecules in tissues and stimulation of the activity of antioxidant enzymes. Curcumin is able to inhibit the expression of matrix metalloproteinases (MMP), especially MMP-3, which is expressed by human pterygium fibroblasts (HPF), as well as the suppression effect of reactive oxidative species (ROS) of fibroblasts due to UV exposure, making curcumin a potential adjuvant therapy for the prevention of pterygium recurrence in the future.(7-9)

Fibrin glue (FG) is a blood-derived product that is used as a biological adhesive and was first introduced in 1909. FG acts by imitating the final step of the coagulation cascade when soluble fibrinogen is activated by thrombin. In the field of ophthalmology, it is an alternative therapy option that is often used in pterygium cases to prevent postoperative recurrence, accelerate wound healing by releasing anti-inflammatory cytokines. FG forms a seal on the wound after pterygium excision to accelerate the wound healing process, minimize recurrence, and prevent more postoperative complications.(10,11)

In this study, HPF cell cultures were treated with curcumin and FG *in vitro* to evaluate their ability in suppressing the expression of MMP-3 and fibroblast migration, and whether can be expected as adjuvant therapy to reduce pterygium recurrence.

Methods

Preparation of Curcumin and FG

Curcumin (Sigma-Aldrich, St. Louis, MO, USA) was dissolved with complete media Dulbecco's Modified Eagle's Medium (DMEM) then sonicated and filtered to create stock solutions at the concentrations of 100 and 200 mol/L.

FG was derived from graded centrifugation of peripheral blood with anticoagulants. A total of 40 mL of peripheral blood was taken from the cubital vein using a sterile syringe containing CPDA in a ratio of 9:1, aseptically. Blood was put into a sterile 10 mL centrifuge tube then centrifuged at 3000 rpm for 15 minutes. The plasma was stored in a sterile centrifuge tube and stored at -20°C for 24 hours, then centrifuged at 4°C, 3000 rpm for 15 minutes. Ten mL of the upper 2/3 of plasma was stored for the preparation of the fibrinogen component, and the platelet-rich plasma (PRP) was stored in sterile micro-tubes to be prepared as material for thrombin production. The plasma was then added with 1 mL ethanol, then incubated at 4°C for 30 minutes followed by centrifugation at 4°C at 3000 rpm for 15 minutes. The supernatant was discarded and the sediment was used as a component of fibrinogen. Thrombin was made by mixing the PRP component with 0.05 mL of 10% CaCl₂. FG was made by mixing the fibrinogen and thrombin in a ratio of 1:1.(12,13)

Sample Collection

Pterygium tissue was taken in the surgical installation of the Rumah Sakit Mata Masyarakat Jawa Timur. The tissue then transported directly to Biomedical Laboratory of Universitas Brawijaya for further *in vitro* evaluation and analysis. The protocol of this study was approved by Health Research Ethics Committee, Faculty of Medicine Universitas Airlangga (No: 131/EC/KEPK/FKUA/2021).

Culture and Passage of HPF

Local anesthesia with subconjunctival injection of 2% lidocaine was performed before removal of the pterygium tissue. A sterile cloth was placed on the operating area and the eyeball was disinfected using 5% povidone iodine and then rinsed with balance salt solution. Pterygium tissue retrieval was carried out with bare sclera technique according to the protocol developed by previous study (Figure 1).(7)

After the pterygium tissue was removed, the top of the specimen was taken about 2 mm from the edge and then cut into small pieces (<0.5 mm²). After being washed




Figure 1. Pterygium excision technique with bare sclera. A: administration of lidocaine 2% before tissue excision; B: pterygium excision using a pterygium spoon.

using phosphate buffer saline (PBS), then the specimen was given red blood cell lysis buffer for 5 minutes and placed on a 100 mm culture dish. One mL of DMEM containing 15% fetal bovine serum (FBS), and penicillin-streptomycin were added to cover the tissue then placed overnight in an incubator at 37°C, and humidified. The medium was replaced daily with fresh culture medium + FBS. When the cells began to reach 80% confluency, FBS was reduced to 10% concentration and then sub-passed. P3-7 cultures were used in this study. Figure 2 showed the primary culture on pterygium tissue.

Cell's Characteristics and Vimentin Staining

The confluent cells were transferred to 24-well plate with a density of 2.5×10^4 cells/plate and then waited until the cells were adhered to the bottom of the plate, then washed 3 times with PBS for 5 minutes. Cells were washed 3 times with PBS for 5 minutes and with 0.1% PBS Triton-X 100 for 1 minute before incubated with 1% bovine serum albumin (BSA) for 30 minutes. The BSA solution was then discarded and incubated with primary antibody overnight. Cells were incubated with Goat Anti Rabbit IgG & LTIRTCVimentin (Cat No. ab6718, BD Biosciences, San Jose, CA, USA) and incubated with DAPI 1: 1000 for 5 minutes. Cells were then photographed with a fluorescence microscope at 40x magnification.




Figure 2. Primary culture on pterygium tissue. Fibroblast growth reached a confluence rate of 90-100% on day 14. White bar: 100 μ m.

Measurement of MMP-3 by Immunofluorescence Staining

Cells that were already confluent were harvested and planted on 24 well plate with a density of 2.5×10^4 for 24-48 hours. Then the wells were grouped into untreated, treated with MMC 0.2 and 0.4 mg/mL, treated with curcumin 100 and 200 mol/mL, and treated with FG. Untreated wells only contained cells and medium; MMC 0.2 and 0.4 mg/mL were given for 5 minutes to cells with medium, then washed once with PBS; curcumin was given at doses of 100 and 200 mol/mL to cells with medium; and FG was given until clots formed, then removed and washed once with PBS. Each well was then added 3-4% formaldehyde for 15 minutes and washed with PBS.

Every well were washed again with 0.1% of PBS Triton-X 100 (Cat No. T8787, Sigma Aldrich) for 5 minutes and incubated with 1% BSA for 30 minutes at room temperature. The fluid was discarded and then incubated with primary antibody MMP-3 Rabbit Polyclonal Antibody (Cat No. bs-0431R, Bioss Antibody, Woburn, MA, USA) overnight at 4°C. Antibody was washed with PBS, incubated with MMP-3 secondary antibody 1:1000 for 1 hour. Results were read using a fluorescence microscope, and the expression levels were analyzed using ImageJ software version 1.51 (National Institutes of Health, Bethesda, MA, USA). The MMP-3 expression level was counted as corrected total cell fluorescence (CTCF), determined using the formula: integrated density - (area of selected cells x mean fluorescence of background values).⁽²⁾

Cell Migration and *in vitro* Scratch Wound Assay

Confluent pterygium fibroblasts were implanted on 24-well plate with a density of 5×10^4 cells per well and incubated for 24-48 hours to achieve 70-80% confluence. The wells were grouped into untreated, treated with 0.4 mg/mL MMC, treated with 200 mol/mL curcumin, and treated with FG.

The cells that had been given the intervention were then scratched perpendicular to the plate using a 200 μ L micropipette and photographed using an Olympus

CKX53 microscope (Olympus Corporation, Tokyo, Japan). Migration distance was measured with imageJ software version 1.51 (National Institutes of Health).

Statistical Analysis

Comparative data between inhibition of fibroblast migration with the interventions and comparison of MMP-3 expression were tested using the one-way Anova test. Tukey Post-Hoc test was carried out to determine the significant difference between one group and another. The p-value is considered significant if $p < 0.05$. All statistical data was processed using SPSS 26.0 software (IBM Corporation, Armonk, NY, USA).

Results

Expression of MMP-3 After the Intervention of MMC, Curcumin, and FG

The results of analysis showed that the highest expression of MMP-3 was found in the untreated group, and the lowest was in the 0.4 mg/mL MMC group. Groups treated with curcumin showed lower MMP-3 expression compare to the untreated group as the negative control, where 200 mol/mL curcumin group showed lower MMP-3 level than 100 mol/mL curcumin group (Table 1). This showed that 200 mol/mL curcumin had a stronger MMP-3 suppression effect that is quite comparable with FG (1002.51 ± 25.22 vs. 1131.55 ± 17.71).

One-Way Anova test followed by Post-Hoc Tukey analysis showed that the comparison of all treatment groups was $p < 0.001$, which suggested that the comparison of MMP-3 expression in all groups was statistically significant. Figure 3 showed pterygium fibroblast cells were stained with MMP-3 antibody and DAPI to assess the comparasion of MMP-3 expression in all treatment groups. MMP-3 was obviously seen in untreated group, meanwhile was found less in 0.4 mg/mL MMC group.

Inhibition of Fibroblast After the Intervention of MMC, Curcumin, and FG

The migration distance showed that the fastest migration happened in the untreated group, while the slowest migration happened in the 0.4 mg/mL group after 48 hours of intervention (Table 2). The 200 mol/mL curcumin and FG groups had less significant differences after 24 hours of intervention. The group that was given 200 mol/mL curcumin had slower migration inhibition after 48 hours when compared to FG group (figure 4).

Statistical test using One-Way Anova followed by Post-Hoc Tukey showed a significant comparison ($p < 0.05$) in all intervention groups. This suggested that the comparison of HPF migration rate after stratching with and without intervention were statistically significant.

Discussion

Pterygium additionally exhibits several tumor-like features including invasion, metaplasia of epithelial cells, presence of oncogenic viruses, inactivation of tumor suppressor genes, and absence of heterozygosity. The pathogenesis of pterygium and its recurrence after excision is still not completely understood, identification of effective drugs that able to suppress the proliferative system of pterygium is urgently required.(2,14)

MMPs have crucial function in pterygium formation because of the excessive fibroblast proliferation and invasion happens at the apex of the pterygium with destruction of the corneal stroma and basal lamina. Pterygium cells will produce MMP which dissolves the basal lamina and triggers the growth of stromal fibroblasts.(2,14) Overexpression of MMP on basal limbal epithelial cells is the primary reason of pterygium development. Extracellular matrix (ECM) modulation through MMP occurs in the early phase of pterygium invasion, numerous studies have proven a sturdy affiliation among MMP and tumor development/

Table 1. MMP-3 expression in human pterygium fibroblasts after intervention.

| Group | Mean±SD | Minimum | Maximum | p-value |
|---------------------|----------------|---------|---------|---------|
| Untreated | 4703.49±108.98 | 4532.22 | 4892.56 | <0.001 |
| 0.2 mg/mL MMC | 593.99±24.4 | 546.72 | 641.32 | |
| 0.4 mg/mL MMC | 270.60±15.88 | 245.41 | 288.32 | |
| 100 mol/mL Curcumin | 2205.84±86.1 | 2072.31 | 2303.21 | |
| 200 mol/mL Curcumin | 1002.51±25.22 | 973.42 | 1034.45 | |
| FG | 1131.55±17.71 | 1113.56 | 1152.36 | |

p-value tested with One-Way Anova.




Figure 3. Pterygium fibroblast cells stained with MMP-3 antibody and DAPI. Observed with inverted fluorescence microscope; White bar: 30 μ m.

invasion. Invasion of pterygium lesions will growth MMP activity.(15-19).

Pterygium will result in activation of fibroblast tissue, specifically on the top of the pterygium, ensuing in cleavage of fibrillar collagen within the basal lamina, particularly because of MMP-1 and MMP-3. Degradation of the ECM results in the release of stimulating cytokines which includes vascular endothelial growth factor (VEGF) and basic fibroblast growth factor (bFGF) that cause angiogenesis and MMP-3 has a major effect on angiogenesis because it triggers angiostatin. MMP-3 additionally has large outcomes due to the fact it may activate pro-MMPs such as MMP-1, MMP-9 and MMP-13 so that they play a role in cell migration, proteolysis and angiogenesis.(2,14,19) In this

study, it was proven that the highest MMP-3 expression was found in the untreated group, while the lowest expression was found in the 0.4 mg/mL MMC group, followed by the 0.2 mg/mL MMC group. The curcumin intervention also had a suppressive effect on MMP-3 that was more than 50% while compared to the untreated group, especially higher in the 200 mol/mL curcumin group. The suppressive impact of FG was quite similar to of curcumin, so it may be suggested that in this study curcumin and FG have been able to exert a suppressive effect on the expression of MMP-3 in pterygium fibroblast tissue.

Curcumin has been extensively studied as an antifibrosis agent. Previous *in vitro* and *in vivo* study showed curcumin is able to inhibit liver fibrosis and prevent liver

Table 2. Migration distance of human pterygium fibroblast cells after scratching within 48 hours with intervention.

| Group | Treatment Period | Mean±SD | Minimum | Maximum | p-value |
|---------------------|------------------|-------------|---------|---------|---------|
| Untreated | 0H | 221.33±3.57 | 216.23 | 225.34 | 0.000 |
| | 24H | 141.59±3.12 | 137.32 | 147.22 | 0.003 |
| | 48H | 32.46±1.25 | 30.28 | 34.35 | 0.001 |
| 0.4 mg/mL MMC | 0H | 233.60±2.87 | 230.38 | 237.32 | 0.000 |
| | 24H | 210.34±2.57 | 206.52 | 215.08 | 0.000 |
| | 48H | 132.05±1.65 | 129.88 | 135.06 | 0.002 |
| 200 mol/mL Curcumin | 0H | 228.89±3.75 | 222.99 | 233.87 | 0.000 |
| | 24H | 178.67±2.85 | 174.34 | 183.21 | 0.001 |
| | 48H | 88.83±1.48 | 85.28 | 91.21 | 0.001 |
| FG | 0H | 231.62±3.17 | 227.16 | 235.88 | 0.000 |
| | 24H | 180.40±2.56 | 175.32 | 183.52 | 0.001 |
| | 48H | 72.45±1.25 | 70.77 | 74.67 | 0.002 |

p-value tested with One-Way Anova.

disease progression. Curcumin dampen the inflammation and reduce fibrosis through inhibition of the transforming growth factor (TGF)- β 1/Sma and drosophila MAD (Smad) signaling pathway.(19) Other studies reporting the use of curcumin on skin fibroblast cells to prevent photoaging showed that ROS are the basis for overexpression of UVB-modulated MMP-3. Curcumin has a robust antioxidant effect that may suppress ROS by means of suppressing the effects of NF- κ B and AP-1 DNA binding activity. UVB

induces phosphorylation of the janus kinase pathway and p38 and curcumin has been shown to regulate the activation of NF- κ B and AP-1 through the MAPK signaling pathway. (8,20)

The migration rate showed that the untreated group had the fastest migration, while the slowest migration rate was found in the of 0.4 mg/mL MMC group after 48 hours. The curcumin and FG groups had much less substantial differences at 24 hours and 200 mol/L curcumin




Figure 4. Fibroblast migration distance after scratching analysis. Comparison of pterygium fibroblast cells were scanned to assess cell migration at 0, 24 and 48 hours. Observed with inverted fluorescence microscope; Yellow line: the scratch border; Red line: as fibroblast migration; White bar: 100 μ m.

group had slower migration inhibition after 48 hours when compared to fibrin glue. This indicates that the comparison of the HPF migration rate with and without intervention is statistically significant. The scratching method performed in this study is an easy and effective step to trigger inflammation.(21) This is in accordance with the outcomes of previous study of fibroblast cells in the tenon conjunctival tissue model that was scratched with the conclusion the less the rate of migration of tenon's fibroblast cells to the wound place, the decrease the opportunity of extracellular matrix manufacturing resulting in fibrosis in that place.(12,22)

Another study confirmed the relationship among MMP expression and activity on pterygium fibroblast migration and its suppressive effect on Bevacizumab and cyclosporine A (CsA). The fibroblast migration ratio was assessed to investigate the recovery migration process which confirmed the migration speed suppression of MMP-3 and MMP-13 after bevacizumab and/or CsA intervention.(12)

An *in vitro* trial on cardiac fibroblasts and stated that curcumin had a higher inhibitory impact on fibroblast migration when compared to the control group and the MTT assay showed that the curcumin group also had a decrease proliferative effect than the control group, which is in line with the study.(23) This indicates that the rate of migration in the curcumin group might be slower when compared to the control group. This occurs most probably due to the suppressing effect of the mediators including TGF- β , MMP and other profibrotic mediators via the NF- κ B through tumor necrosis factor (TNF)- α dependent induction and mitogen-activated protein kinase (MAPK) signaling pathway.(20,23,24)

FG significantly inhibits collagen synthesis and induces collagen degradation in human tenons, in which the wound healing manner is regulated by inflammation. A decrease in the amount of collagen will eliminate the anti-inflammatory response and fibrin glue substantially inhibits collagen in the injured tenon model and is able to reduce wound formation and contractures.(24,25) In line with this study, which showed that in addition to reduce MMP-3 expression, FG was also able to reduce the speed of HPF migration although it was not as significant as the standard for adjuvant pterygium treatment, particularly MMC. Current study is the first study to analyze the inhibition of MMP-3 expression and HPF migration after the intervention of curcumin and FG. However, additional learning about other wound healing mechanisms is necessary to fully define and understand their effects.

Conclusion

Our findings suggest that curcumin and FG are able to suppress pterygium proliferation by inhibiting HPF migration and suppressing MMP-3 expression. Therefore, FG and curcumin have the potential to be an adjuvant therapy for MMC replacement in the management of pterygium.

Acknowledgments

This study was funded by Indonesian Ministry of Research and Technology and Community Development and Service Institution of Universitas Airlangga. Our gratitude to Biomedical Laboratory of Universitas Brawijaya in facilitating the data collection. We also thank to staffs and residents of Ophthalmology Department, Faculty of Medicine, Universitas Airlangga/Dr Soetomo General Hospital who had support us in conducting this study.

Authors Contribution

FR, LI and EK were involved in conceptualizing and planning the research; FR and LI performed the data acquisition/ collection; JA interpreted the result of data and designed the figures and table; EF and IJ calculated the experimental data and performed the analysis; FR, IZ and EK drafted the manuscript and took parts in giving critical revision of the manuscript; FR and EK also collected the research funding.

References

1. Ery, Uly AM, Dwi S. Distribusi dan karakteristik pterigium di Indonesia. Buletin Penelitian Sistem Kesehatan. 2011; 14(1): 84-9.
2. Lu CW, Hao JL, Yao L, Li HJ, Zhou DD. Efficacy of curcumin in inducing apoptosis and inhibiting the expression of VEGF in human pterygium fibroblasts. Int J Mol Med. 2017; 39(5): 1149-54.
3. Singh SK. Pterygium: epidemiology prevention and treatment. Community Eye Health. 2017; 30(99): S5-6.
4. Liu T, Liu Y, Xie L, He X, Bai J. Progress in the pathogenesis of pterygium. Curr Eye Res. 2013; 38(12): 1191-7.
5. Ginting CN, Lister INE, Girsang E, Riastawati D, Kusuma HSW, Widowati W. Antioxidant activities of Ficus elastica leaves ethanol extract and its compounds. Mol Cell Biomed Sci. 2020; 4 (1): 27-33.
6. Girsang E, Lister INE, Ginting CN, Bethasari M, Amalia A, Widowati W. Comparison of antiaging and antioxidant activities of protocatechuic and ferulic acids. Mol Cell Biomed Sci. 2020; 4 (2): 68-75.

5. Li DQ, Sao BL, Zeenat GS, Yunqi L, Abraham S, Daniel M, *et al.* Overexpression of collagenase (MMP-1) and stromelysin (MMP-3) by pterygium head fibroblasts. *Arch Ophthalmol.* 2001; 119(1): 71-80.
6. Sadoughi D, Edalatmanesh MA, Rahbarian R. Protective effect of curcumin on quality parameters of sperm and testicular tissue alterations in alloxan-induced diabetic rats as animal model. *Indones Biomed J.* 2019; 11(3): 225-337
7. Qin Z, Qiuli F, Lifang Z, Houfa Y, Xiuming J, Qiaomei T, *et al.* Proliferative effects of histamine on primary human pterygium fibroblasts. *Mediators Inflamm.* 2016; 2016: 9862496. doi: 10.1155/2016/9862496.
8. Panda A, Sandeep K, Abhiyan K, Raseena B, Shibal B. Fibrin glue in ophthalmology. *Indian J Ophthalmol.* 2009; 57(5): 371-9.
9. Yang MB, Melia M, Lambert SR, Chiang MF, Simpson JL, Buffenn AN. Fibrin glue for closure of conjunctival incision in strabismus surgery: a report by the american academy of ophthalmology. *Ophthalmology.* 2013; 120(9): 1935-41.
10. Komaratih E, Rindiastuti Y, Susilowati H, Wijayanti NT, Dranindi S, Primitasari Y, *et al.* Antifibrotic effects of limbal mesenchymal stem cells-conditioned media (LMSCS-CM) on human tenon's fibroblasts (HTFs) in glaucomatous eyes: comparison with mitomycin c. *Sch Acad J Biosci.* 2019; 7(1): 4-12.
11. Ferris D, Frisbie D, Kisiday J, McIlwraith CW. In vivo healing of meniscal lacerations using bone marrow-derived mesenchymal stem cells and fibrin glue. *Stem Cells Int.* 2012; 2012: 691605. doi: 10.1155/2012/691605.
12. Kim YH, Jae CJ, Sang IG, Su BP, Jin YM, Yong IK, *et al.* Inhibition of pterygium fibroblast migration and outgrowth by bevacizumab and cyclosporine A involves down-regulation of matrix metalloproteinases-3 and -13. *PLoS One.* 2017; 12(1): 1-24.
13. An MX, Kai LW, Shao CL. Detection and comparison of matrix metalloproteinase in primary and recurrent pterygium fibroblast. *Int J Ophthalmol.* 2011; 4(4): 353-6.
14. Cárdenas-Cantú E, Zavala J, Valenzuela J, Valdez-García JE. Molecular basis of pterygium development. *Semin Ophthalmol.* 2016; 31(6): 567-83.
15. Girolamo ND, Jeanie C, Minas TC, Denis W. Pathogenesis of pterygia: role of cytokines, growth factors, and matrix metalloproteinases. *Prog Retin Eye Res.* 2004; 23: 195-228.
16. Solomon A, De-Quan L, Sao BL, Scheffer CGT. Regulation of collagenase, stromelysin, and urokinase-type plasminogen activator in primary pterygium body fibroblasts by inflammatory cytokines. *Invest Ophthalmol Vis Sci.* 2000; 41: 2154-63.
17. Supriono, Nugraheni A, Kalim H, Eko MH. The effect of curcumin on regression of liver fibrosis through decreased expression of transforming growth factor- β 1 (TGF- β 1). *Indones Biomed J.* 2018; 11(1): 52-8.
18. Hwang BM, Noh EM, Kim JS, Kim JM, You YO, Hwang JK, *et al.* Curcumin inhibits UVB-induced matrix metalloproteinase-1/3 expression by suppressing the MAPK-p38/JNK pathways in human dermal fibroblasts. *Exp Dermatol.* 2013; 22(5): 371-4.
19. Stamm A, Reimers K, Straub S, Vogt P, Scheper T, Pepelanova. In vitro wound healing assays – state of the art. *Bio Nano Mat.* 2015; 17(1-2): 79-87.
20. Brilliyanto FS, Suhendro G, Wahyuni I, Rochmanti M, Pramono W. Metformin reduced collagen deposition and contractility, but increased collagen degradation in in vitro posterior capsule opacification model. *Indones Biomed J.* 2021; 13(2): 186-91.
21. Chung CC, Kao YH, Liou JP, Chen YJ. Curcumin suppress cardiac fibroblasts activities by regulating proliferation, migration, and the extracellular matrix. *Acta Cardiol Sin.* 2014; 30(5): 474-82.
22. Wulandari F, Ikawati M, Kirihata M, Kato J, Meiyanto E. Curcumin analogs, PGV-1 and CCA-1.1 exhibit anti-migratory effects and suppress MMP9 expression on WiDr cells. *Indones Biomed J.* 2021; 13(3): 271-80.
23. Rindiastuti Y, Komaratih E, Susilowati H, Soebagyo HD, Rantam FA. Fibrin glue (FG) attenuates fibrosis on human tenon's fibroblasts (HTFS) of glaucomatous eyes : comparison with mitomycin C. *Biochem Cell Arch.* 2019; 19(2): 4713-20.

01. Matrix Metalloproteinase-3 Down

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|--|-----|
| 1 | eprints.ucm.es Internet Source | 1% |
| 2 | www.researchgate.net Internet Source | 1% |
| 3 | tessera.spandidos-publications.com Internet Source | 1% |
| 4 | discovery.ucl.ac.uk Internet Source | 1% |
| 5 | s3-ap-southeast-2.amazonaws.com Internet Source | 1% |
| 6 | uvadoc.uva.es Internet Source | <1% |
| 7 | Baiq Ratna Kumaladewi, Willy Sandhika, Heryawati Heryawati. "Role of CD44 and CD8 in Colorectal Adenocarcinoma metastatic", Medical Laboratory Technology Journal, 2020 Publication | <1% |
| 8 | bura.brunel.ac.uk Internet Source | <1% |

| | | |
|----|--|------|
| 9 | kuscholarworks.ku.edu Internet Source | <1 % |
| 10 | f1000research.com Internet Source | <1 % |
| 11 | www.mdpi.com Internet Source | <1 % |
| 12 | coek.info Internet Source | <1 % |
| 13 | www.linguistics.ualberta.ca Internet Source | <1 % |
| 14 | Hong Sun, Li Li, Cécile Vercherat, Neriman Tuba Gulbagci et al. "Stra13 regulates satellite cell activation by antagonizing Notch signaling", <i>The Journal of Cell Biology</i> , 2007 Publication | <1 % |
| 15 | biomedpharmajournal.org Internet Source | <1 % |
| 16 | iovs.arvojournals.org Internet Source | <1 % |
| 17 | www.medsci.org Internet Source | <1 % |
| 18 | Michelle Vang Mouritzen, Håvard Jenssen. "Optimized Scratch Assay for <i>In Vitro</i> Testing of Cell Migration with an Automated Optical | <1 % |

Camera", Journal of Visualized Experiments, 2018

Publication

19 cris.maastrichtuniversity.nl <1 %
Internet Source

20 elifesciences.org <1 %
Internet Source

21 www.e-journal.unair.ac.id <1 %
Internet Source

22 www.lib.kobe-u.ac.jp <1 %
Internet Source

23 Shigenobu Kawashima, Atsushi Mizota, Emiko Adachi-Usami, Tsuyoshi Kimura. "Effects of mitomycin C on the rat retina", Documenta Ophthalmologica, 1996 <1 %
Publication

24 bmcvetres.biomedcentral.com <1 %
Internet Source

25 dx.doi.org <1 %
Internet Source

26 garuda.kemdikbud.go.id <1 %
Internet Source

27 ies.ijo.cn <1 %
Internet Source

28 nursekey.com
Internet Source

<1 %

29

pubs.rsc.org

Internet Source

<1 %

30

www.hindawi.com

Internet Source

<1 %

31

www.nature.com

Internet Source

<1 %

32

Hyun Jin Park, Garam Choi, Seongmin Ha, Yesl Kim et al. "MBP-11901 Inhibits Tumor Growth of Hepatocellular Carcinoma through Multitargeted Inhibition of Receptor Tyrosine Kinases", *Cancers*, 2022

Publication

<1 %

33

Jia Li, Yan An, Jia-Ning Wang, Xiao-Ping Yin, Huan Zhou, Yong-Sheng Wang. " Curcumin targets vascular endothelial growth factor activating the PI3K/Akt signaling pathway and improves brain hypoxic-ischemic injury in neonatal rats ", *The Korean Journal of Physiology & Pharmacology*, 2020

Publication

<1 %

34

Jianwu Fan, Xin Zhang, Yaping Jiang, Li Chen, Minjie Sheng, Yihui Chen. "SPARC knockdown attenuated TGF- β 1-induced fibrotic effects through Smad2/3 pathways in human

<1 %

pterygium fibroblasts", Archives of Biochemistry and Biophysics, 2021

Publication

35

Kim, H.S.. "TGF- β 1 stimulates production of gelatinase (MMP-9), collagenases (MMP-1, -13) and stromelysins (MMP-3, -10, -11) by human corneal epithelial cells", Experimental Eye Research, 200408

Publication

<1 %

36

Ofira Zloto, Vicktoria Vishnevskia-Dai, Joseph Moisseiev, Michael Belkin, Ido Didi Fabian. "A Biological Tissue Adhesive and Dissolvent System for Intraocular Tumor Plaque Brachytherapy", Ophthalmic Surgery, Lasers and Imaging Retina, 2016

Publication

<1 %

37

Qiao Zhou, Nanxue Cui, Shihai Zhang, Miaomiao Zhou, Younian Xu. "TSLP reduced neuroinflammation by blocking the activation of microglia in the JAK2/STAT5 pathway", Research Square Platform LLC, 2023

Publication

<1 %

38

Robert T. Chang. "Early Postoperative Bleb Maintenance", Cataract Surgery in the Glaucoma Patient, 2009

Publication

<1 %

39

daneshyari.com

Internet Source

<1 %

40

ddd.uab.cat

Internet Source

<1 %

41

downloads.hindawi.com

Internet Source

<1 %

42

ia601702.us.archive.org

Internet Source

<1 %

43

journals.lww.com

Internet Source

<1 %

44

Cheng-Wei Lu, Ji-Long Hao, Lei Yao, Hai-Jun Li, Dan-Dan Zhou. "Efficacy of curcumin in inducing apoptosis and inhibiting the expression of VEGF in human pterygium fibroblasts", International Journal of Molecular Medicine, 2017

Publication

<1 %

45

Gabriela Lopezcarasa-Hernandez, Jose-Francisco Perez-Vazquez, Jose-Luis Guerrero-Naranjo, Maria A. Martinez-Castellanos. "Versatility of use of fibrin glue in wound closure and vitreo-retinal surgery", International Journal of Retina and Vitreous, 2021

Publication

<1 %

46

Henny J.M. Beckers, Florent Aptel, Carroll A.B. Webers, Elisa Bluwol et al. "Safety and Effectiveness of the PRESERFLO® MicroShunt

<1 %

in Primary Open-Angle Glaucoma",
Ophthalmology Glaucoma, 2022

Publication

47

Hwang, Bo-Mi, Eun-Mi Noh, Jong-Suk Kim, Jeong-Mi Kim, Yong-Ouk You, Jin-Ki Hwang, Kang-Beom Kwon, and Young-Rae Lee. "Curcumin inhibits UVB-induced matrix metalloproteinase-1/3 expression by suppressing the MAPK-p38/JNK pathways in human dermal fibroblasts", Experimental Dermatology, 2013.

Publication

48

Josine S. Quaedackers. "Cardiovascular and endocrine effects of a single course of maternal dexamethasone treatment in preterm fetal sheep", BJOG An International Journal of Obstetrics and Gynaecology, 2/2005

Publication

49

Maninder Kaur, Narpinder Singh. "Relationships Between Selected Properties of Seeds, Flours, and Starches from Different Chickpea Cultivars", International Journal of Food Properties, 2006

Publication

50

Omnia Elshreef, Adel Abdelwahab, Muhammad Abu Elaish. "RECURRENCE RATE AFTER CONJUNCTIVAL AUTOGRAFT VESUS INTRAOPERATIVE MITOMYCIN C FOR

<1 %

<1 %

<1 %

<1 %

TREATMENT OF PRIMARY PTERYGIUM",
Journal of Recent Advances in Medicine, 2020

Publication

51

Prathita Amanda, Erik Prabowo, Muflihatul Muniroh, Abdul Mughni Rozy, Neni Susilaningsih. "The Effectiveness of Curcumin on Transforming Growth Factor- β and the Proportion of Liver Fibrosis in Deutschland Denken Yoken Mice with Common Bile Duct Ligation", Open Access Macedonian Journal of Medical Sciences, 2022

Publication

<1 %

52

Shun-Fa Yang, Ching-Yang Lin, Pei-Yu Yang, Shih-Chun Chao, Yi-Zhen Ye, Dan-Ning Hu. "Increased Expression of Gelatinase (MMP-2 and MMP-9) in Pterygia and Pterygium Fibroblasts with Disease Progression and Activation of Protein Kinase C", Investigative Ophthalmology & Visual Science, 2009

Publication

<1 %

53

Yeoun-Hee Kim, Jae-Chang Jung, Sang Il Gum, Su-Bin Park, Jin Yeul Ma, Yong Il Kim, Kyoo Won Lee, Young Jeung Park. "Inhibition of Pterygium Fibroblast Migration and Outgrowth by Bevacizumab and Cyclosporine A Involves Down-Regulation of Matrix Metalloproteinases-3 and -13", PLOS ONE, 2017

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

01. Matrix Metalloproteinase-3 Down

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8