

04. Beneficial Effect of Resveratrol on

by Medhi Denisa Alinda

Submission date: 16-May-2023 02:07PM (UTC+0800)

Submission ID: 2094409942

File name: 04._Beneficial_Effect_of_Resveratrol_on.pdf (221.78K)

Word count: 3852

Character count: 21786

ORIGINAL ARTICLE

Beneficial Effect of Resveratrol on Conditioned Medium Human Adipose-derived Mesenchymal Stem Cell for Cutaneous Wound Healing Related Growth Factors

Medhi Denisa Alinda^{1,2}, Muhammad Yulianto Listiawan², Anang Endaryanto³, Heri Suroto^{4,5}, Evelyn Komaratih⁶, Fedik Abdul Rantam⁷, Bagus Haryo Kusumaputra², Regitta Indira Agusni², Yuyun Rindiastuti⁶, Camilla Amanda Prakoeswa⁸, Cita Rosita Sigit Prakoeswa²

¹ Post Graduate Doctoral Program, Faculty of Medicine, Universitas Airlangga, Surabaya Jawa Timur 60115, Indonesia

² Department of Dermatology and Venereology, Faculty of Medicine Universitas Airlangga-Dr. Soetomo General Academic Hospital, Surabaya Jawa Timur 60115, Indonesia

³ Department of Pediatric, Faculty of Medicine Universitas Airlangga-Dr. Soetomo General Academic Hospital, Surabaya, Jawa Timur 60286, Indonesia

⁴ Cell and Tissue Bank, Faculty of Medicine Universitas Airlangga-Dr. Soetomo General Academic Hospital, Surabaya, Jawa Timur 60115, Indonesia

⁵ Department of Orthopaedic & Traumatology, Faculty of Medicine Universitas Airlangga-Dr. Soetomo General Academic Hospital, Surabaya, East Java 60286, Indonesia

⁶ Department of Ophthalmology, Faculty of Medicine Universitas Airlangga-Dr. Soetomo General Academic Hospital, Surabaya, Jawa Timur 60115, Indonesia

⁷ Stem Cell Research and Development Center, Universitas Airlangga, Surabaya, Jawa Timur 60115, Indonesia

⁸ Faculty of Medicine Universitas Airlangga-Dr. Soetomo General Academic Hospital, Surabaya, Jawa Timur 60286, Indonesia

ABSTRACT

Introduction: Cutaneous wound healing, a type of soft tissue injury, needs the interaction between multiple growth factors (epidermal growth factor [EGF], transforming growth factor [TGF], platelet-derived growth factor [PDGF], and vascular endothelial growth factor [VEGF]) in order to attain physiological healing. Nevertheless, in certain conditions, i.e., chronic wound, certain growth factors might be insufficient. VEGF is among the most significant pro-angiogenic mediators during wound healing to increase migration and invasion of endothelial cells. Conditioned medium adipose-derived mesenchymal stem cells (CM-AdMSCs) are a new hope to ameliorate impaired healing. Resveratrol (RV), a small molecule that can increase the effect of CM-AdMSCs, is needed to help stimulates growth factors from stem cells for culture. This study aims to evaluate growth factors in CM-AdMSCs with resveratrol. **Methods:** Isolated and characterized MSCs from CM-AdMSCs treated with RV supplemented Serum Deprived Medium (SDM) were used in this study. Growth factors level were measured using ELISA. **Results:** A decrease of mean were noted in EGF (290.19±53.19 VS 271.64±52.58) and VEGF (788.76±204.82 VS 117.92±49.22), while an increase of mean in TGF (633.14±11.38 VS 824.73±121.15) and PDGF (2999.00±203.54 VS 3961.40±378.68) between CM-AdMSCs without resveratrol and with resveratrol group, respectively. PDGF and VEGF were statistically significant ($p < 0.05$). **Conclusion:** In conclusion, RV promotes PDGF secretion and decreases VEGF in CM-AdMSCs.

Keywords: Wound healing; CM-AdMSCs; Resveratrol; VEGF; PDGF

Corresponding Author:

Medhi Denisa Alinda, MD
Email: medhidenisaalinda@gmail.com
Tel: +6285645050746

INTRODUCTION

Wound healing, a biophysiological process in the human body, is an intricate multi-stage process orchestrated reconstitution in the skin's dermal and

epidermal layers (1,2). This multi-stage process is reached through four dynamic and integrated programmed phases: hemostasis, inflammation, proliferation, and remodeling/resolution where cells (i.e., thrombocyte, granulocytes, macrophages, fibroblast, and keratinocytes), cytokines, growth factors, and protease are involved (1,3). After a dermal injury, aggregation of platelet initiate the coagulation cascade and formation of clots. Subsequently, the wound bed will be infiltrated with leukocytes producing pro-

inflammatory cytokine (i.e., neutrophils and monocytes/macrophages). In later healing stages, fibroblasts will be recruited to the site of the wound to deposit extracellular matrix and provide the basis for regeneration of new tissue(4). Reported data suggest that essential wound-care products ought to have the composition that are similar to the skin; a mixture of distinct growth factors and extracellular matrix (ECM) proteins endogenous to the skin, viable epithelial cells, fibroblasts, and coupled with mesenchymal stem cells (MSCs) (5). Stem cells are distinguished by their multipotency and self-renewal capability, while MSCs are a group of stem cells with self-renewal ability, multiple differentiation potential, and strong immunoregulatory effects (6,7). Due to this nature, MSCs are desirable subjects for a number of cell therapies in previous literature (2,8,9).

The various MSCs' differentiation lineages induced by growth factors are as follows, 1) chondrocytes, 2) osteoblasts, 3) adipocytes, 4) hepatocytes, 5) cardiomyocytes, 6) neurons, 7) epithelium, 8) endothelial cells, 9) pancreatic β cells, and 10) epidermis (10). Adipose-derived mesenchymal stem cells (ADMSCs), stem cells derived from adipose tissue stroma, are characterized by its considerable self-renewal ability, ability to differentiate into numerous types of functional cells, and possible paracrine function of the cells (11,12). Studies have reported ADMSCs to enhance cutaneous wound healing, with significantly quicker reepithelization, greater granulation tissue formation (day 3 to day 7), and angiogenesis (6,13). This mechanism has shown its beneficial effect in wound healing, especially in chronic or non-healing wounds.

Resveratrol (RV), a naturally occurring polyphenol, is produced by various plants, such as grapes and peanuts. This compound has attracted research in cancer due to its ability to restrict growth, stimulate apoptosis in numerous malignancies, and increase tumor susceptibility to chemotherapeutic agents (14). To an additional extent, various studies published over the last few years have reported RV's advantage on the behavior of several stem cells. Nonetheless, only few literatures have reported RV's effect on stem cell behavior, especially in cell secretion products/secretomes. In this study, we examined the effects of resveratrol on CM-AdMSCs for cutaneous wound healing related growth factors *in vitro*.

MATERIALS AND METHODS

Mesenchymal stem cells isolation and expansion

MSCs are characterized by their capacity to attach to a plastic surface in standard culture conditions, their surface markers expression, their lack of hematopoietic markers expression, and their potential to differentiate along osteoblastic, adipocytic, and chondrocyte pathways *in vitro* (15).

Firstly, a 1 cm³ fat tissue sample that is not contaminated with blood clots or connective tissue is prepared in alpha MEM transport medium inside a 50 ml conicle. Next, the sample is removed from the conicles and rinsed with PBS solution until it is clean and the erythrocytes are removed. Then, the fat tissue is chopped and mixed with collagenase to be incubated at 37°C hot plate for 30 minutes until it dissolves completely. Next, the solution is filtered to separate the remaining insoluble fatty tissue. The filtered products are then centrifuged at 3000 rpm for 5 minutes until pellets are formed. The pellets then re-suspended with 10 ml of alpha MEM medium to form a homogeneous solution that needs to be incubated until colonies are formed, and the cells achieve 80% confluence.

After the cells reach a confluence of 80%, various doses of resveratrol are added to the media of the cells (23). Propagation is done by collecting the conditioning medium from the petri dish and storing it in a 50 ml conicle, washing it with 5 ml of PBS solution, and adding triple express enzyme. If the monolayer is already released, a medium stopper is added and it will develop into a single cell. The single-cell solution is centrifuged at 3000 rpm for 5 minutes until pellets are formed and given Alpha MEM medium to be re-suspended and planted in 2 new Petri dishes afterward.

Before being characterized using CD 105, the mesenchymal stem cells must be effectively separated from fat tissue first. A fluorescent color on the membrane's surface indicates a positive test for CD 105.

To harvest the metabolites, 50 ml of conditioned medium is inserted into the dialysis tubing membrane that needs to be spun at 500 rpm on a hot plate magnetic stirrer. Let it sit overnight until the conditioning medium's color in the dialysis tubing is muted. After that, pour the contents from the dialysis tubing, filter the metabolite products with a size of 0.22 microns, and then pack in 50 ml conicles to be stored in a sealed sterile medipack at a -200°C temperature. It is ready to use when needed and as much as 1 ml is taken for ELISA examination purposes. This research was approved by the Ethical Committee in Health Research of Dr. Soetomo General Academic Hospital (reference number: 0052/LOE/301.4/VIII/2020).

Statistical analysis

Statistical analyses were carried out using the IBM Statistical Package for the Social Sciences (SPSS) Statistics 25 for Windows (International Business Machines Corporation, New York, United States). Data are expressed as mean \pm standard deviation (SD). Groups were compared using independent T-test or one-way ANOVA. A value of $p < 0.05$ was considered statistically significant.

RESULTS

Characterization of adipocyte-derived mesenchymal stem cells.

Comparison of growth factors in conditioned adipocyte-derived mesenchymal stem cells regarding resveratrol (RV)

Four growth factors were measured; namely, epidermal growth factor (EGF), transforming growth factor (TGF), platelet-derived growth factor (PDGF), and vascular endothelial growth factor (VEGF). A decrease of mean was noted in EGF (290.19 ± 53.19 VS 271.64 ± 52.58 [Figure 1a]) and VEGF (788.76 ± 204.82 VS 117.92 ± 49.22 [Figure 1b]) between CM-AdMSCs without RV and with RV group, respectively. On the contrary, an increase of mean in TGF (633.14 ± 11.38 VS 824.73 ± 121.15 [Figure 1c]) and PDGF (2999.00 ± 203.54 VS 3961.40 ± 378.68 [Figure 1d]) between CM-AdMSCs without RV and with RV group, respectively, were noted. To further clarify the mean difference of the growth factors investigated regarding RV, we analyzed the growth factors. We found that PDGF and VEGF were statistically significant (p-value of 0.018 and 0.005, respectively).




Figure 1: Comparison of mean 1a) EGF levels, 1b) VEGF levels, 1c) TGF levels, 1d) PDGF levels in CM-AdMSCs without RV and with RV, respectively.

DISCUSSION

Adipose tissue stroma, an important secretory and endocrine organ, can be conveniently obtained through liposuction technique (16). This tissue can be harvested autologously and does not present the ethical, tumorigenic, or immunogenic issues presented by pluripotent stem cells with immunosuppressive characteristic (17,18). Adipose tissue volume is constituted almost 90% of adipocytes. It produces a diverse population of cell types (preadipocytes, fibroblasts, vascular smooth muscle cells, endothelial cells, resident monocytes/macrophages, and lymphocytes) when broken down enzymatically (19). Once isolated, it is referred to as the stromal vascular

fraction (SVF) (19). There are several ways of ADMSCs' mechanism of action in wound healing: 1) collagen synthesis and degradation, 2) site-specific differentiation and improved survival, 3) immunomodulation, and 4) neovascularization (20).

The alterations in the skin's cell composition and the various epithelial cells' ability to secrete specified growth factors (i.e., TGF- β , VEGF, matrix metalloproteinase [MMP]-1, MMP-2, MMP-9, and extracellular matrix [ECM]) offer the potential of determining an equilibrium between cell regeneration as well as rejuvenation to the ADMSCs' microenvironment (21). This milieu can be further enhanced through the addition of RV. Preconditioned RV promotes MSCs proliferation, sustains differentiation, and hinders MSCs senescence. Through this accepted approach of MSCs with RV, we hoped it would facilitate wound healing related growth factors secretion. Wound healing related growth factors are EGF, TGF, PDGF, VEGF, and fibroblast growth factor (FGF), to name a few (22). A Previous study has reported that 0.1 μ M is the most potent RV dose to promote secretion of EGF, PDGF, TGF- β 1, and hepatocyte growth factor (HGF) (23).

Resveratrol (3,4',5 trihydroxysilbene), a phytochemical, is a well-known biologically active compound synthesized by plants undergoing ionizing radiation or infection (24). RV was stated to display numerous therapeutic advantages, such as anti-inflammatory, antioxidant, immunomodulator, vasorelaxant, and neuroprotective effect, to name a few. In vitro, RV modulates angiogenesis through VEGF expression and the formation of vascular networks (25). Due to its properties, RV potential target is the adult stem cells that dwell in different tissues and organs through oral or intravenous injection (23).

The EGF family, also known as ErbB1 or Her1, consists of various members, four of which – EGF, heparin-binding EGF-like growth factor (HB-EGF), epiregulin, and TGF- α . The EGF family members are synthesized in membrane-associated forms and need MMPs or ADAMs (a disintegrin and metalloproteinase) activation (22). EGF, secreted by platelets, macrophages, fibroblasts, endothelial cells, keratinocytes, and bone marrow-derived MSCs, is a powerful inducer of epithelialization, angiogenesis, fibroblast proliferation, granulation tissue formation, and survival (22). This role is not restricted to straightforward consequences on keratinocytes, fibroblasts, and endothelial cells; a considerable lot of these factors are powerful stimulator of inflammatory mediators and their receptors (22). For example, the increase of antimicrobial peptides and proinflammatory interleukin (IL)-8 production due to TGF- α induction of Toll-like receptors (TLR5 and TLR9) (22, 26).

The TGF- β superfamily, particularly the TGF- β 1-3 (a bone morphogenetic proteins [BMPs]), is involved in wound healing. Various cell types, such as macrophages,

platelets, keratinocytes, and fibroblasts, produced this family of growth factors. TGF- β is activated by MMP-2, MMP-9, thrombospondin 1, and integrin $\alpha v\beta 6$ along with membrane-type MMP (22). The influence of TGF- $\beta 1$ on cells is concentration-dependent; low TGF- $\beta 1$ concentration promotes endothelial proliferation and migration, while high concentration increases matrix production (22). In a chronic wound, decreased level of TGF- β ligands has been documented, implying that an incorporation of exogenous TGF- β s may be advantageous for wound healing (22, 27).

Platelet-derived growth factor (PDGF) is multifunctional. It is also among the first growth factors produced in response to injury and stimulates cellular responses during the entirety of the repair processes (22). It is secreted by platelets, macrophages, keratinocytes, fibroblasts, and endothelial cells. Generally, activation of several pathways (i.e., phospholipase C γ pathway, phosphatidylinositol-3-kinase (PI3K), and several mitogen-activated protein kinase (MAPK) pathways) results in the enhancement of cell migration and proliferation and the increase of VEGF and insulin-like growth factor production. IGF PDGF levels are of a low value in chronic wounds, possibly due to insufficient production and/or exceeding protease-mediated degradation. This underlying condition was advantageous for patients with chronic wounds (22, 28). The VEGF family includes six members. The VEGF is a heparin-binding glycoprotein, it needs to be bound to cell-surface tyrosine kinase receptors (VEGFR-1, VEGFR-2, and VEGFR-3) to conduct their functions (22). Angiogenesis is predominantly mediated by VEGFR1 and VEGFR2, while VEGFR-3 mediates lymphangiogenesis. This family of growth factors plays a part during injury-induced hypoxia leading to their up-regulation. In wound healing, VEGF secreted by the platelets, macrophages, fibroblasts, and keratinocytes, acts in a paracrine approach on endothelial cells, stimulating and promoting wound angiogenesis (22). This growth factors also acts as an immunostimulator for the migration and activation of monocyte. It also assists in MMPs production by smooth myocytes, induces their migration and proliferation throughout hypoxia, and stimulates fibroblast proliferation, scars formation, and keratinocyte motility essential for wound re-epithelization. In chronic wound, a paradox occurred between increase VEGF mRNA and decrease VEGF protein levels due to proteolytic activity within wound bed (22). To our attention, RV suppressed the expression of VEGF as seen as shown in other studies. This is achieved through several proposed mechanism of 1) restrictive outcome in VEGF-induced phosphorylation of VEGFR2 without any effect on total VEGFR2 expression, 2) promotion of VEGF degradation, subsequently reducing the interaction with its receptor, and 3) glycolysis inhibition during latent endothelial cells activation (29, 30).

Our study showed that ADMSCs in conditioned RV medium could increase the secretion of PDGF from ADMSCs. This finding is supported through a proposed mechanism of RV to help preserve the human MSCs self-healing and multi-lineage differentiation abilities beyond passage 10 through a SIRT1-dependent pathway, especially adipogenic potential. SIRT1 (a NAD $^{+}$ -dependent lysine deacetylase) deacetylates SOX2 to prevent its degradation, subsequently preserving the MSCs stemness after a lengthy passaging (31). In addition, SIRT1 manages OCT4 and NANOG expression indirectly through p53 deacetylation in human embryonic and cancer stem cells (31). This is further enhanced through RV mechanism of reducing p53 level by stimulating its deacetylation in adult stem cells. Collectively, RV-activated signalling pathway is revealed as a straightforward relation of RV-SIRT1-SOX2 complex, and the activation of SIRT1 enables an approach for wound therapy acceleration (23, 31).

CONCLUSION

The present study indicated a favorable therapeutic potential for CM-AdMSCs in cutaneous wounds healing, especially in chronic wound. Increase production of PDGF in CM-AdMSCs plays a major role in chronic wounds where they are in a lower value. Lower VEGF should be put into consideration when considering utilizing this growth factor from a CM-AdMSCs. RV promotes PDGF secretion and decreases VEGF in CM-AdMSCs. Future research should be explored to the specific mechanism of CM-AdMSCs in cutaneous wound healing in clinical trials settings.

ACKNOWLEDGEMENTS

We would like to thank the Cell and Tissue Bank of Dr. Soetomo General Academic Hospital, Surabaya, Indonesia for supporting this research.

REFERENCES

1. Guo S, DiPietro LA. Critical review in oral biology & medicine: Factors affecting wound healing. *J Dent Res*. 2010; 89(3):219–29.
2. Nuschke A. Activity of mesenchymal stem cells in therapies for chronic skin wound healing. *Organogenesis*. 2014;10(1):29–37.
3. Harper D, Young A, McNaught CE. The physiology of wound healing. *Surg (United Kingdom)*. 2014;32(9):445–50. Available from: <http://dx.doi.org/10.1016/j.mpsur.2014.06.010>
4. Hanson SE, Bentz ML, Hematti P. Mesenchymal Stem Cell Therapy for Nonhealing Cutaneous Wounds. *Plast Reconstr Surg*. 2010;125(2):510–516.

5. Julianto I, Rindastuti Y. Topical Delivery of Mesenchymal Stem Cells "Secretomes" in Wound Repair. *Acta Med Indones*. 2016;48(3):217–20.
6. Duscher D, Barrera J, Wong VW, Maan ZN, Whittam AJ, Januszzyk M, et al. Stem Cells in Wound Healing: The Future of Regenerative Medicine? A Mini-Review. *Gerontology*. 2016;62(2):216–25.
7. Zhang Y, Han F, Gu L, Ji P, Yang X, Liu M, et al. Adipose mesenchymal stem cell exosomes promote wound healing through accelerated keratinocyte migration and proliferation by activating the AKT/HIF-1 α axis. *J Mol Histol* [Internet]. 2020;51(4):375–83. Available from: <https://doi.org/10.1007/s10735-020-09887-4>
8. Han Y, Li X, Zhang Y, Han Y, Chang F, Ding J. Mesenchymal Stem Cells for Regenerative Medicine. *Cells*. 2019;8(8):886.
9. Neveen AS. Mesenchymal Stem Cell Based Therapy for Parkinson's Disease. *Int J Stem Cell Res Ther*. 2019;6(1).
10. Afflerbach AK, Kiri MD, Detinis T, Maoz BM. Mesenchymal stem cells as a promising cell source for integration in novel in vitro models. *Biomolecules*. 2020;10(9):1–30.
11. Li Q, Guo Y, Chen F, Liu J, Jin P. Stromal cell-derived factor-1 promotes human adipose tissue-derived stem cell survival and chronic wound healing. *Exp Ther Med*. 2016;12(1):45–50.
12. Na YK, Ban JJ, Lee M, Im W, Kim M. Wound healing potential of adipose tissue stem cell extract. *Biochem Biophys Res Commun* [Internet]. 2017;485(1):30–4. Available from: <http://dx.doi.org/10.1016/j.bbrc.2017.01.103>
13. Kim H, Hyun MR, Kim SW. The Effect of Adipose-Derived Stem Cells on Wound Healing: Comparison of Methods of Application Stem Cells *Int*. 2019;2019.
14. Mikula-Pietrasik J, Kuczmarska A, Kucińska M, Murias M, Wierchowski M, Winckiewicz M, et al. Resveratrol and its synthetic derivatives exert opposite effects on mesothelial cell-dependent angiogenesis via modulating secretion of VEGF and IL-8/CXCL8. *Angiogenesis*. 2012;15(3):361–76. Available from: doi: 10.1007/s10456-012-9266-0.
15. Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini FC, Krause DS, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* [Internet]. 2006;8(4):315–7. Available from: <http://dx.doi.org/10.1080/14653240600855905>
16. Lee YJ, Baek SE, Lee S, Cho YW, Jeong YJ, Kim KJ, et al. Wound-healing effect of adipose stem cell-derived extracellularmatrix sheet on full-thickness skin defect rat model: Histological and immunohistochemical study. *Int Wound J*. 2018;2018:1–11.
17. Francis SL, Duchi S, Onofrillo C, Di Bella C, Choong PFM. Adipose-derived mesenchymal stem cells in the use of cartilage tissue engineering: The need for a rapid isolation procedure. *Stem Cells Int*. 2018;2018:13–6.
18. Hassanshahi A, Hassanshahi M, Khabbazi S, Hosseini-Khah Z, Peymanfar Y, Ghalamkari S, et al. Adipose-derived stem cells for wound healing. *J Cell Physiol*. 2019;234:7903-7914.
19. Bertozzi N, Simonacci F, Grieco MP, Grignaffini E, Rapisio E. The biological and clinical basis for the use of adipose-derived stem cells in the field of wound healing. *Ann Med Surg*. 2017;20:41–8.
20. Gadelkarim M, Abushouk AI, Ghanem E, Hamaad AM, Saad AM, Abdel-Daim MM. Adipose-derived stem cells: Effectiveness and advances in delivery in diabetic wound healing. *Biomed Pharmacother*. 2018;107(April):625–33.
21. Mazini L, Rochette L, Admou B, Amal S, Malka G. Hopes and limits of adipose-derived stem cells (ADSCs) and mesenchymal stem cells (MSCs) in wound healing. *Int J Mol Sci*. 2020;21(4).
22. Manuscript A. Potential and Methods of Delivery. *Adv Ski Wound Care*. 2013;25(8):349–70.
23. Prakoeswa CRS, Rindiastuti Y, Wirohadidjojo YW, Komaratih E, Nurwasis, Dinaryati A, et al. Resveratrol promotes secretion of wound healing related growth factors of mesenchymal stem cells originated from adult and fetal tissues. *Artificial Cells, Nanomedicine and Biotechnology*. 2020;48:1159-66.
24. Salehi B, Mishra AP, Nigam M, Sener B, Kilic M, Sharifi-Rad M, et al. Resveratrol: A double-edged sword in health benefits, *Biomedicines*. 2018;6(3):1–20.
25. Shaito A, Posadino AM, Younes N, Hasan H, Halabi S, Alhababi D, et al. Potential adverse effects of resveratrol: A literature review. *Int J Mol Sci*. 2020;21(6).
26. de Lima PO, Joseph S, Panizza B, Simpson F. Epidermal Growth Factor Receptor's Function in Cutaneous Squamous Cell Carcinoma and Its Role as a Therapeutic Target in the Age of Immunotherapies. *Curr Treat Options Oncol*. 2020;21(1).
27. Liarte S, Bernabiy-Garcna Б, Nicol6s FJ. Role of TGF- β in Skin Chronic Wounds: A Keratinocyte Perspective. *Cells*. 2020;9(2):306.
28. Nardini M, Perteghella S, Mastracci L, Grillo F, Marrubini G, Bari E, et al. Growth factors delivery system for skin regeneration: An advanced wound dressing. *Pharmaceutics*. 2020;12(2).
29. Hu WH, Duan R, Xia YT, Xiong QP, Wang HY, Chan GKL, et al. Binding of Resveratrol to Vascular Endothelial Growth Factor Suppresses Angiogenesis by Inhibiting the Receptor Signaling. *J Agric Food Chem*. 2019;67(4):1127–37.
30. Wu H, He L, Shi JJ, Hou X, Zhang H, Zhang X, et al. Resveratrol inhibits VEGF-induced angiogenesis in human endothelial cells associated with

- suppression of aerobic glycolysis via modulation of PKM2 nuclear translocation. *Clin Exp Pharmacol Physiol*. 2018;45(12):1265–73.
31. Choi Y, Yoon DS, Lee KM, Choi SM, Lee MH, Park KH, et al. Enhancement of mesenchymal stem cell-driven bone regeneration by resveratrol-mediated SOX2 regulation. *Aging Dis*. 2019;10(4):818–33.

04. Beneficial Effect of Resveratrol on

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | Mark A. Babizhayev, Yegor E. Yegorov. "Tissue formation and tissue engineering through host cell recruitment or a potential injectable cell-based biocomposite with replicative potential: Molecular mechanisms controlling cellular senescence and the involvement of controlled transient telomerase", Journal of Biomedical Materials Research Part A, 2015
Publication | 1% |
| 2 | www.panafrican-med-journal.com
Internet Source | 1% |
| 3 | www.spandidos-publications.com
Internet Source | 1% |
| 4 | mts.intechopen.com
Internet Source | 1% |
| 5 | "Stem Cell Therapy for Vascular Diseases", Springer Science and Business Media LLC, 2021
Publication | 1% |
-

6	Didi Erwandi Mohamad Haron, Minoru Yoneda, Rafidah Hod, Muhammad Ikram A. Wahab, Mohd Yusmaidie Aziz. "Perfluoroalkyl and polyfluoroalkyl substances, bisphenol and paraben compounds in dust collected from residential homes in Klang Valley, Malaysia", Human and Ecological Risk Assessment: An International Journal, 2022 Publication	1 %
7	lbeifits.files.wordpress.com Internet Source	1 %
8	www.jstage.jst.go.jp Internet Source	1 %
9	www.researchsquare.com Internet Source	1 %
10	cyberleninka.org Internet Source	1 %
11	www.science.gov Internet Source	1 %
12	pure.rug.nl Internet Source	1 %
13	www.thieme-connect.com Internet Source	1 %
14	Jianxia Hu, Xiaoyi Liu, Jingwei Chi, Kui Che, Xiaolong Ma, Mingyue Qiu, Zhengju Fu, Yahao	<1 %

Wang, Yangang Wang, Wei Wang. "Resveratrol Enhances Wound Healing in Type 1 Diabetes Mellitus by Promoting the Expression of Extracellular Vesicle-Carried MicroRNA-129 Derived from Mesenchymal Stem Cells",
Journal of Proteome Research, 2022

Publication

15

apicareonline.com

Internet Source

<1 %

16

rfasper.fasper.bg.ac.rs

Internet Source

<1 %

17

academic.oup.com

Internet Source

<1 %

18

watermark.silverchair.com

Internet Source

<1 %

19

jcbior.com

Internet Source

<1 %

20

www.mdpi.com

Internet Source

<1 %

21

"1-A1: Lung Cancer 1 : Poster Sessions",
Respirology, 2013.

Publication

<1 %

22

Wei-Hui Hu, Ran Duan, Yi-Teng Xia, Qing-Ping Xiong et al. "Binding of Resveratrol to Vascular Endothelial Growth Factor Suppresses Angiogenesis by Inhibiting the Receptor

<1 %

Signaling", Journal of Agricultural and Food Chemistry, 2018

Publication

23

jneuroinflammation.biomedcentral.com

Internet Source

<1 %

24

arthritis-research.com

Internet Source

<1 %

25

coek.info

Internet Source

<1 %

26

Domergue, Sophie, Claire Bony, Marie Maumus, Karine Toupet, Eric Frouin, Valérie Rigau, Marie-Catherine Vozenin, Guy Magalon, Christian Jorgensen, and Danièle Noël.
"Comparison between Stromal Vascular Fraction and Adipose Mesenchymal Stem Cells in Remodeling Hypertrophic Scars", PLoS ONE, 2016.

Publication

<1 %

27

Ferroni, Letizia, Chiara Gardin, Ilaria Tocco, Roberta Epis, Alessandro Casadei, Vincenzo Vindigni, Giuseppe Mucci, and Barbara Zavan.
"Potential for Neural Differentiation of Mesenchymal Stem Cells", Advances in Biochemical Engineering/Biotechnology, 2012.

Publication

<1 %

28

Jen-Her Lu, Kai Hsia, Chih-Kuan Su, Yi-Hsiang Pan, Hsu Ma, Shih-Hwa Chiou, Chih-Hsun Lin.

<1 %

"A Novel Dressing Composed of Adipose Stem Cells and Decellularized Wharton's Jelly Facilitated Wound Healing and Relieved Lymphedema by Enhancing Angiogenesis and Lymphangiogenesis in a Rat Model", Journal of Functional Biomaterials, 2023

Publication

29

Silvia Mihalceanu, Matthias M. Aitzetmüller, Hans-Günther Machens, Dominik Duscher. "Chapter 18 Stem Cell Therapies for Tissue Regeneration and Wound Healing: Strategies to Enhance Therapeutic Effectiveness", Springer Science and Business Media LLC, 2019

Publication

<1 %

30

Wita Saraswati, Erry Gumilar Dahlan, Koosnadi Saputra, Thomas Cahyono Sutrisno. "Effect of Electroacupuncture on Natural-Killer Cells and Tumor Size in Patients with Cervical Squamous-Cell Carcinoma: A Randomized Controlled Trial", Medical Acupuncture, 2019

Publication

<1 %

31

Xiufen Zhuang, Xiao Hu, Shuren Zhang, Xingmin Li, Xiaoying Yuan, Yanhong Wu. "Mesenchymal Stem Cell-Based Therapy as a New Approach for the Treatment of Systemic Sclerosis", Clinical Reviews in Allergy & Immunology, 2022

Publication

<1 %

32	f1000researchdata.s3.amazonaws.com Internet Source	<1 %
33	ijph.in Internet Source	<1 %
34	www.juri.urologi.or.id Internet Source	<1 %
35	www.oncotarget.com Internet Source	<1 %
36	hdl.handle.net Internet Source	<1 %
37	Brian W. Booth, Gilbert H. Smith. "Roles of transforming growth factor- α in mammary development and disease", Growth Factors, 2009 Publication	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

04. Beneficial Effect of Resveratrol on

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
