# ISSN-0973-9122 (Print) • ISSN-0973-9130 (Electronic)

Volume 16 / Number 1 / January-March 2022

# **Indian Journal of Forensic Medicine & Toxicology**

Website: www.ijfmt.com

Official Organ of Indian Association of Medico-Legal Experts (Regd.)



Indian Journal of Forensic Medicine & Toxicology

Home / Archives / Vol. 16 No. 1 (2022): Indian Journal of Forensic Medicine & Toxicology

# Vol. 16 No. 1 (2022): Indian Journal of Forensic Medicine & Toxicology



DOI: https://doi.org/10.37506/ijfmt.v16i1

Published: 2021-11-30

# Articles

# Antibacterial Activity of (+) Usnic Acid against Multi Drug Resistant Acinetobacter baumannii from Clinical Isolates

B. Nagaraju, K. Fathimunnisa, R. Vijayaraghavan, B. Sreekanth 11-17

ዾ Pdf

# A Comparative study to Evaluate the Efficacy of Supervised Exercise Program and Cyriax Physiotherapy on Pain and Function in Lateral Epicondylitis

Gouri Kalaskar, Pratik Phansopkar 18-24



# Deleted

25**-**28

# A Study to Assess Respiratory Health Status among Bricks Workers

Jyot B. Darji, Siddaram Sarate 29-32

🖪 Pdf

# Study of Ossification Centres Around the Elbow and Wrist of Adolescent Age Group 15-18 Years

Kishore Kumar.G, R.Jenifer Jacquiline 33-40



# **Cheiloscopy - A Tool For Identification in Twins**

M.Sugatha, R.Jenifer Jacquiline, M.Taqiuddin Khan

41-46



# Synthesis, Characterization, Molecular Docking of Sulphamethoxazole Schiff Base Metal Complexes and Its Antibacterial, Anti-Inflammatory and Anti Depressant Activity

R.Gandhimathi, Magdivari Sangeetha 47-57



# Aortic Atherosclerosis and Risk Factors: A Clinico- Demographic Autopsy-Based Study

Mainak Tarafder, Suparna Datta, Ranjan Biswas, Prabir Chakraborty, Saptarshi Chatterjee 58-65

🕒 Pdf

# Estimation of Stature & Gender from Thumb Indices in Indian Population

Mohite Hemlata S, Kakade Satish V, Mohite Sandeep S, Roy Priya P, Doshi Megha A 66-75

# Sudden Death with Dual Organ Pathologies–An Autopsy -based Case Report

Mukul Sharma, Kimi Soumya Padhi, Govinda Balmuchu, Prachi Nemani, ManasRanjan Sahu 76-82



# Demographic Study of Unnatural Deaths in Paediatric Age Group in General Hospital, Khammam

# Self-Efficacy with the Quality of Life of Pulmonary Tb Patients

Usastiawaty C.A.S Isnainy, Ridwan Ridwan, Rias Tusianah, M. Arifki Zainaro, Albet Maydiyantoro, Tubagus Ali Rachman Puja Kesuma 872-884

872-884

🖾 Pdf

# Ultrasound Assessment for Thyroid Examination in Patients with Hypothyroidism

Yours Abdel Hassan, Safwan Saeed, Maha Taha Idrees 885-892



# Alpha-Mangostin and Gamma-Mangostin Isolated from Mangosteen (Garcinia mangostana L.) as Promising Candidates against SARS-CoV-2: A Bioinformatics Approach

Arif Nur Muhammad Ansori 893-899



# Comparative Evaluation of Dexmeditomidine & Fentanyl in Terms of Cardiovascular Stress Response During Anesthetic Airway Management in Major Surgical Procedures

Niraj Rathod, Sunil Valand, Seema Rawat, Komal Makwana 900-907



# A Tear Inflammatory Biomarker in Dry Eye Disease

Ni Made Inten Lestari, Evelyn Komaratih, Yuyun Rindiastuti, Cita Rosita Sigit Prakoeswa 908-912



# Serum Preptin Level in Iraqi Beta Major Thalassemic Patients

Noor Haidar Talib, Hedef .D. Al-Yaseen, Ali Mohammed Jwad 913-919

🖾 Pdf

# **Effect of KI on SDF Treated Cavities**

Ola Mohamed Ismail Sakr, Mashail Abdullah Alsaikhan 920-928

🖾 Pdf

# The Cytotoxic Effect of Iraqi Rumex Acetosella against Breast and Esophagus Cancer Cells

Omar Hussein Ahmed, Ali Jabbar Abdulhussein, Enas Jawad Kadhim 929-940

🖾 Pdf

# A molecular Study of the Microsporum Canis and Trichophyton Mentagrophytes Associated Fungal Infection: Athlete's Foot among Farmers

Omar Sadik Shalal, Qahtan Adnan Rasheed, Dunya Abdulrazzaq Alkurjiya 941-950

B P	df
-----	----

Features of Teleroentgenographic Indicators of the Position of the Teeth and the Profile of the Soft Tissues of the Face in Adolescents with Different Profiles and Types of Faces According to Schwarz A.M.

Prokopenko O.S, Gunas I. V, Beliaiev E. V, Kotsyura O. O, Kovalchuk V. V 951-959

🖾 Pdf
-------

# Study of Genetic Variation of the gene NOS3 and Cadmium Concentrations in a Sample of Iraqi Patients with Essential Hypertension

Qusay A. Abdulameer, Ismail H. Aziz, Abdul-jabbar A Ali, Ismail A. Abdulhassan 960-966



# Environmental Analysis of Massive Mask Waste due to the Covid-19 Pandemic in Indonesia

Rahmad Agus Dwianto, Tulus Haryono, Rianto 967-972



# Immunohistochemical Characterization of Hepatic Nuclear Factor 4 Alpha Expression in the Choroid Plexus of the lateral and 4th ventricles of adult Male Rat Brain

Rasha A. Salman, Taghreed Abdulrsool Ali, Duaa AL Musawi 973-978



# Profile Study of Motorcyclists Victims in Road Traffic Accidents at Jaipur Region- An Observational Antemortem Study

Ravindra Kumar, R.K.Punia 979-985



# Urban Slum Settings: A Crosssectional Study

M.C. Yadavannavar, Praveen Ganganahalli, Soumya Mirji, Sneha Arkeri, Sadafara Janwekar 1709-1717

🖾 Pdf

# Comparative Study of Various Human Ossification Centers From 10 Years to 20 Years of Age

Y.V.Ratna Raju, M. Rajeev Varma, Raja sekhar Uppu, P Vijaya Kumar 1718-1726

ß	Pdf
---	-----

# An Examination of the Severity of Periodontal Disease in Elderly Men with Varying Degrees of Physical Activity

Smitha Shetty, Smeet Gusani, Mranali K Shetty

1727**-**1735



# Effect of Gingival Biotype on the Progression of Periodontal Disease

Mranali K Shetty, Neetha J Shetty 1736-1742



Methanol Toxicity: An Emergency Department and Toxicologist Dilemma – Methanol Case Series, Investigation of Medical Professionals' Knowledge, and Review of The Literature

Sahar Y. Issa, Maha K. Al Mazroua, Naglaa F Mahmoud 1743-1758



# The Effect of Local Application of Magnesium Oxide Powder on the Blood Parameters During Nerve Regeneration of Injured Sciatic Nerve in Rat

Helal Montaser, Hussein Aseel 1759-1765

🖾 Pdf

Current Issue
ATOM 1.0
RSS 2.0
RSS 1.0
Make a Submission

**Browse** 

**Open Journal Systems** 

Information

For Readers

For Authors

For Librarians

# Platform & workflow by OJS / PKP

Indian Journal of Forensic Medicine & Toxicology

Home / Editorial Team

# **Editorial Team**

Editor in Chief Prof S K Dhattarwal

Forensic Medicine, PGIMS, Rohtak, Haryana E-mail: <u>editor.ijfmt@gmail.com</u>

# EDITORIAL ADVISORY BOARD

- 1. Prof Sudhir K Gupta, Head, FMT. AIIMS, New Delhi , India
- 2. Prof Mete Gulmen , Cukurova University, TURKEY
- 3. Prof. Leandro Duarte De Carvalho , Minas Gerais, Belo Horizante, BRAZIL
- 4. Dr. Valery Gunas, National Pirogov Memorial Medical University, Vinnytsya, UKRAINE
- 5. Dr. Rahul Pathak Forensic Science, Dept of Life Sciences ,Anglia Ruskin University, Cambridge, UNITED KINGDOM
- 6. Prof Emilo Nuzzalese, University of Turin , Italy
- 7. Dr Noha A. Magdy Elrafie, Forensic Toxicology, Ain Shams University, Cairo, EGYPT
- 8. Dr Rituja Sharma, Associate Prof, Law, Banasthali Vidyapeeth Jaipur
- 9. Dr Shankar Bakkanwar *(Associate Professor)* Forensic Medicine, Kasturba Medical College, Manipal, Karnataka
- 10. Dr K. Ravikumar , Raksha Shakti University, Ahmedabad, Gujrat.
- 11. Dr C. Behera (Addl. Prof) Dept of FMT, AIIMS, New Delhi
- 12. Dr. Kanak Lata Verma, Deputy Director, Toxicology ,RFSL, Chanakyapuri New Delhi
- 13. Dr. Asha Srivastava *(Senior Scientific Officer)* Forensic Psychology,Central Forensic Science Laboratory, CBI, Delhi
- 14. Dr. Raghvendra Kumar Vidua, (Associate Prof), FMT, AIIMS Bhopal
- 15. Dr. Vaibhav Saran (*Asst.Prof.*) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences, Allahabad
- 16. Ms Aparna R. Asst. Prof. Forensic Serology & Biology, Jain University, Bengaluru
- 17. Dr. Deepali Jain, Asst Prof, Forensic Science ,BB Ambedkar University, Lucknow
- 18. Prof. NK Aggrawal Forensic Medicine, UCMS, Delhi
- 19. Prof. Manoj Kumar Mohanty, Forensic Medicine, AIIMS, Bhuvneshwar, Odissha
- 20. Prof. Amar Jyoti Patowary, Forensic Medicine, NEIGRIHMS, Shillong, Meghalaya
- 21. Prof S. Venkata Raghava , Forensic Medicine, Banglore Medical College, Bengaluru
- 22. Prof. Shalini Gupta Oral Pathology and Microbiology,Faculty of Dental Sciences, King George Medical University, Lucknow
- 23. Prof. Virender Kumar Chhoker Forensic Medicine, Santosh Medical College, Ghaziabad, UP
- 24. Prof. Dayanand G Gannur , Forensic Medicine , Shri BM Patil Medical College, Hospital &

Research centre, Bijapur, Karnataka

- 25. Prof Praveen Arora, Forensic Medicine, SAIMS, Indore, MP
- 26. Prof Barkha Gupta , Saraswathi Institute of Medical Sciences Hapur, Uttar Pradesh India
- 27. Prof M Prashant Apollo Medical College Hyderabad
- 28. Prof Dimple Patel , Anatomay, AMC MET Medical College , Ahmedabad , Gujarat
- 29. Dr Mohammed Nader Shalaby, Associate Professor of Biological Sciences and Sports Health Department, Faculty of Physical Education, Suez Canal University, Egypt
- 30 Dr. Avinash Harishchandra Waghmode

Professor and Head,Dept of Forensic Medicine and Toxicology,BKL Walawalkar Rural Medical College Chiplun Ratnagiri

31 Dr. Anita Yadav Assistant Professor Forensic Science, SBAS Galgotias University, Greater Noida, UP

32 Dr. Risha Jasmine Nathan Assistant Professor Forensic Science, SBAS Galgotias University, Greater Noida, UP

# **Current Issue**



Make a Submission

Browse

**Open Journal Systems** 

Information

For Readers

For Authors

For Librarians

# Platform & workflow by OJS / PKP

# A Tear Inflammatory Biomarker in Dry Eye Disease

## Ni Made Inten Lestari<sup>1</sup>, Evelyn Komaratih<sup>1</sup>, Yuyun Rindiastuti<sup>1</sup>, Cita Rosita Sigit Prakoeswa<sup>2</sup>

<sup>1</sup>Department of Ophthalmology, <sup>2</sup>Department of Dermatovenerology, Faculty of Medicine Universitas Airlangga/ Dr Soetomo General Academic Hospital, Jl. Mayjen. Prof. Dr. Moestopo 4-7, Surabaya, Indonesia

# Abstract

**Background:** Dry eye disease is a chronic ocular condition and significantly impacts visual function with multifactorial origin. It is characterized by tear instability and inflammation on the ocular surface. The inflammatory response initiates by synthesis and release of cytokines, Recently, there has been increasing scientific interest using tear film biomarkers that play a role in pathophysiology of dry eye disease. The objective of this study to explain relationship of tear film biomarker in dry eye.

**Methods:** The research design of this study used literature review. The data was collected from electronic database in PubMed, Google Scholar, and website of science and predominantly published in the last 10 years.

**Result:** The defect on the tear film component due to oxidative stress lead to ocular surface epithelial exposure, and intracellular signaling pathways are triggered, involving an inflammatory mediator which plays a role in the pathophysiology of dry eye. Using biomarker lead to better diagnosis, drug development, and effective management for dry eye disease.

**Conclusion:** In conclusion, TNF-a, IL-6, IL-8, IL-1 by number studies to consistently reflect disease severity and strong correlation with schirmer tear test and another test for the ocular surface in dry eye disease.

Keywords: Dry eye disease, Tear Biomarker, Inflammation

# Introduction

In recent years, the epidemiology of DED is changing. The new paradigm impacting younger patients, mostly related to the rise of smartphone

# Corresponding author:

Cita Rosita Sigit Prakoeswa.

Department of Dermatovenerology, Faculty of Medicine Universitas Airlangga/ Dr. Soetomo General Academic Hospital, Surabaya. Jl. Mayjen. Prof. Dr. Moestopo 4-7, Surabaya, 60132. Telephone & Fax: (031) 5501605. Email: cita-rosita@fk.unair.ac.id; risetpublikasi@ gmail.com use. In America, an estimated 16 million had been diagnosed with DED and many countries worldwide with a range of 9.5-90 %. The prevalence is higher among women than man and increases with age<sup>1</sup>. In Indonesia, the majority of DED is 37,6 % by 40-49year age group and 1,4 times higher for men than women<sup>2,6</sup>.

Defect on tear film components due to oxidative stress lead ocular surface epithelial exposed and triggered intracellular signaling pathways. The inflammatory response initiates the synthesis and release of cytokines, matrix metalloproteinases (MMPs), and chemokines. Pro-inflammatory cytokine could activate T-cell to raise inflammation, and MMPs promote extracellular degradation, causing exposure to epithelial barrier<sup>3,6</sup>.

Recently, there has been increasing scientific interest in the use of tear biomarkers in DED. Biomarker or biological marker is an objective indicator medical sign or biological stated of normal and pathogenic processes<sup>4</sup>. Using biomarker will lead to better diagnosis, drug development, and effective management for dry eye disease. However, it is challenging to access ocular specimen—alternatively, many studies identifying protein biomarker in tear fluid. A biomarker originated anatomically close to the disease and noninvasive for dry eye disease is crucial to increase specificity. This article aims to describe an overview of the use of tear fluid biomarker in diagnosing and managing DED.

### Pathophysiology Dry Eye Disease

Dry eye disease can divide into several significant subtypes, such as aqueous deficient and evaporative. The etiology of aqueous deficiency is disruption of the sensory drive to the lacrimal gland, decreased reflex tearing due to nerve damage (refractive surgery), and systemic drug (such as beta-blocker, diuretic, antihistamine), and aging. Aqueous deficiency most common is related to systemic inflammation, such as Sjogren syndrome. Sjogren syndrome is a chronic autoimmune disorder characterized by lymphocytic infiltration in the exocrine gland, resulting in dry mouth and dry eye<sup>3,5</sup>.

Evaporative dry eye resulting in low covering of lipid to the surface of the eye. The most common cause by Meibomian gland dysfunction (MGD). The obstruction of the Meibomian gland leads to tear film instability, hyperosmolarity, and increased tear evaporation. Disruption Meibomian gland function impacts the quantity and quality of Meibomian secretion, affecting changes in tear film composition. The risk factor of MGD including hormonal aspect (imbalance androgen and estrogen), systemic medication (administration of 13-cis retinoid acid), and topical medication (e.g. beta-blocker, prostaglandin analogs, topical epinephrine)<sup>4</sup>.

# **Diagnosis Dry Eye Disease**

To assess DED required a combination of symptom and sign. Dryness symptom assessed on validated questionnaires such as ocular surface disease index (OSDI), MacMonnies dry eye questionnaire, and dry eye questionnaire (DEQ). Clinically, DED routinely assessed using a Schirmer tear test to evaluate production aqueous, tear break up time (TBUT) to measure tear film stability, defect on ocular surfacespecific corneal and conjunctival staining. The tear film quality could be assessed using an osmometer that measures tear film osmolarity <sup>5</sup>.

### **Tear Film Composition**

The tear film is composed of three main layers. The outermost layer is formed predominantly of lipid, which function is to prevent water evaporation and reduce the tension of the ocular surface. The middle is an aqueous layer and plays a role in protecting against pathogens and particle and hydration ocular surface. Its contain insoluble and soluble component such as proteins, electrolytes, peptides, and small molecules. The innermost layer is a mucin layer that resides directly at the surface of the cornea. Mucin originated from the goblet cell. The primary function is to maintain the hydration of the ocular surface<sup>7</sup>.

Dysfunction of the lacrimal functional unit cause tear film instability and change any component of the tear fluid. The lacrimal functional unit consists of the lacrimal gland, ocular surface, ocular nerves, goblet cell, and accessory gland (e.g. Meibomian). Hyperosmolarity would activate innate immune response by generating a pro-inflammatory microenvironment on the ocular surface<sup>9</sup>. Initially, natural immune response mediated by pattern recognition receptor such as a toll-like receptor. Receptor recognition and triggers activation of inflammasomes to secretion pro-inflammatory cytokines such as interleukin -1b<sup>10,11</sup>. The murine model demonstrated that hyperosmolar condition activated protein kinase and stimulates cytokines (IL-8, IL-6, TNF-a), matrix metalloproteinases (MMP9, MMP1)<sup>10,11</sup>. Another study showed that hyperosmolarity induces apoptosis of corneal epithelial cell in vitro condition.

Hyperosmolar stress also activates adaptive immune response through activation antigenpresenting cells (APCs), mainly corneal dendritic cells, and recruitment of inflammatory mediators. Hyperosmolar state on the ocular surface also increased the level of chemokines such as CCL20 and CXCL9 in tear film<sup>9,10</sup>. Mature APCs migrate the regional lymph nodes and prime CD4+ T cell, including T helper 1 (Th1) and Th17. These inflammatory mediators interferon-gamma (IFN-g) through angiogenesis and lymphangiogenesis. IFN-g is secreted by T helper 1 and NK cell related loss of conjunctival goblet cell. Goblet cell produces mucins. In vitro study demonstrated goblet cell is susceptible to IFN-g with significantly reduced proliferation<sup>12,13,14</sup>.

### **Tear Inflammatory Biomarkers**

### Interleukin-6

Interleukin-6 (IL-6) is an important proinflammatory cytokine that is produced by T-lymphocytes and activated macrophages. IL-6 promptly and transiently acute responses to infection, inflammation, immune response, and hematopoiesis. However, dysregulated continue synthesis of IL-6 play a significant effect on chronic inflammation. Interleukin 6 is a representative cytokine with increased expression in tear fluid and conjunctival epithelium and has been known as one of the key molecules in DED.

In many studies, the expression of IL-6 in tear fluid up-regulated in chronic condition<sup>17</sup>. Another study revealed that IL-6 was significantly related to the schirmer tear test. IL-6 levels can be used as indicators to determine the severity and efficacy of anti-inflammatory drugs for DED<sup>18</sup>. Using enzymelinked immunosorbent assay in tear sample patients with DED that are reported IL-6 level is elevated<sup>19,22</sup>. Another study showed IL-6 level was significantly increased in evaporative DED and correlated with meibography and schirmer tear test<sup>18</sup>.

### **Interleukin 1**

The IL-1 family consist of three forms, namely IL-1a, IL-1b, and IL-1 receptor antagonist. The mechanism action of IL-1a and IL-1b activate an immune response that trigger activation to a cascade of inflammation by infiltration, the proliferation of lymphocyte and macrophage, and increasing chemokine production. In ocular surface, IL-1 has been related to corneal neovascularization, bullous keratopathy and reported increasing level in Sjogren Syndrome and ocular rosacea<sup>19,20</sup>. This study noted the pro-inflammatory cytokines IL-1 associated with the intensity of corneal defect epithelium and density of goblet cell <sup>21</sup>. IL-1 receptor antagonist is a form anti-inflammatory of interleukin-1, which has a role to be a natural homeostatic mechanism for preventing undesirable activation of IL-1 mediated inflammatory events on ocular surface $^{20,21}$ .

## Interferon-gamma

Interferon-gamma (IFN- $\gamma$ ) is secreted mainly through lymphocyte T helper type 1, cells, cytotoxic T cells, and natural killer. IFN- $\gamma$  have a role in the innate and adaptive immune response. In tear fluid, the increase IFN- $\gamma$  level inducing conjunctival goblet cell loss and apoptosis of lacrimal gland cell. In the ocular surface, elevated IFN- $\gamma$  in tear correlated with tear hyperosmolarity and corneal fluorescein. Another study revealed the IFN- $\gamma$  level associated with TBUT<sup>28</sup>. In murine model, high IFN- $\gamma$  expression correlates with severity of conjunctival pathology<sup>23</sup>

## Interleukin 8

The chemokines IL-8 is a potent pro-inflammatory cytokine that are chemotactic for lymphocytes, neutrophils, and basophils. IL-8 significantly

increases in the tear of DED. The IL -8 contributes to the occurrence of dry eye disease through infiltration and activation of T lymphocytes, leading to damage lacrimal gland and ocular surface<sup>18</sup>.

# Tumor necrosis factor alfa

Tumor necrosis factor alfa (TNF-a) is a potent mediator inflammatory produced mainly through macrophages during the acute phase of inflammation. The primary role is responsible for the regulation immune system and selectively cytotoxic, leading to necrosis or apoptosis<sup>24</sup>. TNF-a significantly raised in DED and correlated with MMP activation and limit fibrosis. Multiple studies have concluded that TNF-a related to decreased tear production, disease severity, and correlation with OSDI score and schirmer tear test <sup>8,24-26</sup>. In the dry eye mouse model, TNF-a, IL-1, IL-6 level were detected in the tear fluid and conjunctival epithelial<sup>27</sup>.

# Conclusion

In summary, the identification of molecules in tear fluid as biomarker provide evidence that inflammation is a key feature of the pathophysiology of DED. For dry eye disease, IL-6, TNF-a, IL-8, IL-1 by number studies to consistently reflect disease severity and strong correlation with schirmer tear test and another test for the ocular surface.

# Conflict of Interest: Nil

# Ethical Clearance: Nil

**Funding :** The funding is supported by Dana Hibah Mandat Project, Universitas Airlangga Surabaya, Indonesia.

# References

- Farrand KF, Fridman M, Stillman IO, Schaumberg DA. Prevalence of diagnosed dry eye disease in the United States Among adult aged 18 years and older. Am J Ophthalmol. 2017; 182:90-8
- Lee AJ, Lee J, Saw SM, Gazzard G, KOh D, Widjaja D, Tan DT. Prevalence and risk

factor associated with dry eye symptoms: a population-based study in Indonesia. Br J Ophthalmol.2002.86(12):1347-51

- Akpek EK, Bunya VY, Saldanha IJ. Sjogren syndrome: more than just dry eye. Cornea. 2019.38(5):658-661.
- 4. Chahadva P, Goldhardt R, Galor A, Meibomian gland disease: the role of gland dysfunction in dry eye disease. Ophthalmology.2017.124(11):20-26.
- Shanti Y, Shehada R, Bakkar MM, Qaddumi. Prevalence and associated risk factor of dry eye disease in 16 northern west bank towns in Palestine: a cross-sectional study. BMC Ophthalmology. 2020.20:26.
- Phadatare Sp, Momin M, Nighojar P, Askarkar S, Sing KK, A comprehensive review on dry eye disease: diagnosis, medical management, recent development, and future challenges. Advance in Pharmaceutics. 2015.article ID 704946.
- Cwiklik L. Tear film lipid layer: A molecularlevel view. Biochimic Biophys Acta. 2016.1858(10):2421-30.
- Huang JF, Zhang Y, Kay DR, Pickering EH, McDowell MT,. Evaluation of tear protein markers in dry eye disease: repeatability of measurement and correlation with disease. Invest Ophthalmol Vis Sci. 2012: 53. 4556-4564.
- Nijm LM, Dunbar GE. Understanding the science behind the inflammatory cascade of dry eye disease. US Ophthalmic Review. 2019.12(1):15-16.
- Lee HS, Hattori T, Park EY, Expression of toll-like receptor 4 contributes to corneal inflammation in experimental dry eye disease. Investigation Ophthalmology Visual Science. 2012. 53:5632-5640.
- Chi W, Huan X, Chen X. Mitocondrial DNA oxidation induces imbalances activity of NLRP3/ NLRP6 inflammasomes by activation of caspase -8 and BRCC36 in dry eye. Journal Autoimmune. 2017.80:65-67.

- De Paiva CS, Corrales RM, Villareal AL.Farley WJ, Li DQ, Stern ME, Pflugfelder SC. Corticosteroid and doxycycline suppress MMP-9 and inflammatory cytokines expression, MAPK activation in the corneal epithelium in experimental dry eye. Experimental Eye Research.2006.83:526-535.
- Coursey TG, Henriksson JT, Barbosa FL, De-Paiva CS, Pflugfelder SC. Interferon-gamma induced unfolded protein response in conjunctival goblet cell as a cause of mucin deficiency in Sjogren syndrome. 2016:186(6):1647-1558.
- Baudouin C, Irkec M, Messmer EM, Castilo JB, Bonini S, Figueiredo F, Geerling G, Labetoulle M, Lemp M, Rolando M, Settten GV, Aragona P. Clinical impact of inflammation in dry eye disease: proceedings of the ODISSEY group meeting. Acta Ophthalmol. 2018;96(2):111-119.
- Na KS, Mok JW, Kim JY, Rho CR, Jo CK. Correlations between tear cytokines, chemokines, and soluble receptors and clinical severity of dry eye disease. Investigative Ophthalmology & Visual Science. 2012:53:5443-5450.
- Ju Hj, Byun YS, Mok JW, Joo CK. The blockade of IL-6 counterparts the osmolar stress-induced apoptosis in human conjunctival epithelial cells. Journal of Ophthalmology. 2016:8350134.
- Pflugfelder SC, de Paiva CS. The pathophysiology of dry eye disease: what we know and future direction for research. Ophthalmology. 2017;124(11):4-13.
- Wu X, Chen X, Ma Y, Lin X, Yu X, He S, Luo C, Xu W. Analysis of inflammatory tear molecules and clinical correlations in evaporative dry eye disease caused by Meibomian gland dysfunction. 2020;40:3049-3058.
- Yoon KC, Jeong IY, Park YG, Yang SY. Interleukin-6 and tumour necrosis factor-a levels in tears of patients with dry eye syndrome. Cornea. 2007;26(4):431-437.
- 20. Solomon A, Dursun D, Liu Z, Xie Y, Macri

A, Pflugfelder SC. Pro-and anti-inflammatory forms of interleukin -1 in the tear fluid and conjunctiva of patient with dry eye disease. Investigation Ophthalmology Visual Science. 2001;42(10):2283-2292.

- 21. Amparo F, Dana R. IL-1 receptor antagonist in the treatment of dry eye disease. Expert Review of Ophthalmology. 2003;8(6):581-586.
- Tanaka T, Narazaki M, Kishimoto T. IL-6 in inflammation, Immunity, and Disease. Cold Spring Harbor Perspectives In Biology. 2014;6(10):a016295.
- Jackson DC, Zeng W, Wong CY, Mifsud EJ, Williamson NA, Ang CS, Vingrys AJ, Downie LE. Tear interferon-gamma as a biomarker for evaporative dry eye disease. Investigate Ophthalmology & Visual Science. 2016;57:4824-4830.
- Idriss HT, Naismith JH. TNF-a and the TNF Receptor superfamily: Structure-function relationship. Microscopy Research And Techique.2000;50:184-195.
- Lam H, Bleiden L, de Paiva CS, Farley W, Stern M, Pflugfelder SC. Tear cytokine profiles in dysfunctional tear syndrome. Am J Ophthalmol. 2009;147(2):198-201.
- Fong PY, Shih KC, Lam PY, Chan TC, Jhanji V, Tong L. Role of tear film biomarker in the diagnosis and management of dry eye disease. Taiwan J Ophthalmol.2019;9(3):150-159.
- 27. Ji YW, Byun YJ, Choi W, Jeong E, Kim JS, Noh H, Kim ES, Song YJ, Park SK, Lee HK. Neutralization of ocular surface TNF-a reduces ocular surface and lacrimal gland inflammation induced by in vivo dry eye. Invest Ophthalmol Vis Sci. 2013;54:7557-7566.
- Jackson DC, Zeng W, Wong CY, Mifsud EJ, Williamson NA, Ang CS, Vinggrys AJ, Downie LE. Tear interferon-gamma as a biomarker for Evaporative dry eye disease. Invest Ophthalmol Vis Sci. 2016;57:4824-4830.