

Genetic Identification of Shiga Toxin Encoding Gene from Cases of Multidrug Resistance (MDR) Escherichia coli Isolated from Raw Milk

by Hani Plumeriastuti

Submission date: 12-Jun-2023 02:08PM (UTC+0800)

Submission ID: 2114263542

File name: ug_Resistance_MDR_Escherichia_coli_Isolated_from_Raw_Milk_1.pdf (581.94K)

Word count: 4955

Character count: 26470

Genetic Identification of Shiga Toxin Encoding Gene from Cases of Multidrug Resistance (MDR) *Escherichia coli* Isolated from Raw Milk

Ansharieta^a, M. H. Effendi^{b*}, & H. Plumeriastuti^c

^aMaster Program in Animal Diseases and Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga,

^bDepartment of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga,

^cDepartment of Veterinary Pathology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, 60115, Indonesia

*Corresponding author: mheffendi@yahoo.com

(Received 29-03-2020; Revised 08-07-2020; Accepted 22-07-2020)

ABSTRACT

Escherichia coli is one of bacteria which have resistant to three or more classes of antimicrobial agents. *E. coli* having resistant to three or more classes of antimicrobial drugs can be defined as multidrug-resistant (MDR) bacteria. The aim of the study was to evaluate the expression of Shiga toxin gene in MDR *E. coli*. A total of 250 raw milks samples were taken from dairy farms in Kediri, Probolinggo, Pasuruan, Blitar, and Batu Region, East Java Province, Indonesia. Each sample was cultured into enrichment media Brilliant Green Bile Lactose Broth and Eosin Methylene blue agar, then identified with TSIA agar and IMVIC biochemistry test. Antibiotic sensitivity testing was done using Kirby-Bauer disc diffusion assay on medium Mueller-Hinton agar (Oxoid, CM0337). Antibiotics disks used were 30 µg of Tetracycline (Oxoid, CT0054), 10 µg of Streptomycin (Oxoid, CT0047), 30 µg of Chloramphenicol (Oxoid, CT0013), 5 µg of Trimethoprim (Oxoid, CT0057), and 30 µg of Aztreonam (Oxoid, CT0264). Isolate showing resistance to at least 3 antibiotics disk were then continued with PCR assay to identify Shiga toxin *E. coli* (STEC) encoding *stx2* gene. The study was designed to evaluate the nucleotide analysis of STEC gene. The result showed that 6.46% (1/16) of STEC encoding gene was found in MDR *E. coli*. This report of molecular identification on the presence of STEC gene in MDR *E. coli* confirmed a wider spread of MDR *E. coli* that can threaten animal health and human health.

Keywords: *stx2* gene; MDR; *Escherichia coli*; raw milk; public health

INTRODUCTION

Escherichia coli which produces Shiga toxin (STEC), is defined as a group of *E. coli* which can produce a toxin called Shiga toxin (stx). There are two main types of Shiga toxins found in this STEC group. The two toxins are *stx1* and *stx2* (Hunt, 2010). According to several reports of findings in the field, *stx2* toxin is more common and more prevalent than *stx1* toxin in *E. coli* isolates obtained from livestock feces (Tahamtan *et al.*, 2010). Cattle are believed to be the main reservoir of the STEC bacteria (Hussein & Sakuma, 2005). STEC is often identified with foodborne outbreaks throughout the world with mild symptoms such as mild diarrhea to severe symptoms such as hemolytic uremic syndrome (HUS) (Noris *et al.*, 2012). Foods related to STEC infection are foods without cooking processes, such as raw meat, cheese, non-pasteurized milk or raw milk, fruit and vegetable juices, and other natural ingredients (Mohammadi *et al.*, 2013).

Antimicrobial resistance is a global problem, in which scientific research has been carried out and indicates that this problem has a negative impact on

human and animal health (Ventola, 2015). Irrational treatment and administration of antibiotics greater than the recommended dosage in agriculture increase the incidence of antimicrobial resistance in daily life, and this condition has a great impact on the social and economic aspects (Barriere, 2014). Antibiotic resistance makes health workers lose many antibiotic choices in the treatment of bacterial diseases, causing the high potential for treatment failure in cases of clinical complications (Llor & Bjerrum, 2014). The applications of good hygiene and sanitation practices in animal farms are something that should be emphasized to prevent the distribution of MDR microbes in the populations and the transfer of MDR microbes from animals to humans.

Although antibiotic treatment is not the first choice in cases of infections caused by STEC, multidrug-resistant (MDR) STEC is a public health problem that is quite dangerous because the strain acts as a reservoir of resistant genes (Cointe *et al.*, 2018). MDR STEC bacteria can easily transfer the resistant genes to the other Enterobacteriaceae family bacteria in the body of living organisms or in the environment. Some bacteria

in the gut of living organisms are friends and have the potential to transmit resistance genes to one another (Colavecchio *et al.*, 2017). A number of researches have been conducted on STEC isolates from humans, but only a few have been identified from foodstuffs of animal origins. Report on the status of STEC bacterial resistance to antibiotics is also limited. Therefore, this study was conducted to identify the STEC gene, *stx2*, which is a prevalent form of MDR *E. coli* bacteria isolated from several dairy farms in East Java, Indonesia.

MATERIALS AND METHODS

The sample size in this study was 250 raw milk samples taken from the cow bulk milk of each dairy farm located in Kediri (K), Probolinggo (A), Pasuruan (G), Blitar (S), and Batu (H), East Java, Indonesia, during the period of September-December 2019. The study was conducted by purposive sampling with sample criteria of poor enclosure sanitation and dirty environment (Dairy, 2019). The number of farms calculated on average reached 1500-1800 lactation cows per location. Samples taken at each dairy farm location were 50 samples (cow bulk milk). A total of 25 mL of milk samples from each milk can be taken and placed in sterile plastics. Milk samples were put into a cool box at 4°C for transportation (Arya *et al.*, 2012; Kristianingtyas *et al.*, 2020; Putra *et al.*, 2019).

Each sample was cultured into enrichment media Brilliant Green Bile Lactose Broth (Merck, 105454) and incubated at 37°C for 18-24 hours. The positive results were characterized by the change of media from green to cloudy green color and the presence of gas in Durham tube (Effendi *et al.*, 2019). The positive samples were then cultured in Eosin Methylene Blue (EMB) (Merck, 101347) and incubated at 37°C for 18-24 hours (Putra *et al.*, 2019). The colonies showing metallic green colors were purified and continued to the identification steps with Triple Sugar Iron Agar (TSIA) and Indol-Motility, Methyl Red, Voges Proskauer, and Citrate (IMViC) tests. The positive samples were then stored in LB Broth (Himedia, M1245) for further use in the next testing.

Antibiotic sensitivity testing was done using Kirby-Bauer disc diffusion assay on Mueller-Hinton agar medium (Oxoid, CM0337). Antibiotics disks used were 30 µg of Tetracycline (Oxoid, CT0054), 10 µg of Streptomycin (Oxoid, CT0047), 30 µg of Chloramphenicol (Oxoid, CT0013), 5 µg of Trimethoprim (Oxoid, CT0057), and 30 µg of Aztreonam (Oxoid, CT0264). Interpretation of results was con-

ducted by measuring the diameter of the inhibitory zone formed, based on Clinical and Laboratory Standards Institutions, and the resistance to Aztreonam that is the monobactam can be referred as a presumptive extended spectrum beta lactamase (ESBL) (CLSI, 2018).

Isolate which shown resistance to at least 3 antibiotics disk was then taken from the LB Broth and transferred onto Nutrient Agar (Merck, 105450). The whole cell suspensions were made by mixing approximately 8-10 colonies from Nutrient Agar to 0.3 mL Tris-HCl containing EDTA (TE) buffer. The cell lysate was made by heating the suspension for 10 min with the boiling method. The lysates were centrifuged for 10 min at 8000 rpm to shed the cellular debris. A volume of 5 µL of the supernatant was used as a template for amplification by PCR. The specific primer used in this study was *stx2* according to Brenjchi *et al.* (2011), F: 5'- CCA TGA CAA CGG ACA GCA GTT-3' and R: 5'- CCT GTC AAC TGA GCA CTT-3'.

Each PCR reaction was performed in a 25 µL amplification mixture consisting of 2.5 µL 10 X PCR buffer (500 mM KCl, 200 mM Tris HCl), 0.5 µL dNTPs (10 mM), 1 µL MgCl₂ (50 mM), 1.25 µL of each primer (0.5 µM), 0.2 µL of Taq DNA polymerase (5 unit/µL), and 2 µL of template. The thermocycler (Bio Rad, Hercules, CA) program was started with initial incubation at 94°C for 5 min, followed by 35 cycles of denaturation at 94°C for 60 sec, annealing at 52°C for 30 sec and elongation at 72°C for 60 sec, and a final extension at 72°C for 10 min. The PCR products were separated by electrophoresis in 1.5% agarose gel at 100 V for 40 min in the tris-acetate buffer, visualized by ethidium bromide staining, illuminated by UV-transilluminator, and documented by a gel documentation apparatus. DNA ladder with 100 bp was used as a marker for m-PCR assay. The expected size of products for *stx2* is 779 bp (Brenjchi *et al.*, 2011).

RESULTS

A total of 250 samples were taken from several dairy farms in East Java Province, Indonesia. *E. coli* isolates were found in 176 samples of fresh cow's milk. The *E. coli* isolates were then tested using Kirby Bauer disc diffusion assay with several antibiotics, namely Tetracycline, Streptomycin, Trimethoprim, Chloramphenicol, and Aztreonam. The number of isolates that were resistant to Tetracycline, Streptomycin, Trimethoprim, Chloramphenicol, and Aztreonam was 17.05% (30/176), 14.2% (25/176), 9.66% (17/176), 7.95% (14/176), and 1.7% (3/176), respectively (Table 1).

Table 1. Multidrug resistant *Escherichia coli* from raw cow's milk samples

Location	Sample Size	Confirmed <i>E. coli</i>	Multidrug resistant	Presumptive ESBL
Kediri (K)	50	35	3	0
Probolinggo (A)	50	36	2	0
Pasuruan (G)	50	30	4	1
Blitar (S)	50	38	1	2
Batu (H)	50	37	6	0
Total	250	176	16	3

Note: Presumptive ESBL= presumptive extended spectrum beta lactamase.

A total of 16 isolates were multidrug-resistant *E. coli*, and 3 isolates were isolates of presumptive ESBL producing *E. coli*. Seventeen isolates of MDR *E. coli* and presumptive ESBL that were evaluated in this study had patterns of resistance that were almost the same to the pattern of resistance to aminoglycoside class drug, i.e., Streptomycin (Table 41 and Figure 1).

After finding *E. coli* MDR isolates that were resistant to 34 antibiotics and presumptive ESBL producing *E. coli*, the presence of *stx2* gene was tested by PCR method. From 17 isolates tested by PCR, there was 1 *E. coli* MDR isolates (5.88%) that was found in the sample obtained from Probolinggo (Sample Code A-26), which showed a PCR product of *stx2* gene at the size of 779 bp (Figure 2). However, on Figures 3 and 4 there was no sample tested with a positive result of *stx2* gene.

DISCUSSION

Tetracycline has the highest resistance level because this antibiotic is used most often for dairy farming, followed by the other antibiotics such as those used in this study, namely the aminoglycoside (Streptomycin), macrolide (Chloramphenicol), and sulphonamide (Trimethoprim) groups (Hunter *et al.*, 2010). The use of broad-spectrum antibiotics such as the tetracycline and beta-lactam classes are more common in clinical mastitis cases of dairy cows in Europe, due to their effective treatment results. For respiratory and digestive tract problems, the tetracycline and aminoglycoside groups are the first choice antibiotics, while the second choice is the macrolide group and the combination of sulfonamide-trimethoprim drugs, which have a significant

Table 2. Antimicrobial resistance of multidrug resistant (MDR) isolates

No.	Sample code	Resistant to					Multidrug resistant (MDR)	Presumptive ESBL	Positive to <i>stx2</i> gene
		TE 30 µg	S 10 µg	W 5 µg	C 30 µg	ATM 30 µg			
1.	K-03	√	√	√	√	-	+	-	
2.	K-04	√	√	√	√	-	+	-	
3.	K-36	√	√	-	√	-	+	-	
4.	A-26	√	√	√	√	-	+	+	
5.	A-44	-	√	√	√	-	+	-	
6.	G-12	√	√	√	-	-	+	-	
7.	G-31	-	√	√	-	√	+	-	
8.	G-35	√	√	√	-	-	+	-	
9.	G-43	√	√	-	√	-	+	-	
10.	S-25	√	-	-	√	√	+	-	
11.	S-38	-	√	-	-	√	-	+	
12.	H-11	√	√	√	√	-	+	-	
13.	H-15	√	√	√	-	-	+	-	
14.	H-28	√	√	-	√	-	+	-	
15.	H-37	√	√	-	√	-	+	-	
16.	H-44	√	√	-	√	-	+	-	
17.	H-45	√	√	-	√	-	+	-	

Note: TE= Tetracycline; S= Streptomycin; W= Trimethoprim; C= Chloramphenicol; ATM= Aztreonam; √ = positive resistant. MDR is resistant to three or more classes; ESBL is resistant to ATM; presumptive ESBL= presumptive extended spectrum beta lactamase; Kediri (K), Probolinggo (A), Pasuruan (G), Blitar (S), and Batu (H); += positive; -= negative.




Figure 1. Multidrug resistant (MDR) *Escherichia coli* in Kirby-Bauer disk diffusion test




Figure 2. PCR analysis of *stx2* gene on multidrug resistant (MDR) *Escherichia coli* from raw milk of cows; *stx2* gene indicated by the DNA band at 779 bp. M= marker 100 bp; 1= Negative Control; 2= Positive Control; 3= K-03; 4= K-04; 5= K-36; 6= A-26; 7= A-44




Figure 3. PCR analysis of *stx2* gene on multidrug resistant (MDR) *Escherichia coli* from raw milk of cows; *stx2* gene is indicated by the DNA band at 779 bp. M= marker 100 bp; 1= Negative Control; 2= Positive Control; 3= G-12; 4= G-31; 5= G-35; 6= G-43; 7= S-25; 8= S-38.




Figure 4. PCR analysis of *stx2* gene on multidrug resistant (MDR) *Escherichia coli* from raw milk of cows; *stx2* gene is indicated by the DNA band at 779 bp. M= marker 100 bp; 1= Negative Control; 2= Positive Control. Sample: 3= H-11; 4= H-15; 5= H-28; 6= H-37; 7= H-44; 8= H-45.

impact on microbial activity in the rumen, and the last choice is the cephalosporin class three and four antibiotics (the sulfonamide-trimethoprim drug combination which has a significant impact on microbial activity in the rumen, and the last choice is the cephalosporin class three (Economou and Gousia, 2015).

This resistance to Streptomycin can be attributed to the effectiveness of the streptomycin drug in the treatment of mastitis in dairy cows in Indonesia. According to Riyanto *et al.* (2015), the administration of a combination of Penicillin-Streptomycin antibiotics is still considered as a good therapeutic treatment for dairy cows suffering from mastitis. Even though the level of resistance to the drug Streptomycin observed in this study is quite low (14.2%), there is a need to monitor the use of antibiotics Streptomycin in dairy cows in Indonesia.

Chloramphenicol is not approved for use in food animals in Indonesia refer to Law No. 18 of 2009, concerning Animal Husbandry and Health. The persistence of chloramphenicol resistance in *E. coli* has been observed by the other authors (Tadesse *et al.* 2012), showing that more than 90% of chloramphenicol-resistant *E. coli* isolates were concurrently resistant to tetracycline. In addition, Tadesse's data showed not only the persistence of chloramphenicol but an increasing trend of tetracycline resistance over time among animal *E. coli* isolates. These observations could be explained by co-selection of mobile resistance elements or by the unknown substrate(s) for the chloramphenicol-resistance determinants that serve as a selection pressure.

One method that is often used by various researchers to characterize organisms as Multidrug-Resistant (MDR) is based on the results of antimicrobial susceptibility testing *in vitro*, when researchers test resistance to several antimicrobial agents, classes or subclasses of antimicrobial agents (Kallen *et al.*, 2010). MDR-organism is an organism that is resistant to three or more classes of antimicrobials (Magiorakos *et al.*, 2012), or MDR is Gram-negative bacteria that are resistant to three

or more classes of antimicrobials (Gould, 2008), as is shown in Figure 1. A review of the variability of this definition is given in a comprehensive MDR review by Falagas *et al.* (2006), which is used as a reference by some researchers that a large number of studies do not propose specific definitions for MDR, as were shown in Table 2.

The literature shows that food-producing animals represent the most important source for the inclusion of STEC in the food chain. Most infections in humans are caused by the consumption of food contaminated with STEC. Food-producing animals are important reservoirs of STEC and function as sources of food contamination. Several studies found the presence of STEC, especially in fresh milk products and their processed products (Marshall & Beutin, 2011).

Shiga toxin-producing *Escherichia coli* (STEC) is a subgroup of *Escherichia coli* that are able to produce one or two strong toxins called Shiga 1 and Shiga 2 toxins (*stx1* and *stx2*), which are thought to have additional virulence factors such as intimin responsible for attaching STEC to the cells. Intestinal epithelium and causes the tight attachment and effect of lesions on the intestinal mucosa (Melton-Clegg *et al.*, 2014). This bacterial pathotype is a causative agent of gastroenteritis and can be exacerbated by hemorrhagic colitis (HC) or hemolytic uremic syndrome (HUS), which is a major cause of acute kidney failure in children (Etcheverria & Padola, 2013). Foodborne diseases related to STEC have been documented throughout the world. STEC serotype O157: H7 was reported as the causative agent of a series of outbreaks that were occurred mainly in Canada (Chui *et al.*, 2011), Japan (Kanayama *et al.*, 2013), the United States (Mc Collum *et al.*, 2012), and the United Kingdom (Adams *et al.*, 2016). In Indonesia, STEC was detected in feces (8.3%) and meats (5.3%), both isolated from cattle slaughtered in Qurban festival (Ningrum *et al.*, 2016). In addition, from samples with positive STEC tested in the slaughterhouse in Surakarta, fifteen *E. coli* isolates were detected to contain *stx2* gene (Goma *et al.*, 2019).

Observation of raw milk in Surabaya (11) detected two *E. coli* (2.7%) having stx2 gene (Effendi *et al.*, 2018).

One of the most debated topics in the management of STEC infections lies in the possible risks of using antibiotics in natural infections. Because antibiotics can lyse bacterial cell walls, then release Shiga toxin and cause an increase in Shiga toxin gene expression *in vivo*, generally antibiotics are not recommended for treating STEC infections (Paton & Paton, 1998). Antibiotics have long been used in animal and human medical fields for the treatment, control, and prevention of infectious diseases. However, excessive use and not according to the rules of use can have unanticipated side effects, including the development of antibiotic resistance in bacteria against modern β -lactam antibiotics (Tekiner and Özpınar, 2016) or commonly known as an Extended Spectrum β -lactamase (ESBL). Although STEC infections are not treated aggressively with antibiotic therapy, there have been many STEC isolates that are susceptible to many antibiotics, and recent reports indicate that antimicrobial resistance from STEC also continues to increase (Ahmed & Shimamoto, 2015).

Detection of STEC is labor-intensive and the total time required for strain characterization is usually 72 h. On the other hand, molecular methods are sensitive, specific, and a quick approach to detection and characterization of microbiological contaminants in food. The molecular characterization of STEC is performed by means of PCR. The study showed that among cases of MDR from *E. coli* samples, 5.88% contained STEC encoding gene. Environmental contamination, herd management, and poor milking practices are important causes of milk degradation. It has been shown that food animals are important sources of STEC's entry into the food chain (Martin and Beutin, 2011).

The finding of STEC bacteria as contaminants from raw cow milk samples in this study can be considered as important and worthy results, referring to the study of Newell and La Ragione (2018), which states that with only small amounts, STEC colonies found in food samples of animal origin can cause digestive and urogenital disorders. For this reason, housing management and good sanitation practices during the process of milking cows, as well as their storage and distribution, should be improved even better so as to keep consumers away from foodborne diseases originating from raw cow's milk or other dairy products. Therefore, hygiene practices and strict management for dairy herds, processing, and storage of milk should be adopted to avoid undesirable illness due to contaminated milk, and it is important to encourage people to pasteurize milk for consumption.

CONCLUSION

Molecular identification showed that Shiga toxin encoding gene was found in raw cow milk from several regions with cases of Multidrug-Resistance (MDR) of *Escherichia coli* in East Java, Indonesia. These results show that raw cow's milk has a potential for spreading and poses a threat to public health from *E. coli* isolates.

CONFLICT OF INTEREST

The author states that there are no conflicts of interest with financial, personal, or other relationships with other people or organizations related to the material discussed in the manuscript.

ACKNOWLEDGEMENT

This study was supported in part with the Penelitian Hibah Mandat funding from Airlangga University, Indonesia in fiscal year 2019.

REFERENCES

- Adams, N. L., L. Byrne, G. A. Smith, R. Elson, J. P. Harris, R. Salmon, R. Smith, S. J. O'Brien, G. K. Adak, & C. Jenkins. 2016. Shiga toxin-producing *Escherichia coli* O157, England and Wales, 1983-2012. *Emerg. Infect. Dis.* 22 : 590-597. <https://doi.org/10.3201/eid2204.151485>
- Ahmed, A. M. & T. Shimamoto. 2015. Molecular analysis of multidrug resistance in Shiga toxin-producing *Escherichia coli* O157:H7 isolated from meat and dairy products. *Int. J. Food Microbiol.* 193: 68-73. <https://doi.org/10.1016/j.ijfoodmicro.2014.10.014>
- Arya, R., B. Antonisamy, & S. Kumar. 2012. Sample size estimation in prevalence studies. *Indian J. Pediatr.* 79: 1482-1488. <https://doi.org/10.1007/s12098-012-0763-3>
- Barriere, S. L. 2014. Clinical, economic and societal impact of antibiotic resistance. *Expert Opin. Pharmacother.* 16: 151-153. <https://doi.org/10.1517/14656566.2015.983077>
- Brenjchi, M., A. Jamshidi, N. Farzaneh, & MR.Bassami. 2011. Identification of Shiga toxin-producing *Escherichia coli* O157:H7 in raw milk samples from dairy farms in Mashhad using multiplex PCR assay. *Iranian Journal of Veterinary Research.* 12: 145-149.
- Chui, L., M. C. Lee, K. Malejczyk, L. Lim, D. Fok, & P. Kwong. 2011. Prevalence of Shiga toxin-producing *Escherichia coli* as Detected by Enzyme-Linked immunoassays and real-time PCR during the summer months in Northern Alberta, Canada. *J. Clin. Microbiol.* 49: 4307-4310. <https://doi.org/10.1128/JCM.05211-11>
- CLSI. 2018. M100 Performance Standards for Antimicrobial Susceptibility Testing. 28th ed. Clinical and Laboratory Standards Institute, USA.
- Cointe, A., A. Birgy, P. Mariani-Kurkdjian, S. Liguori, C. Courroux, J. Blanco, S. Delannoy, P. Fach, E. Loukiadis, P. Bidet, & S. Bonacorsi. 2018. Emerging multidrug-resistant hybrid pathotype Shiga toxin-producing *Escherichia coli* O80 and related strains of clonal complex 165, Europe. *Emerg. Infect. Dis.* 24: 2262-2269. <https://doi.org/10.3201/eid2412.180272>
- Colavecchio, A., B. Cadieux, A. Lo, & L. D. Goodridge. 2017. Bacteriophages contribute to the spread of antibiotic resistance genes among foodborne pathogens of the *Enterobacteriaceae* family - A Review. *Front. Microbiol.* 8: 1108. <https://doi.org/10.3389/fmicb.2017.01108>
- Dairy, N. Z. 2019. Dairy Cow Housing-A Good Practice Guide for Dairy Housing in New Zealand, Version 2. DairyNZ and Ministry for Primary Industries, Hamilton, New Zealand.
- Economou, V. & P. Gousia. 2015. Agriculture and Food Animals as A Source of Antimicrobial-Resistant Bacteria. *Infect. Drug Resist.* 8: 49-61. <https://doi.org/10.2147/IDR.S55778>
- Effendi, M. H., N. Harijani, S. M. Yanestria, & P. Hastutie. 2018. Identification of Shiga toxin-producing *Escherichia coli* in raw milk samples from dairy cows in Surabaya,

- Indonesia. Philipp. J. Vet. Med. 55: 109-114.
- Effendi, M. H., N. Harijani, Budiarto, N. P. Triningtya, W. Tyasningsih, & H. Plumeriastuti. 2019. Prevalence of pathogenic *Escherichia coli* isolated from subclinical mastitis in East Java Province, Indonesia. Indian Vet. J. 96: 22-25.
- Etcheverria A. I. & N. L. Padola. 2013. Shiga toxin-producing *Escherichia coli*, factors involved in virulence and cattle colonization. Virulence 4: 366-372. <https://doi.org/10.4161/viru.24642>
- Falagas, M.E., P.K. Koletsis, & I. A. Bliziotis. 2006. The diversity of definitions of multidrug-resistant (MDR) and pandrug-resistant (PDR) *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. J. Med. Microbiol. 55: 1619-1629. <https://doi.org/10.1099/jmm.0.46747-0>
- Goma, MKE, A. Indraswari, A. Haryanto, & D.A. Widiasih. 2019. Detection of *Escherichia coli* O157:H7 and Shiga toxin 2a gene in pork, pig feces, and clean water at Jagalan slaughterhouse in Surakarta, Central Java Province, Indonesia. Vet. World, 12(10): 1584-1590. <https://doi.org/10.14202/vetworld.2019.1584-1590>
- Gould, I. M. 2008. The epidemiology of antibiotic resistance. Int. J. Antimicrob. Agents. 32: 2-9. <https://doi.org/10.1016/j.ijantimicag.2008.06.016>
- Hunt, J. M. 2010. Shiga toxin-producing *Escherichia coli* (STEC). Clin. Lab. Med. 30: 21-45. <https://doi.org/10.1016/j.cll.2009.11.001>
- Hunter, P. A., S. Dawson, G. L. French, H. Goossens, P. M. Hawkey, E. J. Kuijper, D. Nathwani, D. J. Taylor, C. J. Teale, R. E. Warren, M. H. Wilcox, N. Woodford, M. W. Wulf, & L. J. Piddock. 2010. Antimicrobial-resistant pathogens in animals and man: Prescribing, Practices and Policies. J. Antimicrob. Chemother. 65: i3-i17. <https://doi.org/10.1093/jac/dkp433>
- Hussein, H. S. & T. Sakuma. 2005. Invited Review: Prevalence of Shiga Toxin-producing *Escherichia coli* in dairy cattle and their products. J. Dairy Sci. 88: 450-465. [https://doi.org/10.3168/jds.S0022-0302\(05\)72706-5](https://doi.org/10.3168/jds.S0022-0302(05)72706-5)
- Kallen, A. J., A. I. Hidron, J. Patel, & N. A. Srinivasa. 2010. Multidrug resistance among gram-negative pathogens that caused healthcare-associated infections reported to The National Healthcare Safety Network, 2006-2008. Infect. Control Hosp. Epidemiol. 31: 528-531. <https://doi.org/10.1086/652152>
- Kanayama, A., Y. Yahata, Y. Arima, T. Takahashi, T. Saitoh, K. Kanou, K. Kawabata, T. Sunagawa, T. Matsui, & K. Oishi. 2015. Enterohemorrhagic *Escherichia coli* outbreaks related to childcare facilities in Japan, 2010-2013. BMC Infectious Diseases. 15: 539. <https://doi.org/10.1186/s12879-015-1259-3>
- Kristianingtyas, L., M. H. Effendi, W. Tyasningsih, & F. Kurniawan. 2020. Genetic identification of bla_{CTX-M} gene and bla_{TEM} gene on extended spectrum beta lactamase (ESBL) producing *Escherichia Coli* from Dogs. Indian Vet. J. 97:01:17-21.
- Llor, C. & L. Bjerrum. 2014. Antimicrobial resistance: risk associated with antibiotic overuse and initiatives to reduce the problem. Ther. Adv. Drug Saf. 5: 229-241. <https://doi.org/10.1177/2042098614554919>
- Magiorakos, A. P., A. Srinivasan, R. B. Carey, Y. Carmeli, M. E. Falagas, C. G. Giske, S. Harbarth, J. F. Hindler, G. Kahlmeter, B. Olsson-Liljequist, D. L. Paterson, L. B. Rice, J. Stelling, M. J. Struelens, A. Vatopoulos, J. T. Weber, & D. L. Monnet. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: An international expert proposal for interim standard definitions for acquired resistance. Clin. Microbiol. Infect. 18: 268-281. <https://doi.org/10.1111/j.1469-0691.2011.03570.x>
- Martin, A. & L. Beutin. 2011. Characteristics of Shiga toxin-producing *Escherichia coli* from meat and milk products of different origins and association with food producing animals as main contamination sources. Int. J. Food Microbiol. 146: 99-104. <https://doi.org/10.1016/j.ijfoodmicro.2011.01.041>
- Mc Collum, J. T., N. J. Williams, S. W. Beam, S. Cosgrove, P. J. Ettestad, T. S. Ghosh, A. C. Kimura, L. Nguyen, S. G. Stroika, R. L. Vogt, A. K. Watkins, J. R. Weiss, I. T. Williams, & A. B. Cronquist. 2012. Multistate outbreak of *Escherichia coli* O157:H7 infections associated with in-store sampling of an aged raw-milk Gouda cheese. J. Food Prot. 75: 1759-1765. <https://doi.org/10.4315/0362-028X.JFP-12-136>
- Melton-Celsa, A. R. 2014. Shiga Toxin (Stx) classification, structure and function. Microbiol. Spectr. 2: EHEC-0024-2013. <https://doi.org/10.1128/microbiolspec.EHEC-0024-2013>
- Mohammadi, P., R. Abiri, M. Rezaei, & S. Salmanzadeh-Ahrabi. 2013. Isolation of Shiga toxin-producing *Escherichia coli* from raw milk in Kermanshah, Iran. Iran. J. Microbiol. 5: 233-238.
- Newell, D. G. & R. M. La Razione. 2018. Enterohaemorrhagic and other Shiga toxin-producing *Escherichia coli* (STEC): where are we now regarding diagnostics and control strategies?. Transbound. Emerg. Dis. 65: 49-71. <https://doi.org/10.1111/tbed.12789>
- Ningrum, S. G., R. D. Soejoedono, H. Latif, W. Arnafia, & I. W. T. Wibawan. 2016. Prevalence and characterization of Shiga toxin-producing *Escherichia coli* isolated from slaughtered qurban animal in Jakarta Province. Med. Pet., 39: 90-94. <https://doi.org/10.5398/medpet.2016.39.2.90>
- Noris, M., F. Mescia, & G. Remuzzi. 2012. STEC-HUS, Atypical HUS and TTP are all diseases of complement activation. Nat. Rev. Nephrol. 8: 622-633. <https://doi.org/10.1038/nrneph.2012.195>
- Paton J. C. & A. W. Paton. 1998. Pathogenesis and diagnosis of Shiga toxin-producing *Escherichia coli* infections. Clin. Microbiol. Rev. 11: 450-479. <https://doi.org/10.1128/CMR.11.3.450>
- Putra, A. R. S., M. H. Effendi, S. Koesdarto, & W. Tyasningsih. 2019. Molecular identification of Extended Spectrum Beta-Lactamase (ESBL) producing *Escherichia coli* isolated from dairy cows in East Java Province, Indonesia. Indian Vet. J. 96: 26-30.
- Riyanto, J., Sunarto, B. S. Hertanto, M. Cahyadi, R. Hidayah, & W. Sejati. 2016. Produksi dan kualitas susu sapi perah penderita mastitis yang mendapat pengobatan antibiotik. Sains Peternak. 14: 30-41. <https://doi.org/10.20961/sainspet.14.2.30-41>
- Tadesse, D. A., S. Zhao, E. Tong, S. Ayers, A. Singh, M.J. Bartholomew, and P.F. McDermott. 2012. Antimicrobial drug resistance in *Escherichia coli* from humans and food animals, United States, 1950-2002. Emerging Infectious Diseases. 18: 741-749. <https://doi.org/10.3201/eid1805.111153>
- Tahamtan, Y., M. Hayati, & M. Namavari. 2010. Prevalence and distribution of the stx1, stx2 genes in Shiga toxin producing *E. coli* (STEC) isolates from cattle. Iran. J. Microbiol. 2: 8-13.
- Tekiner, I. H. & H. Özpınar. 2016. Occurrence and characteristics of extended spectrum Beta-Lactamases-producing *Enterobacteriaceae* from foods of animal origin. Braz. J. Microbiol. 47: 444-451. <https://doi.org/10.1016/j.bjm.2015.11.034>
- Ventola, C. L. 2015. The antibiotic resistance crisis: part 1: causes and threats. Pharm. Ther. 40: 277-283.

Genetic Identification of Shiga Toxin Encoding Gene from Cases of Multidrug Resistance (MDR) Escherichia coli Isolated from Raw Milk

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 www.researchgate.net 1%

Internet Source
- 2 Ahmed A. Ramadan, Neveen A. Abdelaziz, Magdy A. Amin, Ramy K. Aziz. "Novel blaCTX-M variants and genotype-phenotype correlations among clinical isolates of extended spectrum beta lactamase-producing Escherichia coli", Scientific Reports, 2019 1%

Publication
- 3 researcherslinks.com 1%

Internet Source
- 4 Rituparna Tewari, Susweta Mitra, Feroze Ganaie, Sangita Das et al. "Dissemination and characterization of extended spectrum β -lactamase, AmpC β -lactamase and metallo β -lactamase producing Escherichia coli from livestock and poultry in Northeastern India: A molecular surveillance approach", Journal of Global Antimicrobial Resistance, 2019 1%

Publication

5	bmccardiovascdisord.biomedcentral.com Internet Source	1 %
6	slovetres.si Internet Source	1 %
7	Azucena Mora, Jesús E. Blanco, Miguel Blanco, M. Pilar Alonso et al. "Antimicrobial resistance of Shiga toxin (verotoxin)-producing Escherichia coli O157:H7 and non-O157 strains isolated from humans, cattle, sheep and food in Spain", Research in Microbiology, 2005 Publication	1 %
8	Zoonoses - Infections Affecting Humans and Animals, 2015. Publication	1 %
9	www.scielo.br Internet Source	1 %
10	mspace.lib.umanitoba.ca Internet Source	1 %
11	Maria Cecília Cergole-Novella, Lucilia S. Nishimura, Kinue Irino, Tãçnia Mara I. Vaz et al. " genotypes and antimicrobial resistance profiles of Shiga toxin-producing strains isolated from human infections, cattle and foods in Brazil ", FEMS Microbiology Letters, 2006 Publication	<1 %

12

QIAN WU, MEILI XI, XIAOYING LV, YUNFENG XU, YUQING FENG, QIONG LI, QINNAN YANG, XIAODONG XIA. "Presence and Antimicrobial Susceptibility of Escherichia coli Recovered from Retail Chicken in China", Journal of Food Protection, 2014

Publication

<1 %

13

agritrop.cirad.fr

Internet Source

<1 %

14

Salwa MMasoud, Reham Ibrahim, Rehab Abd El-Baky, Sherine Abd El-Rahman. "Prevalence of Blactx-M Gene Among MDR Escherichia Coli Isolated from Wound Infections in Minia University Hospital, Egypt", Minia Journal of Medical Research, 2022

Publication

<1 %

15

Rituparna Tewari, Susweta Mitra, Feroze Ganaie, Sangita Das et al. "Dissemination and characterisation of Escherichia coli producing extended-spectrum β -lactamases, AmpC β -lactamases and metallo- β -lactamases from livestock and poultry in Northeast India: A molecular surveillance approach", Journal of Global Antimicrobial Resistance, 2019

Publication

<1 %

16

www.animbiosci.org

Internet Source

<1 %

17

www.coursehero.com

Internet Source

<1 %

18

Ashraf M. Ahmed, Tadashi Shimamoto. "Molecular analysis of multidrug resistance in Shiga toxin-producing Escherichia coli O157:H7 isolated from meat and dairy products", International Journal of Food Microbiology, 2015

Publication

<1 %

19

profdoc.um.ac.ir

Internet Source

<1 %

20

Grzegorz Węgrzyn, Maite Muniesa. "Editorial: Shiga Toxin-Converting Bacteriophages", Frontiers in Microbiology, 2021

Publication

<1 %

21

Monalisa S. M. Souto, Fernanda M. Coura, Elaine M. S. Dorneles, Ana Paula R. Stynen et al. "Antimicrobial susceptibility and phylotyping profile of pathogenic Escherichia coli and Salmonella enterica isolates from calves and pigs in Minas Gerais, Brazil", Tropical Animal Health and Production, 2016

Publication

<1 %

22

ejurnal.undana.ac.id

Internet Source

<1 %

23

www.nrcresearchpress.com

Internet Source

<1 %

24	d-nb.info Internet Source	<1 %
25	mail.scialert.net Internet Source	<1 %
26	ugspace.ug.edu.gh Internet Source	<1 %
27	www.foodstandards.gov.scot Internet Source	<1 %
28	"Escherichia coli in the Americas", Springer Science and Business Media LLC, 2016 Publication	<1 %
29	A. Bentancor, M.V. Rumi, M.V. Gentilini, C. Sardoy, K. Irino, A. Agostini, A. Cataldi. " Shiga toxin-producing and attaching and effacing in cats and dogs in a high hemolytic uremic syndrome incidence region in Argentina ", FEMS Microbiology Letters, 2007 Publication	<1 %
30	Himmatul Khasanah, Hidayat Bambang Setyawan, Roni Yulianto, Desy Cahya Widianingrum. "Subclinical mastitis: Prevalence and risk factors in dairy cows in East Java, Indonesia", Veterinary World, 2021 Publication	<1 %
31	S. Bandyopadhyay, C. Lodh, H. Rahaman, D. Bhattacharya et al. "Characterization of shiga	<1 %

toxin producing (STEC) and enteropathogenic Escherichia coli (EPEC) in raw yak (Poephagus grunniens) milk and milk products", Research in Veterinary Science, 2012

Publication

32

Siobhán C. McCarthy, Guerrino Macori, Gina Duggan, Catherine M. Burgess, Séamus Fanning, Geraldine Duffy. "Prevalence and Whole-Genome Sequence-Based Analysis of Shiga Toxin-Producing Escherichia coli Isolates from the Recto-Anal Junction of Slaughter-Age Irish Sheep", Applied and Environmental Microbiology, 2021

Publication

<1 %

33

Yasuko Kondo, Seiji Kondo. "Chapter 9 Telomerase RNA Inhibition Using Antisense Oligonucleotide Against Human Telomerase RNA Linked to a 2',5'-Oligoadenylate", Springer Science and Business Media LLC, 2007

Publication

<1 %

34

is.muni.cz
Internet Source

<1 %

35

journal.ipb.ac.id
Internet Source

<1 %

36

journals.plos.org
Internet Source

<1 %

37	rd.springer.com Internet Source	<1 %
38	www.dovepress.com Internet Source	<1 %
39	www.theses.fr Internet Source	<1 %
40	<p>Jay N. Worley, Kristopher A. Flores, Xun Yang, Jennifer A. Chase, Guojie Cao, Shuai Tang, Jianghong Meng, Edward R. Atwill. "Prevalence and Genomic Characterization of Escherichia coli O157:H7 in Cow-Calf Herds throughout California", Applied and Environmental Microbiology, 2017</p> Publication	<1 %
41	<p>NAHLA O. ELTAI, HADI M. YASSINE, TAHRA EL-OBEID, SARA H. AL-HADIDI, ASMAA A. AL THANI, WALID Q. ALALI. " Prevalence of Antibiotic-Resistant Isolates from Local and Imported Retail Chicken Carcasses ", Journal of Food Protection, 2020</p> Publication	<1 %
42	<p>Rokhaya Gueye, Viviane Chatchueng Sandefo, Babacar Beye, Elhadji Ousmane Faye et al. "Assessment of Poultry Feed Contamination Level by Aflatoxin B1: Quantification by Two Chromatographic Analysis Methods", Food and Nutrition Sciences, 2022</p>	<1 %

43

Sümeyye Betül BOZATLI, Abdullah DİKİCİ, Bülent ERGÖNÜL. "Determination of the changes in the gastric fluid endurance of O157 and non-O157 Shiga toxin-producing Escherichia coli during storage of experimentally produced beef frankfurter", *Journal of Food Science and Technology*, 2020

Publication

<1 %

44

Yashini Naidoo, Angel Valverde, Errol D. Cason, Rian E. Pierneef, Don A. Cowan. "A clinically important, plasmid-borne antibiotic resistance gene (β -lactamase TEM-116) present in desert soils", *Science of The Total Environment*, 2020

Publication

<1 %

45

epublications.marquette.edu

Internet Source

<1 %

46

hdl.handle.net

Internet Source

<1 %

47

repository.library.du.ac.bd:8080

Internet Source

<1 %

48

studentsrepo.um.edu.my

Internet Source

<1 %

49

www.ajol.info

Internet Source

<1 %

50

www.inabj.org

Internet Source

<1 %

51

www.islandscholar.ca

Internet Source

<1 %

52

Aty Widyawaruyanti, Muhammad Asrory, Wiwied Ekasari, Dwi Setiawan, Achmad Radjaram, Lidya Tumewu, Achmad Fuad Hafid. "In vivo Antimalarial Activity of Andrographis Paniculata Tablets", *Procedia Chemistry*, 2014

Publication

<1 %

53

Bill Clinton Aglomasa, Cynthia Kyerewaa Adu-Asiamah, Samuel Opoku Asiedu, Priscilla Kini et al. "Multi-drug resistant bacteria isolates from lymphatic filariasis patients in the Ahanta West District, Ghana", *BMC Microbiology*, 2022

Publication

<1 %

54

Trott, Darren. " β -lactam Resistance in Gram-negative Pathogens Isolated from Animals", *Current Pharmaceutical Design*, 2013.

Publication

<1 %

55

Bartłomiej Grygorcewicz, Wioleta Chajęcka - Wierzchowska, Adrian Augustyniak, Agata Wasak et al. "In - milk inactivation of O157:H7 by the environmental lytic bacteriophage ECPS - 6 ", *Journal of Food Safety*, 2019

Publication

<1 %

56

Ibraheem Ghali-Mohammed, Ismail Ayoade Odetokun, Ibrahim Adisa Raufu, Nma Bida Alhaji, Victoria Olusola Adetunji. "Prevalence of Escherichia coli O157 isolated from marketed raw cow milk in Kwara State, Nigeria", Scientific African, 2023

Publication

<1 %

57

Kamelia M. Osman, Ashgan M. Mustafa, Magdy A.K. Aly, Ghada S. AbdElhamed. " Serotypes, Virulence Genes, and Intimin Types of Shiga Toxin-Producing and Enteropathogenic Isolated from Mastitic Milk Relevant to Human Health in Egypt ", Vector-Borne and Zoonotic Diseases, 2012

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Genetic Identification of Shiga Toxin Encoding Gene from Cases of Multidrug Resistance (MDR) Escherichia coli Isolated from Raw Milk

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
