

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/355546379>

# Detection of Knockdown-Resistance Mutations (V1016G and F1534C) in Dengue Vector from Urban Park, Surabaya, Indonesia

Article in *Journal of Tropical Biodiversity and Biotechnology* · October 2021

DOI: 10.22146/jtbb.65357

CITATIONS

0

READS

59

8 authors, including:



**Shifa Fauziyah**

Akademi Analisis Kesehatan Delima Husada Gresik

23 PUBLICATIONS 38 CITATIONS

[SEE PROFILE](#)



**Sri Subekti**

Airlangga University

85 PUBLICATIONS 249 CITATIONS

[SEE PROFILE](#)



**Budi Utomo**

Airlangga University

83 PUBLICATIONS 104 CITATIONS

[SEE PROFILE](#)



**Teguh Hari Sucipto**

Airlangga University

45 PUBLICATIONS 267 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Relationship between d-dimer level and clinical severity of sepsis [View project](#)



Keanekaragaman kupu-kupu [View project](#)

## Research Article

# Detection of Knockdown-resistance Mutations (V1016G and F1534C) in Dengue Vector from Urban Park, Surabaya, Indonesia

Shifa Fauziyah<sup>1</sup>, Sri Subekti<sup>2,3\*</sup>, Budi Utomo<sup>4</sup>, Teguh Hari Sucipto<sup>5</sup>, Hebert Adrianto<sup>6</sup>, Aryati<sup>5,7</sup>, Puspa Wardhani<sup>5,7</sup>, Soegeng Soegijanto<sup>5</sup>

1) Institute of Tropical Disease, Universitas Airlangga, Mulyorejo Street, Mulyorejo 60115, Surabaya, East Java, Indonesia

2) Faculty of Fisheries and Marine Science, Universitas Airlangga, Mulyorejo Street, Mulyorejo 60115, Surabaya, East Java, Indonesia

3) Entomology Study Group, Institute of Tropical Disease, Universitas Airlangga, Mulyorejo Street, Mulyorejo 60115, Surabaya, East Java, Indonesia

4) Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga, Prof. Dr. Moestopo Street, Tambaksari 60132, Surabaya, East Java, Indonesia

5) Dengue Study Group, Institute of Tropical Disease, Universitas Airlangga, Mulyorejo Street, Mulyorejo 60115, Surabaya, East Java, Indonesia

6) School of Medicine, Universitas Ciputra, CitraLand CBD Boulevard, Sambikerep 60219, Surabaya, East Java, Indonesia

7) Clinical Pathology Department, Faculty of Medicine, Universitas Airlangga, Prof. Dr. Moestopo Street, Tambaksari 60132, Surabaya, East Java, Indonesia

\* Corresponding author, email: denguelpt@gmail.com

Submitted: 23 April 2021; Accepted: 21 August 2021; Published online: 22 October 2021

## ABSTRACT

An urban park is potentially a source of vector-borne disease transmission due to it being a natural and artificial mosquito breeding habitats combined with people's continuous presence. Thus, this study aims to screen the occurrence of knockdown-resistance (*kdr*) mutant alleles (V1016G and F1534C) in mosquito populations collected from urban parks in Surabaya, Indonesia. Cross sectional study was conducted in July 2019. A total of 28 ovitraps were installed in seven urban parks, having four ovitraps installed in each park. In total, 1,662 eggs were collected, and only 187 emerged into adult mosquitoes, consisting of 97 *Aedes (Stegomyia) aegypti* and 90 *Aedes (Stegomyia) albopictus*. All-female adult mosquitoes (n=55) were tested using allele-specific polymerase chain reaction assay (AS-PCR) to detect voltage gated sodium channel (VGSC) gene mutations. This study found no mutations in Valine to Glycine mutation in point 1016 (V1016G) and Phenylalanine to Cysteine in point 1534 (F1534C) alleles in both two species. All of mosquito samples have wild type genotype of *kdr* alleles (V1016V and F1534F). Data were analysed using R Studio 1.4 Version by *Genetics* package. Results showed that the frequency of resistant alleles (G1016 and C1534) was zero, and the frequency of susceptible allele was 1 (V1016 and F1534). Insecticide bioassay could not be established due to the limited number of adult mosquitoes, so insecticide resistance status could not be determined. However, this study can be used as preliminary monitoring for the vector control program.

**Keywords:** dengue, insecticide, *kdr* allele, mosquito, Surabaya, urban parks

## INTRODUCTION

Mosquitoes can spread and carry diseases so that it can making them as one of the deadliest animals. Many mosquito-borne diseases still show an increasing number, including dengue, Zika, chikungunya, West-Nile Virus, malaria, and yellow fever. Half of the population worldwide have lived in an area where mosquitoes are present. Four genus are commonly found as the vector of mosquito-borne diseases, such as *Aedes*, *Culex*, *Mansonia*, and *Anopheles*. Genus of *Aedes* can transmit chikungunya virus, Zika virus, dengue virus, lymphatic filariasis, and yellow fever virus (World Health Organization 2020b). Two species of *Aedes*, including *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus*, are the important vector of arboviral disease (Reinert et al. 2009). Dengue infection was a disease transmitted by both of them and became a burden for public health (Simmons et al. 2012).

Data from WHO in 2020 shows that approximately 390 million people have been estimated infected by dengue infection and distributed across 128 countries (World Health Organization 2020a) Indonesia was also endemic to dengue infection, increasing its annual incidence rate from 0.05/100,000 in 1968 to 78.8/100,000 in 2016. However, the case fatality rate was decreased from 41% in 1968 (Nathin et al. 1988) to 1.21% in 2004 (World Health Organization 2006).

Currently, there is no effective vaccine that can be a prophylaxis for all age groups. Dengvaxia (CYD-TDV) is a live recombinant tetravalent dengue vaccine by Sanofi-Pasteur that was first licensed and can be used for individuals 9 to 45 years old of age individuals in endemic areas. The efficacy of this vaccine was varied, with the highest efficacy being against dengue serotypes 3 and 4 (71.6% and 76.9%) followed by dengue serotypes 1 and 2 (54.7% and 43%) (World Health Organization 2020c). Following that, the primary method to prevent dengue infection transmission is through vector control. A meta-analysis study showed that varieties of dengue control variation significantly reduces dengue risk, such as house screening, water-container cover, and community-based environmental management. Interestingly, indoor residual spraying (IRS) did not significantly reduce dengue risk, while the use of insecticide aerosol and mosquito coils was associated with increased dengue risk. In line with that, skin repellent, insecticide-treated bed nets were also had no effects (Bowman et al. 2016). Thus, the efficacy of vector control remained in question and maybe can vary in different geographical areas.

The primary method of dengue infection control in Indonesia is the combination of environmental management (eradicating larval mosquito, covering the water-container, draining the bathtub regularly) and thermal fogging. In Indonesia, Organophosphates have been used for a long decade to control adult mosquitoes. Malathion, as the derivative of Organophosphate, was used to control adult mosquitoes, while temephos was used as larvicides. Malathion was firstly introduced in Indonesia in 1969, while temephos was introduced in 1980 for dengue control (Hardjanti et al. 2015). The intensive and massive use of insecticide can lead to the sensitivity decreasing of mosquito population against commonly-used insecticide. This phenomenon can be called as insecticide-resistance mechanism (World Health Organization 2012). There are four types of insecticide-resistance: metabolic resistance, target-site resistance, cuticular resistance, and behavioural resistance (Corbel & Guessan 2013).

Metabolic resistance is a mechanism in insects enzyme system to naturally detoxify from insecticides exposure. The over-expression of enzymes that can detox and amino acid substitution within the enzyme can increase the enzyme's affinity, resulting in metabolic resistance (David et al. 2007; Hemingway et al. 2004). The decreasing effectivity of the site of action

in mosquito to bind insecticide called target-site resistance, such as the target site of carbamates and organophosphate insecticides is acetylcholinesterase (AChE) in the nerve cell (Fournier 2005). The insect can also develop cuticular resistance as the impact of the reduced uptake of insecticide and the increasing thickness of the cuticle, but the study of this resistance is still limited in *Anopheles* genus (Djouaka et al. 2008). Another mechanism of insecticide resistance contributing to vector control failure is behavioural resistance, that is the mosquito's ability to avoid insecticide exposure. The investigation of this mechanism was also limited and only found in *Anopheles* population (Russell et al. 2011).

Detection of insecticide resistance in mosquito populations can be conducted through various methods, such as biochemical assay and bioassays using WHO diagnosed doses, dose-response bioassays, and molecular assays. Molecular assays are the most sensitive way to predict the possibility of insecticide-resistance in the future. This method can determine the frequency of resistant allele in a population (Ranson et al. 2011). Point mutations of *kdr* have been revealed in some countries worldwide, such as point mutations G923V (Glisin to Valin) and I1011M (Isoleusin to Metionin) found in Brazil, Guyana, and Martinique (Bregues et al. 2003), L982W (Leusin to Tryptofan) found in Vietnam (Bregues et al. 2003), F1534C (Phenylalanin to Cystein) found in Indonesia and Thailand (Kawada et al. 2014; Wuliandari et al. 2020), and V1016G (Valin to Glisin) widely distributed in Indonesia and Thailand (Bregues et al. 2003; Rajatileka et al. 2008; Wuliandari et al. 2015). The mutations are associated with the exposure of permethrin and DDT. In several regions of Indonesia, mutant allele was reported in Kuningan (Jakarta), Padang, Samarinda, Pontianak, Mataram, Denpasar, Dompu, West Manggarai, and East Sumba with the frequency 0.73; 0.6; and 0.02 respectively for V1016G, S989P, and F1534C (Amelia-Yap et al. 2019). As the first place where dengue infection was found in 1968, Surabaya also used malathion as the insecticide for vector control, but the resistant alleles report in mosquito from these populations is still yet detected. This study aims to screen the occurrence of knockdown-resistance (*kdr*) mutant alleles in Surabaya, Indonesia. The public urban park is the chosen study site due to its importance value as the public facility.

## MATERIALS AND METHODS

### Materials

Mosquito eggs from the field were reared until they emerged to be adult mosquitoes, consisting of two species, namely *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus*. Rearing processes were in accordance to the guidelines mentioned in procedure paragraph. Materials that were used in this research were QIAamp Viral RNA Extraction Kit (Qiagen, Hilden, Germany), NEXscript<sup>TM</sup> RT-PCR 2x Master Mix (NEX<sup>TM</sup> Diagnostic, USA), Nuclease Free Water (NFW), 70% ethanol, agarose gel, ethidium bromide, TBE buffer, sugar solution, cotton, and labelled paper. Instruments used were micropipette, tray, plastic pipette, board marker, microcentrifuge, freezer -80°C, vortex mixer, biosafety cabinet level-2, thermal cycler Bio Rad, Erlenmeyer, gel documentation imaging system, and stereo microscope.


### Methods

#### Study Area

Surabaya is in Java Island and is highlighted as the most populated city in East Java, resided by two million citizens of various ethnicities. This situation can lead to urbanization and affect the transmission of mosquito-borne diseases, including dengue infection. This study was conducted in seven

**Table 1.** Details of parks that have successfully had mosquito hatched eggs, collection date, and their coordinates.

Collection Site	District	Coordinates	Collection Date
Apsari	Genteng	7°15'52.34"S 112°44'34.22"E	15072019
Harmoni	Sukolilo	7°17'40.04"S 112°48'13.16"E	08072019
Lansia	Gubeng	7°16'18.85"S 112°45'17.74"E	22071019
Pelangi	Gayungan	7°19'39.23"S 112°43'52.34"	15072019
Persahabatan	Wonokromo	7°16'39.93"S 112°45'07.44"E	15072019
Prestasi	Genteng	7°15'41.30"S 112°44'34.25"E	22072019
Wira Agung	Wonokromo	7°17'56.29"S 112°44'16.73"E	15072019



**Figure 1.** Geographical map of Surabaya City, seven urban parks were marked with circle (Ha: Harmoni Park, Pr: Prestasi Park, La: Lansia Park, Pe: Persahabatan Park, Wi: Wira Agung Park, Pl: Pelangi Park). Figure was created with QGIS 3.10.4 Version.

urban parks in Surabaya, Indonesia. Above are the coordinate of the sampling site (Table 1) and the geographical map was shown in Figure 1.

**Procedures**

*Mosquito Collection and Rearing*

Ovitrap surveillance method was adopted from the following guidelines (Imam et al. 2014). Ovitrap were installed in some of the public urban parks distributed in Surabaya (Figure 1) and designed with black colour to attract mosquitoes to lay their eggs, but only some of the eggs from several parks were successfully attached (Table 1). A total of 28 ovitraps were installed in which 4 ovitraps were installed in each park. Observation of attached eggs in filter paper was done every seven days. All filter papers are labelled based on the date of collection and the collection site. After that, they would be

brought to Entomology Laboratory, Institute of Tropical Disease, Universitas Airlangga. Mosquito rearing was set based on the mosquito rearing guidelines (Imam et al. 2014). Adult mosquitoes were identified using an identification key from the Indonesian Ministry of Health (Indonesian Ministry of Health 1989). All of study design in present study was approved by Ethical Committee Faculty of Medicine, Universitas Airlangga with the referee number of 24-934/UN3.14/PPd/2013.

*RNA Extraction*

Detection of the dengue virus on the mosquito that were caught was also conducted, but the result will be reported in another study. In regards to the double purpose, RNA extraction was conducted. After mosquito identification was made, mosquitos were extracted using QIAamp® Viral RNA by Qiagen, Germany, based on the manufacturer's instruction to extract RNA from mosquito specimens.

*AS-PCR Assays for the Detection of kdr Mutant Alleles (V1016G and F1534C)*

After cDNA from the procedure of 2.2.2.2 were obtained, genotyping process was conducted. Genotyping of *kdr* mutant alleles to detect mutant in codon 1016 and 1534 were performed using Allele-specific PCR assays following previous guidelines (Stenhouse et al. 2013; Yanola et al. 2011) AS-PCR was conducted by using forward and reverse primer below (Table 2).

Reactions were performed using PCR Thermocycler with following stage: 94°C in 2 min, 35 cycles of 30 sec in 94°C, 55°C in 30 sec, 72°C in 30 sec, and final elongation 72°C in 2 min for the detection of V1016G. For the detection of F1534C, PCR was performed with the stage of 94°C in 2 min, 35 cycles of 30 sec in 94°C, 30 sec in 60°C, 30 sec in 72°C, and elongation step with 72°C in two minutes. PCR products were then loaded into 1.5% agarose gel.

*Data Analysis*


Data were analysed using R Application of 4.0.4 Version, using packages ‘HardyWeinberg’ to analyse the frequency of resistant allele.

**RESULTS**

A total of 55 female *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus* were tested, consisting of 37 *Aedes (Stegomyia) aegypti* and 18 *Aedes (Stegomyia) albopictus*. The composition of mosquitoes in every park was shown in Figure 2. The electrophoresis results were shown in Figure 3. The comparison between our result and the reference was shown in Figure 4 and Figure 5.

**Table 2.** The sequences of oligonucleotides used to amplify fragments of the VGSC gene (Stenhouse et al. 2013; Yanola et al. 2011).

	Primer sequence	Product (bp)
Gly1016f	5'-ACCGACAAATTGTTTCCC-3'	
Gly1016r	5'GCGGGCAGGGCGGCGGGGGCGGGG CCAGCAAGGCTAAGAAAAGGTTAACTC-3'	60
Val1016r	5'-GCGGGCAGCAAGGCTAAGAAAAGGT TAATTA-3'	80
Cys1534f	5'GCGGGCAGGGCGGCGGGGGCGGGGCTCTACTTT- GTGTTCTTCATCATGTG3'	113
Cys1534r	5'TCTGCTCGTTGAAGTTGTCGAT3'	
Phe1534f	5'GCGGGCTCTACTTTGTGTTCTTCATCATATT3'	93




**Figure 2.** The distribution of mosquito species, dark grey indicated the occurrence of *Aedes (Stegomyia) albopictus*, meanwhile light grey indicated the occurrence of *Aedes (Stegomyia) aegypti*. Ha: Harmoni Park, Pr: Prestasi Park, La: Lansia Park, Pe: Persahabatan Park, Wi: Wira Agung Park, Pl: Pelangi Park). (Figure was created with QGIS).

It was genotyped to only female mosquitoes due to the important role of female mosquitoes as the vector of mosquito-borne diseases. We estimated that random mating was occurred in this population and VGSC is not sex-linked. The results showed that no point mutation was detected in


**Table 3.** The distribution of species in every park, AS-PCR result (SS: Susceptible allele/V1016V/F1534F/Homozygous wildtype, SR: Susceptible resistant/V1016G/F1534C/Heterozygous mutant, RR: Resistant resistant/G1016G/C1534C/Homozygous mutant).

Sampling Site	<i>Aedes (Stegomyia) aegypti</i>	<i>Aedes (Stegomyia) albopictus</i>	n (pool)	Genotype			Allele Frequency	
				SS (%)	SR (%)	RR	S	R
Apsari Park	5	0	3	3 (100)	0 (0)	0 (0)	1	0
Harmoni Park	12	8	2	2 (100)	0 (0)	0 (0)	1	0
Lansia Park	3	0	1	1 (100)	0 (0)	0 (0)	1	0
Pelangi Park	7	1	1	1 (100)	0 (0)	0 (0)	1	0
Persahabatan Park	5	2	1	1 (100)	0 (0)	0 (0)	1	0
Prestasi Park	1	7	1	1 (100)	0 (0)	0 (0)	1	0
Wira Agung Park	4	0	2	2 (100)	0 (0)	0 (0)	1	0
<b>Total</b>	37	18	11	11	0	0		

all of the samples. In other words, all the mosquitoes collected have homozygous wildtype of V1016V and F1534F, as shown in Table 3 below. If the AS-PCR results shows the homozygous wildtype/V1016V/F1534F, it can be symbolized as genotype SS (means that this pool represents susceptible allele), heterozygous mutant/V1016G/F1534C symbolized as genotype SR (means that this pool represents susceptible and resistant allele), homozygous mutant/G1016G/C1534C symbolized as genotype RR (means that this pool represents resistant allele).




**Figure 3.** The electrophoresis results of the detection of kdr mutant allele (A: V1016G and B: F1534C). Amplified products could be differentiated by size (60 bp for V1016, 80 bp for G1016), 93 bp for F1534 and 113 bp for C1534.



**Figure 4.** The representative of the electrophoresis of V1016G. Study result compared with the reference, left figure was the study result, right figure was the figure from reference (Stenhouse et al. 2013). M=marker of 25 bp; V1016V = Homozygous wildtype; V1016G =Heterozygous Mutant; G1016G = Homozygous Mutant.





**Figure 5.** The representative of the electrophoresis of F1534C. Study result compared with the reference, left figure was the study result, right figure was the figure from reference (Yanola et al. 2011). M=marker of 25 bp; F1534F= Homozygous wildtype; F1534C=Heterozygous Mutant; C1534C= Homozygous Mutant.

## DISCUSSIONS

This study highlighted that no *kedr* mutant alleles (V1016G and F1534C) have been found from the study site. This study was the first report of *kedr* mutation screening in Surabaya, East Java Province, Indonesia. Some of *kedr* mutant alleles in Indonesia have been reported from many regions and showed various mutations on V1016G, F1534C, and S989P. A study from Amelia-Yap shows the occurrence of mutation on V1016G and S989P in mosquitoes collected from Kuningan, Padang, Denpasar, Samarinda, Mataram, and Sumba Timur (Amelia-Yap et al. 2019). Pointing out and detecting mutation of Val1016Gly/V1016G and Phe1534Cys/F1534C is important, including in national and local areas. The Allele-Specific PCR Assay (AS-PCR) can be implemented to detect *kedr* mutant and provide a rapid result, accurate, and cost-effective genotyping (Lee et al. 2016). Rapid results and the precise target can be used to figure out the VGSC gene of mosquitoes in a population. However, this method cannot stand alone and must be combined with WHO tube test Bioassay to capture information about the susceptibility status of mosquito against insecticides (Corbel & Guessan 2013). World Health Organization (WHO) Tube Test Bioassay against insecticides require too much number of samples. Instead, AS-PCR was used for the screening of *kedr* mutant alleles in the mosquito, using the previous method (Stenhouse et al. 2013). *Kdr* screening using Allele-Specific PCR assay is a very sensitive method that can provide early warning for future resistance detection that is suitable to use given the limited number of mosquitoes (Ranson et al. 2011). In this study, not all the mosquito eggs that were rearing successfully emerge into adult mosquitoes. Thus, AS-PCR assay screening method was adopted.

Surabaya is a dengue-endemic area and the first place where dengue cases were found in 1968. Management of dengue infection in Indonesia, including in Surabaya, was focused on vector control, which can be broken down into three methods: physical control, chemical control, and biological control. Physical control can be done by washing bathtubs regularly every seven days before the mosquito eggs become adult mosquitoes, larval

elimination through program one house one larval observer, and closing water containers. Fogging activity (hot-fogging) was applied in some Indonesian regions as chemical control, except in Surabaya where they used Ultra Low Volume spray (cold-fogging). Some chemical insecticides in fogging and ULV were from the organophosphate group (Malathion and methylpirimiphos), Pyrethroid (Cypermethrine, Lambda-cyhalothrine, Cyflutrine, Permethrine, S-Bioalethrine, and its derivation). In comparison, chemical control in goal for killing mosquito larvae was temephos from the organophosphate group and piriproxifen ([Indonesian Ministry of Health 2017](#)).

Since insecticides have been used for many decades, long exposure to insecticides has been a major public health problem. Continuously using the same insecticide for a long period led to mosquitoes' development resistance to insecticide exposure. In this case, every country should be responsible for their vector control program. World Health Organization (WHO) defined resistance as an insect's ability to survive against the effect of insecticides through natural selection and mutations ([World Health Organization 2012](#)). Other countries in South-East Asia were also facing the same problem. In the South-East Asia region, the number of V1016G mutations have been reported from Cambodia, Laos, Myanmar, Malaysia, Singapore, Thailand, and Vietnam. While the report from Philippine and Timor Leste are still not available ([Amelia-Yap et al. 2018](#)). Other mutations were also reported, such as mutations in Thailand that show mutations in codon 1011, where isoleucine become valin (I1011V) ([Rajatileka et al. 2008](#)). Some resistant alleles frequencies, including F1534C and V1016G have been reported in various range. A cross sectional study in India in 2015, shows the frequency of F1534C(C) and V1016G(G) were 0.51 and 0.18 respectively ([Kushwah et al. 2020](#)). In other part of India, it also has been reported the frequency of F1534C(C) was around 0.41-0.79, combined with the new point mutation T1520I(I) with the frequency of 0.13 ([Kushwah et al. 2015](#)). Researches about point mutation in the voltage-gated sodium channel of dengue vector were rapidly increase, a report from Taiwan found a novel point mutation, D1794Y that occurs with the V1023G mutation ([Chang et al. 2009](#)). Meanwhile, a study in Mexico successfully revealed the co-occurrence of point mutation V1016I and F1534C was associated with pyrethroid resistance during 16 years of observation ([Saavedra-Rodriguez et al. 2018](#)). The derivative of permethrin ( $\lambda$ -cyhalothrin) was also reported to be the causative of *kn*dr point mutation (V419L) in Colombia, with the frequency ranging from 0.06 to 0.46 ([Granada et al. 2018](#)).

Study of knockdown-resistance (*kn*dr) alleles in Indonesia may give another impact on vector control method. In Yogyakarta, a dengue-endemic city, the frequency of *kn*dr mutant alleles V1016G and F1534C in the area where Wolbachia will be released as vector control method was firstly measured and shows high frequencies of V1016G mutation, but F1534C was low detected. The result of *kn*dr screening can figure out population background, so that the vector control that may be released can be suitable ([Wuliandari et al. 2020](#)). Since having the first case of dengue infection in 1968 in Jakarta and Surabaya, Indonesia has applied some vector control, but it still needs to be improved so that the dengue outbreaks can be avoided. The number of dengue incidence rate (IR) in Indonesia was increased from 0.05 to 40 per 100,000 populations in 2013. The highest epidemic was reported in 2010, with the IR value of 85.7/100,000 population ([Haryanto 2018](#)). Some risk factors of dengue infection in Indonesia were various breeding sites especially during rainy season, the mobility of citizens inside or outside the country, and the low level of awareness toward health and hygiene lifestyle ([Setiati et al. 2006](#)). Indonesia is a suitable place for some

vector's growth and development as a tropical country, including mosquitoes. Although this study had not found *kdr* mutant allele in *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus*, it does not take alleles in another district/region into account, so regular surveillance in other endemic areas are strongly suggested. *Kdr* mutant alleles in points 989, 1534, and 1016 was a leading factor that causes mosquitoes to develop resistance phenotype against insecticide (Harris et al. 2010; Srisawat et al. 2010). AS-PCR assay contributes to the early detection for future warning resistance in the mosquito that cannot be detected by bioassay methods. Thus, the presence of a single/double *kdr* mutant allele should be considered. In other cases, the negative result/the absence of *kdr* mutant allele did not lead to complacency because of the specific target that are being examined (Corbel & Guessan 2013). The absence of *kdr* mutant allele in one point does not reflect another absence of mutation in another point. Hence, this study might be used as complement data for the vector control program.

## CONCLUSION

Result of present study confirmed that no mutations were found in Valine to Glycine in point 1016 (V1016G) and Phenylalanine to Cysteine in point 1534 (F1534C) alleles in both two species, *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus* collected from urban parks, Surabaya, Indonesia. All of mosquito samples have homozygous wild type genotype of *kdr* alleles (V1016V and F1534F).

## AUTHORS CONTRIBUTION

SF was responsible for conceptualization, data collection, and manuscript preparation. SS was responsible for supervision, investigation of the data, and manuscript preparation. BU was responsible for data analysis and manuscript preparation. THS was responsible for data collection and manuscript preparation. HA was responsible for manuscript preparation and grammatical check. AA was responsible for data validation and manuscript preparation. PW was responsible for data validation and manuscript preparation. SS was responsible for investigation of the data and manuscript preparation.

## ACKNOWLEDGMENTS

We would like to thank the urban parks workers for the help during data collection.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest in this research.

## REFERENCES

- Amelia-Yap, Z.H. et al., 2018. Pyrethroid resistance in the dengue vector *Aedes aegypti* in Southeast Asia: Present situation and prospects for management. *Parasites and Vectors*, 11(1), p.17.
- Amelia-Yap, Z.H. et al., 2019. V1016G Point Mutation: The Key Mutation in the Voltage-Gated Sodium Channel (Vgsc) Gene of Pyrethroid-Resistant *Aedes aegypti* (Diptera: Culicidae) in Indonesia. *Journal of Medical Entomology*, 56(4), pp.953–958.
- Bowman, L.R., Donegan, S. & McCall, P.J., 2016. Is Dengue Vector Control Deficient in Effectiveness or Evidence?: Systematic Review and Meta-analysis. *PLoS Neglected Tropical Diseases*, 10(03), e004551.
- Bregues, C. et al., 2003. Pyrethroid and DDT cross-resistance in *Aedes aegypti* is correlated with novel mutations in the voltage-gated sodium channel gene. *Medical and Veterinary Entomology*, 17(1), pp.87–94.

- Chang, C. et al., 2009. A novel amino acid substitution in a voltage-gated sodium channel is associated with knockdown resistance to permethrin in *Aedes aegypti*. *Insect Biochemistry and Molecular Biology*, 39(4), pp.272–278.
- Corbel, V. & Guesson, R.N., 2013. Distribution, Mechanisms, Impact and Management of Insecticide Resistance in Malaria Vectors: A Pragmatic Review. *Springer*.
- David, J., Gallet, C. & Despre, L., 2007. The evolutionary ecology of insect resistance to plant chemicals. *TRENDS in Ecology and Evolution*, 22(6), pp.298–307.
- Djouaka, R. F. et al., 2008. Expression of the cytochrome P450s, CYP6P3 and CYP6M2 are significantly elevated in multiple pyrethroid resistant populations of *Anopheles gambiae* s.s. from Southern Benin and Nigeria. *BMC Genomics*, 13(9), 538.
- Fournier, D., 2005. Mutations of acetylcholinesterase which confer insecticide resistance in insect populations. *Chemico-Biological Interactions*, 158, pp.257–261.
- Granada, Y. et al., 2018. A point mutation V419I in the sodium channel gene from natural populations of *Aedes aegypti* is involved in resistance to  $\lambda$ -cyhalothrin in Colombia. *Insects*, 9(1), 23.
- Hardjanti, A., Donanti, E. & Jakarta, E., 2015. Detection of Insecticide Resistance in *Aedes Aegypti* to Organophosphate in Pulogadung, East Jakarta Detection of Insecticide Resistance in *Aedes aegypti* to Organophosphate in. *Makara Journal of Health Research*, 19(3), pp.117–120.
- Harris, A.F., Rajatileka, S. & Ranson, H., 2010. Pyrethroid resistance in *Aedes aegypti* from Grand Cayman. *American Journal of Tropical Medicine and Hygiene*, 83(2), pp.277–284.
- Haryanto, B. 2018, 'Indonesia Dengue Fever: Status, Vulnerability, and Challenges', in A. J. Rodriguez-Moralez (eds.), *Current Topics in Tropical Emerging Diseases and Travel Medicine*, IntechOpen.
- Hemingway, J. et al., 2004. The molecular basis of insecticide resistance in mosquitoes. *Insect Biochemistry and Molecular Biology*, 34, pp.653–665.
- Imam, H. et al., 2014. The basic rules and methods of mosquito rearing (*Aedes aegypti*). *Tropical Parasitology*, 4(1), p.53.
- Indonesian Ministry of Health, 1989. *Kunci identifikasi Aedes di Jawa*. Direktorat Jendral Pemberantasan Penyakit Menular dan Penyehatan Lingkungan.
- Indonesian Ministry of Health. 2017. *InfoDatin-Situasi-Demam-Berdarah-Dengue*. Pusat Data dan Informasi Kementerian Kesehatan RI.
- Kawada, H. et al., 2014. Co-occurrence of Point Mutations in the Voltage-Gated Sodium Channel of Pyrethroid-Resistant *Aedes aegypti* Populations in Myanmar. *PLoS Neglected Tropical Diseases*, 8(7), e3032.
- Kushwah, R.B.S. et al., 2015. Pyrethroid-Resistance and Presence of Two Knockdown Resistance (kdr) Mutations, F1534C and a Novel Mutation T1520I, in Indian *Aedes aegypti*. *PLoS Neglected Tropical Diseases*, 9(1), e3332.
- Kushwah, R.B.S. et al., 2020. A new knockdown resistance (kdr) mutation, F1534L, in the voltage-gated sodium channel of *Aedes aegypti*, co-occurring with F1534C, S989P and V1016G. *Parasites and Vectors*, 13(1), 327.
- Lee, H.B. et al., 2016. Allele-specific quantitative PCR for accurate, rapid, and cost-effective genotyping. *Human Gene Therapy*, 27(6), pp.425–435.
- Nathin, M.A., Harun, S.R. & Sumarmo, 1988. Dengue haemorrhagic fever and Japanese B encephalitis in Indonesia. *Southeast Asian J Trop Med Public Health*, 19(3), pp.475–481.

- Rajatileka, S. et al., 2008. Development and application of a simple colorimetric assay reveals widespread distribution of sodium channel mutations in Thai populations of *Aedes aegypti*. *Acta Tropica*, 108(1), pp.54–57.
- Ranson, H. et al., 2011. Pyrethroid resistance in African anopheline mosquitoes : what are the implications for malaria control? *Trends in Parasitology*, 27(2), pp.91–98.
- Reinert, J.F., Harbach, R.E. & Kitching, I.A.N.J., 2009. Phylogeny and classification of tribe Aedini (Diptera : Culicidae). *Zoological Journal of the Linnean Society*, 157, pp.700–794.
- Russell, T.L. et al., 2011. Increased proportions of outdoor feeding among residual malaria vector populations following increased use of insecticide-treated nets in rural Tanzania. *Malaria Journal*, 10(1), 80.
- Saavedra-Rodriguez, K. et al., 2018. Parallel evolution of vgsc mutations at domains IS6, IIS6 and IIIS6 in pyrethroid resistant *Aedes aegypti* from Mexico. *Scientific Reports*, 8(1), 6747.
- Setiati, T.E. et al., 2006. Changing epidemiology of dengue haemorrhagic fever in Indonesia. *Dengue Bulletin*, 30, pp.1–14.
- Simmons, C.P. et al., 2012. Dengue. *The New England Journal of Medicine*, 366, pp.1423–1432.
- Srisawat, R. et al., 2010. Point mutations in domain II of the voltage-gated sodium channel gene in deltamethrin-resistant *Aedes aegypti* (diptera: culicidae). *Applied Entomology and Zoology*, 45(2), pp.275–282.
- Stenhouse, S.A. et al., 2013. Detection of the V1016G mutation in the voltage-gated sodium channel gene of *Aedes aegypti* (Diptera: Culicidae) by allele-specific PCR assay, and its distribution and effect on deltamethrin resistance in Thailand. *Parasites and Vectors*, 6(1), 253.
- World Health Organization, 2006. *Dengue*. World Health Organization. <http://www.searo.who.int/>
- World Health Organization, 2012. *GLOBAL PLAN FOR INSECTICIDE RESISTANCE MANAGEMENT IN MALARIA VECTORS*. WHO Library Cataloguing in Publication Data.
- World Health Organization, 2020a. *Dengue and Severe Dengue*. World Health Organization. <https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue>
- World Health Organization, 2020b. *Mosquito borne diseases*. World Health Organization. [https://www.who.int/neglected\\_diseases/vector\\_ecology/mosquito-borne-diseases/en/](https://www.who.int/neglected_diseases/vector_ecology/mosquito-borne-diseases/en/)
- World Health Organization, 2020c. *Questions and Answers on Dengue Vaccines*. World Health Organization. [https://www.who.int/immunization/research/development/dengue\\_q\\_and\\_a/en/](https://www.who.int/immunization/research/development/dengue_q_and_a/en/)
- Wuliandari, J.R. et al., 2015. Association between Three Mutations, F1565C, V1023G and S996P, in the Voltage-Sensitive Sodium Channel Gene and Knockdown Resistance in *Aedes aegypti* from. *Insects*, 6, pp. 658–685.
- Wuliandari, J.R. et al., 2020. Frequency of kdr mutations in the voltage-sensitive sodium channel (V SSC) gene in *Aedes aegypti* from Yogyakarta and implications for Wolbachia-infected mosquito trials. *Parasites and Vectors*, 13(1), 429.
- Yanola, J. et al., 2011. High-throughput assays for detection of the F1534C mutation in the voltage-gated sodium channel gene in permethrin-resistant *Aedes aegypti* and the distribution of this mutation throughout Thailand. *Tropical Medicine and International Health*, 16(4), pp.501–509.