

The Potensial Predictive Role

by Kumboyono Kumboyono

Submission date: 22-May-2023 07:58PM (UTC+0800)

Submission ID: 2099173505

File name: The_Potensial_Predictive_Role.pdf (1.37M)

Word count: 5083

Character count: 27156

The Potential Predictive Role of Tumour Necrosis Factor- α , Interleukin- 1β , and Monocyte Chemoattractant Protein-1 for COVID-19 Patients Survival

Kumboyono Kumboyono¹, Indah Nur Chomsy², Agustin Iskandar³, Aryati Aryati⁴, Ida Parwati⁵, Titin Andri Wihastuti⁶

¹Nursing Department, Faculty of Health Sciences, University of Brawijaya, Malang, 65151, Indonesia; ²Doctoral Program of Medical Science, Faculty of Medicine, University of Brawijaya, Malang, 65145, Indonesia; ³Department of Clinical Pathology, Faculty of Medicine, University of Brawijaya, Malang, 65145, Indonesia; ⁴Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga, Surabaya, 60131, Indonesia; ⁵Department of Clinical Pathology, Faculty of Medicine, Universitas Padjadjaran, Bandung, 40161, Indonesia; ⁶Basic Nursing Department, Faculty of Health Sciences, University of Brawijaya, Malang, 65151, Indonesia

Correspondence: Titin Andri Wihastuti, Basic Nursing Department, Faculty of Health Sciences, University of Brawijaya, Malang, 65151, Indonesia, Email titinwihastuti@gmail.com

Background: Tumour necrosis factor- α (TNF- α), interleukin- 1β (IL- 1β), and monocyte chemoattractant protein-1 (MCP-1) are early phase cytokines often encountered when the body is exposed to severe acute respiratory syndrome-associated-coronavirus-2. TNF- α , IL- 1β , and MCP-1 are pro-inflammatory cytokines critical in the defence response against systemic infection and injury. Therefore, TNF- α , IL- 1β , and MCP-1 are the most aggressive responses to viral infections in the acute phase, so they can be used to determine the survival of coronavirus disease 2019 (COVID-19) patients.

Purpose: The study aimed to determine the levels of TNF- α , IL- 1β , and MCP-1 as predictors of survival for COVID-19 patients.

Patients and Methods: A prospective cohort study was conducted on confirmed COVID-19 by a reverse-transcriptase-polymerase-chain-reaction (RT-PCR) in 84 adults admitted to the hospital in Indonesia. TNF- α , IL- 1β , and MCP-1 level were measured from serum subjects using the enzyme-linked immunosorbent assay.

Results: The results from logistic regression modelling of the survival status of COVID-19 patients based on TNF- α , IL- 1β , and MCP-1 levels were significant (p-value=0.024). The predictors of all cytokines had P Wald <0.05, so the three cytokines could be used simultaneously to predict the survival status of COVID-19 patients. MCP-1 has the most dominant risk relative value (2.76; 95% CI; 2.53–4.68) compared to TNF- α and IL- 1β in predicting patient survival.

Conclusion: TNF- α , IL- 1β , and MCP-1 as markers of acute systemic inflammatory cytokines can be measured at the beginning of hospitalisation of COVID-19 patients for early diagnosis of disease severity so that healthcare professionals can determine clinical guidance needs for therapeutic programs.

Keywords: cytokine, survival predictors, COVID-19, SARS COV-2

Introduction

The coronavirus disease 2019 (COVID-19) pandemic is a global health crisis with very high morbidity and mortality rates in a short time. Activation of innate and adaptive immune responses by coronavirus infection causes an uncontrolled inflammatory response resulting in a cytokine storm in severe COVID-19 patients.¹ Crosstalk between immune cells and endothelial cells during coronavirus infection impacts the activation of several inflammatory cascades, causing hyperproduction of pro-inflammatory cytokines. Cytokine storms have been shown to cause manifestations of acute respiratory distress syndrome and thromboembolism leading to vascular occlusions, such as in cases of acute ischemic stroke, myocardial infarction, encephalitis, acute kidney injury, vasculitis and sudden death.^{2,3} Significantly elevated pro-inflammatory

cytokines suggest a poor prognosis in COVID-19 patients. Research shows that patients with severe COVID-19 tend to have higher concentrations of pro-inflammatory cytokines (such as Tumour Necrosis Factor- α (TNF- α), Interleukin-1 β (IL-1 β), and Monocyte Chemoattractant Protein-1 (MCP-1) than those with mild or moderate illness.^{4,5} This phenomenon shows a very close relationship between the severity of COVID-19 and the production of pro-inflammatory cytokines. Therefore, predicting the survival of COVID-19 patients using the three pro-inflammatory cytokines will allow clinicians to take prompt and appropriate treatment to prevent the severity of COVID-19 from reducing mortality.

Cytokines are now in the spotlight in various countries because of their essential roles in the pathogenesis of COVID-19. Proinflammatory cytokines such as IL1 β , IL-2, IL-4, IL-6, IL-8, TNF α , IL-2, IL-4, IP-10 and MCP-1 are associated with an increased risk of mortality from acute respiratory distress syndrome (ARDS). Huang (2020) research stated that COVID-19 patients treated in the ICU had higher IL2, IL7, IL10, GSCF, IP10, MCP1, MIP1A, and TNF α than patients not admitted to the ICU.⁶ Meanwhile, a prospective cohort study in 24 COVID-19 patients revealed that levels of IL-4, IL-6, IL-8, IL-10, FGF, IP-10, MCP-1, and TNF correlated with 30-day mortality.⁷ In addition, Chen et al found that the levels of IP-10, MCP-1, and IL-6 were significantly higher in COVID-19 critically ill patients.⁸ MCP-1 and IL-10 were not significantly associated with mortality but were associated with high levels of d-dimer associated with mortality.

Many studies have reported that acute respiratory distress syndrome (ARDS) is one of the leading causes of death related to COVID-19. ARDS is triggered by a cytokine storm that directly damages the respiratory epithelium, liver, heart, and kidneys, including TNF- α .⁹ The soluble form of TNF- α and the membrane protein form are active forms with different affinities for the TNF receptor. When it binds to the TNF receptor, TNF- will cause a pleiotropic effect or stimulate the secretion of other systemic cytokines, so TNF- α is known as an early-phase cytokine.^{10,11}

Interleukin 1 β is released by macrophages via inflammasomes and contributes to the cytokine storm as the most aggressive response to COVID-19.¹² IL-1 β is known to play a role in immunological dysregulation in severe patients with respiratory failure,¹³ which in cytokine storm of ARDS correlates with disease severity, also may be associated with organ failure.^{14,15} Yang et al (2020) found that increased IL-1 receptor antagonist as a marker of increased viral load and risk of death in 14 severe COVID-19 patients.¹⁶ Therefore, IL-1 β is part of the cytokine storm generated by coronavirus infection.¹⁷ IL-1 functions after binding to the IL-1 receptor (IL-1R) and is known to transduce and activate one of the well-known inflammatory pathways, the NF κ B pathway.¹⁸

MCP-1 is a potent monocyte chemotactic factor constitutively produced or induced by oxidative stress, cytokines, or growth factors.¹⁹ MCP-1 was associated with high levels of d-dimer associated with death.⁸ Furthermore, it has been reported that MCP-1 expression increases rapidly in the early phase of acute infection and then progressively decreases with disease progression.²⁰

IL-1 β , and MCP-1 are pro-inflammatory cytokines critical in the defence response against systemic infection and injury.^{17,21} IL-1 β and MCP-1 respond as an emergent cytokine by initiating acute-phase protein synthesis. In addition to TNF- α , IL-1 β and Monocyte Chemoattractant Protein-1 (MCP-1) are early phase cytokines often encountered when the body is exposed to coronavirus.^{4,21} Although there have been studies using similar cytokines to predict the survival of COVID-19 patients in various countries, there has been no reporting in Indonesia in the period of Delta variants of SARS-CoV-2. So, this study wants to confirm the research results on the survival of COVID-19 patients in Indonesia with various other countries.

Materials and Methods

Study Design and Setting

A prospective cohort study confirmed COVID-19 in adults admitted to the public hospitals to refer coronavirus disease patients in Indonesia. The grouping of patients as a risk factor for predicting the survival of COVID-19 patients is based on the severity level: mild, moderate, severe, and critically ill. The grouping of COVID-19 patients is divided into asymptomatic, mild, moderate, severe and critical.²²

1. Mild: Patients with symptoms without evidence of viral pneumonia or hypoxia. Symptoms include fever, cough, fatigue, anorexia, shortness of breath, myalgia, nasal congestion, headache, diarrhoea, nausea and vomiting, loss of smell (anosmia), or loss of taste (ageusia).

2. Moderate: patients with clinical signs of pneumonia (fever, cough, shortness of breath, rapid breathing) but no signs of severe pneumonia, including $SpO_2 > 93\%$ in room temperature, and no signs of severe pneumonia.

3. Severe/Severe Pneumonia: patients with clinical signs of pneumonia (fever, cough, shortness of breath, rapid breathing), with the addition of one of the following symptoms: respiratory rate > 30 times/minute, severe respiratory distress, or $SpO_2 < 93\%$ on room air respiratory rate > 30 times/minute, severe respiratory distress, or $SpO_2 < 93\%$ in room temperature.

4. Critically ill: Patients with ARDS, sepsis, and or septic shock.

Each group was then observed to determine the research's outcome: survival or non-survival. There are three general hospitals in this study: Saiful Anwar Hospital Malang, Soetomo General Hospital Surabaya, and Hasan Sadikin General Hospital Bandung.

Study Subject and Data Collection

The subjects of this study were 84 inpatients at the three hospitals who were diagnosed as confirmed COVID-19, proven by a Reverse-Transcriptase-Polymerase-Chain-Reaction (RT-PCR) method. The inclusion criteria for this research subject are: patients who confirmed COVID-19 by the RT-PCR (nasopharyngeal or oropharyngeal swab); ages 17–60 years; and have onset time 14 days from the first time the patient has symptoms until confirmed as Covid-19 patient. This study's subject exclusion criteria are pregnant or breastfeeding women, having a coagulation disorder before being diagnosed as COVID-19, and co-morbidities such as cancer, hepatitis, or HIV/AIDS.

Cytokine levels of TNF- α (ng/mL), IL-1 β (pg/mL), and MCP-1 (ng/mL) were measured from serum subjects using the Enzyme-Linked Immunosorbent Assay (ELISA) examination procedure. The ELISA procedure was carried out according to the manual protocol of the Human IL-1 β Bioassay Technology Laboratory (BT-Lab) ELISA kit (Cat. No. E0143Hu) and Human TNF- α BT-Lab ELISA Kit (Cat. No. E0082Hu) and, and Human MCP-1 BT-Lab ELISA Kit (Cat. No. E0124Hu). The ELISA procedure was carried out by adding 40 μ L of a serum sample to the sample well. After that, it added 50 μ L of streptavidin-HRP to the sample wells and standard wells. Covered well and incubated for 60 minutes at 37°C, then washed with Wash Buffer five times. First, 50 μ L of substrate solution. Then was attached to each well, and then 50 μ L of substrate solution B was added to each well. Next, it was incubated in the dark for 10 minutes at 37°C. After that, 50 μ L of Stop Solution was summed for each well. Each optical density (OD) value was appropriately read using a microplate reader at a wavelength of 450 nm within 10 minutes after adding the stop solution.

Ethical Clearance

The institutional ethics committee of University of Brawijaya approved the study (No. 209/EC/KEPK/07/2021) and also ethical committee of Saiful Anwar General Hospital (No. 4000/011/K.3/302/2021). All of the patients participate in this study provided informed consent that has been accordanced with the Declaration of Helsinki.

Data Analysis

The authors conducted the Mann-Whitney *U*-test to examine differences in TNF- α , IL-1B, and MCP-1 levels between survivors and non-survivors. The association between levels of TNF- α , IL-1B, and MCP-1 with survival status was multivariate analysed using logistic regression. In addition, the authors calculated values of the diagnostic test (sensitivity, specificity, positive predictive value, negative predictive value) and accuracy of the test (the area under Receiver Operator Characteristic curve) to describe TNF- α , IL-1B, and MCP-1 levels to predict mortality of COVID-19 patient. All statistical analyses were conducted with IBM[®] Statistical Product and Service Solution[®] for Windows version 25.0 with *p*-value < 0.05 as a statistical significance level.

72

Table 1 Characteristics of the Study Population

Characteristics	Group of COVID-19 Patient		p-value
	Survivors (n=47; %)	Non-Survivors (n=37; %)	
Age (x±SD) [#]	56.1 ± 11.5	58.8 ± 13.4	0.325
Sex (n (%)) [^]			
Male	40.4	37.8	0.809
Female	59.6	62.2	
Comorbid (n (%)) [^]	83.0	86.5	0.659
Hypertension [^]	29.8	27.0	
STEMI [^]	4.3	0.0	
Chronic Kidney Disease	4.3	29.7	
Diabetes Mellitus	58.5	41.5	
Cerebrovascular Accident	66.7	33.3	
Overweight	53.8	46.2	
Malignancy	8.5	2.7	
HIV/AIDS	2.1	2.7	
Cirrhosis of the Liver	0.0	2.7	
Severity [^]			0.004*
Mild	12.8	2.7%	
Moderate	38.8	16.2%	
Severe	29.8	27.0%	
Critically ill	9.1	54.1%	

Notes: [#]Independent t-test; [^]Chi-square test; *Considered significance at p-value<0.05.

Results

Baseline Characteristic of the Research Subject

This study found 84 COVID-19 patients according to the inclusion and exclusion criteria from June 2020 to March 2021. There were 47 (55.9%) survivors of COVID-19 and 37 (44.1%) of COVID-19 patients who were died (non-survivors). Table 1 presents the characteristics of research subjects based on patient outcomes, namely survivors and non-survivors.

65

TNF- α , IL-1 β , and MCP-1 Levels in Survivors and Non-Survivors in COVID-19 Patients

Measurement of TNF- α , IL-1 β , and MCP-1 levels in the serum of COVID-19 patients was carried out on the first day of hospitalisation (Table 2). The median levels of TNF- α , IL-1 β , and MCP-1 in the non-survivor group of COVID-19 patients were significantly higher than those of COVID-19 survivors.

Meanwhile, Table 3 shows logistic regression modelling of the survival status of COVID-19 patients based on TNF- α , IL-1 β , and MCP-1 levels were significant (p-value=0.024). The predictors of all cytokines had P Wald <0.05, so the three cytokines could be used simultaneously to predict the survival status of COVID-19 patients. MCP-1 has the most dominant relative risk (RR) value compared to TNF- α and IL-1 β in predicting patient survival.

28

Table 2 Levels of TNF- and IL-1 β in the Survivors and Non-Survivors of COVID-19

Cytokine	Group of COVID-19 Patient		p-value**
	Survivors (n=47)	Non-Survivors (n=37)	
TNF- α (ng/L)	28.03 (15.30–50.76)*	53.97 (18.16–98.49)*	0.007
IL-1 β (pg/L)	665.36 (440.49–813.53)*	939.0 (468.76–2185.82)*	0.035
MCP-1 (ng/L)	41.27 (31.12–68.46)*	86.13 (68.50–94.82)*	0.004

Notes: *Median (min-max) **Based on Mann-Whitney U-Test with α 5%.

Table 3 Results of Multivariate Logistic Regression Analysis Between TNF- α , IL-1 β , and MCP-1 Level with Survival Status Among COVID-19 Patients

Variables	B	P Wald	Relative Risk (RR)	95% Confidence Interval
TNF- α	2.39	0.018	2.15	1.98–2.98
IL-1 β	12.83	0.042	1.12	1.03–2.04
MCP-1	9.68	0.006	2.76	2.53–4.68
Constant	-2.51	0.003		

Notes: -2 Log Likelihood = 64.718; Goodness of Fit = 28.83; p-value = 0.024.

The accuracy of the three cytokines tests in predicting the death of COVID-19 patients can be seen in the following description.

The Ability of TNF- α as a Predictor of Mortality of COVID-19 Patients

ROC curve analysis of TNF- α levels was carried out to determine the ability of this biomarker as a predictor of mortality. The analysis results show that the area under the curve (AUC) is 0.685 with p-value=0.007 (Figure 1). Thus, the cut-off value of TNF- α to predict patient mortality is 37.88 ng/L, with a sensitivity of 62.07%, specificity 68.89%, positive predictive value (PPV) 56.25% and negative predictive value (NPV) 73.81%.

The Ability of IL-1 β as a Predictor of Mortality of COVID-19 Patients

The ability of IL-1 β was determined as a predictor of COVID-19 patient's mortality, an analysis of the ROC curve of IL-1 β levels was carried out on the mortality of COVID-19 patients (Figure 2). The analysis results show that the area under the curve (AUC) is 0.645, with a significance value of p=0.035. Furthermore, the analysis results show that the area under the curve (AUC) is 0.645, with p-value=0.035. Thus, IL-1 β levels predict the death of COVID-19 patients at the cut-off point of 761.54 pg/L with a sensitivity of 59.46%, specificity 70.59%, PPV 68.75% and NPV 61.54%.




Figure 1 ROC curve of TNF- α levels on death of COVID-19 patients.




Figure 2 ROC curve of IL-1 β levels on death of COVID-19 patients.

The Ability of MCP-1 as a Predictor of Mortality of COVID-19 Patients

MCP-1 ability as a predictor of mortality in COVID-19 patients, an analysis of the ROC curve of MCP-1 levels on the mortality of COVID-19 patients was carried out (Figure 3). The analysis results show that the area under the curve (AUC) is 0.702 with a significance value of $p=0.004$. Therefore, MCP-1 levels predict the death of COVID-19 patients with a cut-off value of 51.31 PPV 52.9% and NPV of 71.1%.

Discussion

Cytokine dysregulation has been identified as one of the major factors for poor prognosis in COVID-19.²³ This cytokine storm produces an exaggerated inflammatory response, particularly in the lungs, leading to ARDS, pulmonary oedema, and multi-organ failure. Thus, pro-inflammatory factors play a central role in the severity of COVID-19, especially in patients with co-morbidities.^{24,25} An overactive and untreated immune response can be lethal in these situations, suggesting that death in COVID-19 cases is likely due to this virus-driven hyper inflammation.²⁶

TNF- α is one of the cytokines that promote inflammation, including during SARS-CoV-2 infection. Of the 41 patients suffering from COVID-19, 13 were admitted to the ICU due to severe COVID-19. The patients treated in the ICU had higher IL2, IL7, IL10, GSCF, IP10, MCP-1, MIP1A, and TNF than non-ICU patients.²⁷ According to a cohort study conducted by Del Valle et al, 1484 COVID-19 patients were followed up to 41 days after admission with a median of 8 days.²³ They found that high serum levels of IL-6, IL-8 and TNF- α at the time of hospitalisation were strong and independent predictors of patient survival. In addition, the investigators obtained a cut-off of 35 ng/L, where values above the cut-off value had a high mortality rate.²⁸

We followed the patient's condition for up to 28 days so that we got the patient outcomes in the form of survivors and non-survivors. TNF- α was calculated on the first day the patient was admitted. The optimal cut-off value of TNF- α obtained statistically was 37.88 ng/L. This cut-off value becomes a reference for calculating sensitivity, specificity, positive and negative predictive values, and relative risk in the TNF- α examination. In this study, the median TNF- α in




Figure 3 ROC curve of MCP-1 levels on death of COVID-19 patients.

non-survivor patients was higher than in survivors. The increase in TNF- α levels in non-survivor patients in this study supports the results of a study conducted by Huang et al and Valle et al, where high levels of TNF- α were found in both ICU-treated and deceased COVID-19 patients. In addition, TNF- α can also be an independent predictor of severity in COVID-19 patients.^{28,29}

Cytokine storm is positively correlated with disease severity. Cytokine storm conditions in COVID-19 patients have been shown to increase the severity of COVID-19 disease so that it can increase the risk of death in COVID-19.³⁰ However, few studies prove the relationship of IL-1 β with the death of COVID-19 patients. In this study, the mean levels of IL-1 β were higher in the non-survivor group of COVID-19 patients than in the survivor group, which is consistent with previous studies. The study by Nile et al on 27 severe COVID-19 patients analysed nine biomarkers (IL-1 β , IL-2R, IL-6, IL-8, IL-10, TNF-, CRP, Ferritin) and lymphocytes as pro-inflammatory biomarkers in COVID-19.³¹ This study found that pro-inflammatory cytokines were increased in patients with severe COVID-19.

In addition, a study by Shang et al in 55 COVID-19 (mild-severe) patients admitted to the general ward and ICU showed an increase in IL-1 β levels.³² The increase in IL-1 β is significant and correlates with worsening the patient's condition, namely damage to lung tissue, respiratory failure, and death.²⁷ Thus, an increase in IL-1 levels in COVID-19 patients can potentially be a predictor of death, wherein this study, IL-1 β levels 761.54 pg/L had a sensitivity of 59.46%, specificity 70.59%, positive predictive value 68.75%, a negative predictive value of 61.54% for death and an increase in IL-1 β levels of 1.12 times can increase the risk of death of COVID-19 patients.

Cytokine storm conditions in COVID-19 patients have been shown to increase the severity of COVID-19 disease so that it can increase the risk of death in COVID-19.^{33,34} However, few studies have proven the relationship between MCP-1 and the mortality of COVID-19 patients. In this study, the mean MCP-1 level was higher in the non-survivor group of COVID-19 patients than in the survivor group, consistent with previous studies. A study by Abers et al, on 175 COVID-19 patients (mild-critical) analysed 66 soluble biomarkers with the severity of symptoms and mortality in COVID-19.³⁵ CCL2/MCP-1 as a biomarker associated with monocyte/macrophage activation and NF κ B signalling increased COVID-19 mortality compared to moderate-critical cases and healthy controls, and increased MCP-1 was associated with mortality independently. In addition, a study by Anderberg et al, on 24 confirmed COVID-19 patients admitted to the ICU showed that MCP-1 levels were correlated with the occurrence of respiratory failure, acute renal failure and death.⁷ Thus, an increase in MCP-1 levels can predict death in COVID-19, wherein this study, MCP-1 levels 51.31 ng/L had

a sensitivity of 62.1%, specificity 62.8%, the positive predictive value of 52.9%, a negative predictive value of 71.1% for mortality and an increase in MCP-1 levels 2.71 times increased the risk of death of COVID-19 patients. This study found that the significant difference was obtained from the level of severity, not based on the risk factors and morbidity in COVID-19 patients.

Conclusion

The levels of TNF- α , IL-1 β , and MCP-1 showed significant differences between survivors and non-survivors in COVID-19 patients. Therefore, the levels of all three have strong potential as predictors of outcome for COVID-19 patients. In addition, TNF- α , IL-1 β , and MCP-1 as markers of acute systemic inflammatory cytokines can be measured at the beginning of hospitalisation of COVID-19 patients for early diagnosis of disease severity so that healthcare professionals can determine clinical guidance needs for therapeutic programs. Finally, the recovery of COVID-19 patients is expected to increase.

Acknowledgments

The authors are thankful for the Ministry of Culture and Education; the Republic of Indonesia sponsored this research financially through "Program Penelitian Kolaborasi Indonesia" World Class University with grant numbers: 342.2/UN10.C10/PN/2021.

Disclosure

The authors report no conflicts of interest in this work.

References

- Fajgenbaum DC, June CH. Cytokine Storm. *N Engl J Med*. 2020;383(23):2255–2273. doi:10.1056/NEJMra2026131
- Abou-Ismaïl MY, Diamond A, Kapoor S, Arafah Y, Nayak L. The hypercoagulable state in COVID-19: incidence, pathophysiology, and management. *Thromb Res*. 2020;194:101–115. doi:10.1016/j.thromres.2020.06.029
- Jose RJ, Manuel A. COVID-19 cytokine storm: the interplay between inflammation and coagulation. *Lancet Respir Med*. 2020;8(6):e46–e47. doi:10.1016/S2213-2600(20)30216-2
- Costela-Ruiz VJ, Illescas-Montes R, Puerta-Puerta JM, Ruiz C, Melguizo-Rodríguez L. SARS-CoV-2 infection: the role of cytokines in COVID-19 disease. *Cytokine Growth Factor Rev*. 2020;54:62–75.
- Darif D, Hammi I, Kihel A. The pro-inflammatory cytokines in COVID-19 pathogenesis: what goes wrong? *Microb Pathog*. 2021;153:104799. doi:10.1016/j.micpath.2021.104799
- Huang C, Wang Y, Li X, Ren L. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497–506. doi:10.1016/S0140-6736(20)30183-5
- Bülow Anderberg S, Luther T, Berglund M, et al. Increased levels of plasma cytokines and correlations to organ failure and 30-day mortality in critically ill Covid-19 patients. *Cytokine*. 2021;138:155389. doi:10.1016/j.cyto.2020.155389
- Chen Y, Wang J, Liu C, et al. IP-10 and MCP-1 as biomarkers associated with disease severity of COVID-19. *Mol Med*. 2020;26(1):97. doi:10.1186/s10020-020-00230-x
- Leija-Martínez JJ, Huang F, Del-Río-Navarro BE, et al. IL-17A and TNF- α as potential biomarkers for acute respiratory distress syndrome and mortality in patients with obesity and COVID-19. *Med Hypotheses*. 2020;144:109935. doi:10.1016/j.mehy.2020.109935
- Kany S, Vollrath JT, Relja B. Cytokines in Inflammatory Disease. *Int J Mol Sci*. 2019;20(23):6008.
- Idriss HT, Naismith JH. TNF alpha and the TNF receptor superfamily: structure-function relationship(s). *Microsc Res Tech*. 2000;50(3):184–195.
- Birra D, Benucci M, Landolfi L, et al. COVID 19: a clue from innate immunity. *Immunol Res*. 2020;68(3):161–168. doi:10.1007/s12026-020-09137-5
- Giamarellos-Bourboulis EJ, Netea MG, Rovina N, et al. Complex Immune Dysregulation in COVID-19 Patients with Severe Respiratory Failure. *Cell Host Microbe*. 2020;27(6):992–1000.e3. doi:10.1016/j.chom.2020.04.009
- Song P, Li W, Xie J, Hou Y, You C. Cytokine storm induced by SARS-CoV-2. *Clin Chim Acta*. 2020;509:280–287. doi:10.1016/j.cca.2020.06.017
- Zhang X, Li S, Niu S. ACE2 and COVID-19 and the resulting ARDS. *Postgrad Med J*. 2020;96(1137):403–407. doi:10.1136/postgradmedj-2020-137935
- Yang Y, Shen C, Li J, et al. Exuberant elevation of IP-10, MCP-3 and IL-1ra during SARS-CoV-2 infection is associated with disease severity and fatal outcome. *Infect Dis*. 2020. doi:10.1101/2020.03.02.20029975
- Ye Q, Wang B, Mao J. The pathogenesis and treatment of the 'Cytokine Storm' in COVID-19. *J Infect*. 2020;80(6):607–613.
- Zhong J, Tang J, Ye C, Dong L. The immunology of COVID-19: is immune modulation an option for treatment? *Lancet Rheumatol*. 2020;2(7):e428–e436.
- Deshmane SL, Kremlev S, Amini S, Sawaya BE. Monocyte chemoattractant protein-1 (MCP-1): an overview. *J Interferon Cytokine Res*. 2009;29(6):313–326. doi:10.1089/jir.2008.0027
- Lin L, Lu L, Cao W, Li T. Hypothesis for potential pathogenesis of SARS-CoV-2 infection-A review of immune changes in patients with viral pneumonia. *Emerg Microbes Infect*. 2020;9(1):727–732. doi:10.1080/22221751.2020.1746199

21. Lorkiewicz P, Waszkiewicz N. Biomarkers of Post-COVID Depression. *J Clin Med.* 2021;10(18):4142.
22. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines; 2020. Available from: <https://www.covid19treatmentguidelines.nih.gov/>. Accessed 26 January 2022.
23. Kim JS, Lee JY, Yang JW, et al. Immunopathogenesis and treatment of cytokine storm in COVID-19. *Theranostics.* 2021;11(1):316–329.
24. Ni M, Tian FB, Xiang DD, Yu B. Characteristics of inflammatory factors and lymphocyte subsets in patients with severe COVID-19. *J Med Virol.* 2020;92(11):2600–2606. doi:10.1002/jmv.26070
25. Heriansyah T, Chomsy IN, Febrianda L, Farahiya Hadi T, Andri Wihastuti T. The Potential Benefit of Beta-Blockers for the Management of COVID-19 Protocol Therapy-Induced QT Prolongation: a Literature Review. *Sci Pharm.* 2020;88(4):55. doi:10.3390/scipharm88040055
26. de la Rica R, Borges M, Gonzalez-Freire M. COVID-19: in the Eye of the Cytokine Storm. *Front Immunol.* 2020;11:1–11.
27. McElvaney OJ, McEvoy NL, McElvaney OF, et al. Characterization of the Inflammatory Response to Severe COVID-19 Illness. *Am J Respir Crit Care Med.* 2020;202(6):812–821.
28. Del Valle DM, Kim-Schulze S, Huang HH, et al. An inflammatory cytokine signature predicts COVID-19 severity and survival. *Nat Med.* 2020;26(10):1636–1643.
29. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet.* 2020;395(10223):497–506.
30. Ragab D, Salah Eldin H, Tacimah M, Khattab R, Salem R. The COVID-19 Cytokine Storm; What We Know So Far. *Front Immunol.* 2020;11:1446.
31. Nile SH, Nile A, Qiu J, Li L, Jia X, Kai G. COVID-19: pathogenesis, cytokine storm and therapeutic potential of interferons. *Cytokine Growth Factor Rev.* 2020;53:66–70. doi:10.1016/j.cytogfr.2020.05.002
32. Shang W, Dong J, Ren Y, et al. The value of clinical parameters in predicting the severity of COVID-19. *J Med Virol.* 2020;92(10):2188–2192. doi:10.1002/jmv.26031
33. Tang Y, Liu J, Zhang D, Xu Z, Ji J, Wen C. Cytokine Storm in COVID-19: the Current Evidence and Treatment Strategies. *Front Immunol.* 2020;11:1708.
34. Wu F, Zhao S, Yu B, et al. Author Correction: a new coronavirus associated with human respiratory disease in China. *Nature.* 2020;580(7803):E7.
35. Abers MS, Delmonte OM, Ricotta EE, et al. An immune-based biomarker signature is associated with mortality in COVID-19 patients. *JCI Insight.* 2021;6(1):e144455.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

The Potential Predictive Role

ORIGINALITY REPORT

18%

SIMILARITY INDEX

9%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Xiao Tang, Ronghui Du, Rui Wang, Tanze Cao et al. "Comparison of Hospitalized Patients with Acute Respiratory Distress Syndrome Caused by COVID-19 and H1N1", Chest, 2020
Publication <1 %
- 2 radar.brookes.ac.uk
Internet Source <1 %
- 3 Nicolas Vabret, Graham J. Britton, Conor Gruber, Samarth Hegde et al. "Immunology of COVID-19: current state of the science", Immunity, 2020
Publication <1 %
- 4 Francesca Coperchini, Luca Chiovato, Gianluca Ricci, Laura Croce, Flavia Magri, Mario Rotondi. "The cytokine storm in COVID-19: Further advances in our understanding the role of specific chemokines involved", Cytokine & Growth Factor Reviews, 2021
Publication <1 %
- 5 Ming Ni, Fang - Bing Tian, Dan - Dan Xiang, Bing Yu. "Characteristics of inflammatory <1 %

factors and lymphocyte subsets in patients with severe COVID - 19", Journal of Medical Virology, 2020

Publication

6

Mohd. Iqbal Yattoo, Zeenat Hamid, Izhar Rather, Qurat Ul Ain Nazir et al.

"Immunotherapies and immunomodulatory approaches in clinical trials - a mini review", Human Vaccines & Immunotherapeutics, 2021

Publication

7

Pau Yen Wu, Ana Isabel Caceres, Jiegen Chen, Jamie Sokoloff et al. "Vagus nerve stimulation rescues persistent pain following orthopedic surgery in adult mice.", Cold Spring Harbor Laboratory, 2023

Publication

8

repository.medri.uniri.hr

Internet Source

9

www.koreascience.or.kr

Internet Source

10

Bo Diao, Chenhui Wang, Yingjun Tan, Xiewan Chen et al. "Reduction and Functional Exhaustion of T Cells in Patients With Coronavirus Disease 2019 (COVID-19)", Frontiers in Immunology, 2020

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

- | | | |
|----|---|------|
| 11 | Luca Gallelli, Leiming Zhang, Tian Wang, Fenghua Fu. "Severe Acute Lung Injury Related to COVID - 19 Infection: A Review and the Possible Role for Escin", The Journal of Clinical Pharmacology, 2020
Publication | <1 % |
| 12 | Na Wang, Lushan Liu, Wei He, Na Shang, Junyu Li, Zhou Qin, Xiaoxia Du. "Circulating Mid-regional Proadrenomedullin is a predictor of mortality in patients with COVID-19: a systematic review and meta-analysis", Research Square Platform LLC, 2022
Publication | <1 % |
| 13 | encyclopedia.pub
Internet Source | <1 % |
| 14 | pesquisa.bvsalud.org
Internet Source | <1 % |
| 15 | www.japtronline.com
Internet Source | <1 % |
| 16 | www.juaps.uoanbar.edu.iq
Internet Source | <1 % |
| 17 | Jordi Rello, Mirko Belliato, Meletios-Athanasios Dimopoulos, Evangelos J. Giamarellos-bourboulis et al. "Update in COVID-19 in the Intensive Care Unit from the 2020 HELLENIC Athens International Symposium", | <1 % |

Anaesthesia Critical Care & Pain Medicine, 2020

Publication

18

Sunita Acharya, Shanti Giri, Asha Panth.
"Knowledge and Perceptions of Corona Virus
Disease (COVID 19) among Nurses Working at
Teaching Hospital, Banke, Nepal", Journal of
Advanced Academic Research, 2023

Publication

<1 %

19

oamjms.eu

Internet Source

<1 %

20

www.biomedcentral.com

Internet Source

<1 %

21

www.imrpress.com

Internet Source

<1 %

22

Domenico Birra, Maurizio Benucci, Luigi
Landolfi, Anna Merchionda et al. "COVID 19: a
clue from innate immunity", Immunologic
Research, 2020

Publication

<1 %

23

Lidya Chaidir, Ida Parwati, Jessi Annisa, Soni
Muhsinin, Intan Meilana, Bacht Alisjahbana,
Reinout van Crevel. "Implementation of LED
Fluorescence Microscopy for Diagnosis of
Pulmonary and HIV-Associated Tuberculosis
in a Hospital Setting in Indonesia", PLoS ONE,
2013

Publication

<1 %

24 Marthe Jøntvedt Jørgensen, Jan Cato Holter, Erik Egeland Christensen, Camilla Schjalm et al. "Increased interleukin-6 and macrophage chemoattractant protein-1 are associated with respiratory failure in COVID-19", Scientific Reports, 2020
Publication

<1 %

25 Muhammad Suhaib Qudus, Mingfu Tian, Summan Sirajuddin, Siyu Liu et al. "The roles of critical pro - inflammatory cytokines in the drive of cytokine storm during SARS - CoV - 2 infection", Journal of Medical Virology, 2023
Publication

<1 %

26 bmcsresnotes.biomedcentral.com
Internet Source

<1 %

27 downloads.hindawi.com
Internet Source

<1 %

28 www.biorxiv.org
Internet Source

<1 %

29 www.ijpmonline.org
Internet Source

<1 %

30 "Correction to 11C-methyl-L-methionine PET measuring parameters for the diagnosis of tumour progression against radiation-induced changes in brain metastases", The British Journal of Radiology, 2023
Publication

<1 %

31

Abdulkarim F. Alhetheel, Ahmed M. Albarrag, Zahid A. Shakoor, Ali M. Somily et al.

"Assessment of Proinflammatory Cytokines Among Patients with Middle East Respiratory Syndrome Coronavirus Infection", *Viral Immunology*, 2023

Publication

<1 %

32

Cedric P. Van Dijck, Abhijit V. Lele, Rajen Nathwani. "Chapter 10 Trauma Care and Coronavirus Disease", Springer Science and Business Media LLC, 2020

Publication

<1 %

33

Elena Sancho Ferrando, Katja Hanslin, Michael Hultström, Anders Larsson, Robert Frithiof, Miklos Lipcsey. "Soluble TNF receptors predict acute kidney injury and mortality in critically ill COVID-19 patients: a prospective observational study", *Cytokine*, 2021

Publication

<1 %

34

Esteban Ortiz-Prado, Katherine Simbaña-Rivera, Lenin Gómez-Barreno, Mario Rubio-Neira et al. "Clinical, molecular and epidemiological characterization of the SARS-CoV2 virus and the Coronavirus disease 2019 (COVID-19), a comprehensive literature review", *Diagnostic Microbiology and Infectious Disease*, 2020

Publication

<1 %

35

Lu Zheng, Lina Zhang, Jiamin Huang, Kutty Selva Nandakumar, Shuwen Liu, Kui Cheng. "Potential treatment methods targeting 2019-nCoV infection", European Journal of Medicinal Chemistry, 2020

Publication

<1 %

36

Lvliang Lu, Wenyu Zhong, Ziwei Bian, Zhiming Li et al. "A Comparison of Mortality-related Risk Factors of COVID-19, SARS, and MERS: A Systematic Review and Meta-analysis", Journal of Infection, 2020

Publication

<1 %

37

Martijn Zoodsma, Aline H. de Nooijer, Inge Grondman, Manoj Kumar Gupta et al. "Targeted proteomics identifies circulating biomarkers associated with active COVID-19 and post-COVID-19", Frontiers in Immunology, 2022

Publication

<1 %

38

Wagner, J., L. Keuky, M. Scully, T. Kuoch, and C. Toch. "PO078 TRAINING VILLAGE HEALTH VOLUNTEERS FOR DIABETES PREVENTION IN CAMBODIA: PRELIMINARY FINDINGS FOR INCREASED KNOWLEDGE WITH THE EAT, WALK, SLEEP CURRICULUM", Diabetes Research and Clinical Practice, 2014.

Publication

<1 %

39 Zheyang Tao, Jing Xu, Wei Chen, Zhitao Yang et al. "Anemia is associated with severe illness in COVID - 19: A retrospective cohort study", Journal of Medical Virology, 2020
Publication <1 %

40 aacijournal.biomedcentral.com
Internet Source <1 %

41 discoveriesjournals.org
Internet Source <1 %

42 ijms.info
Internet Source <1 %

43 jamanetwork.com
Internet Source <1 %

44 pubmed.ncbi.nlm.nih.gov
Internet Source <1 %

45 stpm.si
Internet Source <1 %

46 www.frontiersin.org
Internet Source <1 %

47 www.mdpi.com
Internet Source <1 %

48 www.medicinabuenosaires.com
Internet Source <1 %

49 www.termedia.pl
Internet Source <1 %

50

1library.net

Internet Source

<1 %

51

Conrado Avendaño, Anahí Franchi, Hakan Duran, Sergio Oehninger. "DNA fragmentation of normal spermatozoa negatively impacts embryo quality and intracytoplasmic sperm injection outcome", *Fertility and Sterility*, 2010

Publication

<1 %

52

Dita Aditianingsih, Ratna Farida Soenarto, Artheta Mutiara Puiantana, Raymond Pranata et al. "Dose response relationship between D-dimer level and mortality in critically ill COVID-19 patients: a retrospective observational study", *F1000Research*, 2022

Publication

<1 %

53

Gabriel Siracusano, Claudia Pastori, Lucia Lopalco. "Humoral Immune Responses in COVID-19 Patients: A Window on the State of the Art", *Frontiers in Immunology*, 2020

Publication

<1 %

54

H. Burke, A. Freeman, D. C. Cellura, B. L. Stuart et al. "Inflammatory phenotyping predicts clinical outcome in COVID-19", *Respiratory Research*, 2020

Publication

<1 %

55

Hao-Yu Wu, Chien-Ting Pan, Chiao-Feng Cheng, Chi-Ying Lin et al. "Combined

<1 %

Intravenous Immunoglobulin and Baricitinib Treatment for Severe COVID-19 with Rhabdomyolysis: A Case Report", Journal of the Formosan Medical Association, 2021

Publication

56

Jingfeng Bao, Chenxi Li, Kai Zhang, Haiquan Kang, Wensen Chen, Bing Gu. "Comparative analysis of laboratory indexes of severe and non-severe patients infected with COVID-19", Clinica Chimica Acta, 2020

Publication

<1 %

57

Lea Skak Filtenborg Frederiksen, Yibang Zhang, Camilla Foged, Aneesh Thakur. "The Long Road Toward COVID-19 Herd Immunity: Vaccine Platform Technologies and Mass Immunization Strategies", Frontiers in Immunology, 2020

Publication

<1 %

58

Mehrdad Rostami, Hassan Mansouritorghabeh. "D-dimer level in COVID-19 infection: a systematic review", Expert Review of Hematology, 2020

Publication

<1 %

59

Mutlu Kuluöztürk, Erdal İn, Selda Telo, Ercan Karabulut, Ayşegül Altıntop Geçkil. "Efficacy of copeptin in distinguishing COVID - 19 pneumonia from community - acquired pneumonia", Journal of Medical Virology, 2021

Publication

<1 %

60 Piotr Lorkiewicz, Napoleon Waszkiewicz.
"Biomarkers of Post-COVID Depression",
Journal of Clinical Medicine, 2021
Publication <1 %

61 Weifeng Shang, Junwu Dong, Yali Ren, Ming
Tian, Wei Li, Jianwu Hu, Yuanyuan Li. "The
value of clinical parameters in predicting the
severity of COVID - 19", Journal of Medical
Virology, 2020
Publication <1 %

62 Xian Zhou, Gerald Münch, Hans Wohlmuth,
Sualiha Afzal et al. "Synergistic Inhibition of
Pro-Inflammatory Pathways by Ginger and
Turmeric Extracts in RAW 264.7 Cells",
Frontiers in Pharmacology, 2022
Publication <1 %

63 ann-clinmicrob.biomedcentral.com
Internet Source <1 %

64 medauth2.mdedge.com
Internet Source <1 %

65 usir.salford.ac.uk
Internet Source <1 %

66 wjgnet.com
Internet Source <1 %

67 www.omicsdi.org
Internet Source <1 %

68

www.pure.ed.ac.uk

Internet Source

<1 %

69

www.sysrevpharm.org

Internet Source

<1 %

70

www.tandfonline.com

Internet Source

<1 %

71

H Nataprawira, I Parwati. "PO-0199
Microscopic Observation Drugs Susceptibility
(mods): A Promising Culture-based Assay To
Diagnosing Tuberculosis In Children", Archives
of Disease in Childhood, 2014

Publication

<1 %

72

Joshua Krieger, Flynn McGuire, Erik Risa,
August Longino et al. "Emergency department
characteristics and associations with intensive
care admission among patients with
coronavirus disease 2019", Journal of the
American College of Emergency Physicians
Open, 2021

Publication

<1 %

73

Paulo Ricardo Martins-Filho, Carolina Santos
Souza Tavares, Victor Santana Santos.
"Factors associated with mortality in patients
with COVID-19. A quantitative evidence
synthesis of clinical and laboratory data",
European Journal of Internal Medicine, 2020

Publication

<1 %

74

Piotr Lorkiewicz, Napoleon Waszkiewicz. "Is SARS-CoV-2 a Risk Factor of Bipolar Disorder? —A Narrative Review", Journal of Clinical Medicine, 2022

Publication

<1 %

75

Raffaele Costa, Alberto Castagna, Giovanni Ruotolo. "COVID - 19 and cardiovascular problems in elderly patients: Food for thought", AGING MEDICINE, 2021

Publication

<1 %

76

Sajad Khiali, Elnaz Khani, Taher Entezari - Maleki. "A Comprehensive Review of Tocilizumab in COVID - 19 Acute Respiratory Distress Syndrome", The Journal of Clinical Pharmacology, 2020

Publication

<1 %

77

Sara I. Taha, Sara Samaan, Aalaa Shata, Shereen Baioumy, Shaimaa Abdalgeleel, Mariam Youssef. "Baseline Characteristics and Outcomes of 180 Egyptian COVID-19 Patients Admitted to Quarantine Hospitals of Ain Shams University: A Retrospective Comparative Study", Afro-Egyptian Journal of Infectious and Endemic Diseases, 2021

Publication

<1 %

78

Zheyong Huang, Jiatian Cao, Yumeng Yao, Xuejuan Jin et al. "The effect of RAS blockers on the clinical characteristics of COVID-19

<1 %

patients with hypertension", *Annals of Translational Medicine*, 2020

Publication

79

Aloon Eko Widiono, Sukarni Sukarni, Retno Wulandari, Ardianto Prasetyo, Heru Suryanto, Uun Yanuhar. "Pyrolytic thermal decomposition behavior and kinetic parameters of *Tetraselmis chuii* microalgae", AIP Publishing, 2020

<1 %

80

Jing Guo, Shuting Wang, He Xia, Ding Shi et al. "Cytokine Signature Associated With Disease Severity in COVID-19", *Frontiers in Immunology*, 2021

<1 %

81

Mohd Arish, Wei Qian, Harish Narasimhan, Jie Sun. "COVID - 19 immunopathology: From acute diseases to chronic sequelae", *Journal of Medical Virology*, 2022

<1 %

82

Xiaojing Zou, Minghao Fang, Shusheng Li, Liang Wu et al. "Characteristics of liver function in patients with SARS-CoV-2 and chronic HBV co-infection", *Clinical Gastroenterology and Hepatology*, 2020

<1 %

83

Zili Zhang, Fanjie Lin, Fei Liu, Qiongqiong Li et al. "Proteomic profiling reveals a distinctive

<1 %

molecular signature for critically ill COVID-19 patients compared with asthma and COPD", International Journal of Infectious Diseases, 2022

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

The Potential Predictive Role

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
