

# Effect of Zinc(II)

*by* Aswandi Wibrianto

---

**Submission date:** 23-May-2023 11:17AM (UTC+0800)

**Submission ID:** 2099750111

**File name:** Effect\_of\_Zinc\_II.pdf (1.17M)

**Word count:** 3783

**Character count:** 20632

15

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/346668634>

# Effect of Zinc(II)-2,4,5-triphenyl-1H-imidazole Complex Against Replication DENV-2 in Vero Cell

Article in *Indonesian Journal of Tropical and Infectious Disease* · December 2020

DOI: 10.20473/ijtd.v8i3.11776

CITATION

1

READS

87

9 authors, including:

**Aswandi Wibrianto**

National Taiwan University of Science and Technology

13 PUBLICATIONS 25 CITATIONS

[SEE PROFILE](#)**Teguh Hari Sucipto**

Airlangga University

45 PUBLICATIONS 267 CITATIONS

[SEE PROFILE](#)**Siti Churrotin**

Airlangga University

33 PUBLICATIONS 272 CITATIONS

[SEE PROFILE](#)**Ilham Harlan Amarullah**

Airlangga University

13 PUBLICATIONS 113 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



8

Synthesis of Complex Compounds Ni(II)-Chlorophyll as Dye Sensitizer in Dye Sensitizer Solar Cell (DSSC) [View project](#)Microbiology [View project](#)

6

All content following this page was uploaded by [Aswandi Wibrianto](#) on 07 December 2020.

The user has requested enhancement of the downloaded file.

# Indonesian Journal of Tropical and Infectious Disease

Vol. 8 No. 3 September–December 2020

## Case Report

### Effect of Zinc(II)-2,4,5-triphenyl-1H-imidazole Complex Against Replication DENV-2 in Vero Cell

Aswandi Wibrianto<sup>1</sup>, Fatimah Martak<sup>2</sup>, Teguh Hari Sucipto<sup>3a</sup>, Siti Churrotin<sup>3</sup>, Ilham Harlan Amarullah<sup>3</sup>, Harsasi

1. Setyawati<sup>4</sup>, Puspa Wardhani<sup>3</sup>, Aryati<sup>3</sup>, Soegeng Soegijanto<sup>3</sup>

<sup>1</sup> Undergraduate Student, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Indonesia

<sup>2</sup> Department of Chemistry, Faculty of Natural Science, Institut Teknologi Sepuluh Nopember, Indonesia

<sup>3</sup> Dengue Study Group, Institute of Tropical Disease, Universitas Airlangga, Indonesia

<sup>4</sup> Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Indonesia

Received: 8<sup>th</sup> April 2019; Revised: 29<sup>th</sup> January 2020; Accepted: 23<sup>rd</sup> April 2020

#### ABSTRACT

Dengue virus (DENV) is a significant pathogen emerging worldwide as a cause of infectious disease. DENVs are transmitted to humans through female mosquitoes from *Aedes aegypti* and *Aedes albopictus* species. Indonesia is one of the largest countries in the world in dengue endemic regions worldwide. Dengue fever was occurred for the first time as an outbreak in Surabaya and Jakarta in 1968. Many efforts have been made to prevent and treat DENV infections, and clinical trials of a number of vaccines are currently underway. Antiviral testing of DENV is an important alternative for drug characterization and development. Complex compounds are formed as a result of metal and organic complex reactions. Complex compounds can be used as an anti-inflammatory, antimicrobial antifungal, antibacterial, antiviral. The  $Zn^{2+}$  ion can be used as an antiviral candidate. The purpose of this project was investigated Zinc(II)-2,4,5-triphenyl-1H-imidazole antiviral compound to be further tested for inhibitory effect on the replication of DENV-2 in cell culture. DENV replication was measured by antiviral activity assay and cytotoxicity assay. The inhibitory activity of Zinc(II)-2,4,5-triphenyl-1H-imidazole complex compound was determined by Viral ToxGloTM Assay. The cytotoxicity of Zinc(II)-2,4,5-triphenyl-1H-imidazole complex compound was determined by Cell Titer96® AQ<sub>uoeus</sub> assay. The inhibitory concentration ( $IC_{50}$ ) of Zinc(II)-2,4,5-triphenyl-1H-imidazole against dengue virus type-2 was 34.42  $\mu$ g/ml. The cytotoxic concentration ( $CC_{50}$ ) of compound against Vero cell was <100  $\mu$ g/ml. The results of this study demonstrate the antidengue serotype 2 inhibitory activity of investigated Zinc(II)-2,4,5-triphenyl-1H-imidazole complex and its high toxicity in Vero cells. Further studies are not required before investigated Zinc(II)-2,4,5-triphenylimidazole can be applied in the treatment of DENV-2 infections.

**Keywords:** Zinc (II), complex compound, cytotoxicity, inhibitory activity, DENV-2

#### ABSTRAK

Virus Dengue (DENV) adalah patogen signifikan yang muncul di seluruh dunia sebagai penyebab penyakit menular. DENV ditransmisikan ke manusia melalui nyamuk betina dari spesies *Aedes aegypti* dan *Aedes albopictus*. Indonesia adalah salah satu negara terbesar di dunia di daerah endemik demam berdarah di seluruh dunia. Demam berdarah terjadi untuk pertama kalinya sebagai wabah di Surabaya dan Jakarta pada tahun 1968. Banyak upaya telah dilakukan untuk mencegah dan mengobati infeksi DENV, dan uji klinis sejumlah vaksin saat ini sedang berlangsung. Pengujian antivirus DENV adalah alternatif penting untuk karakterisasi dan pengembangan obat. Senyawa kompleks terbentuk sebagai hasil dari reaksi kompleks logam dan organik. Senyawa kompleks dapat digunakan sebagai anti-inflamasi, antimikroba antijamur, antibakteri, antivirus. Ion  $Zn^{2+}$  dapat digunakan sebagai kandidat antivirus. Tujuan dalam proyek ini adalah menyelidiki senyawa antivirus Zink(II)-2,4,5-trifenil-1H-imidazol yang diuji lebih lanjut untuk efek penghambatan pada replikasi DENV-2 dalam kultur sel. Replikasi DENV diukur dengan uji aktivitas antivirus dan uji sitotoksitas. Aktivitas penghambatan senyawa kompleks Zinc(II)-2,4,5-triphenyl-1H-imidazol ditentukan dengan Viral ToxGloTM Assay. Sitotoksitas senyawa kompleks Zinc(II)-2,4,5-triphenyl-

\* Corresponding Author:  
teguhharisucipto@staf.unair.ac.id

*1H*-imidazol ditentukan dengan uji CellTiter96® AQueous. Konsentrasi penghambatan ( $IC_{50}$ ) Zinc(II)-2,4,5-trifenil-1H-imidazol terhadap virus dengue tipe-2 adalah 34,42 µg/ml. Konsentrasi sitotoksik ( $CC_{50}$ ) senyawa terhadap sel Vero adalah <100 µg/ml. Hasil penelitian ini menunjukkan aktivitas penghambatan serotipe 2 antidengue dari Zinc(II)-2,4,5-trifenil-1H-imidazol yang diteliti dan toksisitasnya yang tinggi dalam sel Vero. Studi lebih lanjut tidak diperlukan sebelum investigasi Zinc(II)-2,4,5-trifenil-1H-imidazol dapat diterapkan dalam pengobatan infeksi DENV-2.

**Kata kunci:** Seng (II), senyawa kompleks, sitotoksitas, aktivitas penghambatan, DENV-2

**How to Cite:** Effect of Zinc(II)-2,4,5-triphenyl-1H-imidazole Complex Against Replication DENV-2 in Vero Cell. Wibrianto, A. Martak, F. Setyawati, H. Sucipto, TH. Churrotin, S. Amarullah, IH. Wardhani, P. Aryati, A. Soegijanto, S. Indonesian Journal of Tropical and Infectious Disease, 8(3), 183–188.

## INTRODUCTION

Dengue virus (DENV) is a virus carried by the *flavivirus* genus of the family Flaviviridae. Dengue virus (DENV) consists of four serotypes which is dengue virus type 1, dengue virus type 2, dengue virus type 3, and dengue virus type 4. Dengue virus is transmitted to humans through female mosquitoes from *Aedes aegypti* and *Aedes albopictus* species. World Health Organization (WHO) reported 390 million dengue infections per year.<sup>1</sup> Indonesia is one of the largest countries in the world with dengue endemic areas. Surabaya and Jakarta were the cities where dengue disease was first reported in Indonesia in 1968.<sup>2</sup> Many studies have been conducted to overcome the threat of dengue virus infections, and clinical trials of a number of vaccines are currently on the way.<sup>3</sup> Antiviral testing on DENV is a very important method in the development and characterization of drugs. Supplementary to vaccines, inhibitors in each natural cycle of viral replication have the potential to cure dengue virus infection and indeed compounds such as RNA replication inhibitors have been tested as such.<sup>4</sup> However, there is no commercially available drug with antiviral activity for DENV.<sup>5</sup>

Ligand 2,3,5-triphenyl-1H-imidazole compound is a derivate of imidazole. Imidazole-containing drugs that have strong therapeutic properties have encouraged scientists to synthesize many novel chemotherapeutic agents consisting of these entities. N<sup>5</sup>-(4-fluorophenyl)-N<sup>4</sup>-(2-(pyridin-4-yl)benzyl)-1H-imidazole-4,5-dicarboxamide, a derivate of imidazole, was reported anti-DENV activity.<sup>6</sup> 4-carbamoyl-5-(4,6-diamino-2,5-dihydro-1,3,5-triazin-2-yl)

imidazole-1-β-D-ribofuranoside is examined for four different types of viruses from the flaviridae family *in vitro*, including hepatitis C virus (HCV), Japanese viral encephalitis (JEV), West Nile virus (WNV), and dengue virus (DENV) *in vitro* against NTPases/helicases. The compound showed activity highly active against WNV with  $IC_{50}$  was 23 µM.<sup>7</sup>

Complex compounds are formed as a result of metal and organic compound reactions. Complex compounds can be used as an anti-inflammatory<sup>8</sup>, antimicrobial<sup>9</sup>, antifungal, antibacterial<sup>10</sup>, and antiviral<sup>11</sup>. Based on previous research, Copper(II)-imidazole derivatives can be used as antiDENV-2, can be used as low toxicity and potential as drug candidates. The compound exhibited adsorption inhibitory activity against DENV-2 at  $IC_{50} = 2.3$  µg/ml.<sup>12</sup>

The Zn<sup>2+</sup> ion can be used as an antiviral candidate.<sup>13</sup> Zn<sup>2+</sup> ions can change the activity of various transcription factors and thus, patterns of cellular and viral gene expression.<sup>14</sup> Thus, the antiviral test of the compound Zinc(II)-2,4,5-triphenyl-1H-imidazole was investigated.

## MATERIALS AND METHODS

### Chemicals and Media

Chemical reagents used in this research were Zinc(II)-2,4,5-triphenyl-1H-imidazole complex compound, Minimum Essential Eagle Medium (Sigma-Aldrich, Germany), dengue virus serotype 2 Surabaya isolate (KT012509), Vero cells (African Green Monkey Kidney), Viral ToxGlo™ assay (Promega, USA), CellTiter96®

AQ<sub>uoeus</sub> One Solution Cell Proliferation Assay (Promega, USA).

#### Antiviral Activity Assay

Confluent monolayers of Vero cells were prepared on a 96-well plate ( $1 \times 10^6$  cells/10 ml) and counted using a hemocytometer, and the titer of DENV-2 ( $2 \times 10^4$  FFU/well) was expressed in Foci-Forming Units (FFU) after incubating at 37°C for 2 days. The concentrations of Zinc(II)-2,4,5-triphenyl-1H-imidazole were 50 µg/mL; 25 µg/mL; 12.5 µg/mL; 6.25 µg/mL; 3.13 µg/mL; 1.57 µg/mL; 0.78 µg/mL; and 0.39 µg/mL with addition 100 µL Viral ToxGlo™ Assay per well. The 50% inhibitory concentration (IC<sub>50</sub>) of DENV-2 replication by each compound was further investigated by using GloMax® Discover System.

#### Cytotoxicity Assay

A cytotoxicity assay was performed using CellTiter96® AQ<sub>uoeus</sub> One Solution Cell Proliferation reagent. The CellTiter96® Assay is a modification of the MTT assay method portrayed by Akter.<sup>15</sup> The concentrations of Zinc(II)-2,4,5-triphenyl-1H-imidazole were 100 µg/mL; 200 µg/mL; 400 µg/mL; 600 µg/mL; 800 µg/mL; and 1000 µg/mL. The medium was allowed to equilibrate for 1 hour; then 20 µl/well of CellTiter 96® AQ<sub>uoeus</sub> One Solution Reagent was added. After 1 hour at 37°C in a humidified, 5% CO<sub>2</sub> atmosphere, the absorbance at 490nm was recorded using GloMax® Discover System.

#### Viral Detection by Reverse Transcriptase-Polymerase Chain Reaction

RNA replication was estimated using the Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR). The purpose of this assay was to known RNA replication after treatment. Briefly, DENV-2 RNA was extracted from the DENV-2 infected cells and cell culture supernatant using RNA extraction kit by Qiagen, Germany. The two-step kit (Toyobo, Japan) was used for cDNA synthesis and Polymerase Chain Reaction (PCR) following the manufacturer's instructions. Primer oligonucleotide sequences were as follows by

Bhatnagar et. al. 2012.<sup>16</sup> Amplification condition was 54 °C for one minute (annealing temperature) and the amplified product was the analyzed on 1.5% agarose gel.

## RESULTS AND DISCUSSION

The cytotoxicity of Zinc(II)-2,4,5-triphenyl-1H-imidazole complex compound was determined by CellTiter96® AQ<sub>uoeus</sub> assay and the recorded CC<sub>50</sub> value is <100 µg/ml to Vero cells. When compared with a previous study, Copper(II) was found to be nontoxic to human erythrocyte cells to concentrations of 500 µg/ml.<sup>17</sup> CC<sub>50</sub> is the cytotoxicity level of [Cu(2,4,5-triphenyl-1H-imidazole)<sub>2</sub>]<sub>n</sub> (compound) to cause death to 50% of Vero cells.<sup>12</sup> The toxicity value of Cobalt(II) complex with 2,4,5-triphenyl-1H-imidazole ligand was 362.24 mg/L, which was not toxic.<sup>18</sup> The toxicity value of 2-methyl-4,5-diphenyl-1H-Imidazole ligand compound was 192,3 µg/ml.<sup>19</sup> The toxicity of [Mn(2-(4-chlorophenyl)-4,5-diphenyl-1H-imidazole)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>].2H<sub>2</sub>O was >200 µg/ml which had less toxicity.<sup>20</sup> Zinc(II)-2-(2,4-dihydroxyphenyl)-3,5,7-trihydroxycromen-4-one complex compound defined cytotoxicity with CC<sub>50</sub> at 3.59 µg/ml.<sup>21</sup> But, the metal-free imidazole more toxic for Vero cells (CC<sub>50</sub> = 5.03 µg/ml).<sup>22</sup> Activity against HIV-1 strain IIIB and HIV-2 strain ROD in MT-4 cells (CC<sub>50</sub>) by zinc(II) complexes with hexyl-Me<sub>2</sub>-cyclam (HMC; 3,14-dimethyl-2,6,13,17-tetraazatricyclo(16.4.0.0<sup>7,12</sup>)-docosane) were >372 µM and >372 µM with selectivity index >35 and >3. Activity against HIV-1 strain IIIB and HIV-2 strain ROD in MT-4 cells (CC<sub>50</sub>) by Zn(II)-HMC diacetate were 110.67 ± 12.67 µM and 110.67 ± 12.67 µM with selectivity index 32 and <1.<sup>23</sup>

The complex stability is highly dependent on both the metallic ion and the ligands. As for the central ion (M<sup>2+</sup>), Zn(II) more unstable than Cu(II), Mn(II), and Co(II). The Zn(II) complex has grater polarizability that that Cu(II), Mn(II), and Co(II) because it contains more *d*-electrons, and the Zn(II) complex produced more product ions soluble in water.<sup>24</sup> This effect causes Zn(II) to be more toxic, because Zn<sup>2+</sup> in the medium are



more numerous, so it damages the cell wall faster than complex compound that have high stability such as Cu(II), Mn(II), and Co(II).


The percentage inhibition of the development of dengue virus type-2 by the test sample of Zinc(II)-2,4,5-triphenyl-1*H*-imidazole complex compound was shown on figure 1. The  $IC_{50}$  value was determined from the concentration–response curve (Figure 1); the  $IC_{50}$  value was 34.42  $\mu\text{g/ml}$ ,  $R^2$  was 0.9196. Based on the value of the  $IC_{50}$  Zinc(II)-2,4,5-triphenyl-1*H*-imidazole complex compound was a medium toxic compound.

Antiviral activity was also shown in Figure 2, these findings were corroborated by results obtained from RT-PCR which indicated significant reduction in the amount of DENV-2 genomic RNA levels. The highest percentage of viral inhibition was observed after treating the infected cells with 50  $\mu\text{g/ml}$ .


Based on the previous study, [Cu(2,4,5-triphenyl-1*H*-imidazole)<sub>2</sub>]<sub>n</sub> complex compound exhibited adsorption inhibitory activity against DENV-2 at  $IC_{50} = 2.3 \mu\text{g/ml}$ . The inhibition at  $IC_{50}$  was not significantly high ( $p < 0.005$ ) compared to that of the metal-free imidazole ( $IC_{50} = 0.13 \mu\text{g/ml}$ ).<sup>12</sup> The maximal inhibitory concentration ( $IC_{50}$ ) of Copper(II)chloride Dihydrate against DENV-2 was 0.13  $\mu\text{g/ml}$ .<sup>22</sup>

Activity against HIV-1 strain IIIB and HIV-2 strain ROD in MT-4 cells ( $IC_{50}$ ) by zinc(II) complexes with hexyl-Me<sub>2</sub>-cyclam (HMC; 3,14-dimethyl-2,6,13,17-tetraazatricyclo(16.4.0.0<sup>7,12</sup>)docosane) were  $10.51 \pm 0.23 \mu\text{M}$  and  $133.78 \pm 14.10 \mu\text{M}$ . Activity against HIV-1 strain IIIB and HIV-2 strain ROD in MT-4 cells ( $IC_{50}$ ) by Zn(II)–HMC diacetate were  $3.50 \pm 0.33 \mu\text{M}$  and  $>110.67 \mu\text{M}$ .<sup>23</sup> Anti-HIV-1 activity ( $IC_{50}$ ) in C8166/IIIB, MT-4/GUN1 and PBLs/IIIB were 8.0  $\mu\text{g/ml}$ , 3.5  $\mu\text{g/ml}$ , and 9.3  $\mu\text{g/ml}$ , respectively. The  $IC_{50}$  value of the Cobalt(II)–Morin complex for DENV-2 was 3.08  $\mu\text{g/ml}$ .<sup>25</sup> MB21, a benzimidazole derivative, was found to be the most potential inhibitor of cloned proteases ( $IC_{50} = 5.95 \mu\text{M}$ ).<sup>26</sup>

This study suggest that of Zinc(II)-2,4,5-triphenyl-1*H*-imidazole complex compound can't be an attractive antiviral option. It would



**Figure 1.** Inhibition of DENV-2, at variation concentrations of Zinc(II)-2,4,5-triphenyl-1*H*-imidazole complex compound



**Figure 2.** Electrophoresis on 1.5% agarose of RT-PCR after treatment, molecular weight marker (100 bp), (a) treatment with 50  $\mu\text{g/ml}$  compound, (b) 25  $\mu\text{g/ml}$ , (c) 12.5  $\mu\text{g/ml}$ , (d) 6.25  $\mu\text{g/ml}$ , (e) 3.13  $\mu\text{g/ml}$ , (f) 1.57  $\mu\text{g/ml}$ , (g) 0.78  $\mu\text{g/ml}$ , and (h) 0.39  $\mu\text{g/ml}$

be interesting to further investigate whether 2,4,5-triphenyl-1*H*-imidazole complex with other metal. The result of this study, Zinc(II)-2,4,5-triphenyl-1*H*-imidazole complex compound more toxic than Cu(II)-2,4,5-triphenyl-1*H*-imidazole, this is caused by the Zn (II) complex being unstable compared to the Cu (II) complex.

## CONCLUSION

Further studies are not required before Zinc(II)-2,4,5-triphenyl-1*H*-imidazole can be applied in the medication of DENV-2 infections. This study did not show the potential of the Zinc(II)-2,4,5-triphenyl-1*H*-imidazole complex as a candidate for antiviral agents against DENV-2 because it was shown to be toxic to Vero cells.

## CONFLICT OF INTEREST

There is no conflict of interest of this paper.

## ACKNOWLEDGEMENT

<sup>32</sup> This research was supported by a Research Grant from Mandat Universitas Airlangga (HRMUA) 2019; the Institute of Tropical Disease (ITD); the Center of Excellence (COE) program by the Ministry of Research and Technology (RISTEK) Indonesia; and the Chemistry Department of Universitas Airlangga.

## REFERENCES

- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, *et al.* The global distribution and burden of dengue. *Nature*. 2013;496(7446):504–7.
- Chi J, Meng L, Zhai X, Guo Y, Pan S, Zhou C, *et al.* Tanshinon II A attenuates diabetic cardiomyopathy by down-regulation of mTOR/P70S6K signaling pathway. *Int J Clin Exp Pathol*. 2016;9(12):12297–305.
- Woodland DL. Vaccines Against Dengue Virus. *Viral Immunol* [Internet]. 2015;28(2):75–75. Available from: <https://www.liebertpub.com/doi/10.1089/vim.2015.1502>
- Lim SP, Noble CG, Shi PY. The dengue virus NS5 protein as a target for drug discovery. *Antiviral Res* [Internet]. 2015;119(April):57–67. Available from: <http://dx.doi.org/10.1016/j.antiviral.2015.04.010>
- Assunção-Miranda I, Cruz-Oliveira C, Neris RLS, Figueiredo CM, Pereira LPS, Rodrigues D, *et al.* Inactivation of Dengue and Yellow Fever viruses by heme, cobalt-protoporphyrin IX and tin-protoporphyrin IX. *J Appl Microbiol*. 2016;120(3):790–804.
- Saudi M, Zmurko J, Kaptein S, Rozenski J, Neyts J, Van Aerschot A. Synthesis and evaluation of imidazole-4,5-and pyrazine-2,3-dicarboxamides targeting dengue and yellow fever virus. *Eur J Med Chem* [Internet]. 2014;87:529–39. Available from: <http://dx.doi.org/10.1016/j.ejmech.2014.09.062>
- Wu, H., Kou, F., Jia, F., Liu, B., Yuan, J., and Bai, Y. Synthesis and Characterization of the Ligand Based on Benzimidazole and Its Copper Complex: DNA Binding and Antioxidant Activity. *Bioinorganic Chemistry and Applications*, 2011; 2011: 1–11.
- Agotegaray MA, Dennehy M, Boeris MA, Grella MA, Burrow RA, Quinzani O V. Therapeutic properties, SOD and catecholase mimetic activities of novel ternary copper(II) complexes of the anti-inflammatory drug Fenopropfen with imidazole and caffeine. *Polyhedron* [Internet]. 2012;34(1):74–83. Available from: <http://dx.doi.org/10.1016/j.poly.2011.12.005>
- El-Sonbatia AZ, Diaba MA, and Morgan ShM. Thermal properties, antimicrobial activity and DNA binding of Ni(II) complexes of azo dye compounds. *Journal of Molecular Liquids*. 2017; 225:195-206.
- Kaczmarek MT, Zabiszak M, Nowak M, and Jastrzab R. Lanthanides: Schiff base complexes, applications in cancer diagnosis, therapy, and antibacterial activity. *Coordination Chemistry Reviews*. 2018; 370: 42–54.
- Shahabadi N, Pourfoulad M, and Taherpour A. Synthesis, characterization and in vitro DNA binding studies of a new copper(II) complex containing an antiviral drug, valganciclovir. *Journal of Coordination Chemistry*. 2016; 70(2): 201–222.
- Sucipto TH, Churrotin S, Setyawati H, Martak F, Mulyatno KC, Amarullah IH, *et al.* A new copper (II)-imidazole derivative effectively inhibits replication of dengue-2 in vero cell. *African J Infect Dis*. 2018;12(Special Issue 1):116–9.
- Sensi SL, Canzoniero LMT, Yu SP, Ying HS, Koh J, Kerchner GA, *et al.* Садыков\_Отрошенко\_Курбатов\_(Синтезы\_Орг\_Препаратов\_Пиридинового\_Ряда)\_ (1974). 1997;17(24):9554–64.
- Ishida T. Antiviral Activities of Zn<sup>2+</sup> Ions for Viral Prevention, Replication, Capsid Protein in Intracellular Proliferation of Viruses. *World Sci News* [Internet]. 2018;97(March):28–50. Available from: <http://yadda.icm.edu.pl/yadda/element/bwmeta1.element.psijd-8ab246c3-8e3f-4707-b97f-0c14fa91b094>
- Akter S, Addepalli R, Netzel ME, Tinggi U, Fletcher MT, Sultanbawa Y, and Osborne SA. Antioxidant-Rich Extracts of Terminalia ferdinandiana Interfere with Estimation of Cell Viability. *Antioxidants*. 2019; 8(6): 191-197.
- Bhatnagar J, Paddock CD, Shieh WJ, Blau DM, Liu L, Patel M, and Zaki SR. Molecular Detection and Typing of Dengue Viruses from Archived Tissues of Fatal Cases by RT-PCR and Sequencing: Diagnostic and Epidemiologic Implications. *The American Journal of Tropical Medicine and Hygiene*. 2012; 86(2): 335–340.
- Turecka K, Chylewska A, Kawiak A, and Waleron KF. Antifungal Activity and Mechanism of Action of the Co(III) Coordination Complexes With Diamine Chelate Ligands Against Reference and Clinical Strains of *Candida* spp. *Frontiers in Microbiology*. 2018; 9: 1594-1597.
- Martak F, Cahyani NWD, Nugraheni Z V, Utomo WP. Properties and toxicity of cobalt(II) complex with 2,4,5-triphenyl-1h-imidazole ligand. *Indones J Chem* [Internet]. 2016;16(3):260–7. Available from: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85002759335&doi=10.22146/2Fijc.1147&partnerID=40&md5=bb30e67ee19232caacec030aee50d3ce>
- Martak F, Herliana N, Yosina SD, Merriana PC, Wahyudi A, Gunawan WH. Synthesis and Toxicity Test

- of Cobalt (II), Manganese (II) and Copper (II) Complex with 2-Methyl-4, 5-Diphenyl-1H-Imidazole. *Indian J Sci Technol* [Internet]. 2018;11(7):1–11. Available from: <http://www.indjst.org/index.php/indjst/article/view/115356>
20. Martak F, Wahyudi A, Limanto D, Tajudin M, Ali M. Sintesis dan Karakterisasi Senyawa Kompleks Difenil - 1H - Imidazol. 2018;3(2):159–74.
  21. Sucipto TH, Setyawati H, Churrotin S, Amarullah IH, Sumarsih S, Wardhani P, and Soegijanto S. Anti-Dengue Type 2 Virus Activities of Zinc (II) Complex Compounds with 2-(2,4-dihydroxyphenyl)-3,5,7-trihydroxycromen-4-one Ligands In Vero Cells. *Indones J Trop Infect Dis*. 2019; 7(5): 105-108.
  22. Sucipto TH, Churrotin S, Setyawati H, Kotaki T, Martak F, Soegijanto S. Antiviral Activity of Copper(II) Chloride Dihydrate Against Dengue Virus Type-2 in Vero Cell. *Indones J Trop Infect Dis* [Internet]. 2017;6(4):84. Available from: <http://e-journal.unair.ac.id/index.php/IJTID/article/view/3806>
  23. Biot C, Castro W, Botté CY, Navarro M. The therapeutic potential of metal-based antimalarial agents: Implications for the mechanism of action. *Dalt Trans*. 2012;41(21):6335–49.
  24. Liu Y, Guo M. Studies on Transition Metal-Quercetin Complexes Using Electrospray Ionization Tandem Mass Spectrometry. *Molecules*. 2015;20:8583-8594
  25. Sucipto TH, Churrotin S, Setyawati H, Mulyatno KC, Amarullah IH, Ueda S, et al. INHIBITORY ACTIVITY OF COBALT ( II )– MORIN COMPLEX AGAINST THE REPLICATION OF DENGUE VIRUS TYPE 2. 2017;6(6):6–9.
  26. Raut R, Beesetti H, Tyagi P, Khanna I, Jain SK, Jeankumar VU, et al. A small molecule inhibitor of dengue virus type 2 protease inhibits the replication of all four dengue virus serotypes in cell culture. *Virol J*. 2015;12(1):1–7.



# Effect of Zinc(II)

---

## ORIGINALITY REPORT

---

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

1%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- |   |  |    |
|---|--|----|
| 1 | <a href="https://downloads.hindawi.com">downloads.hindawi.com</a><br>Internet Source   | 1% |
| 2 | Adib Afandi Abdullah, Yean Kee Lee, Sek Peng Chin, See Khai Lim et al. "Discovery of Dengue Virus Inhibitors", <i>Current Medicinal Chemistry</i> , 2020<br>Publication  | 1% |
| 3 | Chanama, S.. "Analysis of specific IgM responses in secondary dengue virus infections: levels and positive rates in comparison with primary infections", <i>Journal of Clinical Virology</i> , 200411<br>Publication | 1% |
| 4 | <a href="https://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a><br>Internet Source   | 1% |
| 5 | Asmarani, One, Much Zaenal Fanani, and Ni Nyoman Tri Puspaningsih. "Biochemical Potential of $\alpha$ -L-Arabinofuranosidase as Anti-Tuberculosis Candidate", <i>Procedia Chemistry</i> , 2016.<br>Publication       | 1% |
-

6	<p>Maher Muayad Hefdhi, Raad Hamdan Thaher. "Design of Microstrip Patch Antenna for C-band Applications", 2022 International Conference on Artificial Intelligence of Things (ICAIoT), 2022</p> <p>Publication</p>	1 %
7	<p>Zhaowenbin Zhang, Qiuxia Dai, Yu Zhang, Hui Zhuang et al. "Design of a Multifunctional Biomaterial Inspired by Ancient Chinese Medicine for Hair Regeneration in Burned Skin", ACS Applied Materials &amp; Interfaces, 2020</p> <p>Publication</p>	1 %
8	<p>Roudaina Abdel Samad, Karim Raafat, Alissar Al Khatib, Hadi Abou Chacra, nada EL DARRA. "Novel Method for The Valorization of Essential Oils From Orange By-Product", Research Square, 2021</p> <p>Publication</p>	1 %
9	<p><a href="https://content.edgar-online.com">content.edgar-online.com</a></p> <p>Internet Source</p>	1 %
10	<p><a href="http://www.db-thueringen.de">www.db-thueringen.de</a></p> <p>Internet Source</p>	1 %
11	<p><a href="http://acikerisim.sakarya.edu.tr">acikerisim.sakarya.edu.tr</a></p> <p>Internet Source</p>	1 %
12	<p><a href="http://journal.unnes.ac.id">journal.unnes.ac.id</a></p> <p>Internet Source</p>	1 %

13	<a href="http://journals.plos.org">journals.plos.org</a> Internet Source	1 %
14	<a href="http://www.nutriclub.co.id">www.nutriclub.co.id</a> Internet Source	1 %
15	<a href="http://repository.up.ac.za">repository.up.ac.za</a> Internet Source	1 %
16	Liu, Yuanzhen, and Mingquan Guo. "Studies on Transition Metal-Quercetin Complexes Using Electrospray Ionization Tandem Mass Spectrometry", <i>Molecules</i> , 2015. Publication	<1 %
17	Zhu, Y.. "Protein bonding on biodegradable poly(l-lactide-co-caprolactone) membrane for esophageal tissue engineering", <i>Biomaterials</i> , 200601 Publication	<1 %
18	<a href="http://cyberleninka.org">cyberleninka.org</a> Internet Source	<1 %
19	<a href="http://jurnal.stikesperintis.ac.id">jurnal.stikesperintis.ac.id</a> Internet Source	<1 %
20	<a href="http://repository.its.ac.id">repository.its.ac.id</a> Internet Source	<1 %
21	Puspa Wardhani, Aryati Aryati, Benediktus Yohan, Hidayat Trimarsanto et al. "Clinical and virological characteristics of dengue in Surabaya, Indonesia", <i>PLOS ONE</i> , 2017	<1 %

22

[mdpi-res.com](https://www.mdpi-res.com)

Internet Source

<1 %

23

[pdfcoffee.com](https://www.pdfcoffee.com)

Internet Source

<1 %

24

Ferial Ghaemi, Amirhassan Amiri, Mohd Yazid Bajuri, Nor Yuliana Yuhana, Massimiliano Ferrara. "Role of different types of nanomaterials against diagnosis, prevention and therapy of COVID-19", Sustainable Cities and Society, 2021

Publication

<1 %

25

Syamal Raha, Rahul Mallick, Sanjay Basak, Asim K. Duttaroy. "Is copper beneficial for COVID-19 patients?", Medical Hypotheses, 2020

Publication

<1 %

26

[www.fspublishers.org](https://www.fspublishers.org)

Internet Source

<1 %

27

[www.i-scholar.in](https://www.i-scholar.in)

Internet Source

<1 %

28

Hanlei Wang, Ran Zhang, Kun Zhang, Xuelin Chen, Yumei Zhang. "Antioxidant, Hypoglycemic and Molecular Docking Studies of Methanolic Extract, Fractions and Isolated Compounds from Aerial Parts of Cymbopogon citratus (DC.) Stapf", Molecules, 2022

<1 %



29

[tede.ufam.edu.br](http://tede.ufam.edu.br)

Internet Source

<1 %

---

30

[worldwidescience.org](http://worldwidescience.org)

Internet Source

<1 %

---

31

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

<1 %

---

32

Chu, Hone-Jay, Ta-Chien Chan, and Fang-Ju Jao. "GIS-aided planning of insecticide spraying to control dengue transmission", *International Journal of Health Geographics*, 2013.

Publication

<1 %

---

33

Mariela A. Agotegaray, Mariana Dennehy, Mónica A. Boeris, María A. Grela, Robert A. Burrow, Oscar V. Quinzani. "Therapeutic properties, SOD and catecholase mimetic activities of novel ternary copper(II) complexes of the anti-inflammatory drug Fenoprofen with imidazole and caffeine", *Polyhedron*, 2012

Publication

<1 %

---

34

Shivkrupa D. Halbandge, Supriya P. Mortale, Sankunny Mohan Karuppayil. "Biofabricated Silver Nanoparticles Synergistically Activate Amphotericin B Against Mature Biofilm Forms

<1 %

of Candida Albicans", The Open  
Nanomedicine Journal, 2017

Publication

---

35

Dengliang Li, Zhanding Cui, Guohua Li,  
Liangting Zhang et al. "Antiviral effect of  
copper chloride on feline calicivirus and  
Synergy with Ribavirin in vitro", Research  
Square, 2020

Publication

---

<1 %

36

Andrew J. Stevens, Michelle E. Gahan, Suresh  
Mahalingam, Paul A. Keller. "The Medicinal  
Chemistry of Dengue Fever", Journal of  
Medicinal Chemistry, 2009

Publication

---

<1 %

37

Raja Sundararajan, Nazeer Ahamed Haja,  
Kumar Venkatesan, Kakali Mukherjee et al.  
"Cytisus scoparius link - A natural  
antioxidant", BMC Complementary and  
Alternative Medicine, 2006

Publication

---

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

# Effect of Zinc(II)

---

GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---