

# A Review of Salmonellosis on Poultry Farms: Public Health Importance

*by* Hani Plumeriastuti

---

**Submission date:** 05-Jun-2023 09:22AM (UTC+0800)

**Submission ID:** 2108955282

**File name:** w\_of\_Salmonellosis\_on\_Poultry\_Farms\_Public\_Health\_Importance.pdf (926.85K)

**Word count:** 5731

**Character count:** 31889

## A Review of Salmonellosis on Poultry Farms: Public Health Importance

Freshindy Marissa Wibisono<sup>1</sup>, Freshinta Jellia Wibisono<sup>2</sup>, Mustofa Helmi Effendi<sup>3\*</sup>, Hani Plumeriastuti<sup>4</sup>, Akvyan Rafi Hidayatullah<sup>1</sup>, Erwan Budi Hartadi<sup>1</sup>, Eka Dian Sofiana<sup>1</sup>

<sup>1</sup>Postgraduate Student in Veterinary Public Health Study, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

<sup>2</sup>Department of Veterinary Public Health, Faculty of Veterinary Medicine, Kusuma Surabaya University, Surabaya, Indonesia

<sup>3</sup>Department of Veterinary Public Health, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

<sup>4</sup>Department of Veterinary Pathology, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

\*Corresponding author: Mustofa Helmi Effendi, Department of Veterinary Public Health, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia, Postal code: 60115. Telephone: +628175111783.

Email: [mheffendi@yahoo.com](mailto:mheffendi@yahoo.com)

### ABSTRACT

Salmonella spp. is a bacterial causative agent that can cause salmonellosis in humans and animals. Salmonellosis is categorized as an important zoonotic disease in public health. Infection can be acquired by direct contact and indirect contact with animals. Indirect transmission can occur through contact with the environment around animals or with contaminated objects around poultry farms. Salmonella in humans, which is an infectious disease, has three types of infections: typhoid fever, paratyphoid fever, and non-typhoid Salmonella (NTS). Typhoid fever and paratyphoid fever are caused by *S. typhi* and *S. enterica* paratyphi (*S. paratyphi*), in contrast, *S. typhimurium*, *S. enteritidis*, *Salmonella enterica* serovar newport (*S. newport*), and *Salmonella enterica* serovar heidelberg (*S. heidelberg*) causes non-typhoid Salmonella infection. The distribution of Salmonella is very widespread and persistent in dry environments but can persist in water for up to several months. In poultry, Salmonella can cause clinical disease or subclinical infection in asymptomatic animals which are often referred to as carriers. Infectious disease control in poultry farms always uses antibiotics. High antibiotic use can lead to increased antibiotic resistance. Raising awareness is critical to limiting the inappropriate use of antibiotics.

**Keywords:** Public health, Antibiotic resistance, Salmonella spp, Salmonellosis, Poultry farms

### Correspondence:

Mustofa Helmi Effendi

Mustofa Helmi Effendi, Department of Veterinary Public Health, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia, Postal code: 60115. Telephone: +628175111783.

Email: [mheffendi@yahoo.com](mailto:mheffendi@yahoo.com)

### INTRODUCTION

QuillBot is going to re-write your email. Start by writing or pasting here something, and then press the paraphrase button. is a bacterial agent that can cause human and animal salmonellosis (1,2). In terms of public health, salmonellosis is regarded as a significant zoonotic disease (3,4). Salmonellosis is a natural, Salmonella spp enteric disease. Acute and chronic diarrhoea disease and even death in human and animal patients are typically induced (5-7). A total of 16 million cases of inflammatory fever include salmonellosis, 1.3 billion cases of gastroenteritis and 3 million deaths from Salmonella. Worldwide (8) each year. There were 1.2 million cases of non-type salmonellosis, 19,000 in hospital, and 370 deaths last year. Apply salmonella to salmonella. Known as one of the primary causes of foodborne disease (4,9), 11% being due to animal contact (10,11). Direct contact and indirect contact with animals can be used to acquire infection. Indirect transmission may occur through interaction with the animals' environment or contaminated items at farms. Salmonella spp infected poultry. It looks good, but can contaminate the surrounding environment and animals (12). Salmonellosis is one of Salmonella spp's most common enteric pathogenic diseases. Has a high degree of morbidity and preventive issues (10,13). Poultry is an important reservoir of bacterial agents. Infected poultry can be a source of disease transmission. Pathogenic microorganisms can cause infectious disease, which is the main disease and the highest cause of death in animals and humans (14). The increasing incidence of infectious diseases causes the use of antibiotics to be the most dominant in health services (15,16).

The high use of antibiotics can lead to increased antibiotic resistance. Antibiotics as prevention are used routinely in most cases. The combination preparation of amoxicillin and colistin (60.8%) was the most widely used preparation in several farms. Farmers do not even realize the purpose of using antibiotics. The high use of antibiotics without a prescription is due to the perception of farmers that the use of antibiotics has no side effects and at a low cost as an effort to prevent disease (17). During the last two decades, there has been increasing awareness of the factors causing the incidence of antibiotic resistance in livestock as food of animal origin for health. Thus, antibiotic resistance in one country is currently a problem for all countries (18).

The need for safe and quality animal protein is a consumer demand. Incidence of Salmonella sp in ducks (19), cattle (20), broilers (21), fish (22). Pathogenic microorganisms in the food chain are transmitted to humans through a variety of foods including beef, poultry and eggs (23,24). Bacterial contamination of food can occur at all stages of production so that it can affect the quality of poultry products (25). Despite awareness of the risks of Salmonellosis from handling raw poultry, the public is generally unaware that Salmonella can also spread between live birds and humans (12). This emphasizes the need for a global initiative with a control and monitoring system to control antibiotic resistance in order to protect public health related to Salmonella spp. from poultry farms. This review article aims to describe the importance of Salmonella which grows from poultry farms in terms of public health.

## OVERVIEW OF SALMONELLA

Salmonella was initially discovered and isolated by Theobald Smith in 1855 from pigs infected with Hog cholera or Classical Swine Fever (CSF). Dr. Daniel Elmer Salmon, an American pathologist with Smith, identified the bacterial strain. The nomenclature of Salmonella is still very controversial and growing. The Centers for Control and Prevention of Diseases (CDC) currently uses the World Health Organization (WHO) recommended scheme of Salmonella (6,26). The Salmonella genus belongs to the family Enterobacteriaceae, with two species, *S. enterica* and *S. enterica*. (3,4) bongori. *S. Enterica* consists of six sub-species, namely I, S, written in Roman numerals. *enterica* subs. *enterica* subs. *enterica*, II, p. ent. *enterica* subs. *enterica* subs. Petrus; IIIa, *S. enterica* subs. *enterica* subs. 3. Arizonae; 3.24, *S. enterica* subs. *enterica* subs. 18 ptoody; IV, p. *enterica* subs. *enterica* subs. The fucks; and VI, *S. enterica* subs. *enterica* subs. indica. indica. *Enterica* subs. *salmonella* subs. *Enterica* (I) occurs primarily in mammals. In humans and warm blooded animals, about 99 per cent of salmonella infections. In comparison, Salmonella and S were the other five subspecies. Bongori are predominantly found in cold-blooded and environmentally friendly animals and seldom infect humans (6,67).

*Salmonella* spp. is a Gram-negative rod-shaped bacteria which is one of the most common infectious agents in the tropics, especially in areas with poor hygiene (23). The source of infection with *Salmonella* spp. Comes from human or animal feces or urine which contaminates drinking water and food so that it becomes a source of infection, besides fish, flies and dust can also act as intermediaries for salmonellosis (7,8,28).

*Salmonella* spp classification. Serotype dependent on 3 major antigens, i.e., somatic (O), capsule (K), and flagellum (H). Somatic antigen (O) is positioned outside the membrane of the bacterial cell and has stable heat to form the lipopolysaccharide (LPS) oligosaccharide portion of the bacterial cell. A unique salmonella serotype can express more than one O antigen. The thermally-labile H antigen is used to activate the immune response of the host and is found mainly in bacterial flagellas. Much of the spp. *Salmonella*. It has two different genes that flagellar encode. These bacteria can be diphasic (phase I and II), so they can release one protein at a time only. Immunologically responsible Phase I H antigen can be expressed in many serotypes, whereas phase II antigen does not have the same characteristics and is present in several other serotypes (9,29).

The K antigen surface is rarely found between most serotypes of *Salmonella* and heat sensitive polysaccharides mainly located on the surface of the bacterial capsule. Only Paratyphi C, Dublin and Typhi serotypes were found for the K antigen subtype virulence (VI). In literature, the term serovar, synonymous with serotype, is widely used. In general, the subspecies in the specific classification *Salmonella* serotype are omitted. Reading literature like S, for example. Typhi enterica subspecies Typhi is generally reduced to *Salmonella* ser. S, or S. Typhi. Typhi (26,50). Typhi. There were currently 2 500 serotypes, each of which has a somatic (O) antigen combination with flagellar antigens H1 and H2. More than 50 percent of these serotypes are S. S. *enterica*. *enterica*, *enterica*. It accounts for the majority of human *Salmonella* infections (30,31).

## SALMONELLOSIS

In humans, *Salmonella*, which is an infectivity, has three distinct infections: typhoid, paratyphoid and non-typhoid *Salmonella* (NTS). In human beings it is a disease. S is the cause of typhoid and paratyphus fever. S, and typhi. S. Enterica serovar Paratyphi, a fever characteristic and complications, including septicemia, immunological symptoms, leukopenia, and neurology. Typhoid and paratyphoid complications may cause death (32). In the reverse, S. Murinum typhimurum, S. enteritidis, salmonella entica serovar Newport (Newport S.), and salmonella entica serovar Heidelberg (H.), cause gastro-enteritis-limited, non-typhoidal infections of *Salmonella*, such as nausea, vomiting, diarrhoea, and bacteremia. But not (2) lethal.

There can be no distinctive clinical signs of typhoid fever and paratyphoid fever, which renders 'enteric fever' a mixture of both. Also S, too. typhus and s. typhus. Paratyphi called typhoid *Salmonella* (5,6). Two forms of *Salmonella* typhoid are stored in humans. Bacteria are spread by food or water intake. Enteric fever is typically characterised by a period of incubation of a week or more accompanied by symptoms such as headache, stomach pain and oral (or constipation) diarrhoea (6,33). Aside from fever, myalgia, bradycardia, hepatomegaly, splenomegaly and chest and stomach rashes can be encountered in infected men (6,34). *Salmonella* strains other than S. typhi and S. paratyphi are often referred to as non-typhoid *Salmonella* (NTS) and the main reservoir is animals. NTS infection is characterized by gastroenteritis in which there is an inflammatory condition in the digestive tract accompanied by symptoms such as bloodless diarrhea, vomiting, nausea, headache, stomach cramps, and myalgia. Symptoms such as hepatomegaly and splenomegaly are rare in patients infected with NTS (6,35). Gastrointestinal complications from NTS infection such as cholecystitis, pancreatitis, and appendicitis, whereas terminal ileal perforation was not associated with NTS infection (35).

Chronic carriers are characterised as bacteria that are carried through the heat for more than one year following the acute *Salmonella* infection. Since people are the *Salmonella* typhoid source, S. S and typhoid. Paratyphoid can be transmitted to human beings through common routes of transmission, such as water consumption or food contaminated by carrier faeces (33). In addition, people as NTS carriers are rare at 0.1% concentrations, since animals are the main stockpile of NTS (35). NTS is a disease that is prevalent in the entire world and is self-restricted diseases. *Salmonella* infection is the most common cause of this disease. Enteric fever, by comparison, is highly morbid and mortal and is widespread particularly in underdeveloped countries (1) due to salmonella typhoid.

## SALMONELLA TRANSMISSION FROM POULTRY

*Salmonella*'s spread is very normal and persistent in dry environments, but it can last for up to several months in water. S. S. Serobar. Enterica has a complex host / reservoir that can cause both human and animal diseases (30). In animals that are sometimes considered "carriers," *Salmonella* can cause clinical or sub-clinical infection in asymptomatic animals. Previous studies have found that chickens can suffer from subclinical infection over 22 weeks (30,36). Transports in farm and climate, in which bacteria can constantly and routinely be excreted in faeces without clinical symptoms, are of great importance to the transmission of *Salmonella* (30,37). *Salmonella*'s mode of transmission can be horizontal or vertical. The transmission of bacteria from the parent to

the infant is vertical transmission. In Salmonella infection in poultry which is caused by serovar enteritidis with a particular preference for the chicken reproductive system, vertical transmission is of great concern. In this case it happens by transovarian infection when the mother bird has a systemic infection that results in ovary infection and egg production in the oviduct. Bacteria migrating from cloaca into reproductive organs (4) contribute to serovar enteritidis gaining access to eggs. Additional evidence indicates that Salmonella may be transferred in dairy cows vertically from the foetal site in the uterus (39). Horizontal transmission takes place through aerogens and/or fecoal transmission. Salmonella is clearly transmitted horizontally by fomite, polluted drinking water, polluted feed, dirty cages, no signs of infected animals and faeces of infected livestock (38, 40). (38).

Some of the investigations of Salmonellosis outbreaks apart from those related to contact with live poultry there are also indirect contact of infected or contaminated food vendors (28). Another example of an epidemic caused by indirect contact with live poultry is foodborne disease which originates from workers who are infected by direct contact with chickens from the farm (41). Animals are the main reservoir for transmission of NTS infection. Transmission of NTS infection to humans can occur through consumption of food or water contaminated with infected animals, as well as by direct contact with infected animals or consuming infected animals. The incidence of NTS around the world is high, bacteria can be found in the environment. It is this diversity of reservoirs that creates significant difficulties for public health authorities in infection control (42).

The risk of Salmonellosis in humans after contact with live birds is well reported through large outbreaks (12). Since the last report on May 20, 2020, 368 people have been infected with Salmonella. As of now as of June 23, 2020, a total of 465 people were infected with Salmonella in 42 US states (43). According to Loharikar *et al.*, (2012), there were 4 outbreaks of Salmonellosis in the United States with the number of laboratory confirmed cases of Salmonella infection in humans from four outbreaks associated with live poultry contact, by the United States in 2009. Outbreak 1: cases infected with the Salmonella montevideo strain, United States on January 20 - December 1, 2009 (n = 96). Outbreak 2: Cases infected with the Salmonella Thompson strain, United States on February 16 - July 11, 2009 (n = 26). Outbreak 3: cases infected with Salmonella typhimurium, United States on May 21 - August 14, 2009 (n = 36). Outbreak 4: cases infected with the Salmonella johannesburg type outbreak, United States May 2009 (n = 7). The reports represent four human outbreaks of Salmonella infection associated with contact with live poultry during 2009. Poultry on farms are the main source of Salmonella spread, however birds often appear healthy despite Salmonella infection. This outbreak highlights the need to focus on strategic efforts to reduce and prevent salmonellosis in humans associated with contact with live birds. (44).

#### ANTIBIOTIC RESISTANCE

The treatment of infectious disease includes antibiotics and can improve the quality of life by reducing bacterial-related deaths. The selectivity of antibiotics against bacterial interference is able to reduce harm to patients while ensuring the target bacteria are eradicated maximally (45). In the veterinary sector, inaccurate antibiotic use can promote antibiotic resistance in different bacteria types (46, 47) including Gram-positive

bacteria (48, 49, 50, 51) and Gram-negative bacteria (52, 53, 54, 55). NTS infection does not normally need antibiotic therapy, but complications including meningitis and septicemia need antibiotic treatments such as ciprofloxacin, ceftriaxone and ampicillin (56). S-causing infections. S and typhi. Paratyphi usually have complications and need serious treatment with antibiotics such as cefixime, chloramphenicol, amoxicillin, trimethoprim / SMX, azithromycin, aztreonam, cefotaxime or ceftriaxone to avoid death (2,57). Paratyphus are usually complications and need serious treatment.

Over the years, animals have often been debated on the contribution of the cattle, which has been an important storage facility for antimicrobial resistance with public health impacts, but in particular some supporting evidence has been given that poultry farms cause human NTS disease (18). More than 3,000 people in America die every year due to antimicrobial resistance (58, 59), according to the CDC, per year more than 48 million cases of foodborne disease occur in the US where at least 70 per cent of the pathogens are immune with at least one antimicrobial. (58, 59). Antibiotics are used for the prevention, treatment and control of infectious diseases as drugs for people and animals. Excessive use of antibiotics can, however, cause antibiotic resistance. The role of cattle in emerging and propagating medically resistant antimicrobial bacteria and the proliferation of antimicrobial resistance is poorly understood among humans (60, 61). The development of antibiotic-resistant Salmonella primarily results from the use of antibiotics for feed-based animals to facilitate growth and antibiotics in veterinary medicine for the treatment of bacterial infections (62). This can establish an increased risk of the transmission, through direct contact and food intake of infected animals, of zoonotic diseases with the salmonella MDR strain from animals to humans via contaminated animal feces' food or water (6, 63).

Antibiotic resistance of Salmonella spp. isolated from broiler chicken meat in Central Surabaya Traditional Market, showed 27 out of 30 positive samples contaminated with Salmonella spp. with the results of the sensitivity test to antibiotics, 0% resistant to the antibiotics meropenem and amikacin; 3.7% resistant to the antibiotic ampicillin sulbactam; 11.1% were resistant to ofloxacin and 44.4% were resistant to nalidixic acid (21). Addressing this challenge will require efforts from various sectors to successfully control the spread and emergence of AMR in the country. Actions should include active surveillance to monitor the emergence and spread of AMR. Infection prevention and control must also be optimized to limit further spread. Raising awareness is critical to limiting the inappropriate use of antibiotics, and antibiotic stewardship programs in hospitals, outpatients, and community pharmacies, should regulate ongoing use of antimicrobials (64).

#### CONTROL AND PREVENTION OF SALMONELLOSIS

Salmonellosis is a major concern in food safety. The European Union has enacted regulations since January 2006 covering control measures implemented throughout the poultry production chain, including production levels, biosecurity measures, and prohibiting the use of antibiotics as growth promoters (65). Poultry and poultry products are frequently associated with outbreaks of Salmonellosis and therefore, are generally recognized as a major source of disease spread (66). Salmonella prevention and control can be achieved by adopting the

principles of HACCP (Hazard Analysis Critical Control Point) (67).

Hygiene and biosecurity should be part of the overall management of the farm. These steps are very important in infection control. Incoming poultry must have a high health status and must be purchased from reliable suppliers that have quality-assured breeding and hatchery facilities. Furthermore, Salmonella can be transmitted in chicken farms through vehicles, workers, clothing, footwear, equipment, water, food, garbage, insects, rodents, wild birds, pets, equipment, and many other factors. To prevent the entry of Salmonella into the farm can be done by limiting people who enter the farm, wearing protective clothing, and wearing boots that have been disinfected. In addition, workers must know basic hygienic principles, such as keeping hands and feet clean. In planning the management of the whole farm, cleaning and disinfection must be carried out regularly. The success of disinfection of chicken farms needs to be tested by taking samples on floors, walls, drinking water, eating places, and the environment, it is hoped that public awareness of the dangers of antimicrobial resistant if an antibiotic use approach is used. (68, 69, 70).

Integrated surveillance with collaboration between human health, food safety and animal health and One Health approaches and contamination strategies including livestock, retail, catering and consumers to minimize contamination and reduce transmission of Salmonella. In addition, continuous monitoring of the level of Salmonella resistance globally is very important for clinicians to support the best treatment options for Salmonellosis, especially for patients receiving antibiotic therapy (71, 72).

## CONCLUSION

Salmonella spp. has always been a major public health problem that occurs worldwide. The spread of Salmonella is very widespread and persistent in the environment, it increases the difficulty in reducing the spread of Salmonella spp. Salmonella spp. it can cause death in humans and animals. Apart from this, the emergence of antibiotic resistance in Salmonella is a major challenge in terms of effective treatment for the Salmonella infection. Limiting the use of antibiotics in feed is an effective measure to stop the spread of antibiotic resistance in poultry production.

## REFERENCES

- Hardy A. *Salmonella*: A continuing problem. *Postgrad Med J*. 2004;80(947):541-5.
- Gut AM, Vasiljevic T, Yeager T, Donkor ON. *Salmonella* infection - Prevention and treatment by antibiotics and probiotic yeasts: A review. *Microbiol (United Kingdom)*. 2018;164(11):1327-1344.
- Shinohara NKS, Barros VB de, Jimenez SMC, Machado E de CL, Dutra RAF, Lima Filho JL de. *Salmonella spp.*, importante agente patogênico veiculado em alimentos. *Cien Saude Colet*. 2008;13(5):1675-1683.
- Anamaria MP, Rafaela S, Junior CAC. Virulence Factors in *Salmonella Typhimurium*: The Sagacity of a Bacterium. *Curr Microbiol*. 2018; 76(6):762-773
- Connor BA, Schwartz E. Typhoid and Paratyphoid Fevers in the Tropics. *J Trop Med Hyg*. 2005; 5 (10) 623-628
- Eng S, Pusparajah P, Mutalib NA, Ser H, Chan K, Lee L. *Salmonella* : A review on pathogenesis, epidemiology and antibiotic resistance. *Front Life Sci*. 2015;3769.
- Barbour EK, Ayyash DB, Alturkistni W, Alyahiby A, Yaghmoor S, Iyer A, et al. Impact of sporadic reporting of poultry *Salmonella* serovars from selected developing countries. *J Infect Dev Ctries*. 2015;9(1):001-7.
- Pui CF, Wong WC, Chai LC, Tunung R, Jeyaletchumi P, Noor Hidayah MN, et al. *Salmonella*: A foodborne pathogen. *Int Food Res J*. 2011;18(2):465-73.
- Lee KM, Runyon M, Herrman TJ, Phillips R, Hsieh J. Review of *Salmonella* detection and identification methods: Aspects of rapid emergency response and food safety. *Food Control*. 2015;47:264-76.
- Anderson TC, Nguyen TA, Adams JK, Garrett NM, Bopp CA, Baker JB, et al. Multistate outbreak of human *Salmonella typhimurium* infections linked to live poultry from agricultural feed stores and mail-order hatcheries, United States 2013. *One Heal*. 2016;2:144-9.
- Hale CR, Scallan E, Cronquist AB, Dunn J, Smith K, Robinson T, et al. Estimates of enteric illness attributable to contact with animals and their environments in the United States. *Clin Infect Dis*. 2012;54(SUPPL.5).
- Behraves CB, Brinson D, Hopkins BA, Gomez TM. Backyard poultry flocks and salmonellosis: A recurring, yet preventable public health challenge. *Clin Infect Dis*. 2014;58(10):1432-8.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe R V., Widdowson MA, Roy SL, et al. Foodborne illness acquired in the United States-Major pathogens. *Emerg Infect Dis*. 2011;17(1):7-15.
- Suardana IW, Utama IH, Ayu P, Putriningsih S, Rudyanto D. Antibiotic Sensitivity Test of *Escherichia coli* O157: H7 Isolate from Chicken Feces. *Bul Vet Udayana*. 2014;6(1).
- Noor SM, Poeloenga M. Use of Antibiotics in Livestock and Their Impact on Human Health. National Workshop on Breeder Production Food Safety. 2014;56-64.
- Wiedosari E, Wahyuardani S. Case Study of Broiler Chicken Disease in Sukabumi and Bogor Districts. *J Kedokt Hewan - Indones J Vet Sci*. 2015; 9(1): 9-13.
- Niasono AB, Latif H, Purnawarman T. Antibiotic Resistance to *Escherichia coli* Bacteria Isolated from Broiler Farms in Subang Regency, West Java. *J Vet*. 2019;20(2):187-95.
- Aarestrup FM. The livestock reservoir for antimicrobial resistance: A personal view on changing patterns of risks, effects of interventions and the way forward. *Philos Trans R Soc B Biol Sci*. 2015;370(1670).
- Loisa, Lukman DW, Latif H. *Salmonella* spp. Resistance against several antibiotics on duck meat in Bogor Regency which can affect consumer health. *J Kedokt Hewan*. 2016;10(2):115-20.
- Kurniawati A, Lukman DW, Wibawan IWT. Antibiotic Resistance to *Salmonella* Isolate of Australian Feeding Cattle Imported Through the Port of Tanjung Priok, Jakarta. *J Vet*. 2016;17(3):449-56.
- Aprillian R, Rahardjo D, Koesdarto S. Evaluation of *Salmonella sp* Contamination and its Antibiotics Resistance Patterns Isolated from Broiler Meat Sold at Wet Market in Center of Surabaya. *Indones J Trop Infect Dis*. 2015;5(6):143.
- Monica WS, Mahatmi H, Besung K. Resistance pattern of *Salmonella typhi* isolated from wolf fish (*Hoplias malabaricus*) to antibiotics. *J Ilmu dan Kesehatan Hewan*. 2013;1(2):64-9.

23. Bell RL, Jarvis KG, Ottesen AR, Mcfarland MA, Brown EW. Recent and emerging innovations in *Salmonella* detection: A food and environmental perspective. *Microb Biotechnol.* 2016;9(3):279–92.
24. Ximenes E, Ku S, Hoagland L, Ladisch MR. Chapter 1 Accelerated Sample Preparation for Fast *Salmonella* Detection in Poultry Products. 2019;1918.
25. Herman L, Heyndrickx M, Grijspeerd K, Vandekerchove D, Rollier I, De Zutter L. Routes for *Salmonella* contamination of poultry meat: epidemiological study from hatchery to slaughterhouse. *Epidemiol Infect.* 2003;131(3):1169–80.
26. Popoff MY, Bockemühl J, Gheesling LL. Supplement 2001 (no. 45) to the Kauffmann-White scheme. *Res Microbiol.* 2003;154(3):173–4.
27. Brenner FW, Villar RG, Angulo FJ, Tauxe R, Swaminathan B. *Salmonella* Nomenclature. *J Clin Microbiol.* 2000;38(7):2465–2467
28. Loharikar A, Vawter S, Warren K, Deasy M, Moll M, Sandt C, et al. Outbreak of human *Salmonella typhimurium* infections linked to contact with baby poultry from a single agricultural feed store chain and mail-order hatchery, 2009. *Pediatr Infect Dis J.* 2013;32(1):8–12.
29. McQuiston JR, Fields PI, Tauxe R V., Logsdon JM. Do *Salmonella* carry spare tyres? *Trends Microbiol.* 2008;16(4):142–148.
30. Jajere SM. A review of *Salmonella enterica* with particular focus on the pathogenicity and virulence factors, host specificity and adaptation and antimicrobial resistance including multidrug resistance. *Vet World.* 2019;12(4):504–21.
31. Guibourdenche M, Roggentin P, Mikoleit M, Fields PI, Bockemühl J, Grimont PAD, et al. Supplement 2003–2007 (No. 47) to the White-Kauffmann-Le Minor scheme. *Res Microbiol.* 2010;161(1):26–29.
32. Chong A, Lee S, Yang YA, Song J. The role of typhoid toxin in *Salmonella typhi* virulence. *Yale J Biol Med.* 2017;90(2):283–90.
33. Bhan MK, Bahl R, Bhatnagar S. Typhoid and paratyphoid fever. *Lancet.* 2005; 366: 749–762.
34. Kuvandik C, Karaoglan I, Namiduru M, Baydar I. Predictive value of clinical and laboratory findings in the diagnosis of the enteric fever. *New Microbiol.* 2009;32(1):25–30.
35. Hohmann EL. Nontyphoidal Salmonellosis. *Food Saftey.* 2001;32:263–269.
36. Poppe C, Johnson RP, Forsberg CM, Irwin RJ. *Salmonella enteritidis* and other *Salmonella* in laying hens and eggs from flocks with *Salmonella* in their environment. *Can J Vet Res.* 1992;56(3):226–232.
37. Gow AG, Gow DJ, Hall EJ, Langton D, Clarke C, Papsouliotis K. Prevalence of potentially pathogenic enteric organisms in clinically healthy kittens in the UK. *J Feline Med Surg.* 2009;655–662.
38. Zamora-Sanabria R, Alvarado AM. Preharvest *Salmonella* Risk Contamination and the Control Strategies. *Curr Top Salmonella Salmonellosis.* 2017.
39. Hanson DL, Loneragan GH, Brown TR, Nisbet DJ, Hume ME, Edrington TS. Evidence supporting vertical transmission of *Salmonella* in dairy cattle. *Epidemiol Infect.* 2016;144(5):962–967.
40. Yanestria, S.M., Rahmaniar, R.P., Wibisono, F.J., Effendi, M.H. Detection of *invA* gene of *Salmonella* from milkfish (*Chanos chanos*) at Sidoarjo wet fish market, Indonesia, using polymerase chain reaction technique. *Veterinary World.* 2019; 12(1): 170–175.
41. Hedican E, Miller B, Ziemer B, Lemaster P, Jawahir S, Leano F, et al. Salmonellosis outbreak due to chicken contact leading to a foodborne outbreak associated with infected delicatessen workers. *Foodborne Pathog Dis.* 2010;7(8):995–997.
42. Dione MM, Ikumapayi UN, Saha D, Mohammed NI, Geerts S, Ieven M, et al. Clonal differences between non-typhoidal *Salmonella* (NTS) recovered from children and animals living in close contact in the Gambia. *PLoS Negl Trop Dis.* 2011;5(5):1–7.
43. CDC. US Outbreaks of Zoonotic Diseases Spread between Animals & People. Centers for Disease Control and Prevention. 2020.
44. Loharikar A, Briere E, Schwensohn C, Weninger S, Wagendorf J, Scheffel J, et al. Four Multistate Outbreaks of Human *Salmonella* Infections Associated with Live Poultry Contact, United States, 2009. *Zoonoses Public Health.* 2012;59(5):347–354.
45. Nami Y, Khosroushahi AY, Abdullah N, Haghshenas B, Barzegari A, Radiah D, et al. Probiotics or antibiotics : future challenges in medicine. *J Med Microbiol.* 2014;1–29.
46. Helmi, AM., Mukti, AT., Soegianto, A., Mahardika, K., Mastuti, I., Effendi, MH., Plumeriastuti, H. A Review of Bacterial Zoonoses and Antimicrobial Resistant (AMR) on Grouper fish (*Epinephelus sp.*). *Sys Rev Pharm 2020;11(9):79-88*
47. Witaningrum AM, Wibisono FJ, Permatasari DA, Tyasningsih W, Effendi MH, Kurniawan F. Potential Hazards of Antibiotics Resistance On *Escherichia Coli* Isolated From Cloacal Swab In Several Layer Poultry Farms, Blitar, Indonesia. *Indian Journal of Public Health Research & Development.* 2020; 11 (03): 2429–2435.
48. Effendi, M.H., Oktavianto, A and Hastutiek, P. Tetracycline Resistance Gene in Streptococcus Agalactiae Isolated from Bovine Subclinical Mastitis in Surabaya, Indonesia. *Philipp. Journal of Veterinary Medicine.* 2018; 55 (SI): 115-120.
49. Tyasningsih, W., Effendi, M. H., Budiarto, B., & Syahputra, I. R. Antibiotic Resistance to Staphylococcus aureus and Methicillin Resistant Staphylococcus aureus (MRSA) Isolated from Dairy Farms in Surabaya, Indonesia. *Indian Vet. J.* 2019; 96(11), 27-31.
50. Rahmaniar, R. P., Yunita, M. N., Effendi, M. H., & Yanestria, S. M. Encoding Gene for Methicillin Resistant Staphylococcus aureus (MRSA) Isolated from Nasal Swab of Dogs. *Indian Vet. J.* 2020; 97(02), 37-40.
51. Effendi M. H., Hisyam M. A. M., Hastutiek P., Tyasningsih W. Detection of coagulase gene in *Staphylococcus aureus* from several dairy farms in East Java, Indonesia, by polymerase chain reaction. *Vet World.* 2019; 12(1): 68-71.
52. Wibisono FJ, Sumiarto B, Untari T, Effendi MH, Permatasari DA, Witaningrum AM. The Presence of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* On Layer Chicken Farms In Blitar Area, Indonesia. *Biodiversitas.* 2020; 21 (6): 2667-2671.
53. Kristianingtyas L, Effendi, MH, Tyasningsih W, Kurniawan F. Genetic Identification of *blactx-M* Gene and *blatem* Gene on Extended Spectrum Beta Lactamase (ESBL) Producing *Escherichia Coli* from Dogs. *Indian Vet. J.* 2020; 97 (01): 17 – 21

54. Putra, A.R. Effendi, M.H. Koesdarto, S. Suwarno, S. Tyasningsih, W. and Estoepangestie, A.T. Detection of the extended spectrum  $\beta$ -lactamase produced by *Escherichia coli* from dairy cows by using the Vitek-2 method in Tulungagung regency, Indonesia. *Iraqi Journal of Veterinary Sciences*, 2020; 34 (1): 203-207.
55. Effendi, M.H., Harijani, N., Budiarto, Triningtya, N.P., Tyasningsih, W. and Plumeriastuti, H. Prevalence of Pathogenic *Escherichia Coli* Isolated from Subclinical Mastitis in East Java Province, Indonesia. *Indian Vet. J.*, 2019; 96 (03) : 22 – 25
56. Medalla F, Gu W, Mahon BE, Judd M, Folster J, Griffin PM, et al. Estimated Incidence of Antimicrobial Drug – Resistant Nontyphoidal *Salmonella* Infections, United States, 2004-2012. *Emerg Infect Dis*. 2017;23(1):2004-12.
57. Kumar P, Kumar R. Enteric Fever. *Indian J Pediatr*. 2017; 84, 227–230
58. Cosby DE, Cox NA, Harrison MA, Wilson JL, Jeff Buhr R, Fedorka-Cray PJ. *Salmonella* and antimicrobial resistance in broilers: A review. *J Appl Poult Res*. 2015;24(3):408–426.
59. Scharff RL. Health-related costs from foodborne illness in the United States. *Prod Saf Proj*. 2010;1–28.
60. Fall-Niang NK, Sambe-Ba B, Seck A, Deme SN, Wane AA, Bercion R, et al. Antimicrobial Resistance Profile of *Salmonella* Isolates in Chicken Carcasses in Dakar, Senegal. *Foodborne Pathog Dis*. 2019;16(2):130–136.
61. Muloi D, Ward MJ, Pedersen AB, Fe EM. Are Food Animals Responsible for Transfer of Antimicrobial-Resistant *Escherichia coli* or Their Resistance Determinants to Human Populations? 2018; XX(X):1–8.
62. Hyeon JY, Chon JW, Hwang IG, Kwak HS, Kim MS, Kim SK, et al. Prevalence, antibiotic resistance, and molecular characterization of *Salmonella* serovars in retail meat products. *J Food Prot*. 2011;74(1):161–166.
63. Holmberg SD, Osterholm MT, Senger KA, Cohen ML. Drug-resistant *Salmonella* from animals fed antimicrobials. *N Engl J Med*. 1984; 311(10):617-622.
64. Zowawi HM. Antimicrobial resistance in Saudi Arabia. An urgent call for an immediate action. *Saudi Med J*. 2016;37(9):935–40.
65. Vandeplas S, Dauphin RD, Beckers Y, Thonart P, The A. *Salmonella* in Chicken : Current and Developing Strategies To Reduce Contamination at Farm Level. *J Food Prot*. 2010;73(4):774–85.
66. Saravanan S, Purushothaman V, Ramasamy T, Krishna G, Sukumar K, Srinivasan P, et al. Molecular Epidemiology of Nontyphoidal *Salmonella* in Poultry and Poultry Products in India : Implications for Human Health. *Indian J Microbiol*. 2015; 55:319–326
67. World Organisation for Animal Health (OIE). Chapter 6.6. Prevention, detection, and control of *Salmonella* in poultry. *Terrestrial Animal Health Code*, 2019;Vol. I, p. 1–6.
68. Immerseel FVAN, Zutter LDE, Houf K, Pasmans F. Strategies to control *Salmonella* in the broiler production chain. *Worlds Poult Sci J*. 2009; 65(9):367–92.
69. Effendi, MH., Cicilia, R., Rahmahani, J., Tyasningsih, W. 2020. Public Awareness for Antimicrobial Resistance from *Escherichia coli* Isolated from Beef Sold on Several Wet Market in Surabaya, Indonesia. *Indian Journal of Public Health Research & Development*, 2020; 11( 9): 295-300.
70. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A., Witaningrum, A.M. CTX Gene of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* on Broilers in Blitar, Indonesia. *Sys Rev Pharm* 2020;11(7): 396-403.
71. Antunes P, Mourão J, Campos J, Peixe L. Salmonellosis: The role of poultry meat. *Clin Microbiol Infect*. 2016;22(2):110–121.
72. Widodo, A., Effendi, M.H., Khairullah, A.R. Extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* from livestock. *Sys Rev Pharm*, 2020;11(7):382-392.

# A Review of Salmonellosis on Poultry Farms: Public Health Importance

---

## ORIGINALITY REPORT

---

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1	<a href="http://www.science.gov">www.science.gov</a> Internet Source	2%
2	<a href="http://repositorio-aberto.up.pt">repositorio-aberto.up.pt</a> Internet Source	1%
3	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	1%
4	<a href="http://research.tees.ac.uk">research.tees.ac.uk</a> Internet Source	1%
5	Mamta Pandey, Emmagouni Sharath Kumar Goud. "Non-Typhoidal Salmonellosis: A Major Concern for Poultry Industry", IntechOpen, 2021 Publication	1%
6	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	1%
7	<a href="http://garuda.ristekbrin.go.id">garuda.ristekbrin.go.id</a> Internet Source	1%

---

8	<p>Sumit Sharma, Peter D. Fowler, Dhan Kumar Pant, Subir Singh, Melinda J. Wilkins.</p> <p>"Prevalence of non-typhoidal Salmonella and risk factors on poultry farms in Chitwan, Nepal", Veterinary World, 2021</p> <p>Publication</p>	1 %
9	<p>Kelechi Edward, V.I. Ibekwe, E.S. Amadi, S.I. Umeh. "ISOLATION AND CHARACTERIZATION OF MULTIDRUG RESISTANT SALMONELLA SPECIES FROM ABATTOIR WASTEWATER IN ABIA STATE, NIGERIA", Bacterial Empire, 2021</p> <p>Publication</p>	1 %
10	<p><a href="#">epdf.pub</a></p> <p>Internet Source</p>	1 %
11	<p><a href="#">rr-europe.woah.org</a></p> <p>Internet Source</p>	1 %
12	<p>Seockmo Ku, Thomas Kreke, Eduardo Ximenes, Kirk Foster, Xingya Liu, Christopher J. Gilpin, Michael R. Ladisch. " Protein particulate retention and microorganism recovery for rapid detection of ", Biotechnology Progress, 2017</p> <p>Publication</p>	<1 %
13	<p><a href="#">mts.intechopen.com</a></p> <p>Internet Source</p>	<1 %
14	<p><a href="#">docksci.com</a></p> <p>Internet Source</p>	<1 %

15	<a href="http://ebpj.e-iph.co.uk">ebpj.e-iph.co.uk</a> Internet Source	<1 %
16	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	<1 %
17	<a href="http://wwwnc.cdc.gov">wwwnc.cdc.gov</a> Internet Source	<1 %
18	<a href="http://ijpsr.com">ijpsr.com</a> Internet Source	<1 %
19	<a href="http://journal.unhas.ac.id">journal.unhas.ac.id</a> Internet Source	<1 %
20	<a href="http://sites.google.com">sites.google.com</a> Internet Source	<1 %
21	<a href="http://bsdwebstorage.blob.core.windows.net">bsdwebstorage.blob.core.windows.net</a> Internet Source	<1 %
22	<a href="http://core.ac.uk">core.ac.uk</a> Internet Source	<1 %
23	<a href="http://refubium.fu-berlin.de">refubium.fu-berlin.de</a> Internet Source	<1 %
24	<a href="http://www.britannica.com">www.britannica.com</a> Internet Source	<1 %
25	<a href="http://www.intechopen.com">www.intechopen.com</a> Internet Source	<1 %
26	Bellil Zahra, Mairi Assia, Touati Abdelaziz. "Chapter 5 Distribution and Prevalence of	<1 %

Antimicrobial Resistance of NTS Salmonella Isolated from Farm Animals and Animal Food Products in Africa", Springer Science and Business Media LLC, 2023

Publication

---

27

Dingka Song, Xuebai He, Yiming Chi, Zhao Zhang, Jing Shuai, Hui Wang, Qiuming Li, Mengze Du. "Cytotoxicity and Antimicrobial Resistance of Salmonella enterica Subspecies Isolated from Raised Reptiles in Beijing, China", Animals, 2023

<1 %

Publication

---

28

Irshad M. Sulaiman, Ying-Hsin Hsieh. "Foodborne Pathogens in Milk and Dairy Products", Elsevier BV, 2017

<1 %

Publication

---

29

Molly R. Tobin, Jesse L. Goldshear, Lance B. Price, Jay P. Graham, Jessica H. Leibler. "A Framework to Reduce Infectious Disease Risk from Urban Poultry in the United States", Public Health Reports, 2015

<1 %

Publication

---

30

Sabah A.A. Jassim, Richard G. Limoges. "Bacteriophages: Practical Applications for Nature's Biocontrol", Springer Nature, 2017

<1 %

Publication

---

31

Yue Liu, Dao-Feng Zhang, Xiujuan Zhou, Li Xu, Lida Zhang, Xianming Shi. "Comprehensive

<1 %

# Analysis Reveals Two Distinct Evolution Patterns of Salmonella Flagellin Gene Clusters", *Frontiers in Microbiology*, 2017

Publication

32

[atrium.lib.uoguelph.ca](http://atrium.lib.uoguelph.ca)

Internet Source

<1 %

33

[d.lib.msu.edu](http://d.lib.msu.edu)

Internet Source

<1 %

34

[link.springer.com](http://link.springer.com)

Internet Source

<1 %

35

[livrepository.liverpool.ac.uk](http://livrepository.liverpool.ac.uk)

Internet Source

<1 %

36

[warm.dovepress.com](http://warm.dovepress.com)

Internet Source

<1 %

37

Anagha Loharikar, Shannon Vawter, Kim Warren, Marshall Deasy et al. "Outbreak of Human Salmonella Typhimurium Infections Linked to Contact With Baby Poultry From a Single Agricultural Feed Store Chain and Mail-order Hatchery, 2009", *The Pediatric Infectious Disease Journal*, 2013

Publication

<1 %

38

*Zoonoses - Infections Affecting Humans and Animals*, 2015.

Publication

<1 %

39

[psasir.upm.edu.my](http://psasir.upm.edu.my)

Internet Source

<1 %

40

EE Effa. "Azithromycin for treating uncomplicated typhoid and paratyphoid fever (enteric fever)", Cochrane Database of Systematic Reviews Reviews, 07/19/2006

Publication

<1 %

41

S. Rubino. "Molecular characterization of ciprofloxacin-resistant *Salmonella enterica* serovar Typhi and Paratyphi A causing enteric fever in India", Journal of Antimicrobial Chemotherapy, 10/20/2006

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# A Review of Salmonellosis on Poultry Farms: Public Health Importance

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---