

Detection of Multidrug-Resistant (MDR) Escherichia coli Isolated from Raw Milk in East Java Province, Indonesia

by Hani Plumeriastuti

Submission date: 05-Jun-2023 09:11AM (UTC+0800)

Submission ID: 2108945100

File name: coli_Isolated_from_Raw_Milk_in_East_Java_Province,_Indonesia.pdf (516.33K)

Word count: 2982

Character count: 15485

Detection of Multidrug-Resistant (MDR) *Escherichia coli* Isolated from Raw Milk in East Java Province, Indonesia

Ribby Ansharieta¹, Mustofa Helmi Effendi^{2,3}, Hani Plumeriastuti⁴

¹Postgraduate Student on Veterinary Public Health Program, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia, ²Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia, ³Halal Research Center, Universitas Airlangga, Surabaya, Indonesia, ⁴Department of Veterinary Pathology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia

Abstract

The aim of this study is to figure out the profile of antibiotic resistance in *E. coli*, sourced from raw milk, obtained from five districts in East Java Province, Indonesia. A total of 250 samples were tested using the Kirby-Bauer Test method. The result revealed that the highest percentage of antibiotic resistance was Tetracycline (17.05%), followed by Streptomycin (14.2%), Trimethoprim (9.66%), Chloramphenicol (7.95%) and Aztreonam (1.7%). A total of 16 *E. coli* isolates (9.1%) were detected as MDR, and 3 *E. coli* isolates (1.7%) were suspected as presumptive ESBL. Thus, threat of multidrug resistance *E. coli* possibly sourced from the milk.

Keywords: *Escherichia coli*, multidrug resistant, raw milk, antibiotic susceptibility test.

Introduction

Contamination of milk with microorganisms is something that cannot be completely avoided, therefore microbial contamination in milk is an important point measuring its quality¹. Some microorganisms can be found in milk and other dairy products, one of which is *E. coli*. Pathogenic *E. coli* infections in humans, which originate from milk, are widely reported in several cases in developing countries².

The case of *E. coli* contamination that is resistant to various antibiotics has been widely studied. At present, a focus on public health is being developed regarding antibiotic resistance due to its use in the treatment of livestock, following the conditions of human diseases in developing countries³. Antibiotics are used as prevention and treatment of various cases of *E. coli* bacterial infections that attack livestock⁴. However, inappropriate and irrational administration of antibiotics results in the

emergence of antibiotic resistance in *E. coli* bacteria⁵.

E. coli can be a reservoir of various antibiotic resistance genes, which can be transmitted to other bacteria. Raw milk, including one of the foodstuffs of animal origin which is a potential source in the spread of multi-drug-resistant bacteria, which impacts on human health⁶. Therefore, the objective of this study was to detect the presence of multidrug resistant *E. coli* from raw milk in East Java Province, Indonesia.

Materials and Methods

Raw milks were taken from dairy farm with a total of 250 samples, located in Kediri, Probolinggo, Pasuruan, Batu and Blitar Region, East Java Province, Indonesia, in the period between August-November 2019. Raw milk samples contained in wrapped sterile plastic and were taken to the laboratory using cool box container at 4°C.

Each sample were cultured into enrichment media Brilliant Green Bile Lactose Broth (Merck, 105454) and incubated at 37°C for 18-24 hours. The positive results characterized by change of media from green colour to cloudy green and gas presence in Durham tube⁷. Then were streaked onto Eosin Methylene Blue (EMB)

Corresponding author :

Mustofa Helmi Effendi,

Halal Research Center, Universitas Airlangga,
Surabaya, Indonesia. Postal code: 60115. Tel:
+628175111783. Email: mheffendi@yahoo.com

agar (Merck, 101347) and incubated at 37 °C for 18-24 hours⁸. Colonies that showed metallic green colour (Figure. 1) was purified and continued to identification with IMViC test.

Antibiotic sensitivity testing was done using Kirby-Bauer disc diffusion assay on medium Mueller-Hinton agar (Oxoid, CM0337). Antibiotics disks used was

Tetracycline 30 µg (Oxoid, CT0054), Streptomycin 10 µg (Oxoid, CT0047), Chloramphenicol 30 µg (Oxoid, CT0013), Trimethoprim 5 µg (Oxoid, CT0057), Aztreonam 30 µg (Oxoid, CT0264) (Figure 2 and 3). Interpretation of results by measuring the diameter of the inhibitory zone formed, based on Clinical and Laboratory Standards Institutions⁹.

Table 1. Antimicrobial Resistance Profile

Location (Code)	Sample Size	Confirmed E. coli	Resistant to					Multidrug Resistant	Presumptive ESBL
			TE	S	W	C	ATM		
Kediri (K)	50	35	7	5	1	4	0	3	0
Probolinggo (A)	50	36	3	2	3	2	0	2	0
Pasuruan (G)	50	30	7	5	8	1	1	4	1
Batu (H)	50	38	4	5	1	1	2	1	2
Blitar (S)	50	37	9	8	4	6	0	6	0
Total	250	176	30	25	17	14	3	16	3
Percentage (%)	100	70.4	17.05	14.2	9.66	7.95	1.7	9.1	1.7

Info: TE: Tetracycline, S: Streptomycin, W: Trimethoprim, C: Chloramphenicol, ATM: Aztreonam

Table 2. Measurement of inhibition zone on antibiotic sensitivity test of Multidrug Resistant and Presumptive ESBL *E. coli*

No.	Sample Code	Diameter of inhibition zone of antibiotics in mm					MDR	Presumptive ESBL
		TE 3 µg	20 10 µg	W 5 µg	C 30 µg	ATM 30 µg		
1.	K-03	3 (R)	6 (R)	6 (R)	6 (R)	36 (S)	+	-
2.	K-04	3 (R)	10 (R)	30 (S)	6 (R)	33 (S)	+	-
3.	K-36	3 (R)	6 (R)	30 (S)	6 (R)	34 (S)	+	-
4.	A-26	6 (R)	6 (R)	6 (R)	6 (R)	31 (S)	+	-
5.	A-44	16 (S)	9 (R)	6 (R)	6 (R)	40 (S)	+	-
6.	G-12	6 (R)	6 (R)	6 (R)	22 (S)	30 (S)	+	-
7.	G-31	17 (S)	11 (R)	7 (R)	16 (I)	6 (R)	+	+
8.	G-35	6 (3)	6 (R)	6 (R)	23 (S)	31 (S)	+	-
9.	G-43	3 (R)	6 (R)	26 (S)	6 (R)	30 (S)	+	-
10.	H-11	6 (R)	6 (R)	6 (R)	6 (R)	30 (S)	+	-
11.	H-15	9 (R)	6 (R)	6 (R)	28 (S)	34 (S)	+	-
12.	H-28	10 (R)	6 (R)	25 (S)	7 (R)	28 (S)	+	-
13.	H-37	9 (R)	9 (R)	22 (S)	6 (R)	27 (S)	+	-
14.	H-44	7 (R)	10 (R)	30 (S)	6 (R)	29 (S)	+	-
15.	H-45	7 (R)	10 (R)	31 (S)	6 (R)	30 (S)	+	-
16.	S-25	7 (R)	16 (S)	24 (S)	7 (R)	15 (R)	+	+
17.	S-38	18 (S)	11 (R)	27 (S)	30 (S)	10 (R)	-	+

Info: R: Resistant, I: Intermediate, S: Sensitive.



Figure 1. E coli in EMB Agar




Figure 2. Isolate resistant to 4 antibiotic discs (K-03)



Figure 3. Isolate of Presumptive ESBL (G-31)

Results and Discussion

Among the total 250 raw cow milk samples collected in 24 different dairy farms, 176 samples (70.4%) were found to be positive for *E. coli* (Figure 1). The antimicrobial resistance profiles of the bacterial isolates were summarised in Table 1. *E. coli* showed resistance to antibiotics like Tetracycline (17.05%), Streptomycin (14.2%), Trimethoprim (9.66%), Chloramphenicol (7.95%) and Aztreonam (1.7%). A total of 16 multiple drug resistance (MDR) isolates were found (8.52%) with six different patterns (Figure 2). The highest MDR patterns noted was Tetracycline, Streptomycin, and

Chloramphenicol in 7 out of 16 MDR isolates (Table 2). Three presumptive ESBL isolate detected from Pasuruan and Blitar, which resistant to Aztreonam disc (Figure 3).

The result of this study showing high level for *E. coli* found in milk indicating a poor sanitation practices of farmers during milking process¹⁰. This number has the same level reported by Chey et al.¹¹ in Malaysia, stated that the highest prevalence (72.2%) of *E. coli* originated from the Eastern zone of the research location. Compared to other developing countries, namely Bangladesh, as much as 75% of milk samples studied contain *E. coli* bacteria¹².

Tetracycline has the highest antibiotic resistance because it is commonly used in the veterinary field, as well as other antibiotics used in this study such as aminoglycoside (Streptomycin), macrolide Chloramphenicol and sulphonamide (Trimethoprim)¹³. The use of broad-spectrum antibiotics such as the tetracycline and beta-lactam classes is more common in clinical mastitis cases of dairy cows in Europe, due to their effective treatment results. For respiratory and digestive tract problems, the tetracycline and aminoglycoside groups are the first choice antibiotics, while the second choice is the macrolide group and the combination of sulfonamid-trimethoprim drugs which have a significant impact on microbial activity in the rumen, and the last choice is the cephalosporin class three and four antibiotics (sulfonamide-trimethoprim drug combination which has a significant impact on microbial activity in the rumen, and the final choice is the cephalosporin class three¹⁴.

Seventeen MDR and Presumptive ESBL *E. coli* isolates that have been studied in this study, have almost the same pattern of resistance to one antibiotic, that is resistant to aminoglycoside class drugs, (Streptomycin), except for S-38 isolates which are still intermediate status. This resistance can be attributed to the effectiveness of the drug streptomycin in the treatment of dairy cows affected by mastitis in Indonesia. According to Riyanto et al.¹⁵, a good treatment for dairy cows suffering from mastitis is a combination of the Penicillin-Streptomycin antibiotic. Although in this study the level of resistance of the drug Streptomycin is quite low (14.2%), but there is a need to monitor the use of antibiotics Streptomycin in dairy cows in Indonesia.

Based on this study, Trimethoprim and Aztreonam antibiotics may still be used to treat MDR bacteria that are found because there are still several isolates of *E. coli*

MDR that are sensitive to them. Trimethoprim is usually combined with Sulfonamide which works synergistically with broad spectrum activity. In the case of urinary tract infections, Trimethoprim has been shown to be effective in killing infections caused by *E. coli* bacteria¹⁶.

While monobactam (Aztreonam) has not been defined for veterinary application¹⁸ the present of resistant isolates has the possibility of ESBL producing *E. coli* found in raw milk. The finding of ESBL Enterobacteriaceae sourced from milk, indicates the presence of environmental contamination, and the lack of environmental sanitation when milking is carried out¹⁷. *E. coli* is a bacterium that can be a reservoir of various antibiotic resistance genes, including the beta lactam antibiotic resistance gene, which makes *E. coli* capable of producing the β -lactamase enzyme^{7,18}. Cage sanitation, enclosure base, and drainage of the cage need to be considered by breeders to prevent contamination of milk by the ESBL *E. coli* presumptive bacteria.

The occurrence of antibiotic resistance is known to originate from the bacterial plasmid which is able to decrease the resistance gene and spread it to other bacteria around it. *E. coli* is a bacterium that is well known as the owner of resistance genes for various antibiotics¹⁹. Various resistance genes can accumulate in bacterial plasmids, usually in plasmid R (resistance), which is the cause of the discovery of bacterial isolates that are resistant to various antibiotics. In the R plasmid, there is an integron sequence, which is able to add various kinds of resistance coding genes to the base arrangement in the plasmid, so that a series of resistance genes is created in a bacterium^{20,21}.

E. coli MDR isolates are very common in various countries and are responsible for a series of infections with high severity and difficult to treat. In Canada, there are studies on cases of urinary tract infections due to *E. coli* bacteria, of which 60% are cases of *E. coli* infections with resistance to more than 3 classes of antibiotics. Consumption of food from raw uncooked animals, inter-regional travel habits, and contact with reservoir animals is associated with an increased risk of urinary tract infections caused by *E. coli* MDR²².

The multidrug resistance reported in the current study might be due to high and irrational antimicrobial use in individual cows to treat various diseases affecting the dairy sector. As a presence of resistance to many antimicrobial drugs has become a public health problem

according to the fact that food borne outbreaks would be difficult to treat and this findings of MDR *E. coli* in food supply establish a source for communicable resistant genes²³.

Conclusion

Multidrug resistant *E. coli* is a public health threat that can be sourced from the consumption of foods derived from animals, one of which is raw milk. From this study it was concluded that *E. coli* is prevalent in raw milk and has multidrug resistance (MDR) properties from East Java Province, Indonesia. *E. coli* found in milk is expected to increase public awareness of the dangers it poses. In this way, the application of sanitation and monitoring of antibiotic use must continue to be promoted.

Acknowledgement: The authors would like to thank the Rector of Airlangga University for providing HIBAH PENELITIAN MANDAT research fund with grant numbers; 371/UN3.14/LT/2019. This article is part of the research.

Ethical Clearance: Raw milk were used in this study, hence ethical clearance was not necessary. Raw milk samples were collected from dairy farms East Java province, Indonesia.

Source of Funding: Mandat Research Grant, Universitas Airlangga

Conflict of Interest: Nil.

References

1. Møretro T., Langsrud S. Residential Bacteria on Surfaces in the Food Industry and Their Implications for Food Safety and Quality. *Compr Rev Food Sci Food Saf.* 2017;16: 1022-1041.
2. Tanzin, T., Nazir, K.H.M.N.H., Zahan, M.N., Md. Parvej, S., Zesmin, K, Rahman, M.T. Antibiotic Resistance Profile of Bacteria Isolated from Raw Milk Samples of Cattle and Buffaloes. *J Adv Vet Anim Res.* 2016. 3(1): 62-67.
3. Robinson, T.P., Bu D.P., Carrique-Mas, J., Fèvre, E.M., Gilbert, M., Grace, D., Hay, S.I., Jiwakanon, J., Kakkar, M., Kariuki, S., Laxminarayan, R., Lubroth, J., Magnusson, U., Ngoc, P.T., Van Boeckel, T.P., and Woolhouse M.E.J. Antibiotic Resistance: Mitigation Opportunities in Livestock Sector Development. *Animal* 2016. 11(1): 1–3.

4. Landers, T.F., Cohen, B., Wittum, T.E., Larson, E.L. A Review of Antibiotic Use in Food Animals: Perspective, Policy, and Potential. *Public Health Rep.* 2012.127(1): 4–22.
5. Hinthong, W., Pumipuntu, N., Santajit, S., Kulpeanprasit, S., Buranasinsup, S., Sookkrung, N., Chaicumpa, W., Aiumurai, P., Indrawattana, N. Detection and Drug Resistance Profile of *Escherichia coli* from Subclinical Mastitis Cows and Water Supply in Dairy Farms in Saraburi Province, Thailand. *PeerJ.* 2017.13; 5: 1-16.
6. Odenthal, S., Akineden, Ö., Usleber, E. Extended-spectrum β -lactamase Producing Enterobacteriaceae in Bulk Tank Milk from German Dairy Farms. *Int J Food Microbiol.* 2016. 238: 72-78.
7. Putra, A.R.S., Effendi, M.H., Koesdarto S., Tyasningsih, W. Molecular Identification of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* Isolated from Dairy Cows in East Java Province, Indonesia. *Indian Vet. J.*, 2019. 96(10): 26-30.
8. Effendi, M.H., Harijani, N., Yanestria, S.M. Hastutiek, P. Identification of Shiga Toxin-Producing *Escherichia coli* in Raw Milk Samples from Dairy Cows in Surabaya, Indonesia. *Philipp J Vet Med.* 2018. 55 (SI): 109-114.
9. Clinical and Laboratory Standards Institute. M100 Performance Standards for Antimicrobial Susceptibility Testing. 28th ed. Pennsylvania: Clinical and Laboratory Standards Institute. 2018. p.296
10. Munera-Bedoya, O.D., Cassoli, L.D., Machado, P.F., Cerón-Muñoz, M.F. Influence of attitudes and behavior of milkers on the hygienic and sanitary quality of milk. *PLoS One* 2017. 12(9): 1-13.
11. Chey, F.Y., Abdullah, A., Ayobb. M.K. Bacteriological Quality and Safety of Raw Milk in Malaysia. *Food Microbiol.* 2004. 21: 535–541.
12. Islam, M.A., Kabir, S.M.L., Seel, S.K. Molecular Detection and Characterization of *Escherichia coli* Isolated from Raw Milk Sold in Different Markets of Bangladesh. *Bangl J Vet Med.* 2016.14(2): 271-275.
13. Hunter, P.A., Dawson, S., French, G.L., Goossens, H., Hawkey, P.M., Kuijper, E.J., Nathwani, D., Taylor, D.J., Teale, C.J., Warren, R.E., Wilcox, M.H., Woodford, N., Wulf, M.W., Piddock, L.J. Antimicrobial-Resistant Pathogens in Animals and Man: Prescribing, Practices and Policies. *J Antimicrob Chemother.* 2010. 65(Suppl.1): i3-i17.
14. Economou, V., Gousia, P. Agriculture and Food Animals as A Source of Antimicrobial-Resistant Bacteria. *Infect Drug Resist.*, 2015. 8: 49-61.
15. Riyanto, J., Sunarto, Hertanto, B.S., Cahyadi, M., Hidayah, R., Sejati, W. Milk Quality and Production of Dairy Cow Mastitis Patients Who Get Treatment of Antibiotics. *Sains Peternak.* 2017. 14(2): 30-41.
16. Al Rabiah, H., Allwood, J.W., Correa, E., Xu, Y., Goodacre, R. pH Plays a Role in the Mode of Action of Trimethoprim on *Escherichia coli*. *PLoS ONE*, 2018. 13(7): 1-20.
17. Okeke, I.N., Laxminarayan, R., Bhutta, Z.A., Duse, A.G., Jenkins, P., O'Brien, T.F., Pablos-Mendez, A., Klugman, K.P. Antimicrobial Resistance in Developing Countries. Part I: Recent Trends and Current Status. *Lancet* 2015. 5: 481-493.
18. Effendi, M.H., Harijani, N., Budiarto, Triningtya, N.P., Tyasningsih, W., Plumeriastuti, H. Prevalence of Pathogenic *Escherichia coli* Isolated from Subclinical Mastitis in East Java Province, Indonesia. *Indian Vet. J.* 2019. 96 (03): 22 - 25
19. Ramírez-Castillo, F.Y., Moreno-Flores, A.C., Avelar-González, F.J., Márquez-Díaz F., Harel, J., Guerrero-Barrera, A.L. An Evaluation of Multidrug-Resistant *Escherichia coli* Isolates in Urinary Tract Infections from Aguascalientes, Mexico: Cross-Sectional Study. *Ann Clin Microbiol Antimicrob.* 2018. 17: 1-13.
20. Nikaido, H. Multidrug Resistance in Bacteria. *Annu Rev Biochem.* 2010.78: 119–146.
21. Kristianingtyas, L., Effendi, M.H., Tyasningsih, W., and Kurniawan, F. Genetic Identification of bla_{CTX-M} Gene and bla_{TEM} Gene on Extended Spectrum Beta Lactamase (ESBL) Producing *Escherichia coli* from Dogs. *Indian Vet. J.*, 2020. 97(01): 17-21.
22. Ukah, U.V., Glass, M., Avery, B., Daignault, D, Mulvey, M.R., Reid-Smith, R.J., Parmley, E.J. Portt, A., Boerlin, P., Manges, A.R. Risk Factors for Acquisition of Multidrug-Resistant *Escherichia coli* and Development of Community-Acquired Urinary Tract Infections. *Epidemiol Infect.*, 2017. 146: 46-57.
23. Founou, L.L., Founou, R.C., Essack S.Y. Antibiotic Resistance in the Food Chain: A Developing Country-Perspective. *Front Microbiol.* 2016. 7: 1-19.

Detection of Multidrug-Resistant (MDR) Escherichia coli Isolated from Raw Milk in East Java Province, Indonesia

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

12%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | www.hindawi.com
Internet Source | 3% |
| 2 | docplayer.net
Internet Source | 2% |
| 3 | T. S. P. J. Jayaweera, J. L. P. C. Randika, H. G. C. L. Gamage, N. N. Udawatta et al. "Isolation of Escherichia coli, Klebsiella spp. and Staphylococcus spp. from Bovine Mastitic Milk in Nuwera Eliya District of Sri Lanka and their Sensitivity to Commonly Used Antibiotics", South Asian Journal of Research in Microbiology, 2018
Publication | 2% |
| 4 | pdfs.semanticscholar.org
Internet Source | 1% |
| 5 | Recep Kara, Mukadderat Gokmen, Levent Akkaya, Veli Gok. "Microbiological Quality and Salmonella spp., /Listeria monocytogenes of Spices in Turkey", Research Journal of Microbiology, 2015 | 1% |

6	e-journal.unair.ac.id Internet Source	1 %
7	Fook Yee Chye, Aminah Abdullah, Mohd Khan Ayob. "Bacteriological quality and safety of raw milk in Malaysia", Food Microbiology, 2004 Publication	1 %
8	www.frontiersin.org Internet Source	1 %
9	livrepository.liverpool.ac.uk Internet Source	1 %
10	jurnal.unimus.ac.id Internet Source	1 %
11	researchspace.ukzn.ac.za Internet Source	1 %
12	Christiana Jesumirhewe, Adriana CABAL-ROSEL, Franz ALLERBERGER, Burkhard SPRINGER, Werner RUPPITSCH. "Genetic characterization of Escherichia coli and Klebsiella spp. from Humans and Poultry in Nigeria", Microbiology Society, 2023 Publication	<1 %
13	JungHa Woo, Jae-Ho Guk, Saehah Yi, Junbum Lee, Hyocheon Song, Woo-Hyun Kim, Seongbeom Cho. "Effect of biofilm formation	<1 %

by antimicrobial-resistant gram-negative bacteria in cold storage on survival in dairy processing lines", International Journal of Food Microbiology, 2022

Publication

14

www.mdpi.com

Internet Source

<1 %

15

eprints.lib.hokudai.ac.jp

Internet Source

<1 %

16

pt.scribd.com

Internet Source

<1 %

17

Rajni Sharma, Akash Thakur, Anita Saini, Shiv Kumar Giri, Anil Kumar, Kanu Priya, Gulab Singh. "Antibiotics Stress Response of Bacteria as Mechanism of Development of Drug Resistance", American Chemical Society (ACS), 2023

Publication

<1 %

18

archive.lib.cmu.ac.th

Internet Source

<1 %

19

link.springer.com

Internet Source

<1 %

20

res.mdpi.com

Internet Source

<1 %

21

www.ijcmas.com

Internet Source

<1 %

22

www.inabj.org

Internet Source

<1 %

23

www.scielo.br

Internet Source

<1 %

24

www.vetmedmosul.com

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Detection of Multidrug-Resistant (MDR) Escherichia coli Isolated from Raw Milk in East Java Province, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
