

Effect of Cinnamon (*Cinnamomum burmannii*) Bark Oil on Pancreatic Histopathology of white Rats (*Rattus norvegicus*) Induced with Streptozotocin

by Hani Plumeriastuti

Submission date: 05-Jun-2023 09:13AM (UTC+0800)

Submission ID: 2108946865

File name: of_white_Rats_Rattus_norvegicus_Induced_with_Streptozotocin.pdf (1.4M)

Word count: 3038

Character count: 16706

Effect of Cinnamon (*Cinnamomum burmannii*) Bark Oil on Pancreatic Histopathology of white Rats (*Rattus norvegicus*) Induced with Streptozotocin

Budiarjanti¹, Reni Utami², Mustofa Helmi Effendi³, and Hani Plumeriastuti^{4*}

¹Doctoral Program on Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia, ²Undergraduate student on Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia, ³Department of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia, ⁴Department of Veterinary Pathology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya, Indonesia

Abstract

The purpose of this research was to understand the effect of the cinnamomum (*Cinnamomum burmannii*) bark oil (CBO) on the pancreatic histopathology white rat (*Rattus norvegicus*) with diabetes mellitus. The variables taken were the diameter and the number of Langerhans Island in the pancreatic of white male rats which had been induced by streptozotocin (STZ). This research using diabetic rat induced by streptozotocin intraperitoneally with single dose of 45 mg/kg BW. Samples consist of fifty rats were divided into five groups, negative control group (K-) was not induced by streptozotocin, which treatment by giving CMC Na 1%, positive control group (K+) was induced by streptozotocin without CBO, group P1 was induced by streptozotocin and gave CBO 100 mg/kgBW, group P2 was induced by streptozotocin and gave CBO 200 mg/kgBW, group P3 was induced by streptozotocin and gave CBO 400 mg/kgBW daily for 14 days period. The results showed that cinnamomum bark oil (CBO) with dose 200 mg/kgBW (P2) was significantly higher ($p < 0,05$) than group P1, group P3 and positive control group (K+). It can be concluded that CBO can be used to improve pancreatic function of STZ induced in diabetic rats

Keywords : Streptozotocin, Cinnamomum bark oil, Island of Langerhans, Diabetes mellitus, Rats.

Introduction

Treatment for DM patients is currently mostly done by administering intravenous insulin and oral antidiabetic drugs, such can cause gastrointestinal side effects, hypoglycemic reactions, allergic skin reactions, gastrointestinal disorders and ketoacidosis. Therefore, natural herbal remedies are needed from plants that do not cause side effects. Herbal alternative medicine is one form of treatment therapy that is more affordable to the public, has mild side effects and is easily obtained¹.

Cinnamon is a spice shaped like a bark and is widely used as a spice for cooking cakes and drinks. Cinnamon plant including one of the plants that are often found in the tropics with a number of diverse species. Cinnamon plants that have been studied have antidiabetic effects including *Cinnamomum zeylanicum*, *Cinnamomum burmannii*, and *Cinnamomum cassia*². *Cinnamomum burmannii* is a type of cinnamon that is mostly found in Indonesia³. Cinnamon is a plant that has the effect of hypoglycemia, where the plant contains cinnamic acid and cinnamaldehyde which can increase insulin circulation and are effective in reducing fasting blood glucose levels².

Cinnamon bark oils (CBO) contain cinnamaldehyde and eugenol compounds which have potential as antimicrobial organisms⁴ and cinnamaldehyde compounds resulting from the isolation of cinnamon oil have the potential to enzyme α -glucosidase⁵. This

Corresponding author:

Hani Plumeriastuti
Department of Veterinary Pathology, Faculty of
Veterinary Medicine, Universitas Airlangga, Surabaya,
Indonesia. Postal code: 60115. Tel: +6287711462134.
Email: hani-p@fkh.unair.ac.id

enzyme plays a role in the breakdown of carbohydrates into blood sugar, so that the essential oil content can be developed as an antidiabetic compound. Based on the description above, the effect of cinnamon essential oil in repairing pancreatic β cells as antidiabetic has not been proven. Therefore, the authors wish to conduct research on the effects of cinnamon essential oil on the histopathology of the pancreatic Langerhans island in white rats (*Rattus norvegicus*) induced by streptozotocin.

Materials and Methods

The experimental animals used were white male rats (*Rattus norvegicus*), healthy wistar strains aged 2-3 months with a body weight between 150-250 grams totaling 50 animals. The white rat (*Rattus norvegicus*) was first adapted for seven days in the Pharmacology Laboratory of the Faculty of Medicine, Airlangga University. Diabetic white mice were made by streptozotocin-induced at a dose of 45 mg / kg body weight (BW) injected intraperitoneally. This injection was carried out in all treatments except negative controls. Streptozotocin (STZ) is dissolved in a citrate buffer solution with a pH of 4.5. STZ induction was done immediately after making a single dose of 45 mg / kg BW⁶.

White rats (*Rattus norvegicus*) male Wistar strain of 50 animals divided by five groups, each group of ten. Adaptation was done for seven days with the aim to avoid stress in experimental animals. The study were conducted with five treatments including: (K-): A group of white mice that were not induced by STZ and given CMC Na 1%, (K +): A group of STZ-induced white rats at a dose of 45 mg / kgBW intraperitoneally and give 1% CMC Na, (P1): A group of STZ-induced white rats at a dose of 45 mg / kgBW intraperitoneally and treated with CBO at a dose of 100 mg / kgBW orally, (P2): STZ-induced white rats at a dose of 45 mg / kgBW intraperitoneally and treated with CBO at a dose of 200 mg / kgBW orally, (P3): A group of STZ-induced white rats at a dose of 45 mg / kgBW intraperitoneally and treated with CBO at a dose of 400 mg / kgBW orally^{7,8,9}.

White rat samples that have been made with histopathological preparations using Hematoxylin and eosin (HE) staining, followed by examination of white rat pancreatic histopathological preparations at the Veterinary Pathology Laboratory

of the Faculty of Veterinary Medicine, Airlangga University, Surabaya.

Results and Discussion

Histopathological observations of Langerhans Island white rats in group K (-) were only given CMC Na 1% solvent, K (+) was induced by streptozotocin 45 mg / kg BW, groups P1, P2 and P3 were treated with CBO, once for 14 days. The dose of CBO given is 100 mg / kg BW, 200 mg / kg BW and 400 mg / kg BW.

The data analyzed are the number of cells and the diameter of the island of Langerhans. Data obtained from observations and calculations on histological preparations in accordance with the treatment, repetition and dosage use. The results of statistical tests using the ANOVA (Analysis of Variance) test showed that there were significant differences ($p < 0.05$) between treatments followed by Duncan's test.

The results were obtained from the calculation of the number of cells and the measurement of the diameter of the island of Langerhans which was carried out using a 400x magnification microscope. the results of observations in the control and treatment groups can be seen in Table 1 and Table 2. The histopathological picture of Langerhans Island can be observed in Figure 1. The results of observations in Table 1 mean the highest number of Langerhans island cells are group P2 and the lowest in group K (+).

Analysis of Variance (ANOVA) statistical test results showed a significant difference ($p < 0.05$). Further test is the Duncan test with the results obtained by the treatment of K (-) significantly different from K (+), P1, and P3. The treatment of K (+) is significantly different from K (-), P2 and P3.

In Table 1 and Table 2 it can be seen that P1 and P3 show significant differences with P2, so to determine the correct dosage in the use of cinnamon essential oil need to compare the three treatments with a negative control group K (-). P1 and P3 statistical test results were significantly different from K (-), but not significantly different from K (+).

The results of microscopic histopathological observations can be seen in Figure 1. P2 group was diabetic group of rats that gave cinnamon essential oil

at a dose of 200 mg / kg BW orally and produced the highest number of cells and diameter compared to other treatment groups, namely 62.50 ± 21.54 and 91.96 ± 12.38 . The cell looks neatly arranged and fills the island of Langerhans. Group P3 is a group of white rats with DM patients who were given orally cinnamon essential oil at a dose of 400 mg / kg body weight or 48.40 ± 16.24 cells. The obtained diameter is 76.54 ± 16.21 . Cells appear to be rare on the island of Langerhans. The number of cells and diameter of the Langerhans island produced was higher than the treatment groups K (+) and P1. The variance analysis test gave significantly different results ($p < 0.05$) on the number and diameter of the Langerhans island K (-), P2 with the treatment groups K (+), P1, and P3.

Hyperglycemia is a condition in which the body cannot control blood glucose levels caused by disruption of insulin secretion by β cells in the pancreatic islets of Langerhans. The process of glucose metabolism is very important to produce energy sources for the survival of cells in the body. However, in a state of hyperglycemia, the body's cells cannot metabolize glucose so the body lacks an energy source. It responds to the body to search for alternative energy, namely from glycogenesis and gluconeogenesis. Both of these processes produce by-products, namely free radicals, which cause damage to cells such as pancreatic β cells^{10, 11}.

B cell death in the Langerhans island can automatically change the structure of the Langerhans island in terms of shape and size. Therefore, the size of the diameter or size of the area can be presented as an indicator of damage and repair of pancreatic organs in hyperglycemic conditions¹². B cell death is very influential on the area of the island of Langerhans, because there are 60% -80% of β cells in the total endocrine cells in the island of Langerhans. This shows that almost all of the endocrine cells contained in the island of Langerhans are β cells¹³. This study was conducted to determine the effect of CBO on increasing the number of cells and the diameter of the pancreatic Langerhans island in white rats induced by streptozotocin. The results of histopathological examination and observation of the number of cells and the diameter of the pancreatic Langerhans island showed that the positive control group (K +) obtained the lowest average compared to other treatments. This is due to this group of streptozotocin-induced white rats 45mg

/ kgBW and only given 1% Carboxymethyl Cellulose Sodium (CMC Na) therapy. CMC Na is inert, safe and does not react with other groups¹⁴.

It has been proven that cinnamaldehyde compound from cinnamon oil isolation against the α -glucosidase enzyme so that it is very potential as an α -glucosidase enzyme inhibitor. IC50 value is a number that shows the concentration of extract (ppm) that can inhibit 50% of the activity of the α -glucosidase enzyme⁵. Compounds are said to be active to inhibit the activity of the α -glucosidase enzyme¹⁵.

The enzyme α -glucosidase plays a role in the breakdown of carbohydrates into blood sugar. Inhibition of α -glucosidase in mammalian intestine can reduce blood glucose levels from metabolism of polysaccharides or oligosaccharides. Slowing the absorption of blood glucose causes a reduction in postprandial hyperglycemia to prevent chronic complications of diabetes mellitus such as retinopathy, neuropathy, neuropathy and microvascular as well as macrovascular problems¹⁶. The group that plays a role in inhibiting the enzyme α -glucosidase is the cinamoil group. The component of cinnamon oil containing cinamoil group is cinnamaldehyde¹⁷. Cinnamaldehyde is a compound that has a functional group of aldehydes and benzene ring conjugated alkenes. The study of Ping et al (2010) states that cinnamon oil has an antidiabetic effect in mice¹⁸. Components contained in cinnamon oil are cinnamaldehyde (62-75%), eucalyptol, copaene, naphthalene, eugenol, benzaldehyde. Cinnamaldehyde contained is useful as antidiabetic¹⁹.

Zhu R et al (2017) states that cinnamaldehyde shows a decreased effect of sugar on test animals through increased sugar release and improvement of insulin sensitivity in adipose tissue and muscle tissue, increases glycogen synthesis in the liver, improves pancreatic islets, slows down gastric emptying time, and improves gastric emptying kidney disorders due to diabetes and brain damage²⁰. The mean cell count and diameter of the white rat Langerhans island increased in P2 group compared to P1 and P3. This is likely due to metabolic disorders or biotransformation of the body. The aim of cinnamaldehyde metabolism is to change non-polar cinnamaldehyde into polar, so that they can be secreted by the kidneys or bile. This process can cause active

cinnamaldehyde to become inactive. However, some become more active, less active and even become toxic²¹. The speed of metabolism can increase the intensity and length of service of the cinnamaldehyde. This metabolic rate may vary in each individual. A decrease in metabolic rate will increase the intensity and prolong the service life of the cinnamaldehyde and possibly increase the toxicity of the cinnamaldehyde²².

Table 1. Mean cell counts of Langerhans island in white rats (*Rattus norvegicus*)

Treatment	Mean Cell Number ± SD
K(-)	60.27 ^b ± 7.96
K(+)	34.20 ^a ± 6.64
P1	37.05 ^a ± 11.96
P2	62.50 ^b ± 21.54
P3	48.40 ^{ab} ± 16.24

11
 Note: a, b different superscripts in the same column show significant differences (p <0.05).

Table 2. Mean diameter of Langerhans island in white rats (*Rattus norvegicus*).

Treatment	Mean Diameter (µm) ± SD
K(-)	87.79 ^b ± 3.61
K(+)	58.83 ^a ± 8.11
P1	60.91 ^a ± 17.18
P2	91.96 ^b ± 12.38
P3	76.54 ^{ab} ± 16.21

7
 Note: a, b different superscripts in the same column show significant differences (p <0.05).




Figure 1. Histopathological picture of Langerhans Island white rat with 40 0x magnification with HE staining at treatment K (-), K (+), P1, P2, P3. on Langerhans Island Cell of Pancreas.

Conclusion

After conducting research on the effect of cinnamon bark oil (CBO) at a dose of 100 mg / kg, 200 mg / kg and 400 mg / 8 BW for 14 days, it can be summarized that CBO can increase the number of cells and expand the diameter the pancreatic Langerhans island of white rat (*Rattus norvegicus*) induced by streptozotocin. Therefore, the effect of administration of CBO can be

used as an alternative medicine to increase the number of Langerhans island that produce insuline hormone.

Designation :

Budiastuti as a Doctoral candidate on Faculty of Pharmacy, Universitas Airlangga

Reni Utami as an undergraduate student on Universitas Airlangga,

Mustofa Helmi Effendi as an associate professor on Universitas Airlangga,

Hani Plumeriastuti as a researcher on Universitas Airlangga

Ethical approval : Ethical clearance was taken and passed from the Animal Care and Use Committee of the Faculty of Veterinary Medicine, Airlangga University. Certificate ethic number : 1.KE.127.07.2019.

Source of Funding : Penelitian Unggulan Terapan Perguruan Tinggi Grant, Universitas Airlangga, in fiscal year 2019

Conflict of Interest : Nil

References

1. Utami, P., Puspaningtyas, D.E. The Miracle of Herbs. *Agro Media Pustaka. Jakarta*. 2013.8 (2): 27-30.
2. Roswien, P.A. Potential of Water and Ethanol Extract of Cinnamon (*Cinnamomum burmannii*) Bark on the A-Glucosidase Enzyme Activity. *Jurnal Kedokteran YARSI*. 2015. 23 (2): 91-102.
3. Emilda. Effects of *Cinnamomum burmannii* Nees Cinnamon Compounds Against Diabetes Mellitus: Literature Review. *Jurnal Fitofarmaka Indonesia*, 2018. 5 (1): 1-15.
4. Niu, C., dan Gilbert, E.S. Colorimetric method for identifying plant essential oil components that affect biofilm formation and structure. *Appl Environ Microbiol*, 2004.70(12): 6951-6956.
5. Ngadiwiyana., Iswiyarto., Basid, N., dan Purbowatiningrum. The potential of Cinnamaldehyde from isolation of cinnamon oil as an antidiabetic compound. *Majalah Farmasi Indonesia*, 2011.22(1): 9-14.
6. Sharafeldin, K., and Rizvi, MR. Effect of traditional plant medicines (*Cinnamomum zeylanicum* and *Syzygium cumini*) on oxidative stress and insulin resistance in streptozotocin induced diabetic rats. *The Journal of Basic & Applied Zoology*. 2015. 72: 126-134
7. Kumar, S., Vasudeva, N., and Sharma, S. GC-MS analysis and screening of antidiabetic, antioxidant and hypolipidemic potential of *Cinnamomum*

17. *tamala* oil in streptozotocin induced diabetes mellitus in rats. *Cardiovascular Diabetology*, 2012.11:95.
8. Akbarzadeh, A., Norouzian, D., Mehrabi, N.R. Induction of Diabetes by streptozotocin in rats. *Indian J Clin Biochem*. 2007. 22(2): 60-64.
9. Elsner, M., Guldbakke, B., M. Tiedge, Munday, and S. Lenzen. Relative importance of transport and alkylation for pancreatic beta-cell toxicity of streptozotocin. *Diabetologia*. 2007.43:1528-1533.
10. Firdaus, R., Marliyati, S.A., dan Roosita, K. A Diabetes Mice Model Induced by Sucrose Streptozotocin for the Gestational Diabetes Mellitus Research Approach. *Jurnal MKMI*, 2016. 12 (1): 25-29.
11. Guo, X. Effect of Cinnamaldehyde on Glucose Metabolism and Vessel Function. *Medical Science Monitor*. 2017.23 : 3844-3853.
12. Aisyatussoffi, N., Abdulgani, N. Effect of Giving Extract of Cork Fish (*Channa striata*) on the Histological Structure of Pancreas and Blood Glucose Levels of Mice (*Mus musculus*) Hyperglycemic. *Jurnal Sains dan Seni POMITS*. 2013.2: 10-14
13. Mohamed, NA., and Nassier, OA. Effects Of Cinnamon (*Cinnamomum Cassiae*) and /or Barley (*Hordeum Vulgare L*) On Some Physiological Parameters In Streptozotocin Diabetic Rats. *Egypt. J. Exp. Biol. (Zool.)*, 2013.9(1): 133 -139
14. Ananda, B.S. Potential of Onion (*Allium cepa L.*) Ethanol Extract Against the Area of Langerhans Island in White Rat (*Rattus norvegicus*) Induced by Alloxan. (Thesis). Fakultas Kedokteran Hewan. Universitas Airlangga. 2018.
15. Szkudelski, T. The Mechanism of Alloxan and Streptozotocin Action in B Cells of The Rat Pancreas. *Physiological Research*, 2001.50(6): 537-546.
16. Shibano, M., Kakutani, K., Taniguchi, M., Yasuda, M., and Baba, K. Antioxidant Constituents in the Dayflower (*Commelina communis L.*) and their α -Glucosidase Inhibitory Activity. *J. Nat Med*. 2008. 62 : 349-353.
17. Adisakwattana, S., N., Chavasiri, Deesamer, S., and Yibchok-anun, S. Structure Activity Relationships

- of trans-Cinnamic Acid Derivates on Alpha Glukosidase Inhibition. *Bioorg. Medical Chemical Lett.* 2007.14: 2893-2896.
18. Ping, H., Guijun, Z., dan Guixing, R. Antidiabetic Effects of Cinnamon Oil in Diabetic KK-Ay Mice. *Food and Chemical Toxycology.* 2010.48(1). 2344-2349.
 19. Plumeriastuti H, Budiastuti, Effendi MH, Budiarto. Identification of bioactive compound of the essential oils of *Cinnamomum burmannii* from several areas in Indonesia by gas chromatography–mass spectrometry method for antidiabetic potential. *Natl J Physiol Pharm Pharmacol*, 2019;9(4):279-283.
 20. Zhu, R. Review Cinnamaldehyde In Diabetes : A Review of Pharmacology, Pharmacokinetics and Safety. *Pharmacological Research.* 2017.122:78-89.
 21. Mardjono, M. *Pharmacology and Therapy.* Jakarta. Universitas Indonesia Press. 2007. pp 8-17.
 22. Inoue, T., Murakami, N., Ayabe, T., Oto, Y., Nishino, I., Goto, Y., Koga, Y., and Sakuta, R. Pyruvate Improved Insulin Secretion Status in a Mitochondrial Diabetes Mellitus Patient. *Jurnal Clinics Endocrinol Metabolism.* 2016.101(5): 1924-1926.

Effect of Cinnamon (*Cinnamomum burmannii*) Bark Oil on Pancreatic Histopathology of white Rats (*Rattus norvegicus*) Induced with Streptozotocin

ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

14%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

repository.tma.uz

Internet Source

3%

2

Salmon Charles PT Siahaan, Budi Santoso, - Widjiati. "Effectiveness of *Moringa oleifera* Leaves on TNF- α Expression, Insulin Levels, Glucose Levels and Follicle Count in *Rattus norvegicus* PCOS Model", *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 2022

Publication

2%

3

worldwidescience.org

Internet Source

1%

4

S M Sinaga, M Batubara, M H Pulungan, H Cintya, H Dhermawan. "Study of antiinflammation activity in ethanol extract from coriander leaf (*coriandrum sativum* L.) induced by carrageenan in male rats", *Journal of Physics: Conference Series*, 2021

Publication

1%

5	www.hindawi.com Internet Source	1 %
6	pinpdf.com Internet Source	1 %
7	repo.unand.ac.id Internet Source	1 %
8	e-journal.unair.ac.id Internet Source	1 %
9	journal.uad.ac.id Internet Source	1 %
10	cyberleninka.org Internet Source	1 %
11	download.atlantis-press.com Internet Source	1 %
12	www.sysrevpharm.org Internet Source	1 %
13	assets.researchsquare.com Internet Source	1 %
14	sciendo.com Internet Source	1 %
15	doc-pak.undip.ac.id Internet Source	<1 %
16	D A Susbandi, F Swastawati, S Suharto. " Flavor characteristics and quality of mackerel	<1 %

() "otak-otak" affected by various liquid smoke addition methods ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

17

Yan Zhao, Xinli Zhang, Jiezhong Chen, Chao Lin, Renfu Shao, Chunxia Yan, Chen Chen. "Hexarelin Protects Rodent Pancreatic B-Cells Function from Cytotoxic Effects of Streptozotocin Involving Mitochondrial Signalling Pathways In Vivo and In Vitro", PLOS ONE, 2016

Publication

<1 %

18

e-journal.undikma.ac.id

Internet Source

<1 %

19

Manna, P.. "Protective role of arjunolic acid in response to streptozotocin-induced type-I diabetes via the mitochondrial dependent and independent pathways", Toxicology, 20090304

Publication

<1 %

20

www.ajol.info

Internet Source

<1 %

21

www.bjorl.org

Internet Source

<1 %

22

www.wvj.science-line.com

Internet Source

<1 %

23

pure.rug.nl

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Effect of Cinnamon (*Cinnamomum burmannii*) Bark Oil on Pancreatic Histopathology of white Rats (*Rattus norvegicus*) Induced with Streptozotocin

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
