

OVOZOA

Journal of Animal Reproduction

The addition of vitamin C in tris-egg yolk extender maintained Sapera goat semen quality in 5° C storage <i>Jolly Rifqi Pahlevy, Hermin Ratnani, Faisal Fikri, Tjuk Imam Restiadi, Amung Logam Saputro, Bodhi Agustono</i>	1 – 8
Correlation of parity and age to services per conception, conception rate, and gestation length in Holstein Friesian cross cows <i>Norma Amalia Ulfah, Abdul Samik, Sri Hidanah, Mas'ud Hariadi, Tri Wahyu Suprayogi</i>	9 – 14
Moringa oleifera leaf extract restored the diameter and epithelium thickness of the seminiferous tubules of rat (<i>Rattus norvegicus</i>) injected with gentamicin <i>Meidy Mardatillah, Wurlina Wurlina, Ira Sari Yudaniayanti, Hani Plumeriastuti, Hardany Primarizky, Iwan Sahrial Hamid</i>	15 – 21
pH and ferning score of cervical mucus related to gestation and calf gender of Holstein Friesian crossbreed cows <i>Muhammad Billy Ferdiansyah, Iwan Sahrial Hamid, Herry Agus Hermadi, Abdul Samik, Tatik Hernawati</i>	22 – 26
Pre-freezing at 10 cm above liquid nitrogen surface for eight minutes resulted the best Sapera goat semen quality <i>Zehansyah Algifari Pasha, Bodhi Agustono, Amung Logam Saputro</i>	27 – 33
Virgin coconut oil protected the diameter and the epithelium thickness of seminiferous tubules of mice (<i>Mus musculus</i>) from oral ethanol induction <i>Vishantheni S Ravi, Soeharsono Soeharsono, Hani Plumeriastuti, Kadek Rachmawati</i>	34 – 40
Uterine torsion in Simmental crossbreed cow <i>Sri Rahayu</i>	41 – 48



Ovozoa: Journal of Animal Reproduction by Unair is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.



Publisher: Department of Veterinary Reproduction Faculty of Veterinary Medicine Universitas Airlangga
in collaboration with the Indonesian Association of Department of Veterinary Reproduction
and the Indonesian Animal Reproduction Association
Available at: <https://e-journal.unair.ac.id/OVZ/index>

This journal is supported by



EDITOR IN CHIEF

[Prof. Dr. Imam Mustofa, drh., M.Kes,](#)

EDITORS

[suzanita Utama M.Phil., PhD., drh,](#)
[Dr. Erma Safitri, drh. M.Si,](#)
[Dr. Tita Damayanti Lestari, drh. M.Sc,](#)
[Dr. Rimayanti drh. M.Kes,](#)
[Dr. Sri Mulyati, drh., M.Kes,](#)

FOCUS AND SCOPE

Ovozoa Journal of Animal Reproduction (Ovozoa J Anim Reprod) published original articles, review articles, and case reports in Indonesian or English, in the scope of veterinary reproduction and animal reproductive biotechnology from researchers, lecturers, students, and practitioners around Indonesia and worldwide.

The main objective of the Ovozoa J Anim Reprod is to disseminate the results of scientific researches in the field of animal reproduction.

Ovozoa J Anim Reprod invites manuscripts in the areas:

- ♣ Development of reproductive diagnostic techniques
- ♣ Development of reproductive technologies
- ♣ Infectious and non-infectious reproductive diseases
- ♣ Comparative pathology of reproductive diseases
- ♣ Reproductive endocrinology
- ♣ Infertility
- ♣ Environmental and stress effects on reproduction
- ♣ Better understanding of the in vivo fertilization process
- ♣ Better understanding of embryonic development
- ♣ Artificial insemination and embryo transfer
- ♣ In vitro fertilisation and embryo culture
- ♣ Sex selection of sperm and embryo
- ♣ Semen and embryo preservation
- ♣ Reproductive cloning
- ♣ Stem cell in reproduction
- ♣ Development of assisted reproductive technologies
- ♣ Molecular biology, genetics, proteomics, metabolomics application in reproductive biology

Authors have to prepare the manuscripts according to OVOZOA Journal of Animal Reproduction guidelines. Manuscript and statement of originality and copyright release form should be submitted via online submission through journal website: <https://drive.google.com/file/d/1PvnbOI6kQPm6NXNIV5Z04s0-ej4N5Iq4/view>.

JOURNAL HISTORY

OVOZOA journal is an Online Journal (e-journal), first published (Volume 1) in 2012 on the page <http://journal.unair.ac.id/OVOZOA@ovoza-media-123.html>. This journal was issued based on the Decree of the Dean of the Faculty of Veterinary Medicine, Universitas Airlangga, as a scientific communication media that contained the scientific work of lecturers and students of the Faculty of Veterinary Medicine, Universitas Airlangga in the field of Animal Reproduction. OVOZOA was published under the e-Journal Universitas Airlangga: <http://journal.unair.ac.id/>.

Based on the decision of Asosiasi Departemen Reproduksi Veteriner Indonesia (ADERVI) or the Indonesian Association of Department of Veterinary Reproduction (IADVR) meeting on December 11th, 2019 in Surabaya, the publication of the OVOZOA e-journal starting in 2020 in collaboration with ADERVI. The Memorandum of Agreement between the Dean of the Faculty of Veterinary Medicine, Universitas Airlangga and the Chairperson of ADERVI, Number (FKH unair): 1445/UN3.1.6/KS/2020 and Number (ADERVI): 04/ADERVI/V/2020 signed on May 4th, 2020. Since March 27th, 2021 OVOZOA Journal of Animal Reproduction has also collaborated with the Indonesian Animal Reproduction Association (Asosiasi Reproduksi Hewan Indonesia, ARHI). Starting from Volume 9 – 2020, OVOZOA completes its name as the OVOZOA Journal of Animal Reproduction. It contains the publication of scientific work of lecturers, researchers, and practitioners of ADERVI members in the field of Veterinary Reproduction and Animal Reproduction Biotechnology. Ovozoa Journal of Animal Reproduction is published at <https://e-journal.unair.ac.id/OVZ/index> periodically three times a year: April, August, and December.

SECRETARIAT OF OVOZOA JOURNAL ANIMAL REPRODUCTION

Faculty of Veterinary Medicine Universitas Airlangga, Kampus C, Mulyorejo, Surabaya 60115

Phone: +62-31-5992785, 5993016, Email: ovoza@journal.unair.ac.id, Website: <https://e-journal.unair.ac.id/OVZ/index>

ARTICLE PROCESSING CHARGES

Author Contributions

Article Processing Charges for each accepted article is IDR 450,000. There is no other charges. For the year 2022 article processing charges will not be levied, after a year the charges will be applied as written above.



This journal is supported by

OVOZOA

Journal of Animal

LIST OF CONTENT

- The addition of vitamin C in tris-egg yolk extender maintained Sapera goat semen quality in 5° C storage**
Jolly Rifqi Pahlevy, Hermin Ratnani, Faisal Fikri, Tjuk Imam Restiadi, Amung Logam Saputro, Bodhi Agustono 1 – 8
- Correlation of parity and age to services per conception, conception rate, and gestation length in Holstein Friesian cross cows**
Norma Amalia Ulfah, Abdul Samik, Sri Hidanah, Mas'ud Hariadi, Tri Wahyu Suprayogi 9 – 14
- Moringa oleifera leaf extract restored the diameter and epithelium thickness of the seminiferous tubules of rat (*Rattus norvegicus*) injected with gentamicin**
Meidy Mardatillah, Wurlina Wurlina, Ira Sari Yudaniayanti, Hani Plumeriastuti, Hardany Primarizky, Iwan Sahrial Hamid 15 – 21
- pH and ferning score of cervical mucus related to gestation and calf gender of Holstein Friesian crossbreed cows**
Muhammad Billy Ferdiansyah, Iwan Sahrial Hamid, Herry Agus Hermadi, Abdul Samik, Tatik Hernawati 22 – 26
- Pre-freezing at 10 cm above liquid nitrogen surface for eight minutes resulted the best Sapera goat semen quality**
Zehansyah Algifari Pasha, Bodhi Agustono, Amung Logam Saputro 27 – 33
- Virgin coconut oil protected the diameter and the epithelium thickness of seminiferous tubules of mice (*Mus musculus*) from oral ethanol induction**
Vishantheni S Ravi, Soeharsono Soeharsono, Hani Plumeriastuti, Kadek Rachmawati 34 – 40
- Uterine torsion in Simmental crossbreed cow**
Sri Rahayu 41 – 48

**Publisher: Division of Veterinary Reproduction Faculty of Veterinary Medicine Universitas Airlangga
in collaboration with the Indonesian Association of Department of Veterinary Reproduction
and the Indonesian Animal Reproduction Association
Available at <https://e-journal.unair.ac.id/OVZ/index>**



Ovozoa: Journal of Animal Reproduction by Unair is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.



This journal is supported by



Original article

***Moringa oleifera* leaf extract restored the diameter and epithelium thickness of the seminiferous tubules of rat (*Rattus norvegicus*) injected with gentamicin**

Meidy Mardatillah^{1*}, Wurlina Wurlina², Ira Sari Yudaniayanti³, Hani Plumeriastuti⁴, Hardany Primarizky³, Iwan Sahrial Hamid⁵

¹ Pet and Wild Animal Veterinary Professional Interest Group,
Dr. Ir. H. Soekarno Street, Mulyorejo, Surabaya, East Java Province, Indonesia

² Division of Veterinary Reproduction, ³ Division of Veterinary Clinic,

⁴ Division of Veterinary Pathology, ⁵ Division of Veterinary Basic Medicine,
Faculty of Veterinary Medicine, Universitas Airlangga

* Corresponding author, e-mail: meidymardatillah15@gmail.com

Open access under CC BY – SA license, DOI: 10.20473/ovz.v11i1.2022.15-21
Received October 29 2021, Revised January 14 2022, Accepted March 11 2022
Published online April 11 2022

How to cite this article: Mardatillah M, Wurlina W, Yudaniayanti IS, Plumeriastuti H, Primarizky H, Hamid IS. 2022. *Moringa oleifera* leaf extract restored the diameter and thickness of the seminiferous tubules of rat (*Rattus norvegicus*) injected with gentamicin. *Ovozoa : Journal of Animal Reproduction* 11: 15-21.

ABSTRACT

This research aims to determine the effect of *Moringa oleifera* leaf extract (MLE) on the diameter and epithelium thickness of the seminiferous tubules of rats induced with gentamicin. This study used 25 male rats (*Rattus norvegicus*) aged 8-12 weeks with 150-200 grams body weight. Negative control group (C-) was injected with aquadest and given 0.5% Na-CMC orally, positive control group (C+) was induced with gentamicin injection of 5 mg/kg BW/day and given 0.5% Na-CMC orally, while groups T1, T2, and T3 were induced with gentamicin at the same dose and treated orally with MLE of 200, 316, 500 mg/kg BW/day. Analysis of variance showed significant differences and further analysis using Duncan Multiple Range Test showed difference in each group. The results showed that treatment with MLE of 500 mg/kg BW/day improved the diameter and epithelium thickness of the seminiferous tubules of rats injected with gentamicin. It could be concluded that MLE restored the diameter and epithelium thickness of the seminiferous tubules of rats induced with gentamicin.

Keywords: gentamicin, *Moringa oleifera* leaves, rat (*Rattus norvegicus*), seminiferous tubules

INTRODUCTION

Gentamicin is an effective aminoglycoside antibiotic widely used to treat life-threatening Gram-negative bacteria infections (Edeogu *et al.*, 2020), including mycobacterium infections, septicemia, complications of urinary tract infections, endocarditis, peritonitis, and others in humans and animals (Chaves and Tadi, 2021). So far, it is known that gentamicin had toxic effects, including nephrotoxic effects

(Saleh *et al.*, 2016) and neurotoxic (Rezaei *et al.*, 2018) as well as toxic effects on male reproductive organs (Aly, 2019).

Male fertility is driven by spermatogenic stem cells, which renew themselves and give rise to differentiated spermatogonia. The spermatogonial transition is accompanied by a shift in gene expression, and changes in metabolism. The process consisted of the upregulation of genes involved in mitochondrial function, biogenesis, oxidative phosphorylation,

and glycolysis (Lord and Nixon, 2020). Spermatogenesis was a complex process of proliferation and maturation of male germ cells from diploid spermatogonia, through meiosis, into mature haploid spermatozoa. The process involved dynamic interactions between the developing germ cell and its supporting Sertoli cells. The gonadal tissues, with their high abundance of unsaturated fatty acids, high rates of cell division, and various testicular enzymes produce were particularly susceptible to overexpression of reactive oxygen species (ROS). The testis had developed a sophisticated array of antioxidant systems with enzymes and scavenging free radicals (Guerriero *et al.*, 2014). Gentamicin was an aminoglycoside-class antibiotic that could increase ROS and decrease antioxidant reserves (Rahayu *et al.*, 2019). Aminoglycosides passed through the gram-negative membrane in an oxygen-dependent active transport (Chaves and Tadi, 2021). ROS exposure can be neutralized using antioxidants (He *et al.*, 2017).

Moringa oleifera had many health benefits (Gopalakrishnan *et al.*, 2016). One of the most prominent contents of the *Moringa oleifera* was antioxidants, especially in the leaves. *Moringa oleifera* leaves contained antioxidants such as flavonoids, tannins, steroids and triterpenoids, alkaloids, and saponins based on phytochemical tests. *Moringa oleifera* leaves were rich in minerals, vitamins, and phytochemicals, and antioxidants (Falowo *et al.*, 2018).

Study on the effect of *Moringa oleifera* leaf extract (MLE) on male animal fertility has not been reported. Therefore, this study aimed to determine the effectiveness of MLE on the diameter and epithelium thickness of the seminiferous tubules of the testes of white rats (*Rattus norvegicus*) induced by gentamicin.

MATERIALS AND METHODS

Ethics commission approval

This study used 25 heads of white male rats (*Rattus norvegicus*) Wistar strain aged ten weeks with an average body weight of 200 grams. This research procedure has been approved by the Animal Care and Use Committee of the Faculty of Veterinary

Medicine Universitas Airlangga No. 1.KE.013.01.2020.

Moringa leaf extraction

Moringa oleifera leaves were dried in a dark room at room temperature (27°C) for one week. Then *Moringa* leaves were pulverized using a powder making machine (Primtech, PDK-2000 N, Indonesia). *Moringa* leaf powder (150 grams) was put in an erlenmeyer and added with 96% ethanol. Maceration was carried out for 3 x 24 hours and stirred everyday. Macerate was filtered and dried in a rotary evaporator (Rotavapor Buchi RE301 Malaysia) at 40°C 50 rpm until all the solvent and water content of the *Moringa* leaf were evaporated, leaving the *Moringa* leaf extract (MLE). MLE was resuspended in 0.5% sodium carboxymethyl cellulose (Na-CMC) for the treatment of the experimental animals (Das *et al.*, 2012).

Determination of the dosage of ingredients

The dose of gentamicin was determined based on previous study (Khaki, 2009), which was 5 mg/kg BW/day for 14 days. Determination of the dose of MLE referred to the research of Ogunsola *et al.* (2017), with the minimum dose of 200 mg/kg BW/day and the maximum dose of 500 mg/kg BW/day. The dose between the minimum and maximum doses was determined by multiplying the dosing interval with the lowest dose, thereby a dose of 316 mg/kg BW/day was obtained.

Experimental animals

White rats (*Rattus norvegicus*) were placed in individual cages, where husks were added as a bedding to the cage. Feed pellets and drinking water were given in the morning and evening. During the study, the health of rats was observed concurrently with feeding. Rats were considered healthy if there was no sign of weight loss or changes in body weight of not more than 10% of the sample criteria, the color and condition of the fur were clean and not shedding, and the mouse movement was active, the color of the ears and extremities was red and not pale.

Treatment of rats

Rats were adapted in their respective cages for 7 days. After the adaptation, the rats in group C+, T1, T2, and T3 were injected (sc) with Gentamicin 5 mg/kg BW/day for 14 days, while rats in group C- were injected with 0.5 ml aquadest (sc). At day-15 the rats in group T1, T2, and T3 were treated with 0.5 ml MLE of respectively 200, 316, and 500 mg/kg BW/day, while the rats in group C- and C+ were treated with 0.5 ml of 0.5% Na-CMC. Treatment was given orally for 14 days.

Measurement of variables

After the completion of treatment, all rats were sacrificed by cervical dislocation and dissected for testicle collection. The testes were immersed in 10% formalin for 24 hours, and prepared for Hematoxylin Eosin (HE) stained histological slides (Machmudia et al., 2021). The research variables were examined using a light microscope with a magnification of 100x and photographed using Optilab. Diameter and epithelium thickness of the seminiferous tubules were measured using Image Raster Program.

Diameter of seminiferous tubules

The diameter of the seminiferous tubules was measured from the farthest distance from two opposite points that pass through the midpoint, then a diameter perpendicular to the first diameter that have to pass the midpoint of the tubule was measured, then two perpendicular diameters between the previous two diameters were measured. The four measurements obtained were then averaged (Wurlina et al., 2021).

Epithelium thickness of seminiferous tubules

The epithelium thickness of the seminiferous tubules was measured by drawing a line from the closest distance at the boundary between the layers of spermatogonia cells to the lumen surface of each seminiferous tubule. Epithelial thickness measurement was repeated four times at different places in each seminiferous tubule. The results of epithelial thickness measurements in five seminiferous tubules of each preparation were averaged (Wurlina et al., 2021).

Data analysis

The data on diameter and epithelium thickness were analyzed using one-way Analysis of Variance and Duncan's Multiple Range Test. The statistical analysis was conducted at a 95% level of confidence by using Statistical Product and Service Solutions (SPSS) version 20 International Business Machines (IBM) Corporation (Armonk, New York, USA).

RESULTS

Subcutaneous injection of 0.5 ml of gentamicin 5 mg/kg BW/day for 14 days without MLE administration (C+ group) caused a decrease ($p < 0.05$) in diameter and epithelium thickness of the seminiferous tubules compared to those of the normal rats (C- group). The diameter and the epithelium thickness of the seminiferous tubule in the T3 group (rats injected with 0.5 ml of gentamicin 5 mg/kg BW/day for 14 days and treated with 500 mg/kg BW/day MLE orally for the next 14 days) were higher ($p < 0.05$) than those of the C+ group, and not significantly different ($p > 0.05$) than those of C- group (Table 1).

Table 1 Diameter and epithelium thickness (μm , means \pm SD) of the seminiferous tubules of white rats (*Rattus norvegicus*) injected with gentamicin and treated with Moringa leaf extract

	diameter (μm)	epithelium thickness (μm)
C-	381.80 \pm 22.48 ^a	99.01 \pm 4.63 ^a
C+	328.33 \pm 15.06 ^c	67.77 \pm 8.35 ^c
T1	338.34 \pm 20.40 ^{bc}	75.69 \pm 9.31 ^c
T2	352.58 \pm 28.92 ^{ab}	85.67 \pm 8.31 ^b
T3	361.48 \pm 18.76 ^{ab}	92.00 \pm 4.62 ^{ab}

Different superscripts in the same column show significant differences ($p < 0.05$); C-: rats were injected with 0.5 ml of distilled water subcutaneously for 14 days, followed by administration of 0.5% Na-CMC orally 0.5 ml/day/head for the next 14 days; C+, T1, T2, and T3 groups: rats were injected with 0.5 ml gentamicin (5 mg/kg BW/day for 14 days), followed by the oral administration of 0.5% Na-CMC, 0.5 ml MLE (in 0.5% Na-CMC) (200,

316, and 500 mg/kg BW/day respectively for the next 14 days.

DISCUSSION

Gentamicin could increase the number of reactive oxygen species (ROS) in the testes (El-maddawy, 2014). ROS was one of the free radicals that trigger lipid peroxidation, including in the testicular tissue (Su *et al.*, 2019). The functional development of the testes was played by spermatogenic cells, Sertoli as nursing cells, Leydig cells as testosterone producers, and other testicular stromal tissues. Like other cells, cells in testicular tissue were also rich in polyunsaturated fatty acids (PUFA) (Van Tran *et al.*, 2017). High levels of ROS could damage cells in testicular tissue (Aly and Hassan, 2018).

ROS were short-lived and highly reactive molecules. Low levels of ROS played a role in activating cell survival signaling pathways. However, ROS activated cell death signaling pathways from apoptosis and necroptosis at high levels. ROS activated mitochondria, death receptors, and endoplasmic reticulum apoptotic pathways (Redza-Dutordoir *et al.*, 2016). ROS were free radicals derived from oxygen during normal cellular metabolism by nature.

ROS played a vital role in physiological processes and signaling pathways associated with male fertility. ROS acted as molecular mediators of signal transduction pathways at physiological concentrations involved in regulating the hypothalamic-pituitary-gonadal axis, spermatogenesis, and steroidogenesis. In contrast, oxidative stress occurred when ROS concentrations are higher than the body's antioxidant. Therefore, maintaining a balanced redox state was essential for normal male reproductive function (Baskaran *et al.*, 2021).

Oxidative stress caused by gentamicin injection could directly or indirectly affect testicular tissue. The direct effect of oxidative stress on functional cells and supporting cells of testicular tissue would cause tissue damage, resulting in a decrease in the diameter and thickness of the seminiferous tubules epithelium. Exposure to substances that were thought to increase ROS resulted in a decrease in the number of spermatogenic cells, and

Leydig cells, and Sertoli cells (Pratama *et al.*, 2021). The death of some Sertoli cells reduced the capacity of nursing functions for the process of spermatogenesis (Griswold, 2018). The death of some Leydig cells also decreased spermatogenesis. Leydig cells were cells that produce testosterone, which was essential for spermatogenesis. Testosterone was transported by androgen-binding proteins to Sertoli cells, binding to androgen receptors to regulate spermatogenesis (Ge *et al.*, 2021). Treatment which was also thought to trigger an increase in ROS in rats, caused a decrease in the diameter of the seminiferous tubules and the epithelium thickness of the seminiferous tubules (Wurlina *et al.*, 2021).

Apoptosis and necrosis caused by ROS exposure (Redza-Dutordoir *et al.*, 2016) due to gentamicin injection would be followed by a decrease in epithelium thickness of the seminiferous tubules, which in turn caused shrinkage and a decrease in the diameter of the seminiferous tubules. Gentamicin induction could result in testicular atrophy, degeneration, and loss of spermatogenesis. This was because gentamicin induced oxidative stress associated with spermatogenic disorders (Aly and Hassan, 2018). Sperm viability, sperm motility, total sperm count, testicular weight, seminal vesicle weight, epididymal weight, percentage of total apoptotic cells, and seminiferous tubule diameter were significantly decreased with gentamicin induction (Khaki, 2015). Animals treated with gentamicin decreased testosterone and luteinizing hormone (LH) levels, sperm count, viability, motility and increase sperm abnormalities (Yahya *et al.*, 2019).

Oxidative stress could also occur in cells on the hypothalamic-pituitary-testis axis, thereby inhibiting the secretion of the follicle-stimulating hormone (FSH) and LH and ultimately reducing the synthesis of the hormone testosterone (Al-Damegh, 2014). FSH produced from the pituitary played a role in supporting spermatogonia's structural and metabolic development into mature spermatids through receptors bound to the Sertoli cell membrane. FSH also played a vital role in determining the number of Sertoli cells to regulate the production capacity of spermatogenesis (Oduwole *et al.*, 2018).

The seminiferous tubules could make up to 90 percent of the testes. The walls of the tubules were composed of stratified germinal epithelium that contained spermatogenic cells and Sertoli cells, nutritive cells that have a mature sperm head embedded in them. Sertoli cells helped facilitating the process of spermiogenesis and thus the production of viable sperm. Sertoli cells also secreted many vital molecules, including androgen binding protein (ABP), Inhibin B, and Activin. This secretion facilitated spermatogenesis directly or indirectly through a hormonal negative feedback system. Sertoli cells also responded to pituitary hormones such as FSH to initiate the process of spermatogenesis, complementing adjacent spermatogonia (Wong and Khan, 2021). If the seminiferous tubule tissue was damaged, it could affect the process of spermatogenesis, the inhibition of cell division to differentiate, affecting the number of spermatogenic cells and the quality of the spermatozoa produced.

Damage to the seminiferous tubules of the testes after gentamicin injection was the result of ROS causing oxidative stress. Moringa leaves contain antioxidants, such as flavonoids, polyphenols (Pizzino *et al.*, 2017), and quercetin (Dong *et al.*, 2019). Moringa leaves are also known to contain vitamin C (Dafaalla *et al.*, 2017) and vitamin E (Fejér *et al.*, 2019). Flavonoids acted as antioxidants in preventing male reproductive system (Ye *et al.*, 2020). The flavonoids in Moringa leaves were antioxidants that prevented oxidative stress, increased testosterone secretion (Ogunsola *et al.*, 2017) and reduced testicular disorders due to ROS exposure (Kathun and Varma, 2017). Administration of quercetin prevented the histopathological changes of testicular tissue (seminiferous tubule diameter, epithelial height) exposed to ROS from bisphenol A (Jahan *et al.*, 2020). Vitamin C in MLE functioned as an antioxidant or as a free radical scavenger (Vergara-Jimenez *et al.*, 2017). Vitamin C was able to improve the epithelium thickness of the seminiferous tubules. The administration of vitamin C could increase the previously reduced number of spermatogenic cells, increasing the epithelium thickness of the seminiferous tubules, which eventually affected the weight of

the testes (Vijayprasad *et al.*, 2014). Administration of vitamin E in the form of α -tocopherol maintained the number of spermatogenic and Leydig cells in white rats (*Rattus norvegicus*) exposed to ROS from 2,3,7,8 Tetrachlorodibenzo-p-dioxin (Machmudia *et al.*, 2021).

Antioxidant flavonoids and polyphenols, quercetin, vitamin C, and vitamin E in *Moringa oleifera* leaves prevented oxidative stress, thereby restoring the health of cells involved in spermatogenesis after the exposure to ROS from gentamicin. The study at this stage was limited to knowing the potential of MLE to restore the diameter and epithelium thickness of the seminiferous tubules in rats receiving gentamicin injection.

CONCLUSION

The administration of MLE restored the diameter and epithelium thickness of the seminiferous tubules of white male rats (*Rattus norvegicus*) induced with gentamicin. The MLE dose of 500mg/kg BW/day was the most effective in restoring the diameter and epithelium thickness of the seminiferous tubules. Further research is needed to determine the effect of MLE on semen quality in male animals exposed to ROS.

REFERENCES

- Al-Damegh MA. 2014. Stress-Induced Changes in Testosterone Secretion in Male Rats: Role of Oxidative Stress and Modulation by Antioxidants. *J Anim Sci.* 4: 70-8.
- Aly HAA. 2019. Testicular toxicity of gentamicin in adult rats: Ameliorative effect of lycopene. *Hum Exp Toxicol.* 38: 1302-13.
- Aly HAA, Hassan MH. 2018. Potential Testicular Toxicity of Gentamicin in adults Rats. *Biochem Biophys Res Commun.* 497: 362-7.
- Baskaran S, Finelli R, Agarwal A, Henkel R. 2021. Reactive oxygen species in male reproduction: A boon or a bane? *Andrologia* 53: 1-12.
- Chaves BJ, Tadi P. 2021. Gentamicin. In: *StatPearls [Internet]*. Treasure Island (FL):

- StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK557550/>. 9 December 2021.
- Dafaalla MM, Hassan AW, Idris OF, Abdoun S, Modawe GA, Kabbashi AS. 2016. Effect of ethanol extract of *Moringa oleifera* leaves on fertility hormone and sperm quality of Male albino rats. *World J Pharm Res.* 5: 1-11.
- Das AK, Rajkumar V, Verma AK, Swarup D. 2012. *Moringa oleifera* leaves Extract: A Natural Antioxidant for Retarding Lipid Peroxidation in Cooked Goat Meat Patties. *J Food Sci Tech.* 47: 585-91.
- Dong X, Meng-Jiaou H, Yan-Qiu W, Yuan-Lu C. 2019. Antioxidant activities of quercetin and its complexes for medical application. *Molecules* 24: 1123.
- Edeogu CO, Kalu ME, Famurewa AC, Asogwa NT, Onyeji GN, Ikpemo KO. 2020. Nephroprotective Effect of *Moringa oleifera* Seed Oil on Gentamicin-Induced Nephrotoxicity in Rats: Biochemical Evaluation of Antioxidant, Anti-inflammatory, and Antiapoptotic Pathways. *J Am Coll Nutr.* 39: 307-15.
- Falowo AB, Mukumbo FE, Idamokoro EM, Lorenzo JM, Afolayan AJ, Muchenje V. 2018. Multi-functional application of *Moringa oleifera* Lam. in nutrition and animal food products: A review. *Food Res Int.* 106: 317-34.
- Fejér J, Kron I, Pellizzeri V, Plúchtová M, Eliašová A, Campone L, Gervasi T, Bartolomeo G, Cicero N, Babejová A, Konečná M, Sedlák V, Poráčová J, Grušová D. 2019. First Report on Evaluation of Basic Nutritional and Antioxidant Properties of *Moringa Oleifera* Lam. from Caribbean Island of Saint Lucia. *Plants (Basel)* 8: 537.
- Ge RS, Li X, Wang Y. 2021. Leydig Cell and Spermatogenesis. *Adv Exp Med Biol.* 1288:111-29.
- Gopalakrishnan L, Doriya K, Kumar DS. 2016. *Moringa oleifera*: A review on nutritive importance and its medicinal application. *Food Sci Hum Wellness* 5: 49-56.
- Griswold MD. 2018. 50 years of spermatogenesis: Sertoli cells and their interactions with germ cells. *Biol Reprod.* 99: 87-100.
- Guerrero G, Trocchia S, Abdel-Gawad FK, Ciarcia G. 2014. Roles of reactive oxygen species in the spermatogenesis regulation. *Front Endocrinol (Lausanne).* 5: 56.
- He L, He T, Farrar S, Ji L, Liu T, Ma X. 2017. Antioxidants Maintain Cellular Redox Homeostasis by Elimination of Reactive Oxygen Species. *Cell Physiol Biochem.* 44: 532-53.
- Jahan S, Ain QU, Ullah H. 2016. Therapeutic effects of quercetin against bisphenol A induced testicular damage in male Sprague Dawley rats. *Syst Biol Reprod Med.* 62: 114-24.
- Kathun S, Varma MC. 2017. Role of *Moringa oleifera* Leaf Extract on Silk Dye Waste Effluent Induced Histopathotoxicity on Liver and Testis of Swiss Albino Male Mice *Mus musculus*. *IOSR J Pharm.* 7: 1-7.
- Khaki A. 2015. Assessment on the adverse effects of Aminoglycosides and Flouroquinolone on sperm parameters and male reproductive tissue: A systematic review. *Iran J Reprod Med.* 13: 125-34.
- Khaki A, Khaki AA, Iraj S, Bazi P, Imani SA, Kachabi H. 2009. Comparative Study of Aminoglycosides (Gentamicin & Streptomycin) and Fluoroquinolone (Ofloxacin) Antibiotics on Testis Tissue in Rats: Light and Transmission Electron Microscopy Study. *Pak J Med Sci.* 25: 624-9.
- Lord T, Nixon B. 2020. Metabolic Changes Accompanying Spermatogonial Stem Cell Differentiation. *Dev Cell.* 52: 399-411.
- Machmudia A, Eliyani H, Widjiati W, Wurlina W. 2021. Efek pemberian α -tokoferol terhadap jumlah sel spermatogenik dan sel Leydig pada tikus putih (*Rattus norvegicus*) yang dipapar 2,3,7,8 Tetrachlorodibenzo-p-dioxin. *Ovozoa* 10:74-79.
- Oda SS, El-Maddawy ZKh. 2012. Protective effect of vitamin E and selenium combination on deltamethrin-induced reproductive toxicity in male rats. *Exp Toxicol Pathol.* 64: 813-9.
- Oduwole OO, Peltoketo H, Huhtaniemi IT. 2018. Role of Follicle-Stimulating

- Hormone in Spermatogenesis. *Front Endocrinol (Lausanne)* 9: 763.
- Ogunsola OA, Owalabi JO, Fabiyi OS, Nwobi NL, Faluyi B, Akinbola AS. 2017. Moringa plant parts consumption had effects on reproductive functions in male and female rat models. *IOSR J Dent Med Sci.* 16: 82-6.
- Pizzino G, Irrera N, Cucinotta M, Pallio G, Mannino F, Arcoraci V, Squadrito F, Altavilla D, Bitto A. 2017. Oxidative Stress: Harms and Benefits for Human Health. *Oxid Med Cell Longev.* 2017: 8416763.
- Pratama API, Susilowati S, Maslachah L, Ratnani H, Suprayogi TW. 2021. The effect of watermelon (*Citrullus lanatus*) rind ethanolic extract on the number of leydig, sertoli, and spermatogenic cells of rat (*Rattus novergicus*) exposed to heat. *Ovozoa* 10: 7-11.
- Rahayu I, Usman E, Reza M. 2019. Effect of vitamin C on testosterone level, sperm count and sperm morphology in gentamicin-induced Wistar rats. *Int J Res Med Sci.* 7: 451-6.
- Redza-Dutordoir M, Averill-Bates DA. 2016. Activation of apoptosis signalling pathways by reactive oxygen species. *Biochem Biophys Acta* 1863: 2977-92.
- Rezaei NJ, Bazzazi AM, Naseri Alavi SA. 2018. Neurotoxicity of the antibiotics: A comprehensive study. *Neurol India* 66:1732-40.
- Saleh P, Abbasalizadeh S, Rezaeian S, Naghavi-Behzad M, Piri R, Pourfeizi HH. 2016. Gentamicin-mediated ototoxicity and nephrotoxicity: A clinical trial study. *Niger Med J.* 57: 347-52.
- Su LJ, Zhang JH, Gomez H, Murugan R, Hong X, Xu D, Jiang F, Peng ZY. 2019. Reactive Oxygen Species-Induced Lipid Peroxidation in Apoptosis, Autophagy, and Ferroptosis. *Oxid Med Cell Longev.* 2019: 5080843.
- Van Tran L, Malla BA, Kumar S, Tyagi AK. 2017. Polyunsaturated Fatty Acids in Male Ruminant Reproduction - A Review. *Asian-Australas J Anim Sci.* 30: 622-37.
- Vergara-Jimenez M, Almatrafi MM, Fernandez ML. 2017. Bioactive Components in Moringa Oleifera Leaves Protect against Chronic Disease. *Antioxidants (Basel)* 6: 91.
- Vijayprasad S, Bb G, Bb N. 2014. Effect of vitamin C on male fertility in rats subjected to forced swimming stress. *J Clin Diagn Res.* 8: 5-8.
- Wong WJ, Khan YS. 2021. Histology, Sertoli Cell. In: *StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing.* <https://www.ncbi.nlm.nih.gov/books/NBK560631/>. 27 December 2021.
- Wurlina W, Mustofa I, Meles DK, Mulyati S, Putri DKSC, Suwasanti N. 2021. Administration of the α -tocopherol for repairing testicle histological damage in rats exposed to dioxin. *Thai J Vet Med.* 51: 293-301.
- Yahya K, Hassan AH, Nadhem H. 2019. Evaluation the Effect of Gentamicin on Fertility of Male Rats & Probable Protective Role of Lipoic Acid. *Indian J Public Health Res Dev.* 10: 1230
- Ye RJ, Yang JM, Hai DM, Liu N, Ma L, Lan XB, Niu JG, Zheng P, Yu JQ. 2020. Interplay between male reproductive system dysfunction and the therapeutic effect of flavonoids. *Fitoterapia* 147: 104756.

Author Index

Volume 11 Number 1, April 2022

Abdul Samik	9, 22	Meidy Mardatillah	15
Amung Logam Saputro	1, 27	Muhammad Billy Ferdiansyah	22
Bodhi Agustono	1, 27	Norma Amalia Ulfah	9
Faisal Fikri	1	Soeharsono Soeharsono	34
Hani Plumeriastuti	15, 34	Sri Hidanah	9
Hardany Primarizky	15	Sri Rahayu	41
Hermin Ratnani	1	Tatik Hernawati	22
Herry Agus Hermadi	22	Tjuk Imam Restiadi	1
Ira Sari Yudaniayanti	15	Tri Wahyu Suprayogi	9
Iwan Sahrial Hamid	15, 22	Vishantheni S Ravi	34
Jolly Rifqi Pahlevy	1	Wurlina Wurlina	15
Kadek Rachmawati	34	Zehansyah Algifari Pasha	27
Mas'ud Hariadi	9		



This journal is supported by

Subject Index

Volume 11 Number 1, April 2022

Abnormality	1-8, 27-33	Moringa leaves	15-21
Age	9-14	Motility	1-8, 27-33
Alcoholism	34-40	Nutrition	1-8
Alive calf	41-49	Parity	9-14
Alkaline pH	22-26	Poverty reduction	27-33
Calf gender	22-26	Prevent undernutrition	27-33
Cervical mucus	22-26	Rat (<i>rattus norvegicus</i>)	15-21
Conception rate	9-14	Reproductive health	34-40
Dystocia	41-49	Seminiferous tubule	15-21
Estrus, gestation	22-26	Service per conception	9-14
Financial income	1-8	Simmental crossbreed dam	41-49
Gentamicin	15-21	Spermatozoa	34-40
Gestation length	9-14	Testicle histologic	34-40
Healthy dam	41-49	Uterine torsion	41-49
Male fertility	34-40	Viability	1-8, 27-33



This journal is supported by



ACKNOWLEDGEMENT

The Editors would like to thank and express appreciation to all peer reviewers who have reviewed the manuscripts for publication of OVOZOA Journal of Animal Reproduction Vol. 11 No. 1, April 2022

Prof. Dr. María De Lourdes Arruzazabala.

Centre of Natural Products, National Centre for Scientific Research, Cuba

Prof. Voster Muchenje, M.Sc., B.Sc (Hons), PgDhet.

University of Fort Hare, South Africa

Dr. Xiaoying Zhang.

Northwest A&F University, China

Prof. Mulyoto Pangestu, Dvm., M.Sc., Ph.D.

Monash Clinical School Monash University, Australia

Prof. P Imrat DVM., Ph.D.

Utrecht University, Netherlands

drh. Erif Maha Nugraha Setyawan, M.Sc., Ph.D.

Gadjah Mada University

Prof. Dr. Wurlina Drh., M.S.

Airlangga University

Dr. Sc. Agr. Siti Rasad, Ir., M.S.

Padjadjaran University

Prof. Dr. Drh. Tjok Gde Oka Pemayun, M.S.

Udayana University



This journal is supported by



Author Guidelines

OVOZOA Journal of Animal Reproduction is a journal that contains scientific articles on animal reproduction and animal reproductive biotechnology published by the Department of Veterinary Reproduction of Airlangga University together with the Association of the Indonesian Veterinary Reproduction Department (ADERVI) 3 times per year on April, August and December. OVOZOA received manuscripts in the form of original research articles, review articles and case reports in Indonesian and English. Manuscripts received must be original, current and have never been published or are being planned to be published in other scientific journals.

Manuscripts must be submitted online through the Open Journal System (OJS) in Word format. The entire text is typed in Times New Roman 12pt double spaced, using line numbers. The title is written with a Sentence case capitalization (bold, 14pt, centered). The full length of the manuscripts is a maximum of 15 pages of A4 paper with a top margin of 2.2 cm, other margins of 2 cm, and consecutive page numbers. Italicize only for species names or terms that have not been standardized as Indonesian. Define abbreviations upon first appearance in the text. Do not use non-standard abbreviations unless they appear at least three times in the text. Keep abbreviations to a minimum. Avoid unnecessary duplication of text.

The first page contains titles in Indonesian and English, followed by full names of all authors without titles and initials (bold, centered), followed by the name and complete address of the respective institution (marked with numeric superscripts) and e-mail of corresponding authors (marked with *superscript).

The second page forward contain **ABSTRACT** in English and Indonesian, followed by **INTRODUCTION, MATERIALS AND METHODS, RESULTS, DISCUSSION, CONCLUSIONS, APPROVAL OF ETHICAL COMMISSION, ACKNOWLEDGEMENTS** and **REFERENCES**.

Title should be concise but informative, as far as possible in no more than 12 words.

Abstract written in Indonesian and English, do not exceed 250 words, containing elements of background, material and methods, results and conclusions.

Keywords maximum of 6 (six) words or phrases, written after the abstract in each languages, alphabetically ordered. As far as possible avoid using keywords from the title.

Introduction should be brief, containing elements of background, problems, objectives and reference sources that support.

Research materials (materials and equipment) do not need to be mentioned separately, but rather integrated in the method used, complete with the brand and catalog number if applicable. Do not include common supplies, such as test tubes, pipette tips, beakers, etc. or standard laboratory equipment.

Method must be concise but sufficiently detailed (with reference or modification) so research can be repeated by other researchers.

Results are displayed in a concise but clear narrative with/without tables or figures.

Tables are made without vertical lines (use only lines at the top and bottom of the table as well as for separating heading from the main table), with table title placed before the table, numbered in Arabic numerals (**Table X**), and have to be referred in the text. The description of the table is placed after the table; it must be concise but clear enough so that the table separately can be understood without referring to the text. The table along with the title and description are placed after the References.

Figures presented are only those that support the findings of the study, and not restatement of data from tables in the form of figures. When resulted data in the form of figure is more informative, interesting or significant, presentation of data in table form is not required. Figure title is placed after the figure, numbered in Arabic numerals (**Figure Y**), and have to be referred in the text. The description of the figure is

placed under the title of the figure; it must be concise but clear enough so that the figure separately can be understood without referring to the text. Image (in JPEG format) is sent in separate file. The title and description of the figure are placed after the References.

Discussion contains explanation of what are found related to **the importance of your study** and how it may be able to answer the research question, comparison of findings (internally, between research data, and externally, compared with findings from other studies) and cause-effect analysis.

Conclusion does not only repeat the results of the study, but summarize the findings into a narrative that impacts on the development of science and/or practitioners in the field of veterinary reproduction.

Approval of Ethical Commission have to be stated (number and institution) if the manuscript is constructed based on a research using live animals.

Acknowledgements are delivered to the research funders (state the name, number and recipient name of the grant, if applicable), and to those who have helped carry out the research.

References are sorted alphabetically based on the author's last name. The *titles of journals* should be *abbreviated* according to the style *used* by each journal. Thirty (30)% of the references have to be from publications in the past 10 years.

Journals

Nakamura K, Kusama K, Ideta A, Imakawa K, Hori M. 2020. IFNT-independent Effects of Intrauterine Extracellular Vesicles (EVs) in Cattle. *Reproduction* 159: 503-11.

Windeyer MC, Gamsjager L. 2019. Vaccinating Calves in the Face of Maternal Antibodies:

Challenges and Opportunities. *Review Vet Clin North Am Food Anim Pract.* 35: 557-73.

Books

Gordon I. 2003. *Laboratory Production of Cattle Embryos*. 2nd Ed. CABI Publishing. UK

Chapter in books

Parkinson TJ, Vermunt Jos J, Noakes DE. 2019. Maternal Dystocia: Causes and Treatment. In: Noakes DE, Parkinson TJ, England GCW (Ed). *Veterinary Reproduction and Obstetrics*. 10th Ed. Elsevier, UK. 236-49.

Thesis/Dissertation

Utama S. 2012. *Monitoring and Assessment of Nuclear Transfer Pregnancies using Maternal*

Pregnancy Recognition Proteins. PhD. Thesis. Monash University. Melbourne, Australia.

Internet

Akhir N. 2020. Title of article from the internet. <http://www.link>. access date.

Review article

Ovazoa received review articles on topics included in the scope of the journal which are of current interest. Reviews are invited by the editor. The review must be a comprehensive analysis and perspective on the state of the field and where it is heading. Reviews will be subject to the same peer review process as is applied for original papers. The manuscript is arranged in the same way as the original article with an unstructured abstract (maximum of 250 words). The number of references is limited to 50-70, with 75% of them have to be from publications in the past 10 years. The total length should not exceed 20 pages (A4 paper, double spaced 12pt Times New Roman, top margin of 2.2 cm, other margins of 2 cm).

Case reports:

Ovozoa receives quality and interesting case reports with topics included in the scope of this journal. All reports are peer reviewed as is applied for original articles. All instructions are the same as for the original articles.

PEER REVIEW PROCESS

Manuscript reviews for publications in the OVOZOA Journal of Animal Reproduction (hereinafter referred to as OVOZOA) are carried out by peer reviewers in a double-blind review. The chief editor handles all correspondence with the corresponding author and the reviewer, and leads the final decision on whether the manuscript is recommended to be accepted, rejected, or needs to be returned to the corresponding author for revision. The corresponding author will automatically receive a confirmation email after successfully submitting the manuscript through OJS.

The chief editor and editors will evaluate the manuscripts submitted at the prequalification stage based on the format suitability and substance qualifications. If there is a format discrepancy, the manuscript will be sent back to the corresponding author to be rewritten under the OVOZOA format. However, if there is a discrepancy in substance qualifications (not included in OVOZOA focus and scope), the manuscript is rejected without further review. The decision will change the status of the manuscript in the OJS. The chief editor will notify the corresponding author about the decision.

Manuscripts that pass the prequalification will be evaluated by two or three reviewers determined by the chief editor and editors. The chief editor will send the manuscript to reviewers via email and OJS. The reviewers answer whether they would like to review the manuscript directly via email or through OJS. The reviewers check the manuscript based on predetermined criteria and returns it with recommendations to the chief editor via email or OJS. If one reviewer recommends being accepted and another reviewer recommends being rejected, then the chief editor will ask the third reviewer or three editors to decide on the acceptance or rejection of the manuscript.

The assessment results from the peer reviewers will be consolidated by two editors who match their expertise to make a decision.

1. Accepted without revisions

The decision to accept the manuscript will change the status of the manuscript in the OJS. The chief editor will notify the corresponding author by e-mail if the manuscript is declared accepted and send the payment bill. At the same time, the manuscript was passed on to the editors for editing and layout. If payment has been made by the corresponding author, and, the editing and administration process has been completed, the chief editor will send a proof to the corresponding author via email for final correction and approval. Proof which has been approved by the corresponding author is determined to be published online in what volume and number (month and year edition).

2. Accepted with minor revisions, or major revisions, or must be rewritten.

The decision to revise the manuscript will change the status of the manuscript in the OJS. Manuscripts that require revision will be returned to the corresponding author via email, and the corresponding author must resubmit the revised manuscript through the OJS within the specified time period. Next, the chief editor sends the revised manuscript from the corresponding author to the editors to check whether the manuscript has been revised according to the reviewers' suggestions. Editors provide recommendations to the chief editor that the manuscript was accepted, or must be returned to the corresponding author to be revised again, or rejected.

3. Rejected

Manuscripts are rejected if based on consolidated comments it is concluded that

- 3.1. the corresponding author did not revise the manuscript as suggested by the reviewers and did not respond to the suggestion
- 3.2. based on the assessment of reviewers stated that
 1. originality and novelty in manuscripts for the advancement of science are considered inadequate,
 2. data is incomplete to prove the research hypothesis or the methodology used is incorrect,
 3. inconsistencies between objectives, research methods, results, and conclusions.

If the manuscript is rejected, the corresponding author will be notified by the chief editor via email and through the OJS with a statement of reasons for the refusal.

Deadlines

1. Prequalification to determine the suitability of the format, as well as compliance with the focus and scope is for a maximum of two days
2. The review process by the peer reviewer is no more than 14 days
3. Editors make consolidated comments in no later than 21 days
4. Revisions by the corresponding author:
 - a. minor revisions: no more than 14 days
 - b. major revisions (must be rewritten): a maximum of 30 days.

Manuscripts that exceed the revision deadline will be canceled. When necessary, the corresponding author can request an extension of time to the chief editor before the revision period ends.

5. Editors assess and determine the revision decision no later than seven days.
6. The editing and layout process is a maximum of seven days.
7. Approval of proofreading of the manuscript final version from the corresponding author: a maximum of four days.

The time interval from the date the OJS manuscript submits until the decision of being rejected or accepted for publication varies, depending on the time required for the review and revision process. Decisions at the prequalification stage will be accepted by the corresponding author within two days after submitting through the OJS. Manuscripts that are accepted without revision will receive a notification no later than 37 days after the manuscript submission through OJS. Whereas, manuscripts with minor revisions, the acceptance or rejection will be decided in no later than 58 days, and for manuscripts with major revision in no later than 74 days after the manuscript submission through OJS. The time from submitting the manuscript to publishing is approximately four months.

PUBLICATION FREQUENCY

Ovozoa Journal of Animal Reproduction (Ovozoa J AnimReprod) is published periodically three times a year (April, August, and December) by the Department of Veterinary Reproduction Faculty of Veterinary Medicine Universitas Airlangga in collaboration with the Indonesian Association of Department of Veterinary Reproduction (IADVR).

OPEN ACCESS POLICY

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

PUBLICATION ETHICS

The Ovozoa Journal publishes scientific articles in the form of original research articles, review articles, and case reports on animal reproduction and animal reproductive biotechnology reviewed by Ovozoa Editorial Team. Published scientific articles expected to meet standards of academic excellence. It is necessary to equate stakeholders' views in an

ethical standard to avoid problems in the future. The publication ethics is needed as a reference for Ovozoa Journal policies in publications, including the authors, editors (reviewers), reviewers (peer reviewers), and publisher.

Author

The author is required to submit the original work (no plagiarism) and not partially or fully published in other journals until the Ovozoa Journal provides an answer to the eligibility of the article. Each name in the article must have a contribution to the research and the writing of scientific work, such as developing the concept, creating the design, collecting data, analyzing data, interpreting data, and making a conclusion. To those who play an essential role in the research but have no function as an author (example: funders), their names can be embedded in acknowledgments. The corresponding author is responsible for the research and writing of the manuscript.

Authors are required to cite the literature that affects the articles. The articles must follow the review process by reviewers and peer reviewers. Manuscripts that submit to the Ovozoa Journal are not being sent to other scientific journals, are not being assessed by a journal, or have never is published or are being planned to be published in other scientific journals. If a small part of the data has been published, the source must be given written thanks in the acknowledgment of the article. If the data is reprocessed from a source, permission from the author is needed. Approval of the Ethical Commission must be stated (number and institution) if the manuscript is constructed based on research using live animals. If the author discovers and is aware of the article's mistakes or errors, they must notify the chief editor to withdraw, clarify, and correct the article. If necessary, the author must be willing to apologize for the mistakes. Requests to reduce, add, or change the author's composition for articles, must be approved by the chief editor and prepared before the article is published.

Reviewer

The reviewer plays a role in providing input and assisting the editors in making policies on the articles reviewed in addition to helping authors improve the quality of their articles. The reviewer should inform the editors about the appropriateness and ability to review the articles sent to them. All articles which are undergoing an editing process must be kept confidential. The editing process should be carried out objectively by giving a reason, and not criticizing the author personally. The reviewer must have no personal interest in the research conducted by the author or the research funder. If the article is not suitable enough to be published, the article's confidentiality must still be maintained, so others will not use it without the author's permission.

Editor

The editors are responsible and have full authority in accepting or rejecting a manuscript sent by the authors. An assessment of a manuscript must still prioritize the scientific value, leaving aside personal and commercial interests.

If the manuscript is not suitable enough to be published, the editors must keep the confidentiality of the manuscript, so that it is not used by others, unless they get permission by the authors. The editors must fully keep the principle of blind review.

Publisher

As the publisher of the Ovozoa Journal, Faculty of Veterinary Medicine, Universitas Airlangga, in collaboration with the Association of the Indonesian Veterinary Reproduction, always encourages editors to obey with predetermined scientific writing procedures.

The publisher, in collaboration with the authors and editors in charge of maintaining academic integrity throughout the publication process. The publisher also maintains the quality of the journal and provides policies that encourage the development of the journal.

Animal Ethics

Research involving experimental animals be carried out following animal ethics and welfare. Experimental animals must be appropriately treated and minimizing the pain that might arise during the treatment. The utilization of experimental animals must be following local, national, and international regulations. The authors must make a statement, including the name of the ethical authority and the consent number; the trial is carried out in compliance with animal ethics and welfare.

ARTICLE PROCESSING CHARGES

Author Contributions

Article Processing Charges for each accepted article is IDR 450,000. There is no other charges. For the year 2022 article processing charges will not be levied, after a year the charges will be applied as written above.

PLAGIRISM SCREENING

Submitted manuscript will be checked using Turnitin software. The manuscript is passed if the similarity is less than 30%.

ADDRESS FOR CORRESPONDENCE

Editors of OVOZOA Journal of Animal Reproduction

Faculty of Veterinary Medicine Universitas Airlangga

Kampus C, Mulyorejo, Surabaya 60115

Phone: (031) 5992785 ext. 5993016

Email: ovozoa@journal.unair.ac.id

Website: <https://e-journal.unair.ac.id/OVZ/index>



This journal is supported by



Kampus C Mulyorejo – 60115, Surabaya Tel. +6231 599 2377 Fax. +6231 599 3015
 Email: ovozaa@journal.unair.ac.id, website: <https://e-journal.unair.ac.id/OVZ>

Copyright Transfer Agreement

I(We), the undersigned, am(are) the author(s) of the article entitled:

.....

In connection with the receipt of the article mentioned above in the OVOZOA Journal of Animal Reproduction, I(We) hereby agree to transfer all copyright ownership of the article in all forms of media (electronic, print, etc.) to be published in the OVOZOA Journal of Animal Reproduction.

The author(s) declare that:

- The article is truthful, original work of the author(s) without fabrication, fraud, or plagiarism.
- This paper contains works that have not previously published in any form and not under consideration for publication in other journals or other scientific media.
- I do it full myself (Each author was contributing) intellectually, substantially, directly, (and significantly) in the research and the writing of the article.
- I take full of public responsibility (Each author has the same public responsibility) for the article.
- This article was not intended for harm or benefit certain parties and does not violate the rights of third parties.
- All permissions needed for the research have been obtained in written form.
- I(All authors) has(have) received the final version of the article and agreed to be published in the OVOZOA Journal of Animal Reproduction.

Put a * sign behind the name of the Corresponding Author.

No	Name	Signature	Date
1			
2			
3			
4			
5			
6			
7			

Please add rows if it's not enough.

(This form must be signed by all authors in order as appeared in the article, and should be returned to the editorial office.)