

# Correlation between Neutrophil-to-Lymphocyte Ratio with Disease Severity in Diabetic Patients with COVID-19 at Tertiary Referral Hospital in Indonesia

*by* Hermina Novida

---

**Submission date:** 19-Nov-2021 12:15PM (UTC+0800)

**Submission ID:** 1707284970

**File name:** Artikel.pdf (598.58K)

**Word count:** 7131

**Character count:** 30523

Open Access Article

## Correlation between Neutrophil-to-Lymphocyte Ratio with Disease Severity in Diabetic Patients with COVID-19 at Tertiary Referral Hospital in Indonesia

Cupuwatie Cahyani<sup>1</sup>, Hermina Novida<sup>2</sup>, Soebagijo Adi Soelistijo<sup>2</sup>, Usman Hadi<sup>3</sup>, Nenci Siagian<sup>1</sup>

<sup>1</sup> Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia

<sup>2</sup> Division of Endocrinology, Diabetes, and Metabolism, Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia

<sup>3</sup> Division of Tropic and Infection, Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia

**Abstract:** Coronavirus disease 2019 (COVID-19) is highly transmissible and spreads quickly with the clinical presentation varies from asymptomatic to severe respiratory symptoms, even leading to death. Diabetes mellitus is one of the main comorbidities contributing to worsening the COVID-19 cases. Increased Neutrophil-to-Lymphocyte Ratio (NLR) is considered as an early warning of COVID-19 severity. This study evaluated the correlation between NLR with disease severity at admission in diabetic patients with COVID-19 at a tertiary referral hospital in Indonesia. The authors performed a retrospective study using secondary data from medical records of diabetic inpatients with COVID-19 from May to September 2020. The demographic data were collected from the medical record and the hematologic parameter from the laboratory. NLR was calculated by dividing the absolute neutrophils counts by the absolute lymphocyte counts. The data was analyzed with a significance level of  $p < 0.05$ . Study subjects consisted of 100 non-severe cases and 128 severe cases. The median NLR of severe and non-severe groups was 9.7 vs. 5.22 ( $p < 0.001$ ). The correlation coefficients of NLR with disease severity were 0.52 ( $p < 0.001$ ). The calculated Area Under the Curve (AUC) of the ROC analysis for the NLR was 0.803 (cut-off:  $> 6.15$ ;  $p < 0.001$ ) with sensitivity 91%, specificity 64%, negative predictive value 16%, and positive predictive value 75.82%. NLR had a significant positive correlation with disease severity at admission in diabetic patients with COVID-19. As simple, rapid, and cost-effective biomarkers, NLR can help clinicians identify potentially severe cases early, conduct early triage, and initiate effective management in time so the progress of disease severity should be possibly prevented.

**Keywords:** COVID-19, diabetic patients, disease severity, neutrophil-to-lymphocyte ratio.

## 三级转诊医院糖尿病患者中性粒细胞与淋巴细胞比率与中性粒细胞与淋巴细胞比率疾病轻重程度的相关性分析在印度尼西亚

**提要:** 2019 年冠状病毒病 (新冠肺炎) 具有高度传染性并迅速传播, 临床表现从无症状到严重的呼吸道症状不等, 甚至导致死亡。糖尿病是导致新冠肺炎病例恶化的主要合并症之一。中性粒细胞与淋巴细胞比率的增加被认为是新冠肺炎严重性的早期预警。本研究旨在评估印度尼西亚三级转诊医院新冠肺炎糖尿病患者入院时中性粒细胞与淋巴细胞与疾病严重程度之间的相关性。我们使用 2020 年 5 月至 9 月新冠肺炎糖尿病住院患者病历的二手数据进行了一项回顾性研究。人口统计学数据来自病历和实验室的血液学参数。中性粒细胞与淋巴细胞是通过将绝对中性粒细胞计数除以绝对淋巴细胞计数来计算的。以  $p < 0.05$  的显著性

Received: May 1, 2021 / Revised: June 28, 2021 / Accepted: July 14, 2021 / Published: September 30, 2021

About the authors: Cupuwatie Cahyani, Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia; Hermina Novida, Soebagijo Adi Soelistijo, Division of Endocrinology, Diabetes, and Metabolism, Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia; Usman Hadi, Division of Tropic and Infection, Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia; Nenci Siagian, Department of Internal Medicine, Faculty of Medicine, Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia

水平分析数据。研究对象由 100 名非重症病例和 128 名重症病例组成。重度和非重度组的中位中性粒细胞与淋巴细胞的比率分别为 9.7 和 5.22 (磷<0.001)。中性粒细胞与淋巴细胞的比率与疾病严重程度的相关系数为 0.52 (磷<0.001)。中性粒细胞与淋巴细胞的比率的接收器操作特性分析的计算机曲线面积为 0.803 (临界值 : >6.15 ; 磷<0.001) , 灵敏度为 91% , 特异性为 64% , 阴性预测值为 16% , 阳性预测值为 75.82%。中性粒细胞与淋巴细胞与新冠肺炎糖尿病患者入院时的疾病严重程度呈显著正相关。中性粒细胞与淋巴细胞作为简单、快速且具有成本效益的生物标志物, 可以帮助临床医生及早识别潜在的严重病例, 进行早期分类, 并及时启动有效管理, 从而可能阻止疾病严重程度的进展。

**关键词:** 新冠肺炎, 糖尿病患者, 疾病严重程度, 中性粒细胞与淋巴细胞的比率。

11

## 1. Introduction

The new coronavirus disease-19 (COVID-19), caused by Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2), emerged in late December 2019 in Wuhan, China, and World Health Organization, on January 30, 2020, called public health emergency [1]. It has spread worldwide rapidly and widely. Meanwhile, new cases in Indonesia are still increasing, with the total number of confirmed COVID-19 cases as of mid-September 2020 228,993 people with 9,100 deaths [2]. COVID-19 infection can be categorized as an asymptomatic, mild, moderate, severe, or critical disease [3, 4]. Clinically, supposed early warning of severe COVID-19 infection can be identified, so timely intervention and treatment may help reduce mortality, improve the cure rate, and shorten the hospital stay.

Diabetes mellitus (DM) is one of the most prevalent chronic conditions with devastating multi-systemic complications and was estimated to have inflicted 463 million people in 2019 [5]. Many reports show that diabetes is a frequent pre-existing condition associated with severe disease in COVID-19 patients, and it is an unholly situation wherein one disease entity compliments the other. How diabetes increases the severity of COVID-19 is unclear, though several factors may be responsible [6]. Diabetes is a chronic inflammatory condition, so a pro-inflammatory state could accentuate the cytokine storm, which is believed to be responsible for acute respiratory distress syndrome (ARDS) as well as multi-organ dysfunction in COVID-19. Poor glycemic control, event transient hyperglycemia may temporarily affect innate immune responses to infection. Immune defects named inappropriate T-cell action, impaired natural killer cell activity, and defects in complement action could reduce viral clearance.

Further, diabetes is associated with increased plasminogen levels, which have been postulated to increase the virulence of SARS CoV-2. Increased viral replication in diabetes may also be due to increased

furin, a type-1-membrane-bound- protease involved in coronavirus entry into the cell. In addition, pre-existing comorbidities associated with diabetes like hypertension, coronary artery disease, chronic kidney disease, and obesity may further impair immunity and predispose to severe infection [7, 8, 9].

Neutrophil-to-Lymphocyte Ratio (NLR) is an easy-to-analyzed inflammation biomarker that is feasible in all hospital settings. Many studies denote elevated NLR as an excellent early warning factor to identify severe disease in COVID-19 [10-15]. However, a few studies analyzing the correlation between NLR with severity in special populations like diabetic patients. According to the background, this study aimed to analyze the relation between NLR with disease severity at admission in diabetic patients with COVID-19.

6

## 2. Methods

### 2.1. Study Design, Participants, and Data Collection

This study was a retrospective, cross-sectional method, which was carried out by taking secondary data from May to September 2020 with patients who have diagnosed diabetes with COVID-19 in the medical records of Dr. Soetomo Hospital Surabaya, a major tertiary referral hospital in Indonesia and has been mainly responsible for the treatments of COVID-19 patients assigned by the government.

Diabetic patients were admitted to the wards with COVID-19 infection confirmed by real-time reverse transcription-polymerase chain reaction (RT-PCR). Inclusion criteria were as follows: (1) diabetic with COVID-19, (2) results of blood routine examination were obtained within 24 hours of admission. Patients were excluded using the following criteria: (1) patients with known hematological illnesses, with known HIV-positive status, those on chemotherapeutic drugs, long-term glucocorticoids, and pregnant women because these conditions affect the NLR, (2) patients with missing data.

The severity<sup>18</sup> of the disease was classified into non-severe cases (mild and moderate symptoms) and severe cases (severe symptoms<sup>25</sup> critical illness) based on the severity of symptoms according to interim guidance of the World Health Organization<sup>16</sup>. Severe cases were defined when one of the following criteria was present: (1) respiratory distress (respiratory rate over 30 breaths per minute<sup>15</sup>), (2) oxygen saturation  $\leq 93\%$  on room air, and (3) arterial blood oxygen partial pressure (PaO<sub>2</sub>)/oxygen concentration (FiO<sub>2</sub>)  $\leq 300$  mmHg (1 mmHg = 0.133 kPa).

Information on demographic data, symptoms, pre-existing comorbidities, and laboratory results were collected. The routine laboratory tests complete blood count (CBC), liver function tests (LFTs), renal function test (RFTs), serum electrolytes, C-reactive protein (CRP), Albumin serum, and blood glucose level. CBC was analyzed using the hospital laboratory system, Sysmex XS-800i. The NLR was calculated using the simple formula absolute number of neutrophils divided by an absolute number of lymphocytes. The time from the onset of illness to the hospital admission was also recorded. The assessment of disease severity and laboratory tests was performed on the day of the patient admission before treatment.

## 2.2. Statistical Analysis

The authors describe the categorical variables as number<sup>10</sup> and percentages (%), and continuous variables as mean standard deviation (SD) if they are typically distributed or median with interquartile ranges (IQR) if they are not. The Kolmogorov-Smirnov test was used to verify the normality of the distribution. Independent-group t-test was used<sup>3</sup> to compare parametric continuous variables or Mann-Whitney U test for non-parametric continuous variables. The proportion for categorical variables was compared using the Chi-square test or Fisher's exact test. Correlation between variables was assessed using Spearman's correlation analysis.

Receiver operating characteristics (ROC) were<sup>24</sup> used to study the accuracy of the various predictive test. All statistical analyses were performed using SPSS version 25.0 software. Two-sided P values of less than 0.05 were considered statistically significant.

## 2.3. Ethical Approval

Health Research Ethics Committee of Dr. Soetomo Hospital (Surabaya, Indonesia) approved this study protocol.

## 3. Results

Two hundred twenty-eight diabetic patients who were confirmed positive for COVID-19 were included in this study. 100 (43.86%) were non-severe<sup>19</sup> cases, and 128 (56.14%) were severe cases. The demographic and

clinical characteristics of all patients were shown in Table 1.

There were many significant differences<sup>21</sup> in the parameters of baseline characteristics between the severe group and the non-severe group. Patients with the severe group had older age  $\geq 60$  years (36.7% vs. 23%,  $P = 0.037$ ), more likely to have underlying comorbidities hypertension (43% vs. 29%,  $P = 0.03$ ) and chronic kidney disease (2.2% vs. 0%,  $P = 0.046$ ), had symptoms dyspnea (74.2% vs. 52%,  $P = 0.001$ ) and sore throat (16.4% vs. 5%,  $P = 0.007$ ) compared to patients with the non-severe group. While in the non-severe group had symptoms anosmia (7% vs. 1.6%,  $P = 0.037$ ), fatigue (30% vs. 18%,  $P = 0.033$ ), and myalgia (8% vs. 1.6%,  $P = 0.018$ ) compared to patients with severe group. There were no significant differences in sex, hospital admission, the onset of diabetes, use of diabetes medication, cardiovascular disease, cerebrovascular disease, the onset of symptoms to hospital admission, and symptoms such as dry cough, fever, anorexia, diarrhea, and runny nose.

Table 2 presents the laboratory findings in the non-severe and severe group on the day of hospital admission in diabetic patients with COVID-19. There were many significant differences in the parameters of laboratory findings between non-severe and severe groups. Severe cases had higher white blood cell (9.93 vs 7.22,  $P = 0.001$ ); higher neutrophil count (8.26 vs 5.61,  $P < 0.001$ ); higher neutrophil-to-lymphocyte ratio (NLR) (9.7 vs 5.22,  $P < 0.001$ ); higher platelet-to-lymphocyte ratio (PLR) (304.1 vs 197.78,  $P < 0.001$ ), higher-level aspartate aminotransferase (AST) (59 vs 46,  $P = 0.002$ ), and higher-level C-reactive protein (CRP) (11.6 vs 7.4,  $P = 0.013$ ), lower lymphocyte count (0.835 vs 1.29,  $P < 0.001$ ) and lower albumin (3.1 vs 3.25,  $P = 0.018$ ) compared to patients with the non-severe group. There were no significant differences in hemoglobin level, platelet count, plasma glucose level, sodium level, potassium level, creatinine level, blood urea nitrogen level, alanine aminotransferase level, activated partial thromboplastin time, and prothrombin time.

Table 3 presents the results<sup>19</sup> of Spearman correlation analysis between NLR with the severity of COVID-19. NLR in diabetic patients with COVID-19 has a significant positive correlation with disease severity ( $P < 0.001$ ) with a moderate correlation strength ( $r = 0.52$ ). The ROC curve in Figure 1 showed that The Area under Curve (AUC) for NLR was 0.803 [95% CI (0.74-0.865),  $P < 0.001$ ]. NLR's cut-off-value as the optimal threshold for predicting the disease severity in diabetic patients with COVID-19 at admission is  $> 6.15$  with the sensitivity of 90.6%, specificity of 64%, the positive predictive value 75.82%, and negative predictive value 16%. In addition, the positive likelihood ratio was  $0.906/1 - 0.64 = 2.52$ , and the negative likelihood ratio was  $1 - 0.906/0.64 = 0.15$ .



Table 1 Characteristics of research subjects

Variable	Total = 228 No(%)	Non-severe (n = 100)	Severe (n = 128)	P-value
<b>Age, Median (IQR), y</b>	<b>55 (50-61)</b>	<b>53.5 (48-58.5)</b>	<b>56 (52-61)</b>	<b>0.019<sup>a*</sup></b>
<60 years	158 (69.3%)	77 (77%)	81 (63.3%)	0.037 <sup>b*</sup>
≥60 years	70 (30.7%)	23 (23%)	47 (36.7%)	
<b>Sex</b>				
Female	112(49.1%)	45 (45%)	67 (52.3%)	0.273 <sup>b</sup>
Male	116 (50.9%)	55 (55%)	61 (47.7%)	
<b>Hospital admission</b>				
Referred	68 (29.8%)	25 (25%)	43 (33.6%)	0.161 <sup>b</sup>
Directly	160 (70.2%)	75 (75%)	85 (66.4%)	
<b>Onset of T2DM</b>				
New Onset	45 (19.7%)	19 (19%)	26 (20.3%)	0.262 <sup>c</sup>
<5 Years	127 (55.7%)	52 (52%)	75 (58.6%)	
5-10 Years	35 (15.4%)	17 (17%)	18 (14.1%)	
>10 Years	21 (9.2%)	12 (12%)	9 (7%)	
<b>Use of diabetes medication</b>				
None	46 (20.2%)	19 (19%)	27 (21.1%)	0.15 <sup>c</sup>
Routine	113 (49.6%)	56 (56%)	57 (44.5%)	
Not Routine	69 (30.3%)	25 (25%)	44 (34.4%)	
<b>Comorbidities</b>	<b>97 (42.54%)</b>	<b>32 (32.99%)</b>	<b>65 (67.01%)</b>	<b>0.009<sup>b*</sup></b>
Hypertension	84 (36.8%)	29 (29%)	55 (43%)	0.03 <sup>b*</sup>
Cardiovascular Disease	11 (4.8%)	6 (6%)	5 (3.9%)	0.466 <sup>b</sup>
Chronic Kidney Disease	9 (3.9%)	0 (0%)	5 (2.2%)	0.046 <sup>b*</sup>
Cerebrovascular Disease	5 (2.2%)	3 (3%)	6 (4.7%)	0.518 <sup>b</sup>
<b>Sign and symptoms</b>				
Dry Cough	170 (74.6%)	71 (71%)	99 (77.3%)	0.277 <sup>b</sup>
Fever	153 (67.1%)	68 (68%)	85 (66.4%)	0.8 <sup>b</sup>
Dyspnea	147 (64.5%)	52 (52%)	95 (74.2%)	0.001 <sup>b*</sup>
Anorexia	88 (38.6%)	43 (43%)	45 (35.2%)	0.229 <sup>b</sup>
Anosmia	9 (3.9%)	7 (7%)	2 (1.6%)	0.037 <sup>b*</sup>
Fatigue	53 (23.2%)	30 (30%)	23 (18%)	0.033 <sup>b*</sup>
Sore throat	26 (11.4%)	5 (5%)	21 (16.4%)	0.007 <sup>b*</sup>
Diarrhea	24 (10.5%)	11 (11%)	13 (10.2%)	0.838 <sup>b</sup>
Runny nose	16 (7%)	6 (6%)	10 (7.8%)	0.599 <sup>b</sup>
Myalgia	10 (4.4%)	8 (8%)	2 (1.6%)	0.018 <sup>b*</sup>
Illness onset to hospital admission, Median (IQR), days	4 (3-7)	4 (3-7)	4.5 (3-7)	0.536 <sup>a</sup>

Note: P-values comparing severe and non-severe cases are divided from <sup>a</sup> Mann-Whitney U test, <sup>b</sup> Chi-square test, and <sup>c</sup> Fisher's exact test. P < 0.05 was considered statistically significant and marked by \*

Abbreviations: IQR - interquartile range

Table 2 Laboratory findings at admission in diabetic patients with COVID-19

	Median (IQR) Total (N = 228)	Non-severe (n = 100)	Severe (n = 128)	P value
Hemoglobin (g/L, normal range 11-15)	13.45 (12-14.6%)	13.7 (12-15.1)	13.35 (12-14.4)	0.091 <sup>a</sup>
White blood cell count (x10 <sup>9</sup> /L, normal range 3.5-9.5)	8.96 (6.96-12.47)	7.93 (6.43-10.76)	9.94 (7.6-14.12)	0.001 <sup>b*</sup>
Neutrophil count (x10 <sup>9</sup> /L, normal range 1.8-6.3)	7.4 (5.27-11.33)	5.61 (4.45-8.92)	8.26 (6.37-12.83)	0.000 <sup>b*</sup>
Lymphocyte count (x10 <sup>9</sup> /L, normal range 1.1-3.2)	0.97 (0.75-13.47)	1.29 (0.91-1.61)	0.84 (0.64-10.88)	0.000 <sup>b*</sup>
Platelet count (x10 <sup>9</sup> /L, normal range 125-350)	256.5 (206-337)	260.5 (193.75-349.75)	248.5 (208.5-328.25)	0.747 <sup>b</sup>
The neutrophil-to-Lymphocyte ratio (NLR)	8.12 (5.48-12.35)	5.22 (2.99-8.2)	9.7 (7.4-14.34)	0.000 <sup>b*</sup>
Platelet-to-Lymphocyte ratio (PLR)	265.97 (180.45-367.22)	197.78 (144.83-304.17)	304.17 (234.64-404.75)	0.000 <sup>b*</sup>
Plasma glucose, mmol/L (normal range 140-180)	239 (160.5-312.25)	236 (166.75-288.75)	240.5 (14.25-336.25)	0.755 <sup>b</sup>
Serum sodium, mmol/L (normal range 135-145)	135 (130.25-139)	134 (130-138)	135.5 (131-140)	0.128 <sup>b</sup>
Serum potassium, mmol/L (normal range 3.5-5.5)	4.1 (3.5-4.7)	4.1 (3.5-4.6)	4.1 (3.5-4.7)	0.31 <sup>a</sup>
Creatinine, mmol/L (normal range 0.4-1.2)	1.1 (0.8-1.6)	1 (0.8-1.48)	1.2 (0.8-1.88)	0.098 <sup>b</sup>
BUN, mmol/L (normal range 25-67)	19 (12-30)	17 (11-26.75)	19 (13-31)	0.113 <sup>b</sup>
Albumin, g/dL (normal range 3.4-5.4)	3.13 (2.9-3.35)	3.25 (2.9-3.5)	3.1 (2.9-3.3)	0.018 <sup>b*</sup>
AST U/L (normal range 8-40)	52 (37-83)	46 (30-71.25)	59 (42-88.75)	0.002 <sup>b*</sup>
ALT U/L (normal range 5-35)	46 (33-71.75)	44.5 (31-70)	46.5 (34-74.25)	0.347 <sup>b</sup>
APTT, s (normal range 20-40)	26.35 (25.53-29.85)	26.35 (23.63-29.75)	26.35 (24.9-30.8)	0.15 <sup>b</sup>
PT, s (normal range 9-14)	11.2 (10.2-13.5)	11 (10.13-13.15)	11.7 (10.33-13.6)	0.153 <sup>b</sup>
CRP, mg/L (normal range 0-8)	10.65 (4.76-16.6)	7.4 (3.83-14.4)	11.6 (6.22-17.73)	0.013 <sup>b*</sup>

Note: P-values comparing severe and non-severe cases are divided from <sup>a</sup> at the test and <sup>b</sup> Mann-Whitney U-test. P<0.05 was considered statistically significant and marked by \*

Abbreviations: IQR - interquartile range, BUN - blood urea nitrogen, AST - aspartate aminotransferase, ALT - alanine aminotransferase, APTT - activated partial thromboplastin, PT - prothrombin time, CRP - C-reactive protein, and COVID-19 - Coronavirus disease 2019.

Table 3 Correlation NLR with severity COVID-19 in diabetic patients

	R-value	P-value
NLR	0.52	0.000




Fig. 1 The ROC curves of NLR in the predicting of severe COVID-19 in diabetic patients at admission

#### 4. Discussion

This study was conducted at a tertiary referral hospital, where the patients who came were referred patients but in this study, more patients came directly than referred. This study showed significant differences between non-severe and severe median age (53.5 vs. 56 years;  $P=0.019$ ). These conclusions are quite consistent with the previous studies [16]. Advanced chronological age is one of the main risk factors for the adverse outcomes of COVID-19, presumably due to immunological changes (immunosenescence and inflammation) and other organ dysfunction [17, 18].

This study also found that the severe case of the diabetic patients with COVID-19 who had comorbid hypertension and chronic kidney disease were significantly higher than the non-severe cases group. This finding is supported by the results of a previous study [19-20]. The specific pathogenesis of hypertension that may lead to more severe COVID-19 remains to be studied. The imbalance of cytokines may be considered an explanation for the correlation between hypertension and severe COVID-19. Increasing clinical data have shown a relationship between the deterioration of COVID-19 and cytokine storms, such as elevated levels of interleukin-6, interleukin-7, granulocyte-macrophage-colony-stimulating-factor, and tumor necrosis factor- $\alpha$  [21]. Several lines of evidence suggested that CKD patients, especially those at advanced stages, are vulnerable to SARS-CoV-2 infection.

Moreover, a higher baseline serum creatinine level was an independent risk factor for in-hospital death in COVID-19 [22]. The basis for such vulnerability is

likely multifactorial, and both environmental factors and medical factors such as old age, immune cell dysfunction, cardiovascular and pulmonary comorbidities need to be considered [23]. In conclusion, elderly or older people and a higher frequency of comorbidities in diabetic COVID-19 patients are more susceptible to severe or critical conditions.

Symptoms of COVID-19 may appear anytime from 2 to 14 days after exposure. Therefore, 14-day quarantine is recommended [24, 25]. This study showed no difference in onset of symptoms to hospital admission in patients with the severe and non-severe groups. The severe cases mostly had a cough, dyspnea, fever, and anorexia but significantly higher in dyspnea and sore throat than non-severe cases. The non-severe cases had significantly higher anosmia, fatigue, and myalgia than severe cases. This study was in line with a study in China that the most common symptoms observed from the onset include fever, cough, and fatigue [26]. A meta-analysis from 55 studies explained that clinical manifestations such as fever, cough, fatigue, anorexia, dyspnea, chest tightness, hemoptysis, diarrhea, and abdominal pain were significantly associated with the severity of cases. However, it was not for myalgia, pharyngalgia, nausea, vomiting, headache, dizziness, and sore throat [27]. Therefore, sore throat and dyspnea can be used as an early warning towards critical illness in diabetic patients with COVID-19.

This study found that plasma glucose at admission between non-severe and severe cases was not significantly different, but the median of plasma glucose was  $>200$  mg/dL. It was explained based on SARS-CoV-2 mediated damage of the pancreatic  $\beta$ -cell as ACE-2 is also expressed on the pancreatic islets. It could partly explain the worsening glucose control in diabetic patients with some functional  $\beta$ -cell in reserve [28]. Infection also causes a stress response in the body by increasing certain hormones such as cortisol and adrenaline. These hormones work against the action of insulin and, as a result, the body's production of glucose increases, which results in high blood sugar levels [29].

Higher neutrophil and lower lymphocyte count in this study showed a significant difference in severe than non-severe cases. Neutrophil and lymphocyte are two key indexes affected by SARS CoV-2 infection and demonstrated significant differences in the severe and non-severe COVID-19 infection [11, 30, 31]. Neutrophils are one of the human body's vital immune cells. When pathogenic microorganisms invade the body, immune cells tend to rapidly chemotactically gather to the infection site and play the role of host defense and immune regulation. Lymphocytes are the primary effector cells of the human immune response. The number of lymphocytes in the body is closely related to the body's immunity and defense system

against pathogenic microorganisms and is negatively correlated with inflammation.

NLR reflects the balance of the body's neutrophil, lymphocyte count levels, and the degree of systemic inflammation. Therefore, a high NLR is an important marker that indicates an imbalance in the inflammatory response and marker of disease severity [4, 32]. It is in line with this study that higher NLR was found a significant difference in the severe cases than in the non-severe group (9.7 vs. 5.22,  $P < 0.001$ ). The present study also indicated significant positive correlation between NLR with disease severity of COVID-19 in diabetic patients ( $R = 0.52$ ,  $P < 0.001$ ).

The authors performed the ROC analysis to predict the disease severity in hospitalized diabetic patients with COVID-19 and determined the cut-off levels on-time admission. The cut-off value of NLR obtained in this study was  $> 6.15$  to predict severe cases. The ROC analysis performed according to the NLR cut-off values were calculated as AUC 0.803 (95% CI: 0.74-0.865,  $P < 0.001$ ) with sensitivity 90.6%, specificity 64%, positive predictive value 75.82%, and negative predictive value of 16%. In addition, it was shown that the positive likelihood ratio = 2.52 (fair) and the negative likelihood ratio = 0.15 (excellent), indicating that the cut-off for NLR of 6.15 will provide sufficient probability in predicting the severity of diabetic COVID-19 patients. This value is higher than many studies about NLR and disease severity of COVID-19 (NLR values ranging from 3.3 to 5.9 to predict the severity) because, in diabetic conditions, NLR levels were also increased due to chronic low-grade inflammation [10]. However, no NLR consensus cut-off values have been established to determine normal and elevated NLR values, especially for COVID-19 [13-14, 32-34].

This study also had other abnormal indicators that had significant differences between the severe and non-severe cases, such as higher white blood count, PLR, and albumin serum AST and CRP. These abnormalities suggested that SARS CoV-2 infection might be associated with myocardial injury, hepatic injury, and other related organ damage [35].

## 5. Conclusion

This single-center observational study revealed that some findings in this study might be consistent in severe cases with previous studies, including higher age ( $\geq 60$  years), history of hypertension and chronic kidney disease, presence of symptoms of dyspnea and sore throat, decreasing lymphocyte and albumin, and increasing white blood count, neutrophil count, CRP level, and alanine aminotransferase level. Early identification of risk factors for severe patients is vital to afford appropriate supportive care or access to the intensive care unit (ICU) if necessary. At admission, neutrophil to Lymphocyte Ratio (NLR) had a significant positive correlation with disease severity in

diabetic patients with COVID-19. NLR has known as a simple, rapid, and cost-effective biomarker. Evaluating NLR can help clinicians predict severe cases early, conduct early triage, and initiate effective management in time, reducing the overall mortality of diabetic patients with COVID-19.

However, this study had certain limitations, such as having a small sample size, use secondary data from the medical record, and being a single-center study. For more accurate and precise results, wider generalizability of the findings, and larger sample size, clinical studies are required to confirm the findings further.

## References

- [1] HUANG C, WANG Y, LI X. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*, 2020, 395(10223): 497-506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
- [2] COVID-19 kemenkes.go.id. Situasi terkini perkembangan Coronavirus Disease (COVID-19). 2020. Available (accessed 15 September, 2020) from <https://Covid19.kemkes.go.id/situasi-infeksi-emerging/info-corona-virus/situasi-terkini-perkembangan-coronavirus-disease-COVID-19-15-september-2020/#.XykMYIUzBIU>
- [3] DIAZ-BALLVE L, RISSO-VASQUEZ A, RIOS F. Coronavirus disease 2019 (COVID-19) aspects of interest for critical care – a narrative review. *Rev Arg de Ter Int*, 2020, 5(Suppl 1): 1- 11.
- [4] IMRAN MM, AHMED U, USMAN U. Neutrophil/lymphocyte ratio—A marker of COVID-19 pneumonia severity. *Int J Clin Pract*, 2021, 75: e13698.
- [5] SAEEDI P, PETERSOHN I, SALPEA P. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res Clin Pract*., 2019, 157: 107843.
- [6] PAL R, BHADADA SK. COVID-19 and diabetes mellitus: An unholy interaction of two pandemics. *Diabetes Metab Syndr*, 2020, 14(4): 513-517.
- [7] GUPTA R, HUSSAIN A, MISRA A. Diabetes and COVID-19: evidence, status and unanswered research questions. *Eur J Clin Nutr*, 2020, 74: 864–870.
- [8] CRITCHLEY JA, CAREY IM, HARRIS T. Glycemic Control and Risk of Infections Among People With Type 1 or Type 2 Diabetes in a Large Primary Care Cohort Study. *Diabetes Care*, 2018, 41(10): 2127-2135.
- [9] MADDALONI E, BUZZETTI R. Covid-19 and diabetes mellitus: unveiling the interaction of two pandemics. *Diabetes Metab Res Rev*, 2020, e33213321.
- [10] HUSSAIN M, BABAR MZM, AKHTAR L, HUSSAIN MS. Neutrophil lymphocyte ratio (NLR): A good assessment tool of glycemic control in type 2 diabetic patients. *Pak J Med Sci*, 2017; 33(6): 1366-1370.
- [11] LIU L, ZHENG Y, CAI L. Neutrophil-to-lymphocyte ratio, a critical predictor for assessment of disease severity in patients with COVID-19. *Int J Lab Hematol.*, 2021, 43(2): 329-335.
- [12] KONG M, ZHANG H, CAO X. Higher level of neutrophil-to-lymphocyte is associated with severe COVID-19. *Epidemiol Infect*, 2020, 148: e139.
- [13] SIMADIBRATA DM, CALVIN J, WIJAYA AD. The



neutrophil-to-lymphocyte ratio on admission to predict the severity and mortality of COVID-19 patients: A meta-analysis. *Am J Emerg Med*, 2021, 42: 60-69.

[14] LI X, LIU C, MAO Z. Predictive values of neutrophil-to-lymphocyte ratio on disease severity and mortality in COVID-19 patients: a systematic review and meta-analysis. *Crit Care*, 2020, 24(1): 647.

[15] CHAN ABIGAIL SY, ROUT A. Use of Neutrophil-to-Lymphocyte and Platelet-to-Lymphocyte Ratios in COVID-19. *J Clin Med Res*, 2020, 12(7): 448-453.

[16] KSHANTI IA, EPRILIAWATI M, MOKOAGOW I, et al. The Impact of Coronavirus Disease 2019 Pandemic on People with Diabetes in Indonesia: A Cross-Sectional National Scale Web-Survey. 2020. 10.1101/2020.12.01.20241588.

[17] PERROTTA F, CORBI G, MAZZEO G. COVID-19 and the elderly: insights into pathogenesis and clinical decision-making. *Aging Clin Exp Res*, 2020, 32(8): 1599-1608.

[18] LANDI F, BARILLARO C, BELLINI A. The New Challenge of Geriatrics: Saving Frail Older People from the SARS-COV-2 Pandemic Infection. *J Nutr Health Aging*, 2020, 24(5): 466-470.

[19] SHI Y, YU X, ZHAO H. Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care*, 2020, 24(1): 108.

[20] LI X, XU S, YU M. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol*, 2020, 46(1): 110-118.

[21] DRUMMOND GR, VINH A, GUZIK TJ, SOBEY CG. Immune mechanisms of hypertension. *Nat Rev Immunol*, 2019, 19(8): 517-532.

[22] CHENG Y, LUO R, WANG K. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int*, 2020, 97(5):829-838.

[23] COBO G, et al. Chronic inflammation in end-stage renal disease and dialysis. *Nephrology Dialysis Transplantation*, 2018, 33(3): iii35-iii40.

[24] ZHENG Z, PENG F, XU B. Risk factors of critical & mortal COVID-19 cases: A systematic literature review and meta-analysis. *The Journal of infection*, 2020, 81(2): e16-e25.

[25] FU L, WANG B, YUAN T. Clinical characteristics of coronavirus disease 2019 (COVID-19) in China: A systematic review and meta-analysis. *J Infect*, 2020, 80(6): 656-665.

[26] CHANG DE, MO G, YUAN X. Time Kinetics of Viral Clearance and Resolution of Symptoms in Novel Coronavirus Infection. *Am J Respir Crit Care Med*, 2020, 201(9): 1150-1152.

[27] BAREK MA, AZIZ MA, ISLAM MS. Impact of age, sex, comorbidities, and clinical symptoms on the severity of COVID-19 cases: A meta-analysis with 55 studies and 10014 cases. *Heliyon*, 2020, 6(12): e05684.

[28] LIU J, LI Y, LIU Q. SARS-CoV-2 cell tropism and multi-organ infection. *Cell Discov*, 2021, 7(17).

[29] HANTZIDIAMANTIS PJ, LAPPIN SL. Physiology, Glucose. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK545201/>

[30] HAN Q, WEN X, WANG L. Role of hematological parameters in diagnosing influenza virus infection in patients with respiratory tract infection symptoms. *J Clin Lab Anal*, 2020, 34(5): e23191.

[31] LIU Y-P, LI G-M, HE J. Combined use of the neutrophil-to-lymphocyte ratio and CRP to predict 7-day disease severity in 84 hospitalized patients with COVID-19 pneumonia: a retrospective cohort study. *Ann Transl Med*, 2020, 8(10): 635.

[32] YANG A-P, LIU J-P, TAO W-Q, LI H-M. The diagnostic and predictive role of NLR, d-NLR, and PLR in COVID-19 patients. *Int Immunopharmacol*, 2020, 84: 106504.

[33] SUN S, CAI X, WANG H. Abnormalities of the peripheral blood system in patients with COVID-19 in Wenzhou, China. *Clin Chim Acta*, 2020, 507: 174-180.

[34] SONG C-Y, XU J, HE J-Q, LU Y-Q. COVID-19 early warning score: a multi-parameter screening tool to identify highly suspected patients. *MedRxiv*, 2020, 10-3.

[35] PEIRIS S, MESA H, AYSOLA A. Pathological findings in organs and tissues of patients with COVID-19: A systematic review. 2021. *PLoS ONE*, 16(4): e0250708.

#### 參考文:

[1] HUANG C, WANG Y, LI X. 2019 新型冠状病毒感染患者的臨床特徵。柳葉刀, 2020, 395 (10223) : 497-506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)

[2] 新冠肺炎 kemenkes.go.id. 最新發展情況冠狀病毒病(新冠肺炎)。2020. 可從 <https://Covid19.kemkes.go.id/situasi-infeksi-emerging/info-corona-virus/situasi-terkini-perkembangan-coronavirus-disease-COVID-19-15> 獲得 (2020 年 9 月 15 日訪問) - september-2020/#.XykMYIUzbIU

[3] DIAZ-BALLVE L, RISSO-VASQUEZ A, RIOS F. 對重症監護感興趣的 2019 年冠狀病毒病(新冠肺炎) 方面——敘述性評論。牧師精氨酸阻止整數, 2020, 5(補充 1): 1-11。

[4] IMRAN MM, AHMED U, USMAN U. 中性粒細胞/淋巴細胞比率——新冠肺炎肺炎嚴重程度的標誌。國際臨床實踐雜誌, 2021, 75 : e13698。

[5] SAEEDI P, PETERSOHN I, SALPEA P. 2019 年全球和區域糖尿病患病率估計值以及 2030 年和 2045 年預測: 來自國際糖尿病聯合會糖尿病地圖集第 9 版的結果。糖尿病研究臨床實踐, 2019, 157 : 107843。

[6] PAL R, BHADADA SK. 新冠肺炎和糖尿病: 兩種流行病的邪惡相互作用。糖尿病代謝綜合徵, 2020, 14 (4) : 513-517。

[7] GUPTA, R, HUSSAIN, A, MISRA A. 糖尿病和新冠肺炎: 證據、現狀和未解決的研究問題。歐洲臨床營養學雜誌, 2020, 74: 864-870。

[8] CRITCHLEY JA, CAREY IM, HARRIS T. 大型初級保健隊列研究中 1 型或 2 型糖尿病患者的血糖控制和感染風險。糖尿病護理, 2018, 41(10): 2127-2135。

[9] MADDALONI E, BUZZETTI R. 新冠肺炎和糖尿病: 揭示兩種流行病的相互作用。糖尿病代謝研究修訂版, 2020, e33213321。

[10] HUSSAIN M, BABAR MZM, AKHTAR L 和 HUSSAIN MS. 中性粒細胞/淋巴細胞比率: 2 型糖尿病患者血糖控制的良好評估工具。樸醫學雜誌科學, 2017 年: 33 (6) : 1366-1370。



- [11] LIU L, ZHENG Y, CAI L. 中性粒細胞與淋巴細胞的比率，是評估 新冠肺炎患者疾病嚴重程度的關鍵預測指標。 詮釋 J 實驗室血液學, 2021, 43(2): 329-335。
- [12] KONG M, ZHANG H, CAO X. 較高水平的中性粒細胞-淋巴細胞與嚴重的 新冠肺炎相關。 流行病感染, 2020 年, 148 : e139。
- [13] SIMADIBRATA DM、CALVIN J、WIJAYA AD 等。入院時中性粒細胞與淋巴細胞比值預測 新冠肺炎患者的嚴重程度和死亡率：薈萃分析。 美國急救醫學雜誌, 2021, 42: 60-69。
- [14] LI X, LIU C, MAO Z. 中性粒細胞與淋巴細胞比率對新冠肺炎患者疾病嚴重程度和死亡率的預測值：系統評價和薈萃分析。 暴擊護理, 2020 年, 24 (1) : 647。
- [15] CHAN ABIGAIL、SY ROUTH AMIT。在 新冠肺炎中使用中性粒細胞與淋巴細胞和血小板與淋巴細胞的比率。 臨床醫學雜誌, 2020, 12(7): 448-453。
- [16] KSHANTI IA、EPRILIAWATI M、MOKOAGOW I 等。2019 年冠狀病毒病大流行對印度尼西亞糖尿病患者的影響：一項跨部門的全國規模網絡調查。 2020. 10.1101/2020.12.01. 20241588。
- [17] PERROTTA F、CORBI G、MAZZEO G 等。新冠肺炎和老年人：對發病機制和臨床決策的洞察。 老化臨床實驗研究, 2020, 32(8): 1599-1608。
- [18] LANDI F, BARILLARO C, BELLINI A。老年病學的新挑戰：從 嚴重急性呼吸綜合征冠狀病毒 2 大流行感染中拯救虛弱的老年人。 營養健康老齡化雜誌, 2020, 24(5): 466-470。
- [19] SHI Y, YU X, ZHAO H. 宿主對嚴重 新冠肺炎的易感性和宿主風險評分的建立：武漢以外 487 例病例的發現。 暴擊護理, 2020 年, 24 (1) : 108。
- [20] LI X, XU S, YU M. 武漢成人 新冠肺炎住院患者嚴重程度和死亡率的危險因素。 J 過敏臨床免疫學雜誌, 2020 年, 46 (1) : 110-118。
- [21] DRUMMOND GR、VINH A、GUZIK TJ 和 SOBEY CG。高血壓的免疫機制。 納特免疫師, 2019, 19(8): 517-532。
- [22] CHENG Y, LUO R, WANG K. 腎臟疾病與 新冠肺炎患者的院內死亡有關。 腎臟國際, 2020, 97 (5) : 829-838。
- [23] COBO G。終末期腎病和透析中的慢性炎症。 腎病透析移植, 2018, 33(3) : iii35–iii40。
- [24] ZHENG Z, PENG F, XU B. 危重和致命 新冠肺炎病例的危險因素：系統文獻綜述和薈萃分析。 感染雜誌, 2020 年, 81(2) : e16-e25。
- [25] FU L, WANG B, YUAN T. 中國冠狀病毒病 2019 (新冠肺炎) 的臨床特徵：系統評價和薈萃分析。 期刊感染, 2020, 80(6): 656-665。
- [26] CHANG DE, MO G, YUAN X. 新型冠狀病毒感染中病毒清除和症狀解決的時間動力學。 牛 J 和呼吸急症健康醫學, 2020, 201(9): 1150-1152。
- [27] BAREK MA、AZIZ MA、ISLAM MS。年齡、性別、合併症和臨床症狀對 新冠肺炎病例嚴重程度的影響：一項包含 55 項研究和 10014 例病例的薈萃分析。 赫利永, 2020, 6(12): e05684。
- [28] LIU J, LI Y, LIU Q。嚴重急性呼吸綜合征冠狀病毒 2 細胞嗜性和多器官感染。 細胞發現, 2021 年, 7(17)。
- [29] HANTZIDIAMANTIS PJ, LAPPIN SL。生理學，葡萄糖。在：統計珍珠[互聯網]。金銀島：統計珍珠出版； 2021 年 1 月 -。可從：<https://www.ncbi.nlm.nih.gov/books/NBK545201/>
- [30] HAN Q, WEN X, WANG L. 血液學參數在呼吸道感染症狀患者流感病毒感染診斷中的作用。 臨床實驗室分析雜誌, 2020, 34(5): e23191。
- [31] LIU Y-P, LI G-M, HE J. 聯合使用中性粒細胞與淋巴細胞比率和反應堆來預測 84 名 新冠肺炎肺炎住院患者的 7 天疾病嚴重程度：一項回顧性隊列研究。 安翻譯醫學, 2020, 8(10): 635。
- [32] YANG A-P, LIU J-P, TAO W-Q, LI H-M. 國家廣播電台、國家廣播電台和 磷升電阻 在 新冠肺炎患者中的診斷和預測作用國際免疫藥理學。 2020, 84: 106504。
- [33] SUN S, CAI X, WANG H. 中國溫州 新冠肺炎患者外周血系統異常。 臨床學報, 2020, 507 : 174-180。
- [34] SONG C-Y, XU J, HE J-Q, LU Y-Q. 陸遠強新冠肺炎預警評分：一種用於識別高度疑似患者的多參數篩查工具。 醫學, 2020 年, 10-3。
- [35] PEIRIS S、MESA H、AYSOLA A 等。新冠肺炎患者器官和組織的病理學發現：系統評價。 2021. 公共圖書館一號, 16(4): e0250708。

# Correlation between Neutrophil-to-Lymphocyte Ratio with Disease Severity in Diabetic Patients with COVID-19 at Tertiary Referral Hospital in Indonesia

---

## ORIGINALITY REPORT

---

15%

SIMILARITY INDEX

%

INTERNET SOURCES

15%

PUBLICATIONS

%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1 Nitin Gupta, Kavitha Saravu, Sagar S Maddani, Shashikiran Umakanth, Sowmya Joylin. "Neutrophil-lymphocyte Ratio as a Simple Tool to Predict Requirement of Admission to a Critical Care Unit in Patients with COVID-19", *Indian Journal of Critical Care Medicine*, 2021  
Publication 1%
  - 2 Abdullah Alshukry, Mohammad Bu Abbas, Yaseen Ali, Barrak Alahmad et al. "Clinical characteristics and outcomes of COVID-19 patients with diabetes mellitus in Kuwait", *Heliyon*, 2021  
Publication 1%
  - 3 Aliye Bastug, Hurrem Bodur, Serpil Erdogan, Derya Gokcinar et al. "Clinical and laboratory features of COVID-19: Predictors of severe prognosis", *International Immunopharmacology*, 2020  
Publication 1%
-

4

Ai-Ping Yang, Hui-Ming Li, Wen-Qiang Tao, Xue-Jing Yang, Min Wang, Wen-Juan Yang, Jian-Ping Liu. "Infection with SARS-CoV-2 causes abnormal laboratory results of multiple organs in patients", *Aging*, 2020

Publication

1 %

5

Adekunle Sanyaolu, Chuku Okorie, Aleksandra Marinkovic, Risha Patidar et al. "Comorbidity and its Impact on Patients with COVID-19", *SN Comprehensive Clinical Medicine*, 2020

Publication

1 %

6

Chun Gao, Li Zhu, Cheng Cheng Jin, Yi Xin Tong, Ai Tang Xiao, Sheng Zhang. "Proinflammatory cytokines are associated with prolonged viral RNA shedding in COVID-19 patients", *Clinical Immunology*, 2020

Publication

1 %

7

Hao-Qing Shao, Zhou-Jin Tan. "Microbiota-targeted therapeutics in gastrointestinal diseases", *World Chinese Journal of Digestology*, 2021

Publication

1 %

8

Nurlela Damayanti, Resti Yudhawati. "The Comparison of Pleural Fluid TNF- $\alpha$  Levels in Tuberculous and Nontuberculous Pleural Effusion", *Indian Journal of Tuberculosis*, 2018

Publication

1 %



9

应用激光 应用激光. "Biological Effects of ALA-PDT on PBMCs from Healthy and Psoriatic Patients", Applied laser, 2011

Publication

1 %

10

Haizhou Wang, Yongxi Zhang, Pingzheng Mo, Jing Liu, Hongling Wang, Fan Wang, Qiu Zhao. "Neutrophil to CD4+ lymphocyte ratio as a potential biomarker in predicting virus negative conversion time in COVID-19", International Immunopharmacology, 2020

Publication

1 %

11

Abdul Waris, Misbahud Din, Atiqa Khalid, Raees Abbas Lail et al. "Evaluation of hematological parameters as an indicator of disease severity in Covid - 19 patients: Pakistan's experience", Journal of Clinical Laboratory Analysis, 2021

Publication

<1 %

12

Li-Li Meng, Zhan-Jun Wu. "Clinical utility of fecal neutrophil-derived biomarkers in evaluating activity of inflammatory bowel disease", World Chinese Journal of Digestology, 2014

Publication

<1 %

13

Yue-Ping Liu, Gao-Ming Li, Jun He, Ying Liu, Min Li, Rui Zhang, Ya-Lan Li, Ya-Zhou Wu, Bo Diao. "Combined use of the neutrophil-to-lymphocyte ratio and CRP to predict 7-day

<1 %

disease severity in 84 hospitalized patients with COVID-19 pneumonia: a retrospective cohort study", *Annals of Translational Medicine*, 2020

Publication

---

14

Deasy Ardiany, Agung Pranoto, Soebagijo Adi Soelistijo, Libriansyah, Sauli Ari Widjaja.

"Association between neutrophil-lymphocyte ratio on arterial stiffness in type-2 diabetes mellitus patients: a part of DiORS Study", *International Journal of Diabetes in Developing Countries*, 2021

Publication

---

15

Fang Liu, Lin Li, MengDa Xu, Juan Wu, Ding Luo, YuSi Zhu, BiXi Li, XiaoYang Song, Xiang Zhou. "Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with COVID-19", *Journal of Clinical Virology*, 2020

Publication

---

16

Haerani Rasyid, Alvin Sangkereng, Tutik Harjianti, Audrey S. Soetjipto. "Impact of age to ferritin and neutrophil - lymphocyte ratio as biomarkers for intensive care requirement and mortality risk in COVID - 19 patients in Makassar, Indonesia", *Physiological Reports*, 2021

Publication

---

<1 %

<1 %

<1 %

17

Renato Domingues, Alice Lippi, Cristian Setz, Tiago F. Outeiro, Anita Krisko. "SARS-CoV-2, immunosenescence and inflammaging: partners in the COVID-19 crime", Aging, 2020

Publication

&lt;1 %

18

Yangjing Xie, Qinghai You, Chaoran Wu, Shiyu Cao, Guangbo Qu, Xiaoxiang Yan, Xuan Han, Changhui Wang, Hong Zhang. "Impact of Cardiovascular Disease on Clinical Characteristics and Outcomes of Coronavirus Disease 2019 (COVID-19)", Circulation Journal, 2020

Publication

&lt;1 %

19

Shangwen Pan, Huaqing Shu, Yaxin Wang, Ruiting Li et al. "The Abnormal Imaging of SARS-CoV-2: A Predictive Measure of Disease Severity", Frontiers in Medicine, 2021

Publication

&lt;1 %

20

杜祥琬 杜祥琬. "Effect of low-dose electromagnetic radiation on peripheral blood cells in mouse", High Power Laser and Particle Beams, 2011

Publication

&lt;1 %

21

Hamed Akbari, Reza Tabrizi, Kamran B. Lankarani, Hamid Aria et al. "The role of cytokine profile and lymphocyte subsets in the severity of coronavirus disease 2019

&lt;1 %



(COVID-19): A systematic review and meta-analysis", Life Sciences, 2020

Publication

22

Leithead, Joanna A., Neil Rajoriya, Bridget K. Gunson, and James W. Ferguson. "Neutrophil-to-lymphocyte ratio predicts mortality in patients listed for liver transplantation", Liver International, 2014.

Publication

<1 %

23

Guang Chen, Di Wu, Wei Guo, Yong Cao et al. "Clinical and immunologic features in severe and moderate Coronavirus Disease 2019", Journal of Clinical Investigation, 2020

Publication

<1 %

24

Seong Soo A. An, Yu-Ri Kim, Jong-Il Park, Eun Jeong Lee et al. "Toxicity of 100 nm zinc oxide nanoparticles: a report of 90-day repeated oral administration in Sprague Dawley rats", International Journal of Nanomedicine, 2014

Publication

<1 %

25

Ai-Ping Yang, Jianping Liu, Wenqiang Tao, Hui-ming Li. "The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients", International Immunopharmacology, 2020

Publication

<1 %

26

KW Cheung, SF Ngu, CP Lee. "Fetal fibronectin test on Chinese women with symptoms of

<1 %

preterm labour: a pilot study", Hong Kong Medical Journal, 2013

Publication

---

27

Loris Roncon, Marco Zuin, Gianluca Rigatelli, Giovanni Zuliani. "Diabetic patients with COVID-19 infection are at higher risk of ICU admission and poor short-term outcome", Journal of Clinical Virology, 2020

Publication

---

<1 %

28

Muhammad Miftahussurur, Langgeng Agung Waskito, Ari Fahrial Syam, Iswan Abbas Nusi et al. "

Alternative eradication regimens for *Helicobacter pylori* infection in Indonesian regions with high metronidazole and levofloxacin resistance

", Infection and Drug Resistance, 2019

Publication

---

<1 %

29

Tuoyu Zhou. "SARS-CoV-2 triggered excessive inflammation and abnormal energy metabolism in gut microbiota", Cold Spring Harbor Laboratory, 2021

Publication

---

<1 %

30

TH Rainer, PA Cameron, DV Smit. "Should Differential White Cell Counts be Reported as Percentages or Absolute Counts in Patients with Severe Acute Respiratory Syndrome?",

<1 %

Hong Kong Journal of Emergency Medicine,  
2017

Publication

---

31

Bing-Xu Hou, Li-Ying Feng. "Role of JNK signaling pathway-mediated apoptosis in diseases", World Chinese Journal of Digestology, 2011

Publication

---

<1 %

32

Tim Trausch. "10 Contact Zones and/as Combat Zones: Representing the Nanjing Massacre", Walter de Gruyter GmbH, 2020

Publication

---

<1 %

33

Jian Liu. "Study on Feature Extraction and Feature Selection in Confidence Measure of Speech Recognition", 2009 Chinese Conference on Pattern Recognition, 11/2009

Publication

---

<1 %

34

Tissa Wijeratne, Sheila Gillard Crewther, Carmela Sales, Leila Karimi. "COVID-19 Pathophysiology Predicts That Ischemic Stroke Occurrence Is an Expectation, Not an Exception—A Systematic Review", Frontiers in Neurology, 2021

Publication

---

<1 %

35

Rūta Liutkevičienė, Gediminas Mečėjus, Diana Žilovič, Žana Bumbulienė. "Endometrial biopsy and density of nerve fibers in eutopic endometrium. Looking for easier ways to

<1 %



diagnose endometriosis", Gynecological  
Endocrinology, 2019

Publication

---

36

Shao-rui Hao, Ren Yan, Shan-yan Zhang, Jiang-shan Lian et al. "干扰素- $\alpha$ 2b 喷雾吸入未缩短住院患者的 SARSCoV-2 的脱落时间: 一项初步的配对病例对照研究", Journal of Zhejiang University-SCIENCE B, 2020

Publication

---

<1 %

37

Xiaochen Li, Shuyun Xu, Muqing Yu, Ke Wang et al. "Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan", Journal of Allergy and Clinical Immunology, 2020

Publication

---

<1 %

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On

# Correlation between Neutrophil-to-Lymphocyte Ratio with Disease Severity in Diabetic Patients with COVID-19 at Tertiary Referral Hospital in Indonesia

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---