

PO357 The Number of Metabolic Syndrome's Components Associated with Fibrinogenemia in Patients with Type 2 Diabetes

by Hermina Novida

Submission date: 27-May-2021 01:35PM (UTC+0800)

Submission ID: 1595072565

File name: nskh.pdf (117.97K)

Word count: 714

Character count: 3817

PO357

THE NUMBER OF METABOLIC SYNDROME'S COMPONENTS ASSOCIATED WITH FIBRINOGENEMIA IN PATIENTS WITH TYPE 2 DIABETES

H. Novida¹, A. Tjokroprawiro¹, A. Pranoto¹, A. Sutjahjo¹, S.Murtiwi¹, S. Adi¹, S. Wibisono¹.

¹ Surabaya Diabetes and Nutrition Center Dr. Soetomo Teaching Hospital-Faculty of Medicine Airlangga University, Surabaya, Indonesia

Background: Metabolic syndrome (MetS) is a cluster of cardiovascular disease risk factors that the prevalence increased in developing country. Several studies have demonstrated that this syndrome strongly predicts cardiovascular disease. Recently, close association of MetS with haemostatic abnormalities including plasma fibrinogen has been reported. Plasma fibrinogen is a marker of inflammation and it is considered to be one of the predictors of coronary artery disease. The aim of this study was to determine the prevalence of MetS and to analyze the association between the number of MetS components with fibrinogenemia in patients with type 2 diabetes.

Method: We analyzed 100 patients with type 2 diabetes consisting of 60 male and 40 female patients using cross sectional observational design. Blood pressure, body weight, height and waist circumference (WC) were measured. We also obtained fasting plasma glucose (FPG), HbA1c, total cholesterol (TC), low density lipoprotein cholesterol (LDL-C), high density lipoprotein cholesterol (HDL-C), triglyceride (TG) and fibrinogen level from blood venous samples. MetS was defined according National Cholesterol Education Program Adult Treatment Panel (NCEP ATP) III definition when 3 of 5 components were present, the diagnosis MetS could be established. The components were abdominal obesity (men >90 cm, women >80 cm), blood pressure $\geq 130/\geq 85$ mmHg, FPG ≥ 100 mg/dL, TG ≥ 150 mg/dL and HDL-C (men >400mg/dL. Data was statistically analyzed using logistic regression test.

Result: There were 75 patients had at least 3 components of MetS that the prevalence of MetS in this study based on NCEP ATP III definition was 75%. The mean of WC in these 100 patients was 99.21 ± 10.28 cm, FPG was 181.31 ± 79.60 mg/dL, HbA1c was $8.67 \pm 2.32\%$, TC was 199.02 ± 50.62 mg/dL, LDL-C was 123.83 ± 41.16 mg/dL, HDL-C was 44.93 ± 10.69 mg/dL and TG was 196.53 ± 18.30 mg/dL. The overall mean of fibrinogen level was 381.08 ± 123.07 mg/dl, while 69% patients were normofibrinogenemia and 31% were hyperfibrinogenemia. There was only one patient (1%) patient had one MetS component with fibrinogen level 241mg/dL, 24 patients (24%) had two MetS components with the average fibrinogen level was 345.62 ± 105.22 mg/dL, 44 patients (44%) had three MetS components with the average fibrinogen level was 359.32 ± 92.47 mg/dL, 26 patients (26%) had four MetS components with the average fibrinogen level was 423.00 ± 123.92 mg/dL and 5 patients (5%) had five MetS components with the average fibrinogen level was 380.45 ± 224.54 mg/dL. There was significant association between the number of MetS components and fibrinogen level (p 0.023; p < 0.05).

Conclusion: Prevalence of MetS in this study was 75% based on NCEP ATP III definition and there was significant association between the number of MetS components and fibrinogenemia in patients with type 2 diabetes

Reference(s)

- Bosevski M, Bosevska G, Stojanosvka (2013). Influence of Fibrinogen and CRP on Progression of Peripheral Arterial Disease in Type 2 Diabetes: A Preliminary Report. *Cardiovascular Diabetology* 12, 29–34.
- Brenner DR, Arora P, Garcia-Bailo B, Morrison H, El-Sohemy A, Morrison H et al (2011). The Relationship between Inflammation, Metabolic Syndrome and Markers of Cardiometabolic Disease among Canadian Adults. *J Diabetes Metab S: 2*.
- Grundy SM, Brewer B, Cleeman JI, Smith SC, Lenfant C (2004). Definition of Metabolic Syndrome: Report of National Heart, Lung and Blood Institute/American Heart Association Conference on Scientific Issues Related to Definition. *Circulation* 109, 433–438.
- Iyer UM, Desai P (2010). Assessment of C-Reactive Protein and Fibrinogen Levels in Type 2 Diabetes Mellitus. *Biomedical Research* 21, 208–213.
- Kamath S, Lip GYH (2003). Fibrinogen: Biochemistry, Epidemiology and Determinants. *Q J Med* 96, 711–729.
- Maple-Brown LJ, Cunningham J, Nandi N, Hodge A, O’Dea K (2010). Fibrinogen and Associated Risk Factors in A High-Risk Population: Urban Indigenous Australians, the Druid Study. *Cardiovascular Diabetology* 9, 69–76.
- Remkova, DJ (2007). Prothrombotic State in Metabolic Syndrome. *Bratisl Lek Listy* 108, 279–280.

PO357 The Number of Metabolic Syndrome's Components Associated with Fibrinogenemia in Patients with Type 2 Diabetes

ORIGINALITY REPORT

16%

SIMILARITY INDEX

%

INTERNET SOURCES

16%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Mohammad hossein Somi, Zeinab Nikniaz, Mohammad Asghari Jafarabadi, Amir Taher Eftekharsadat et al. "Association Between Diabetic Retinopathy, Dietary Inflammatory Index and Metabolic Syndrome in Azar Cohort Study", Research Square, 2020
Publication 4%
- 2 Milos Maksimovic, Hristina Vlajinac, Djordje Radak, Jelena Marinkovic, Jadranka Maksimovic, Jagoda Jorga. "Association of overweight and obesity with cardiovascular risk factors in patients with atherosclerotic diseases", Journal of Medical Biochemistry, 2019
Publication 4%
- 3 Tjokroprawiro, A., P.Z. Romadhon, S. Murtiwi, S. Adi, A. Pranoto, A. Suthahjo, and S. Wibisono. "PO345 ADIPONECTIN IS INVERSELY CORRELATED WITH LIPOPROTEIN(A) IN TYPE 2 DIABETES 4%

MELLITUS", Diabetes Research and Clinical Practice, 2014.

Publication

4

"OPENING LECTURE", Gynecological Endocrinology, 2009

Publication

2%

5

Soo Bin Lee, Hyeok Chan Kwon, Jung Yoon Pyo, Mi Il Kang, Jason Jungsik Song, Yong-Beom Park, Jun Yong Park, Sang-Won Lee. "Clinical implication of metabolic syndrome in nonobese patients with antineutrophil cytoplasmic antibody-associated vasculitis", Research Square, 2021

Publication

2%

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On