

# The Effect of Kebar Grass (*Biophytum petersianum* Klotzsch) Extract on the Number of Leydig Cells of Mice (*Mus Musculus*) Treated with TCDD

*by* Hani Plumeriastuti

---

**Submission date:** 05-Jun-2023 09:09AM (UTC+0800)

**Submission ID:** 2108943323

**File name:** mber\_of\_Leydig\_Cells\_of\_Mice\_Mus\_Musculus\_Treated\_with\_TCDD..pdf (603.29K)

**Word count:** 2990

**Character count:** 16149

Research Article

## The Effect of Kebar Grass (*Biophytum petersianum* Klotzsch) Extract on the Number of Leydig Cells of Mice (*Mus Musculus*) Treated with TCDD

MILLA NURSADIDA<sup>1</sup>, HANI PLUMERASTUTI<sup>2</sup>, YENI DHAMAYANTI<sup>3</sup>, WIDJIATI<sup>4</sup>, EPY MUHAMMAD LUQMAN<sup>5\*</sup>

<sup>1,3,4,5</sup>Departement of Anatomy Veteriner, Faculty of Veterinary Medicine Universitas Airlangga, Jalan Mulyorejo, Surabaya-60115 Indonesia

<sup>2</sup>Departement of Pathology Veteriner, Faculty of Veterinary Medicine Universitas Airlangga, Jalan Mulyorejo, Surabaya-60115 Indonesia

\*Corresponding Author

Email ID: epy-m-l@fkh.unair.ac.id

Received: 05.04.20, Revised: 05.05.20, Accepted: 05.06.20

### ABSTRACT

This study aimed to examine the effect of kebar grass extract on the Leydig cells number of mice to exposed 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD). These animals were divided into five groups. Twenty five BALB/C mice were used with five groups: Negative Control (C-), Positive Control (C+) with TCDD 7µg/kgBW IP, and treatment groups that treated with TCDD 7µg/kgBW IP and Kebar Grass orally for 53 days were presented T1 0,045mg/gBW/day, T2 0,080mg/gBW/day, and T3 0,135mg/gBW/day. Results showed that TCDD decreased the Leydig cells number, while kebar grass extract was increase the number of Leydig cells exposed by TCDD significantly (p<0,05). But the increase in the number of Leydig cells had not yet reached the normal number of Leydig cells.

**Keywords:** Kebar Grass, *Biophytumpetersianum* Klotzsch, Leydig cells, TCDD, mice

### INTRODUCTION

The environmental contaminant, such as industrial processes like incineration of medical wastes, chlorine bleaching of paper and pulp and manufacture of pesticides can be called endocrine-disrupting by Dioxin [1]. The dioxin 2,3,7,8-tetrachlorodibenzo-dioxin (TCDD) has been intensively studied and is known to be the most toxic compound among these compounds [2]. Those studies demonstrated that TCDD altered the process of testicular steroidogenesis and caused a reduction of Leydig cell volume and number [3].

TCDD has been reported to induce superoxide anion, lipid peroxidation and DNA damage. Induction of oxidative stress upon exposure to TCDD are considered to be an important mechanism for the toxic effects of TCDD [4]. TCDD increases reactive oxygen species (ROS) in endothelial cells by the induction of CYP1A1 [5]. TCDD toxicity is very dangerous because it causes oxidative stress, therefore we need a compound that can break TCDD toxicity such as antioxidant compounds [6]. Antioxidants defend against excessive ROS levels through enzymatic (superoxide dismutase, catalases, and peroxidases) and non-enzymatic (vitamins, steroids etc.) mechanisms [7].

Antioxidant compounds are found in various plants, one of which is Kebar Grass (*Biophytum petersianum* K) which has a role in protecting against free radical attacks. The usage of kebar grass as a fertile fertilizer had been carried out in Papua Indonesia for a long time [8]. Kebar grass contains three types of chemical compounds flavonoids, saponins, and tannins that have the potential to give effect to the process of spermatogenesis. The administration of 5% kebar grass infusion could significant increase spermatogenesis activity [9]. The active compounds of kebar grass act as antioxidants by inhibiting the transcription of CYP1A1 cytosolic enzymes and steroid dehydrogenase genes through inhibition of AhR, as well as reducing the competitive binding of TCDD to AhR [10].

Research on kebar grass extract so far is not widely known, therefore the researchers wanted to observe the effect of kebar grass extract to the Leydig cell number in mice exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin to observe the potential kebar grass in counteracting the toxic effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin. In the present study, we have sought to investigate whether the effect of Kebar grass extract in multiple dosages can increase the number of Leydig cells exposed by 2,3,7,8-tetrachlorodibenzo-p-dioxin.

## MATERIALS AND METHODS

This research was conducted at the Laboratory of Experimental Animal Faculty Veterinary Medicine Universitas Airlangga the preparation of the histopathology was conducted at the Department of Pathology Faculty Veterinary Medicine of Universitas Airlangga. Ethical clearance was obtained from the Faculty Veterinary Medicine Universitas Airlangga.

### Materials

This research used 25 male mice (*Mus musculus*) 12 weeks old with the body weights between 25-35 grams obtained from Veterinaria Farma Surabaya. Apparatus that was required, 5 groups contain 5 mice, food and drink, per oral needle, 1 cc syringe, scalpel, a petri dish, surgical scissors, and microscope. Materials that have been used were 2,3,7,8-tetrachlorodibenzo-p-dioxin, Kebar Grass extract, distilled water, 10% NaCl 0.5%, and formalin. The dosage of the 2,3,7,8-tetrachlorodibenzo-p-dioxin which has been used for the rats was 50 µg/KgBW. Then it was converted to the dosage of mice  $50 \times 0.14 = 7 \mu\text{g/KgBW}$ . The dosage of Kebar Grass extract was by using three different dosage of 0,045 mg/gBW/day, 0,08 mg/gBW/day, and 0,135 mg/gBW/day [9].

### Methods

Experimental animals were obtained using simple random sampling procedure. There are five treatment groups (5 mice each group): C (-) was treated with a placebo, C (+) was exposed by TCDD 7 µg/KgBW single dose intraperitoneal injection. T (1) group was exposed by TCDD

7 µg/KgBW single dose intraperitoneal injection and treated by Kebar grass extract was given orally at a dosage 0,045 mg/kg BW/day. T (2) was exposed by TCDD 7 µg/KgBW single dose intraperitoneal injection and treated by kebar grass extract was given orally at a dosage 0,080 mg/kg BW/day. T (3) was exposed by TCDD 7 µg/KgBW single dose intraperitoneal injection and treated by Kebar grass extract was given orally at a dosage 0,135 mg/kg BW/day. The treatment were conducted for 53 days consecutively. The testicular examination was conducted by hematoxylin and eosin (HE) staining (Hematoxylin Staining for Millicell®-HA, Merck, Germany) and observed from five different fields by using a light microscope with 400x magnification (Olympus® CX-41). Data analysis was done by using the ANOVA test. Then continued with the Duncan test to determine differences in each group. Data were analyzed by using the Statistical Analysis Software program (SPSS) version 22.


## RESULTS

Table 1. showed a decrease in the mean of Leydig cells number in the positive control group (C+) compared to negative control (C-). Furthermore, in groups T1, T2, and T3 there was a gradual increase in Leydig cells number. The increase in Leydig cells number that occurred in all treatment groups reached the highest size in the T3 group, which was 24.24. The result of the effect Kebar grass extract on the number of Leydig cells exposed to TCDD by calculating the normal Leydig cells in the mice's testes listed in Table 1.

**Table 1: Amount of Leydig cells in each group.**

Groups	Amount of Leydig cells (Mean ± SD)
C (-) Placebo	43,04 <sup>e</sup> ± 3.36
C (+) exposed TCDD 7 µg/KgBw single dose intraperitoneal injection	4,08 <sup>a</sup> ± 0.78
T (1) TCDD 7 µg/KgBw single dose intraperitoneal injection and Kebar grass extract at a dosage 0,045 mg/gBw/day	11,8 <sup>b</sup> ± 2.31
T (2) TCDD 7 µg/KgBw single dose intraperitoneal injection and Kebar grass extract at a dosage 0,080 mg/gBw/day	16,2 <sup>c</sup> ± 4.44
T (3) TCDD 7 µg/KgBw single dose intraperitoneal injection and Kebar grass extract at a dosage 0,135 mg/gBw/day	24,24 <sup>d</sup> ± 13.83

Description: Different superscript in the same column indicate significant differences (P<0.05).



**Fig.1: Overview of testicular microscopic image (Hematoxylin Eosin staining; magnification of 400x; Olympus® CX-41). Image of group C (-) indicates a normal Leydig cells. Image of group C (+) which induced by TCDD showed most decrease of number Leydig cells. Image of group T (1) showed improvement of Leydig cells. Image of group T (2) many cells showed normal shape of Leydig cells still look some cells died. Image of group T (3) showed the most normal cells between another treatment groups.**

## DISCUSSION

In group C (-), there were more normal Leydig cells than Leydig cells which occurred necrosis. Contrary to the C (+) group exposed to TCDD, the normal number of Leydig cells experienced a significant decrease. On histopathological features of Leydig cells exposed to TCDD given various doses of kebar grass extract increased significantly compared to the C (+) group. C(+) group that exposed by TCDD 7 $\mu$ /KgBW single dose intraperitoneal injection proved that TCDD exposure affected the number of Leydig cells due to highly toxicity (Figure 1). The results of the research in this group similar with [3] that hazardous chemicals can cause antiandrogenic or estrogenic disorders. Antiandrogenic disorders can cause infertility by suppressing androgen production in Leydig cells, reduce the number of Leydig cells, induce Leydig cell apoptosis, or increase aryl hydrocarbon (AhR) to bind to TCDD thereby blocking androgen activity [11]. TCDD binds, Arylhydrocabon Receptor (AhR), and Aryl Hydrocarbon Receptor Nuclear Translocator (ARNT) will bind to certain DNA, namely Dioxin-Responsive Enhancer Elements (DRE), the bond will change the expression of various genes including cytochrome P450 in the nucleus [10]. Increased cytochrome P450 can affected on Leydig cells through a receptor called Androgen Receptor (AR) which plays a role in the

maturation of the steroidogenesis pathway [12]. Cytochrome P450 bond with Androgen Receptor (AR) is known to increase Reactive Oxygen Species (ROS) production, so that it can trigger an increase in superoxide radicals which could cause membrane lipid peroxidase [13]. However, an increase in cytochrome P450 results in the formation of cytochrome P450 bonds with AR which could reduce the action of AR on Leydig cells and increase ROS formation in cells so that death occurs in Leydig cells [14].

Furthermore, in groups T1, T2, and T3 there was a gradual increase in Leydig cells number. The treatment group that exposed by TCDD single dose 7 $\mu$ /KgBW intraperitoneal injection and treated by kebar grass extract at dosage of 0.135 mg/kg BW/day had the most significant number of normal Leydig cells was 24.24 when compared with T (1) and T (2) that treated with kebar grass extract 0,045 mg/gBw/d and 0,080 mg/gBw/day. This shows that kebar grass extract can increase the number of Leydig cells that exposed by TCDD.

Kebar grass (*Biophytum petersianum* Klotzsch) was a plant belonging to the family group Oxalidaceae found in Kebar District, West Papua Indonesia [15]. Kebar grass contains flavonoids, retinol, tocopherol and saponins [8]. Flavonoids that had a primary antioxidant function because they were free radical acceptors so they could inhibit free



radical chain reactions in lipid oxidation that could prevent membrane damage [16]. Polyphenol compounds such as flavonoids work by changing  $H_2O_2$  into  $H_2O$  and  $O_2$  [17]. The content of retinol (vitamin A) in Kebar grass functions as an antioxidant. Beta carotene works by reacting with free radicals and causing free radicals to become stable. Beta carotene (retinol) works together with tocopherol contained in Kebar grass and ascorbic acid (vitamin C). Furthermore ascorbic acid which turns into radicals is stabilized by the natural antioxidant glutathione [6]. The content of tocopherol (vitamin E) in Kebar grass prevented free radicals by giving H atoms from hydroxyl groups to lipid peroxy radicals. Radicals formed from alpha-tocopherol would be stabilized through electron delocalization in the aromatic ring [18]. Vitamin E can reduce TCDD-induced AhR activation so that it can be used as prevention and treatment of acute or chronic TCDD intoxication [12]. This can be strengthened by research conducted by [19] which proved that vitamin E has an antagonistic effect on TCDD toxicity in the process of spermatogenesis. Saponins included in the steroid group compound in an acidic stomach will break the sugar portion, so that it can have an effect for increasing the in free sterols. These sterol compounds are the basic ingredients of testosterone (pregnenolone) [20]. From this explanation it can be seen that saponins can increase testosterone levels which accelerate the Leydig cells number.

In this study, there was a significant decrease in the positive control group (C+) in Leydig cells number when compared to negative controls (C-) due to TCDD exposure. Whereas in T1, T2, and T3 there was a significant increase in Leydig cells number each group. This proves that the administration of kebar grass extract at a dose of 0.045mg/g BB/day in group T1 can significantly increase Leydig cells number. In the T2 and T3 groups there was a significant increase in Leydig cells number when compared with positive controls. The three treatment groups differed significantly with C- which proved that all the doses cannot maintain the Leydig cells number. An increase in the number of Leydig cells had not yet reached the normal number of Leydig cells. Higher doses and longer doses can be used to achieve optimal results. The implication of oxidative stress in the etiology of several chronic and degenerative diseases suggests that antioxidant therapy represents a promising avenue for treatment. In the future, a therapeutic strategy to increase the antioxidant capacity of cells may be used to fortify the long term effective treatment [21].

## CONCLUSION

Based on research that has been done, it can be concluded that the Kebar Grass extract could increase the amount of Leydig cells which were exposed by 2,3,7,8-tetrachlorodibenzo-p-dioxin, although the increase in Leydig cells had not reached a normal amount (C-).

**Conflict of interests:** The authors declare that there is no conflict of interests regarding the publication of this article.

**Source of funding:** The authors express sincere thanks to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia and Universitas Airlangga for funding research (Number 1408/UN3/2019).

## REFERENCES

1. Dhanabalan S, Mathur PP, Latha P. TCDD and Corticosterone on Testicular Steroidogenesis and Antioxidant System of Epididymal Sperm in Rats. *Toxicol Ind Health*. 2015; 31(9): 811-822.
2. Yoshida R, Ogawa Y. Review Article Oxidative Stress Induced by 2,3,7,8-Tetrachlorodibenzo-p-Dioxin: An Application of Oxidative Stress Markers to Cancer Risk Assessment of Dioxins. *National Institute of Industrial Health*. 2000; 21-1.
3. Lai KP, Wong MH, Wong CKC. Inhibition of CYP450scc Expression in Dioxin-Exposed Rat Leydig Cells. *J. Endocrinol*. 2005; 185: 519-527.
4. Latchoumycandane C, Chitra KC, Mathur PP. 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD) Induces Oxidative Stress in The Epididymis and Epididymal Sperm of Adult Rats. *Arch Toxicol*. 2003;7: 280-284.
5. Kopf PG, Walker MK. 2,3,7,8-tetrachlorodibenzo-p-dioxin increases reactive oxygen species production in human endothelial cells via induction of cytochrome P4501A1. *Toxicol Appl Pharmacol*. 2010; 15;245(1):91-9. DOI: 10.1016/j.taap.2010.02.007.
6. Sayuti K, Yenrina R (2015). *Natural and Synthetic Antioxidants*. Padang, Andalas University Press. Padang Indonesia.
7. Darbandi S, Darbandi M. Lifestyle modifications on further reproductive problems. *Cresco J Reprod Sci*. 2016;1(1):1-2.
8. Sembiring B, Darwati I. Identification of chemical component of kebar grass accession (*Biophytum petersianum*) from Papua and Java. *Bul. Littro*, 25(1): 37-44. DOI: 10.21082/bullittro.v25n1.2014: 37-44.
9. Lefaan PN. Effect of Kebar Grass (*Biophytum petersianum*) Infusion on Spermatogenesis of Mice (*Mus musculus*). *Jurnal Sains Veteriner*. 2014; 32: 55-67.
10. Yin HP, Xu JP, Zhou XQ, Wang Y. Effect of Vitamin E on Reproductive Hormones and Testis Structure in Chronic Dioxin-Treated Mice. *J.*

- Toxicol and Industrial Health, 2012; 28 (2): 152-160.
11. Chen BB, Zirkin BR, Ge RS. the Leydig Cell as a Target for Toxicants. Elsevier. USA. 2010; PP: 133-142.
  12. Chang HJ, Park JS, Lee EK, Kim MH, Baek MK, Kim HR, Jeong HG, Choi SY, and Jung YD. Ascorbic Acid Suppresses the 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-induced CYP1A1 expression in Human HepG2 Cells. J. Toxicology In Vitro, 2009; 23: 972-979. DOI: 10.1016/j.tiv.2009.02.008
  13. Dorbrzynski MI, Calkosinski I, Przywitowska JK, Brzoza AC, Waszkiewicz ES, and Parulska O. Effects of dioxins in environmental Pollution on development of tooth disorders. Polish J. of environ.2009; 18(3): 319-323.
  14. Karmilawati W. Vitamin E Potential Against Spermatogenic Cell Counts, Seminiferous Tubule Diameter, and Testosterone Levels in Mice Exposed 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Tesis Ilmu Biologi Reproduksi Universitas Airlangga. 2013.
  15. Lisangan MM, Syarif R, Rahayu WP, Dharmaputra OS. Effect of kebar grass (*Biophytum petersianum* Klotzsch) leaf extract on the growth and morphological structure of aflatoxigenic *Aspergillus flavus*. Food Research. 2019; 4(1):234-243. DOI: 10.26656/fr.2017.4(1).129.
  16. Choe E, Min DB. Mechanisms of Antioxidants in the Oxidation of Foods. Comprehensive Reviews in Food Science and Food Safety, 2009; 8: 345-358
  17. Sroka Z and Cisowski W. Hydrogen Peroxide Scavenging, Antioxidant and Anti-Radical Activity of Some Phenolic Acids. Food and Chemical Toxicology. 2003; 41(6):753-8. DOI: 10.1016/S0278-6915(02)00329-0
  18. Pokorny J, Yanishlieva N, Gordon M. Antioxidant in Food; Practical Applications, CRC Press. New York. 2001. //https.books.google.co.id. [22 Oktober 2019].
  19. Wati WK, Wurlina, Sarmanu. Potential of vitamin e on spermatogenic cell number in mice treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Veterinaria Medika 2014; 7(3):224 – 231.
  20. Winarni D. Effects of Javanese and Korean Ginseng Root Extracts on Male Mouse Libido in Low Testosterone Preconditions. Berk. Penel. Hayati, 2007;12: 153–159.
  21. Pham-Huy LA, He H, and Pham-Huy C. Free Radicals, Antioxidants in Disease and Health. Int J Biomed Sci. 2008; 4(2): 89–96.

# The Effect of Kebar Grass (*Biophytum petersianum* Klotzsch) Extract on the Number of Leydig Cells of Mice (*Mus Musculus*) Treated with TCDD

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

15%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://hdl.handle.net">hdl.handle.net</a> Internet Source	2%
2	<a href="http://digital.car.chula.ac.th">digital.car.chula.ac.th</a> Internet Source	1%
3	C. Latchoumycandane, K. C. Chitra, P. P. Mathur. "2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) induces oxidative stress in the epididymis and epididymal sperm of adult rats", <i>Archives of Toxicology</i> , 2003 Publication	1%
4	<a href="http://www.jdas.in">www.jdas.in</a> Internet Source	1%
5	Stohs, Sidney J., and Ezdihar A. Hassoun. "Dioxin-Activated AHR: Toxic Responses and the Induction of Oxidative Stress", <i>The AH Receptor in Biology and Toxicology Pohjanvirta/AH Receptor</i> , 2011. Publication	1%

6	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1 %
7	<a href="http://www.ijpsr.info">www.ijpsr.info</a> Internet Source	1 %
8	K. Svechnikov, G. Izzo, L. Landreh, J. Weisser, O. Söder. "Endocrine Disruptors and Leydig Cell Function", <i>Journal of Biomedicine and Biotechnology</i> , 2010 Publication	1 %
9	<a href="http://coek.info">coek.info</a> Internet Source	1 %
10	<a href="http://allie.dbcls.jp">allie.dbcls.jp</a> Internet Source	1 %
11	<a href="http://www.inabj.org">www.inabj.org</a> Internet Source	1 %
12	"Pollutant Diseases, Remediation and Recycling", Springer Science and Business Media LLC, 2013 Publication	1 %
13	<a href="http://journals.sagepub.com">journals.sagepub.com</a> Internet Source	1 %
14	Yasuhito Shimada, Shisei Kuninaga, Michiko Ariyoshi, Beibei Zhang et al. "E2F8 promotes hepatic steatosis through FABP3 expression in diet-induced obesity in zebrafish", <i>Nutrition &amp; Metabolism</i> , 2015	1 %



---

15	<a href="https://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	1 %
16	<a href="http://journal.unair.ac.id">journal.unair.ac.id</a> Internet Source	1 %
17	<a href="http://www.pgpa.ufrpe.br">www.pgpa.ufrpe.br</a> Internet Source	1 %
18	<a href="http://www.guamagentorange.info">www.guamagentorange.info</a> Internet Source	<1 %
19	Apresio K. Fajrial, Adhitya G. Saputro, Mohammad K. Agusta, Febdian Rusydi, Nugraha Nugraha, Hermawan K. Dipojono. "First principles study of oxygen molecule interaction with the graphitic active sites of a boron-doped pyrolyzed Fe-N-C catalyst", Physical Chemistry Chemical Physics, 2017 Publication	<1 %
20	Sayed M. Rawi, Fatma M. Seif Al Nassr. "Zinc sulphate and vitamin E alleviate reproductive toxicity caused by aluminium sulphate in male albino rats", Toxicology and Industrial Health, 2013 Publication	<1 %
21	<a href="https://oaspub.epa.gov">oaspub.epa.gov</a> Internet Source	<1 %
22	<a href="http://repositorio.ufsm.br">repositorio.ufsm.br</a> Internet Source	

---

<1 %

23

B Agustono, D A Agustin, E K Sabdoningrum, M N Yunita, R T Dewi, I R T Ivani. "Potential feed substitute of Cirripedia sp. flour on body weight gain, feed conversion ratio, feed consumption of buck", IOP Conference Series: Earth and Environmental Science, 2022

Publication

<1 %

24

[biofarmaka.ipb.ac.id](http://biofarmaka.ipb.ac.id)

Internet Source

<1 %

25

[doaj.org](http://doaj.org)

Internet Source

<1 %

26

[forikes-ejournal.com](http://forikes-ejournal.com)

Internet Source

<1 %

27

Rr Dian Tristiana, Ah Yusuf, Rizki Fitryasari, Sylvia Dwi Wahyuni, Hanik Endang Nihayati. "Perceived barriers on mental health services by the family of patients with mental illness", International Journal of Nursing Sciences, 2018

Publication

<1 %

28

Aly, H.A.A.. "2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)-induced cytotoxicity accompanied by oxidative stress in rat Sertoli cells: Possible role of mitochondrial fractions

<1 %

of Sertoli cells", Toxicology and Applied  
Pharmacology, 20110501

Publication

---

29

Ramy K. A. Sayed, Doaa M. Mokhtar, Marisol  
Fernández-Ortiz, José Fernández-Martínez et  
al. "Lack of retinoid acid receptor-related  
orphan receptor alpha accelerates and  
melatonin supplementation prevents  
testicular aging", Aging, 2020

<1 %

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# The Effect of Kebar Grass (*Biophtyum petersianum* Klotzsch) Extract on the Number of Leydig Cells of Mice (*Mus Musculus*) Treated with TCDD

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---