

# A Review of Antimicrobial Resistance (AMR) of Escherichia coli on Livestock and Animal Products: Public Health Importance

*by* Hani Plumeriastuti Plumeriastuti

---

**Submission date:** 05-Jun-2023 09:15AM (UTC+0800)

**Submission ID:** 2108948686

**File name:** n\_Livestock\_and\_Animal\_Products,\_Public\_Health\_Importance\_1.pdf (1.08M)

**Word count:** 8250

**Character count:** 46078

# A Review of Antimicrobial Resistance (AMR) of *Escherichia coli* on Livestock and Animal Products: Public Health Importance

Havyun Durrotul Faridah<sup>1</sup>, Erna Kristiana Dewi<sup>1</sup>, Fatimah<sup>1</sup>, Mustofa Helmi Effendi<sup>2\*</sup>, Hani Plumeriastuti<sup>3</sup>

<sup>1</sup>Department of Biology, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia

<sup>2</sup>Department of Veterinary Public Health, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

<sup>3</sup>Department of Veterinary Pathology, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

\*Corresponding author: Mustofa Helmi Effendi, Department of Veterinary Public Health, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia, Email: [mheffendi@yahoo.com](mailto:mheffendi@yahoo.com)

## ABSTRACT

Antimicrobial resistance (AMR) has become a major challenge for the world of health in this century. This resistance event is closely related to the use of antibiotics that are not in accordance with the rules in both the hospital sector and the livestock sector. Antibiotics have an important role in treating infectious diseases. The role of antibiotics is expanding not only as a treatment but also to prevent disease and as a growth promoter in livestock. *Escherichia coli* has been widely used to monitor resistance in livestock and food of animal origin because these microbes can be found in the digestive tract of warm-blooded animals. Several strains of *Escherichia coli* are potential sources of resistant genes that can be transmitted to humans. Livestock and food of animal origin such as poultry, pigs, cattle, sheep, eggs, and milk have become reservoirs for resistant *Escherichia coli*. It is not only resistant to one type of antibiotic but has also become multidrug resistant, and also become public health problem. *Escherichia coli* has different resistance mechanisms depending on the type of antibiotic. This resistance mechanism is also encoded by resistance genes which can be transmitted horizontally. Resistant bacteria encode different resistance genes for each antibiotic class. There needs to be strict regulations and supervision to limit the use of antibiotics in the livestock sector so that it can reduce cases of antibiotic resistance in the world.

**Keywords:** *Escherichia coli*, Antimicrobial resistance, Livestock, Animal products, Public health

## Correspondence:

Mustofa Helmi Effendi

Department of Veterinary Public Health, Faculty of Veterinary Medicine,

Airlangga University, Surabaya, Indonesia.

Email: [mheffendi@yahoo.com](mailto:mheffendi@yahoo.com)

## INTRODUCTION

Antimicrobial resistance (AMR) was introduced more than fifty years ago and is the biggest challenge of this century. Resistance of pathogenic bacteria to antibiotics is one of the major public health problems of the 21st century (1). This resistance is a growing public health threat throughout the world (2). Microbes can simultaneously be resistant to several groups of antibiotics. This event is called multidrug-resistant, ie when the microbes are resistant to at least 3 types of antibiotics (3). This incident is associated with the use of antibiotics that are not according to the rules. Such as excessive use of antibiotics and errors in diagnosing diseases. Not only because of its use in humans but also on farm animals (4). The use of antibiotics is expected to continue to increase rapidly in the coming years due to the intensification of the livestock sector in many developing countries (5). Animal husbandry is one of the sectors that has the potential as a reservoir for resistant microbes (4). The use of antibiotics in animal husbandry is directly related to the cases of resistant bacteria in food of animal origin (6, 7). In the poultry sector, breeders in various countries use antibiotics to raise poultry (8, 9). Antibiotics are given orally to treat livestock infected with pathogenic bacteria, prevent livestock from disease, and to increase productivity (10). The indiscriminate use of antimicrobials in livestock has the opportunity to accelerate the development of resistant pathogenic microbes as well as in commensal organisms (11). Apart from concerns about the emergence of AMR in bacteria from poultry production, there are also human health concerns about the presence of antimicrobial residues in foods of animal

origin such as milk (11), meat (12), and eggs (13). In addition, AMR in poultry pathogens is likely to result in economic losses. Infectious diseases in livestock that are not handled and due to the purchase of antibiotics that are not effective in treating them (14).

The case of resistance to the gram-negative group of bacteria is very worrying because of the limited choice of antibiotics to treat infections caused by several organisms such as Enterobacteriaceae, *Pseudomonas aeruginosa*, and *Acinetobacter* which have become resistant to almost all available antimicrobials, including carbapenems (4). The Enterobacteriaceae group is a microbe that is clinically fast growing agents to become resistant to available antimicrobials (15). Cases of microbial resistance to antibiotics that are widely reported in livestock and food of animal origin include *Escherichia coli*. Resistant *Escherichia coli* was found in chickens in Ghana (16), pork and beef in Mexico (17), chicken in Pakistan (18), chicken in France (19) and chicken in Indonesia (11).

## REVIEW

### *Escherichia coli*

*Escherichia coli* is a member of the Enterobacteriaceae family in the Gammaproteobacteria class (20) which is commonly found living in human and animal digestion as normal microflora (21). *Escherichia coli* includes Gram negative bacteria which are facultative anaerobes (22). These bacteria are Gram-negative bacteria with short rods, measuring 2.4 x 0.4-0.7 µm, motile with peritrichous flagella, and without spores and including opportunistic pathogens which always show increased resistance to various antibiotics. These bacteria have the highest

incidence of urinary tract infections (UTIs) (23). *Escherichia coli* is able to reproduce itself in approximately twenty minutes (20).

*Escherichia coli* is one of the bacteria that has been studied in depth. Sequence analysis of the *Escherichia coli* genome was first reported in 1997. Since then, more than 4800 *Escherichia coli* genomes have been sequenced (23). *Escherichia coli* can replicate in every 20 minutes. This rapid growth makes it suitable for observations of the evolution of microorganisms and research on long-term experimental evolution of more than 50,000 generations (24).

*Escherichia coli* has been widely used to monitor AMR in livestock and food of animal origin. This is because *Escherichia coli* can be found in the digestive tracts of warm-blooded animals (25). In addition, several strains of *Escherichia coli* are potential sources of the AMR gene which can be transmitted to humans through various means (21, 22). *Escherichia coli* carried through feces or treatment of wastewater that is disposed of into waterways can pollute the environment (26). The concentration of *Escherichia coli* per gram of feces varies across host species, typically reaching 10<sup>7</sup>–10<sup>9</sup> in humans and 10<sup>4</sup>–10<sup>6</sup> in domestic animals (27).

#### VIRULENCE FACTORS OF *ESCHERICHIA COIL*

*Escherichia coli* is a normal opportunistic flora in the digestive tract, that is, if the number is within normal limits, the bacteria can be beneficial, but if there is an increase in the number from normal, the bacteria will become pathogenic (28). Pathogenic *Escherichia coli* can cause disease in the intestinal tract as well as outside the intestine (29). There are six intestinal pathotypes of *Escherichia coli* including Shiga toxin-producing *Escherichia coli* (STEC), enterotoxigenic *Escherichia coli* (ETEC), enteropathogenic *Escherichia coli* (EPEC), diffusely adherent *Escherichia coli*, and enteroinvasive *Escherichia coli*. These strains are classified based on their virulence properties and pathogenicity mechanisms that cause gastrointestinal diseases such as diarrhea (30, 31). *Escherichia coli* is the main cause of urinary tract infections (UTIs). Apart from being infected with *Escherichia coli*, UTIs can also be caused by *Proteus* spp., *Staphylococcus saprophyticus*, *Klebsiella* spp. and other Enterobacteriaceae. However, among UTIS-causing bacteria, *Escherichia coli* is considered to be the most dominant cause of community and nosocomial UTI. In addition, having a history of UTI in a patient increases the chance of infection (23). Therapeutic measures for UTI treatment use several antibiotic options including cotrimoxazole (trimethoprim / sulfamethoxazole), nitrofurantoin, ciprofloxacin and ampicillin (32, 33). However, studies of antibiotic resistance using urinary tract isolates from *Escherichia coli* have shown increased resistance to certain antibiotics, for example to ampicillin and cotrimoxazole (32).

*Escherichia coli* causes infection in the bloodstream which can cause morbidity, mortality, and health problems (34). These infections can increase the mortality rate in the hospital and lead to resistance to antibiotics, leading to longer hospital stays (35). Nosocomial bloodstream infections caused by resistant bacteria have been reported in Africa, especially in areas that are vulnerable and at high risk of developing nosocomial infections, such as in intensive care units and pediatric departments (36-38).

#### USE OF ANTIBIOTICS IN ANIMALS

Antibiotics have contributed greatly to human and animal health since their appearance in the 1940's. The development of antibiotics in the 1940's era cannot be separated from the services of Alexander Fleming who discovered penicillin in 1928 (38). The use of antibiotics can be found in several sectors, such as hospitals (39), animal husbandry and agriculture (40), and fisheries (41) which have the impact of increasing AMR on livestock (42-46), pet (47-50), poultry (51-55), fisheries (56-59) and also obtained from animal products (60-65). The use of antibiotics is increasingly widespread, not only for treating infectious diseases (66), but also for disease prevention and as growth promoters in livestock. Unfortunately, the use of antibiotics in subtherapeutic doses for disease prevention and growth enhancement is higher than for the treatment of disease (67-69). This is one of the factors that trigger AMR in livestock (70).

For at least 50 years, farms in the United States and other developed countries have used antibiotics as Antimicrobial Growth Promoters (AGPs). AGP was first introduced in the mid-1950s. The use of antibiotics as AGP were streptomycin, sulfasuxidine, and streptothricin to pig and chicken feed which has been shown to have a beneficial effect on livestock production for first reported (71).

Antibiotics commonly used in humans include penicillins, fluoroquinolones, and macrolides. Meanwhile, sulfonamides, penicillins, and tetracyclines are often used in animals (11). In a survey conducted in China in 2010, it was found that there were residues of ciprofloxacin, oxytetracycline, enrofloxacin and chlortetracycline in 143 manure samples (72). A survey in Chile suggested that antibiotic residues were also found in eggs produced on farms (73). Due to the high concentration of antibiotics in livestock manure, this manure is often used for fish use because it is believed to increase fish growth and maintain pond fertility (74).

Because the use of antibiotics has resulted in increasing cases of resistance, several countries have limited their use. As in the EU, the use of antibiotics as Antibiotic Growth Promoters (AGP) has been banned since 2006 (18). While in the US it has been banned starting in 2017. In Europe, the European Food Safety Authority (EFSA) and the European Medicines Agency (EMA) recommend steps to reduce the use of antimicrobials in animal husbandry in the European Union (79). France also issued a policy of prohibiting the excessive use of antibiotics in chickens (19).

#### AMR OF *ESCHERICHIA COIL* IN FOOD OF ANIMAL ORIGIN

*Escherichia coli* lives in both animals and humans as normal flora which has many uses (21). But on the other hand, it can also harm humans because it causes infectious diseases (28). To avoid these losses, breeders take precautions so that their livestock remain healthy and have high productivity (72, 75). Farmers give antibiotics to prevent infection and add antibiotics as additional feed to increase livestock productivity. The use of antibiotics without supervision can lead to resistance to one antibiotic or even more than 3 antibiotics called multidrug resistant (MDR). A total of 94.9% *Escherichia coli* in sheep isolates showed the presence of MDR during the antibiotic sensitivity test (76). This high percentage illustrates the potential for spreading resistant microbes through animal husbandry (16). The level of microbial resistance to

antibiotics is called antimicrobial resistance (AMR) will continue to increase if farmers continue to use antibiotics excessively.

Cases of resistant *Escherichia coli* were found on farms, both in livestock and breeders (77). Livestock such as chicken, turkey, pig, cow, as well as food of animal origin such as eggs and milk are reported as reservoirs of resistant *Escherichia coli*. The majority of *Escherichia coli* isolated turned out to be resistant to penicillin class antibiotics such as penicillin, ampicillin and amoxicillin. *Escherichia coli* isolates isolated from milk isolates in India were reported to be 100% resistant to penicillin. Likewise in Indonesia, *Escherichia coli* isolates isolated from beef were also reported to be 100% resistant to ampicillin, which is a penicillin antibiotic class (12). For foods of other animal origin, microbial resistance to the penicillin class is also high, for example *Escherichia coli* in eggs in Pakistan reached 83.3% (78), chicken and turkey in Arizona (79). Apart from livestock, cases of resistant *Escherichia coli* were also found in breeders. As Aworh reports, workers on Nigerian farms are also at risk of carrying *Escherichia coli* resistant to ampicillin, tetracycline, sulfamethoxazole-trimethoprim, and streptomycin. This resistant *Escherichia coli* can be transmitted from animals to humans via several transfer mechanisms (76). Transmission of *Escherichia coli* from animals into the human body can occur in at least two ways, namely (a) when humans eat food of animal origin that is contaminated with bacteria, the food is not washed and cooked properly and (b) through direct contact with existing *Escherichia coli*. Livestock is a reservoir of resistant *Escherichia coli* which has a resistance coding gene. The gene can reach humans via the food chain (76). Antimicrobial resistance in chickens is a common problem in Bangladesh and other developing countries due to the indiscriminate use of antibiotics as feed additives and prophylactic treatment of infectious diseases. Commonly used antibiotic agents, such as ampicillin, tetracycline, streptomycin, ciprofloxacin, erythromycin, sulfamethoxazole-trimethoprim, cholistin sulfate, gentamicin, and levofloxacin have relatively high levels of resistance than previously reported. (80). The presence of MDR also cannot be separated from microbial resistance in animal husbandry. The *Escherichia coli* isolate showing very high resistance to multiple antibiotics and a high prevalence of MDR leads to an alarming situation that requires further investigation. There is a need for a framework to address these problems (81). Due to the lack of choice of antibiotics that are sensitive to bacteria, further research on alternative antimicrobial agents is currently urgently needed (80).

#### MECHANISM OF RESISTANCE ON *ESCHERICHIA COLI*

*Escherichia coli* belongs to the Gram-negative bacteria group. Gram negative bacteria are more easily resistant because they have an outer membrane that is not owned by the Gram-positive bacteria group. The outer membrane of Gram negative is an additional structure in protection against antibiotics (81). The inner structure of this outer membrane is composed of phospholipids (PLs) and lipopolysaccharides (LPs) in the outer structure (82). PLs and LPs act as bacterial membrane defenses because they are composed of saturated chains so that they are hydrophobic (83). The molecules making up PLs and LPs each carry a negative charge which allows for intermolecular linking interactions through the binding of divalent cations (84) and the presence of porin as ion

selective channels so as to limit the absorption of antibiotics (85)

*Escherichia coli* has the ability to modify antibiotic targets including the aminoglycoside class of antibiotics by modifying the ribosomal subunit through the acquisition of plasmids carrying 16S rRNA methyltransferases (86-87) against the fluoroquinolone group by means of mutations in the GyrA subunit which is chromosomally coded from gyrase (gyrA gene) or the ParC subunit of topoisomerase IV (parC gene) (88) and against penicillin by modifying penicillin-binding protein (PBP) (89, 90).

$\beta$ -lactam is a class of antibiotics that has 3 carbon rings and 1 nitrogen ring which is responsible for the bacteriolytic mechanism of action against bacteria.  $\beta$ -lactam works by inhibiting the synthesis of peptidoglycan which is an important component of bacterial cell walls. *Escherichia coli* is able to be resistant to  $\beta$ -lactam by producing  $\beta$ -lactamase enzymes. The most widely used classification for the  $\beta$ -lactamase enzyme is the Ambler structural classification, which is based on the similarity of the sequence so that this protein is divided into four classes, namely classes A, C, and D of serine- $\beta$ -lactamase and class B of metallo- $\beta$ -lactamase (91). *Escherichia coli* is capable of producing several types of enzymes such as broad-spectrum  $\beta$ -lactamase (ESBL), AmpC  $\beta$ -lactamase (AmpC) (92) and carbapenemase which have different hydrolytic activity against  $\beta$ -lactam antibiotics (93). ESBL is a major concern in the medical world because it has the highest level of causative factors for  $\beta$ -lactamase resistance (94). ESBL is predominant class A in the Ambler classification and is resistant to penicillins, first, second, and third generation cephalosporins and monobactams (for example, aztreonam), but cannot hydrolyze cephamycins (cefoxitin) or carbapenems (imipenem, meropenem), and can be inhibited by  $\beta$ -lactamase inhibitors such as clavulanic acid, tazobactam and sulbactam (95).

AmpC has the ability to hydrolyze amino- and ureidopenicillins, oxyimino- $\beta$ -lactams such as ceftazidime, ceftiofur, and aztreonam and deactivate cephalosporins such as cephamycins (cefoxitin) (96-97). AmpC-producing *Escherichia coli* strains ranging from 2.0% were isolated from patients in Portuguese hospitals (98) to 16.7% from three university hospitals in Iran (99) and 29.0% from five referral hospitals in Sudan (100) AmpC and ESBL may be owned by a strain of *Escherichia coli* simultaneously which causes the mechanism of action of antibiotics to be more complicated and resistance to increase (82, 101).

Carbapenems bind to penicillin-binding proteins and induce spheroplast formation and cell lysis. The *Escherichia coli* strain that produces carbapenemase has the ability to hydrolyze the antibiotic carbapenem (102). Increased resistance to carbapenem antibiotics has been reported in several countries, one of which is China, which in 2011 was 0% of cases then increased to 1.9% in 2017 (103).

Several types of efflux pumps found in *Escherichia coli* are the ATP-binding cassette (ABC) family, the multidrug and toxic compound extrusion (MATE) family, the major facilitator superfamily (MFS), and the resistance-nodulation cell division (RND) family. *Escherichia coli* has an ABC group efflux pump, namely the MacAB transporter which provides resistance to several macrolides (104); MATE which is capable of transporting fluoroquinolones (105, 106); MFS capable of transporting macrolides (MefB and MdfA pumps), fluoroquinolones (QepA2, EmrAB-ToIC, and MdfA pumps), tetracyclines (EmrAB-ToIC and MdfA pumps), trimethoprim (Fsr pumps), and chloramphenicol

(MdfA pumps) (107, 108); and RND which can release  $\beta$ -lactam, fluoroquinolone, tetracycline, chloramphenicol, and lincosamides (109-111).

#### FACTORS THAT INCREASE RESISTANCE ON *ESCHERICHIA COILI*

High population density goes hand in hand with the increasing need for food derived from livestock. As a result, an intensive strategy is needed to prevent and control pathogenic infections by using antibiotics. The use of antibiotics in livestock requires a prescription from a doctor. However, farm managers often give antibiotics of their own accord even though it is in accordance with the guidelines given by veterinarians (112, 113). In addition, in some countries, antibiotics are available over the counter without a prescription. This absence of regulation makes antibiotics readily available, abundant and cheap, thus encouraging overuse. The ability to purchase such products online has also made them accessible in countries where antibiotics are regulated (114). Administration of antibiotics in livestock that is not in accordance with the needs (misused) which causes increased resistance to pathogens including *Escherichia coli* (115). Antibiotics used in livestock that induce resistance can enter the human body through the consumption of feed from livestock (116). Resistant bacteria that enter the human body will cause adverse health problems (34).

#### GENE CODE FOR RESISTANT TO ANTIBIOTICS ON *ESCHERICHIA COILI*

Bacterial resistance to antibiotics is classified into two types, namely intrinsic resistance and acquired resistance (117). Intrinsic resistance is a congenital resistance due to the insensitivity of bacteria to certain antibiotic classes. This is due to the inability of the target structure for certain antibiotics, for example, vancomycin resistance in Gram negative bacteria due to the inability of vancomycin to penetrate the outer membrane (118). Conversely, acquired resistance is resistance caused by the acquisition of a foreign resistant gene or gene mutation (119). In bacteria, genes can be inherited from relatives or can be obtained from non-relatives of mobile elements such as plasmids. This horizontal gene transfer (HGT) may allow antibiotic resistance to be transmitted between different bacterial species (120). Resistance can also occur spontaneously via mutation (121).

Resistant bacteria encode different resistance genes for each antibiotic class. Gram negative bacteria that are resistant to the aminoglycoside antibiotic class have resistant coding genes, namely *armA* (122) *npmA*, *rmtA*, *rmtB*, *rmtC*, and *rmtD* (123,124). The *rmt* gene provides resistance to gentamicin and amikacin, while *npmA* provides resistance to gentamicin, neomycin, amikacin, and apramycin, but not to streptomycin (125). Meanwhile, the gene coding for *Escherichia coli* resistance can come from humans and animals. blaTEM-1 is an animal-derived *Escherichia coli* resistance gene to  $\beta$ -lactam that can inactivate penicillins and aminopenicillins. Meanwhile, the most common ESBL gene in *Escherichia coli* isolates of human origin is blaCTX-M-15 (126).

The target of the fluoroquinolone antibiotic against *Escherichia coli* is gyrase. *Escherichia coli* developed a fluoroquinolone resistance mechanism by carrying out mutations encoded by the *gyrA* and *parC* genes (127). Other genes at play are *qnrS1*, *qnrB1*, *qnrB4*, and *qnrB10* (128). *Escherichia coli* resistance to tetracycline

antibiotics is coded by tet (A), tet (B), tet (C), tet (D), tet (E), tet (G), tet (J), tet (L), and tet (Y) as the gene coding for the efflux pump and tet (M) and tet (W) as ribosome protectors (129). The coding genes for *Escherichia coli* resistance to phenicol antibiotics include *pac*, *fluorine*, and *cfr* (130). *Escherichia coli* of animal origin, is resistant to sulfonamide antibiotics mediated by *sul1*, *sul2*, or *sul3*. The *sul2* gene is associated with the streptomycin *strA-strB* resistance gene whereas *sul3* is associated with other resistance genes, such as the *mef* (B) macrolide resistance gene (131).

The *dfr* gene modulates resistance to trimethoprim antibiotics in Enterobacteriaceae and other Gram-negative bacteria. Based on its size and structure, the *dfr* gene is divided into two major groups, namely *dfrA* and *dfrB* (132-133). The *dfrA* gene encodes a protein from 152 to 189 amino acids, while *dfrB* encodes a protein of 78 amino acids. Most of the *dfrA* and *dfrB* genes are found in *Escherichia coli* of animal origin. *Escherichia coli* is also resistant to polymyxin antibiotics which are modulated by the *pmrA*, *pmrB*, *mgrB*, *phoP*, and *phoQ* genes isolated from pigs (134).

#### CONCLUSION

Microbial resistance to antibiotics is a health problem that continues to be a global challenge. One of them is *Escherichia coli* which is commonly found in livestock and humans as well as animal products that are resistant to several classes of antibiotics. Animals and food of animal origin such as poultry, pigs, cattle, sheep, eggs, and milk are reservoirs for resistant *Escherichia coli* that have the potential to cause disease in humans. Because there are fewer choices of antibiotics that are sensitive to microbes, it is necessary to look for new antibiotic alternatives that are more effective in treating pathogenic infections, especially *Escherichia coli*. In addition, there is a need for strict regulations governing the use of antibiotics in livestock and poultry to break the chain of antibiotic resistance.

#### ACKNOWLEDGEMENTS

This study was in part funded by the Direktorat Riset dan Pengabdian Masyarakat, Deputy Bidang Penguatan Riset dan Pengembangan, Kementerian Riset dan Teknologi/Badan Riset dan Inovasi Nasional, Indonesia in fiscal year 2020 with grant number: 756/UN3.14/PT/2020.

#### REFERENCES

1. Irrgang A, Hammerl JA, Falgenhauer L, Guiral E, Schmogger S, Imirzalioglu C, et al. Diversity of CTX-M-1-producing *E. coli* from German food samples and genetic diversity of the blaCTX-M-1 region on Inc11 ST3 plasmids. *Vet Microbiol.* 2018;221(6):98-104.
2. World Health Organization. Antimicrobial Resistance: Global Report on Surveillance. 2014.
3. Ramírez-Castillo FY, Moreno-Flores AC, Avelar-González FJ, Márquez-Díaz F, Harel J, Guerrero-Barrera AL. An evaluation of multidrug-resistant *Escherichia coli* isolates in urinary tract infections from Aguascalientes, Mexico: Cross-sectional study. *Ann Clin Microbiol Antimicrob.* 2018;17(1):1-13.
4. Alonso CA, Zarazaga M, Ben Sallem R, Jouini A, Ben Slama K, Torres C. Antibiotic resistance in *Escherichia coli* in husbandry animals: the African perspective. *Lett Appl Microbiol.* 2017;64(5):318-34.

5. Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, Robinson TP, et al. Global trends in antimicrobial use in food animals. *Proc Natl Acad Sci U S A*. 2015;112(18):5649–54.
6. Baron S, Jouy E, Larvor E, Eono F, Bougeard S, Kempf I. Impact of third-generation-cephalosporin administration in hatcheries on fecal *Escherichia coli* antimicrobial resistance in broilers and layers. *Antimicrob Agents Chemother*. 2014;58(9):5428–5434.
7. Chantziaras I, Boyen F, Callens B, Dewulf J. Correlation between veterinary antimicrobial use and antimicrobial resistance in food-producing animals: A report on seven countries. *J Antimicrob Chemother*. 2014;69(3):827–34.
8. Agunos A, Léger D, Carson C. Review of antimicrobial therapy of selected bacterial diseases in broiler chickens in Canada. *Can Vet J*. 2012;53(12):1289–1300.
9. Landoni MF, Albarellos G. The use of antimicrobial agents in broiler chickens. *Vet J*. 2015;205(1):21–7.
10. Page SW, Gautier P. Use of antimicrobial agents in livestock. *OIE Rev Sci Tech*. 2012;31(1):145–88.
11. Chen Y, Liu Z, Zhang Y, Zhang Z, Lei L, Xia Z. Increasing Prevalence of ESBL-Producing Multidrug Resistance *Escherichia coli* From Diseased Pets in Beijing, China From 2012 to 2017. *Front Microbiol*. 2019;10(December):1–12.
12. Effendi MH, Cicilia R, Tyasningsih W. Public Awareness for Antimicrobial Resistance from *Escherichia coli* Isolated from Beef Sold on Several Wet Market in Surabaya, Indonesia. *Indian Journal of Public Health Research & Development*, 2020, 11(9): 295-300
13. Peralta-Sánchez JM, Martín-Platero AM, Ariza-Romero JJ, Rabelo-Ruiz M, Zurita-González MJ, Baños A, et al. Egg production in poultry farming is improved by probiotic bacteria. *Front Microbiol*. 2019;10(5):1–13.
14. Nhung NT, Chansiripornchai N, Carrique-Mas JJ. Antimicrobial resistance in bacterial poultry pathogens: A review. *Front Vet Sci*. 2017;4(8):1–17.
15. Chah KF, Ugwu IC, Okpala A, Adamu KY, Alonso CA, Ceballos S, et al. Detection and molecular characterisation of extended-spectrum  $\beta$ -lactamase-producing enteric bacteria from pigs and chickens in Nsukka, Nigeria. *J Glob Antimicrob Resist*. 2018;15:36–40.
16. Falgenhauer L, Imirzalioglu C, Oppong K, Akenten CW, Hogan B, Krumkamp R, et al. Detection and characterization of ESBL-producing *Escherichia coli* from humans and poultry in Ghana. *Front Microbiol*. 2019;10(1):1–8.
17. Martínez-Vázquez AV, Rivera-Sánchez G, Lira-Méndez K, Reyes-López MÁ, Bocanegra-García V. Prevalence, antimicrobial resistance and virulence genes of *Escherichia coli* isolated from retail meat in Tamaulipas, Mexico. *J Glob Antimicrob Resist*. 2018;14:266–72.
18. Ur Rahman S, Ahmad S, Khan I. Incidence of ESBL-producing-*Escherichia coli* in poultry farm environment and retail poultry meat. *Pak Vet J*. 2019;39(1):116–20.
19. Casella T, Nogueira MCL, Saras E, Haenni M, Madec JY. High prevalence of ESBLs in retail chicken meat despite reduced use of antimicrobials in chicken production, France. *Int J Food Microbiol*. 2017;257(2):271–275.
20. Jang J, Hur HG, Sadowsky MJ, Byappanahalli MN, Yan T, Ishii S. Environmental *Escherichia coli*: ecology and public health implications—a review. *J Appl Microbiol*. 2017;123(3):570–581.
21. Kheiri R, Akhtari L. Antimicrobial resistance and integron gene cassette arrays in commensal *Escherichia coli* from human and animal sources in IRI. *Gut Pathog*. 2016;1–10.
22. Overdeest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, et al. Extended-spectrum  $\beta$ -lactamase genes of *Escherichia coli* in chicken meat and humans, the Netherlands. *Emerg Infect Dis*. 2011;17(7):1216–1222.
23. Jan N, Meshram SU, Kulkarni A. Plasmid profile analysis of multidrug resistant *E. coli* isolated from UTI patients of Nagpur City, India. *Rom Biotechnol Lett*. 2009;14(5):4635–4640.
24. Tenaillon O, Barrick JE, Ribeck N, Deatherage DE, Blanchard JL, Dasgupta A, et al. Tempo and mode of genome evolution in a 50,000-generation experiment. *Nature*. 2016;536(7615):165–70.
25. Nhung NT, Cuong N V, Thwaites G, Carrique-Mas J. Antimicrobial usage and antimicrobial resistance in animal production in Southeast Asia: A review. *Antibiotics*. 2016;5(4):1–24.
26. Berthe T, Ratajczak M, Clermont O, Denamur E, Petit F. Evidence for coexistence of distinct *Escherichia coli* populations in various aquatic environments and their survival in estuary water. *Appl Environ Microbiol*. 2013;79(15):4684–93.
27. Tenaillon O, Skurmik D, Picard B, Denamur E. The population genetics of commensal *Escherichia coli*. *Nat Rev Microbiol*. 2010;8(3):207–17.
28. Issa AH, Almayah AA, Ibrahim HK. New virulence factor of normal Flora *E. Coli*. *Syst Rev Pharm*. 2020;11(2):71–6.
29. Kai A, Konishi N, Obata H. Diarrheagenic *Escherichia coli*. *Nippon rinsho Japanese J Clin Med*. 2010;68 Suppl 6(1):203–207.
30. Kaper JB, Nataro JP, Mobley HLT. Pathogenic *Escherichia coli*. *Nat Rev Microbiol*. 2004;2(2):123–40.
31. Alanazi MQ, Alqahtani FY, Aleanizy FS. An evaluation of *E. coli* in urinary tract infection in emergency department at KAMC in Riyadh, Saudi Arabia: Retrospective study. *Ann Clin Microbiol Antimicrob*. 2018;17(1):1–7.
32. Foxman B. The epidemiology of urinary tract infection. *Nat Rev Urol*. 2010;7(12):653–60.
33. Manges AR, Geum HM, Guo A, Edens TJ, Fibke CD, Pitout JDD. Global extraintestinal pathogenic *Escherichia coli* (ExPEC) lineages. *Clin Microbiol Rev*. 2019;32(3):1–25.
34. Naylor NR, Pouwels KB, Hope R, Green N, Henderson KL, Knight GM, et al. The health and cost burden of antibiotic resistant and susceptible *Escherichia coli* bacteraemia in the English hospital setting: A national retrospective cohort study. *PLoS One*. 2019;14(9):1–19.
35. Saied T, Elkholy A, Hafez SF, Basim H, Wasfy MO, El-Shoubary W, et al. Antimicrobial resistance in pathogens causing nosocomial bloodstream infections in university hospitals in Egypt. *Am J Infect Control*. 2011;39(9):e61–5.
36. Tansarli GS, Poulidakos P, Kapaskelis A, Falagas ME. Proportion of extended-spectrum  $\beta$ -lactamase

- (ESBL)-producing isolates among enterobacteriaceae in Africa: Evaluation of the evidence-systematic review. *J Antimicrob Chemother.* 2014;69(5):1177-84.
37. Ma F, Xu S, Tang Z, Li Z, Zhang L. Use of antimicrobials in food animals and impact of transmission of antimicrobial resistance on humans. *Biosaf Heal.* 2020;
  38. Petrillo F, Folliero V, Santella B, Franci G, Foglia F, Trotta MC, et al. Prevalence and Antibiotic Resistance Patterns of Ocular Bacterial Strains Isolated from Pediatric Patients in University Hospital of Campania "luigi Vanvitelli," Naples, Italy. *Int J Microbiol.* 2020;2020.
  39. Speksnijder DC, Jaarsma ADC, van der Gugten AC, Verheij TJM, Wagenaar JA. Determinants associated with veterinary antimicrobial prescribing in farm animals in the Netherlands: A qualitative study. *Zoonoses Public Health.* 2015;62(s1):39-51.
  40. Manyi-Loh C, Mamphweli S, Meyer E, Okoh A. Antibiotic use in agriculture and its consequential resistance in environmental sources: Potential public health implications. *Molecules.* 2018; 23
  41. Çelebi H, Bahadır T, Şimşek İ, Tulun Ş. Antibiotic Applications in Fish Farms and Environmental Problems. *Turkish J Eng.* 2019;(1):1-10.
  42. Effendi, M.H., Oktavianto, A and Hastutiek, P. Tetracycline Resistance Gene in *Streptococcus Agalactiae* Isolated from Bovine Subclinical Mastitis in Surabaya, Indonesia. *Philipp. Journal of Veterinary Medicine.* 2018; 55 (SI): 115-120.
  43. Effendi M.H, Harijani N, Budiarto, Triningtya N.P, Tyasningsih W, and Plumeriastuti H. Prevalence of Pathogenic *Escherichia Coli* Isolated from Subclinical Mastitis in East Java Province, Indonesia. *Indian Vet. J.* 2019; 96 (03): 22 - 25.
  44. Tyasningsih, W., Effendi, M. H., Budiarto, B., Syahputra, I. R. Antibiotic Resistance to *Staphylococcus aureus* and Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated from Dairy Farms in Surabaya, Indonesia. *Indian Vet. J.* 2019; 96(11), 27-31.
  45. Putra, A.R. Effendi, M.H. Koesdarto, S. Suwarno, S. Tyasningsih, W. and Estoepangestie, A.T. Detection of the extended spectrum  $\beta$ -lactamase produced by *Escherichia coli* from dairy cows by using the Vitek-2 method in Tulungagung regency, Indonesia. *Iraqi Journal of Veterinary Sciences,* 2020; 34 (1): 203-207.
  46. Putra ARS, Effendi MH, Koesdarto S, and Tyasningsih W. Molecular Identification of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* Isolated from Dairy Cows in East Java Province, Indonesia. *Indian Vet. J.* 2019; 96 (10): 26 - 30.
  47. Rahmaniari, R. P., Yunita, M. N., Effendi, M. H., Yanestria, S. M. Encoding Gene for Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated from Nasal Swab of Dogs. *Indian Vet. J.* 2020; 97(02), 37-40.
  48. Kristianingtyas L, Effendi, MH, Tyasningsih W, Kurniawan F. Genetic Identification of *blactx-M* Gene and *blatem* Gene on Extended Spectrum Beta Lactamase (ESBL) Producing *Escherichia Coli* from Dogs. *Indian Vet. J.* 2020; 97 (01): 17 - 21.
  49. Decline, V., Effendi, M. H., Rahmaniari, R. P., Yanestria, S. M., Harijani, N. Profile of antibiotic-resistant and presence of methicillin-resistant *Staphylococcus aureus* from nasal swab of dogs from several animal clinics in Surabaya, Indonesia. *Int J One Health,* 2020; 6(1), 90-94.
  50. Yunita, M. N., Effendi, M. H., Rahmaniari, R. P., Arifah, S., Yanestria, S. M. Identification Of Spa Gene For Strain Typing Of Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated From Nasal Swab Of Dogs. *Biochem. Cell. Arch.* 2020; 20 (1), 2999-3004.
  51. Wibisono FJ, Sumiarto B, Untari T, Effendi MH, Permatasari DA, Witaningrum AM. The Presence of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* On Layer Chicken Farms In Blitar Area, Indonesia. *Biodiversitas.* 2020; 21 (6): 2667-2671.
  52. Rahmahani J, Salamah, Mufasirin, Tyasningsih W, and Effendi MH. Antimicrobial Resistance Profile of *Escherichia coli* From Cloacal Swab of Domestic Chicken in Surabaya Traditional Market. *Biochem. Cell. Arch.* 2020; 20 (1): 2993-2997.
  53. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A., Witaningrum, A.M. CTX Gene of Extended Spectrum Beta-Lactamase (ESBL) Producing *Escherichia coli* on Broilers in Blitar, Indonesia. *Sys Rev Pharm* 2020;11(7): 396-403.
  54. Permatasari, D.A., Witaningrum, A.M., Wibisono, F.J., Effendi, M.H. Detection and prevalence of multidrug-resistant *Klebsiella pneumoniae* strains isolated from poultry farms in Blitar, Indonesia. *Biodiversitas,* 2020; 21 (10): 4642-4647.
  55. Wibisono, F.J., Sumiarto, B., Untari, T., Effendi, M.H., Permatasari, D.A., Witaningrum, A.M. Short Communication: Pattern of antibiotic resistance on extended-spectrum beta-lactamases genes producing *Escherichia coli* on laying hens in Blitar, Indonesia. *Biodiversitas,* 2020; 21 (10): 4631- 4635.
  56. Helmi, AM., Mukti, AT., Soegianto, A., Mahardika, K., Mastuti, I., Effendi, MH., Plumeriastuti, H. A Review of Bacterial Zoonoses and Antimicrobial Resistant (AMR) on Grouper fish (*Epinepholus sp.*). *Sys Rev Pharm* 2020;11(9):79-88.
  57. Effendi MH, Bintari IG, Aksono EB, Hermawan IP. Detection of *blaTEM* Gene of *Klebsiella pneumoniae* Isolated from Swab of Food Producing Animals in East Java. *Tropical Animal Science Journal.* 2018;41(3):174-178.
  58. Yanestria, S.M., Rahmaniari, R.P., Wibisono, F.J., Effendi, M.H. Detection of *invA* gene of *Salmonella* from milkfish (*Chanos chanos*) at Sidoarjo wet fish market, Indonesia, using polymerase chain reaction technique, *Veterinary World,* 2019; 12(1): 170-175.
  59. Effendi, MH, Harijani N, Yanestria SM, Hastutiek P. 2018. Identification of shiga toxin-producing *Escherichia coli* in raw milk samples from dairy cows in Surabaya, Indonesia. *Philippine J Vet Med.*55(SI):109-114.
  60. Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H. 2020. *MecA* gene and methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from dairy farms in East Java, Indonesia. *Biodiversitas,* 21(8): 3562-3568.
  61. Harijani, N., Wandari, A., Effendi, M.H. and Tyasningsih. W. Molecular Detection of Encoding Enterotoxin C Gene and Profile of Antibiotic Resistant on *Staphylococcus Aureus* Isolated from Several Dairy Farms in East Java, Indonesia. *Biochem. Cell. Arch.* 2020; 20 (1): 3081-3085.
  62. Effendi, M. H., Bintari, I. G., Aksoro, E. B. and Hermawan. I. P. Detection of *blaTEM* Gene of

- Klebsiella pneumoniae Isolated from swab of food-producing animals in East Java. *Trop. Anim. Sci. J.* 2018;41:174-178.
63. Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H., Tyasningsih, W. and Rahmahani, J. Detection of Enterotoxin type B gene on Methicillin Resistant *Staphylococcus aureus* (MRSA) isolated from raw milk in East Java, Indonesia. *Sys Rev Pharm, 2020;11(7):290-298.*
64. Ansharieta, R., Effendi, M.H., Plumeriastuti, H. Detection of Multidrug-Resistant (MDR) *Escherichia coli* Isolated from Raw Milk in East Java Province, Indonesia, *Indian Journal of Forensic Medicine & Toxicology, 2020;14(4):3403-3407*
65. Ramandinianto, S.C., Khairullah, A.R., Effendi, M.H., Hestiana, E. P. Profile of Multidrug Resistance (MDR) and Methicillin Resistant *Staphylococcus aureus* (MRSA) on Dairy Farms in East Java Province, Indonesia, *Indian Journal of Forensic Medicine & Toxicology, 2020;14(4):3439-3445*
66. Devasahayam G, Scheld WM, Hoffman PS. Newer antibacterial drugs for a new century. *Expert Opin Investig Drugs.* 2010;19(2):215-34.
67. Maron DF, Smith TJS, Nachman KE. Restrictions on antimicrobial use in food animal production: An international regulatory and economic survey. *Global Health.* 2013;9(1).
68. El-Lethey H, Huber-Eicher B, Jungi TW. Exploration of stress-induced immunosuppression in chickens reveals both stress-resistant and stress-susceptible antigen responses. *Vet Immunol Immunopathol.* 2003;95(3-4):91-101.
69. Silbergeld EK, Graham J, Price LB. Industrial food animal production, antimicrobial resistance, and human health. *Annu Rev Public Health.* 2008;29:151-69.
70. Bouki C, Venieri D, Diamadopoulos E. Detection and fate of antibiotic resistant bacteria in wastewater treatment plants: A review. *Ecotoxicol Environ Saf.* 2013;91:1-9.
71. Moore, P.R., Eenson, A., Luckey, T.D., McCoy, C.A. Elvehjem, EBH. Use of sulfasuxidine, streptothricin, and streptomycin in nutritional studies with the chick. *J Biol Chem.* 1946;165:437-41.
72. Zhao L, Dong YH, Wang H. Residues of veterinary antibiotics in manures from feedlot livestock in eight provinces of China. *Sci Total Environ.* 2010;408(5):1069-1075.
73. Cornejo J, Pokrant E, Figueroa F, Riquelme R, Galdames P, Pillo F Di, et al. Assessing antibiotic residues in poultry eggs from backyard production systems in Chile, first approach to a non-addressed issue in farm animals. *Animals.* 2020;10(6):1-11.
74. Parvez, M. S., Sarker, M. S., Azad, M. S., & Salequzzaman M. Effect on water quality, pond productivity and growth of carps in polyculture system by using homestead organic wastage as pond manure. *Int J Sustain Agril Tech.* 2002;2(2):45-9.
75. Murphy D, Ricci A, Auce Z, Beechinor JG, Bergendahl H, Breathnach R, et al. EMA and EFSA Joint Scientific Opinion on measures to reduce the need to use antimicrobial agents in animal husbandry in the European Union, and the resulting impacts on food safety (RONAFA). *EFSA J.* 2017;15(1).
76. Gozi KS, Froes JR, Ajude LPTD, Da Silva CR, Baptista RS, Peiró JR, et al. Dissemination of multidrug-resistant commensal *Escherichia coli* in feedlot lambs in southeastern Brazil. *Front Microbiol.* 2019;10(6).
77. Aworh MK, Kwaga J, Okolocha E, Mba N, Thakur S. Prevalence and risk factors for multi-drug resistant *Escherichia coli* among poultry workers in the Federal Capital Territory, Abuja, Nigeria. *PLoS One.* 2019;14(11):1-15.
78. Tasbihullah, R., Sadeeq ur, A., Tariq, S., Umer, A., Shakoor, S., Muhammad, A., et al. High Occurrence Rate of Multidrug-Resistant ESBL-Producing *E. coli* Recovered from Table Eggs in District Peshawar, Pakistan. *Pakistan Journal of Zoology.* 2020; 52(4):1231-1238.
79. Davis GS, Waits K, Nordstrom L, Grande H, Weaver B, Papp K, et al. Antibiotic-resistant *Escherichia coli* from retail poultry meat with different antibiotic use claims. *BMC Microbiol.* 2018;18(1):1-7.
80. Al Azad, MAR, Rahman MM, Amin R, Begum MIA, Fries R, Husna A, et al. Susceptibility and multidrug resistance patterns of *Escherichia coli* isolated from cloacal swabs of live broiler chickens in Bangladesh. *Pathogens.* 2019;8(3):1-9.
81. Kumar P, Kumari S, Kaushik P. Prevalence of Multi-Drug Resistant (MDR) *Escherichia coli* in bovine clinical samples. *Int. J. Curr. Microbiol. AppSci.* (2018). 2018;(7):1476-85.
82. Kallau NHG, Wibawan IWT, Lukman DW, Sudarwanto MB. Detection of multi-drug resistant (MDR) *Escherichia coli* and tet gene prevalence at a pig farm in Kupang, Indonesia. *J Adv Vet Anim Res.* 2018;5(4):388-96.
83. Blair JMA, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJV. Molecular mechanisms of antibiotic resistance. *Nat Rev Microbiol.* 2015;13(1):42-51.
84. Henderson JC, Zimmerman SM, Crofts AA, Boll JM, Kuhns LG, Herrera CM, et al. The Power of Asymmetry: Architecture and Assembly of the Gram-Negative Outer Membrane Lipid Bilayer. *Annu Rev Microbiol.* 2016;70(6):255-78.
85. Raetz CRH, Whitfield C. Lipopolysaccharide Endotoxins Endotoxins as Activators of Innate Immunity. *Annu Rev Biochem.* 2008;71:635-700.
86. Nikaido H, Vaara M. Molecular basis of bacterial outer membrane permeability. *Microbiol Rev.* 1985;49(1):1-32.
87. Delcour AH. Outer membrane permeability and antibiotic resistance. *Biochim Biophys Acta - Proteins Proteomics.* 2009;1794(5):808-816.
88. Mingeot-Leclercq MP, Glupczynski Y, Tulkens PM. Aminoglycosides: Activity and resistance. *Antimicrob Agents Chemother.* 1999;43(4):727-737.
89. Schultz C, Geerlings S. Plasmid-mediated resistance in enterobacteriaceae: Changing landscape and implications for therapy. *Drugs.* 2012;72(1):1-16.
90. Hooper DC JG. Mechanisms of drug resistance: quinolone resistance. *Ann N Y Acad Sci.* 2015;1354:12-31.
91. Sauvage E, Kerff F, Terrak M, Ayala JA, Charlier P. The penicillin-binding proteins: Structure and role in peptidoglycan biosynthesis. *FEMS Microbiol Rev.* 2008;32(2):234-58.
92. Georgopadakou. Penicillin-binding proteins and bacterial resistance to  $\beta$ -lactams. *Antimicrob Agents Chemother.* 1993;37:2045-53.
93. Bush K, Jacoby GA. Updated functional classification of  $\beta$ -lactamases. *Antimicrob Agents Chemother.* 2010;54(3):969-76.
94. Jacoby GA. AmpC B-Lactamases. *Clin Microbiol Rev.*



- 2009;22(1):161-82.
95. Psichogiou M, Tassios PT, Avlami A, Stefanou I, Kosmidis C, Platsouka E, et al. Ongoing epidemic of blaVIM-1-positive *Klebsiella pneumoniae* in Athens, Greece: A prospective survey. *J Antimicrob Chemother.* 2008;61(1):59-63.
  96. Samanta I, Joardar SN, Das PK, Sar TK. Comparative possession of Shiga toxin, intimin, enterohaemolysin and major extended spectrum beta lactamase (ESBL) genes in *Escherichia coli* isolated from backyard and farmed poultry. *Iran J Vet Res.* 2015;16(1):90-93.
  97. Philippon A, Labia R JG. Extended-spectrum beta-lactamases. *Antimicrob Agents Chemother.* 1989;33:1131-1136.
  98. Oliveira C, Amador P, Prudêncio C, Tomaz CT, Tavares-ratado P, Fernandes R. ESBL and AmpC  $\beta$ -lactamases in clinical strains of *Escherichia coli* from Serra da Estrela, Portugal. *Med.* 2019;55(6).
  99. Rizi KS, Mosavat A, Yousefi M, Jamehdar SA, Ghazvini K, Safdari H, et al. High prevalence of blaCMY AmpC beta-lactamase in ESBL co-producing *Escherichia coli* and *Klebsiella* spp. clinical isolates in the northeast of Iran. *J Glob Antimicrob Resist.* 2020;22:477-482.
  100. Dirar M, Bilal N, Ibrahim ME, Hamid M. Resistance Patterns and Phenotypic Detection of  $\beta$ -lactamase Enzymes among Enterobacteriaceae Isolates from Referral Hospitals in Khartoum State, Sudan. *Cureus.* 2020;12(3).
  101. Pitout, JDD., Laupland, KB. Extended-spectrum beta-lactamase-producing Enterobacteriaceae: an emerging public-health concern. *Lancet Infect Dis.* 2008;8(3): 159-166.
  102. Nordmann P, Naas T, Poirel L. Global spread of carbapenemase producing Enterobacteriaceae. *Emerg Infect Dis.* 2011;17(10):1791-1798.
  103. Tian X, Zheng X, Sun Y, Fang R, Zhang S, Zhang X, et al. Molecular mechanisms and epidemiology of carbapenem-resistant *Escherichia coli* isolated from Chinese patients during 2002-2017. *Infect Drug Resist.* 2020;13:501-512.
  104. Lubelski J, Konings WN, Driessen AJM. Distribution and Physiology of ABC-Type Transporters Contributing to Multidrug Resistance in Bacteria. *Microbiol Mol Biol Rev.* 2007;71(3):463-476.
  105. Kuroda T, Tsuchiya T. Multidrug efflux transporters in the MATE family. *Biochim Biophys Acta - Proteins Proteomics.* 2009;1794(5):763-768.
  106. Swick MC, Morgan-Linnell SK, Carlson KM, Zechiedrich L. Expression of multidrug efflux pump genes *acrAB-tolC*, *mdfA*, and *norE* in *Escherichia coli* clinical isolates as a function of fluoroquinolone and multidrug resistance. *Antimicrob Agents Chemother.* 2011;55(2):921-4.
  107. Kumar S, Mukherjee MM, Varela MF. Modulation of Bacterial Multidrug Resistance Efflux Pumps of the Major Facilitator Superfamily. *Int J Bacteriol.* 2013;2013:1-15.
  108. Tuckman M, Petersen PJ, Howe AYM, Orlowski M, Mullen S, Chan K, et al. Occurrence of tetracycline resistance genes among *Escherichia coli* isolates from the phase 3 clinical trials for tigecycline. *Antimicrob Agents Chemother.* 2007;51(9):3205-3211.
  109. Blair JMA, Richmond GE, Piddock LJV. Multidrug efflux pumps in Gram-negative bacteria and their role in antibiotic resistance. *Future Microbiol.* 2014;9(10):1165-77.
  110. Anes J, McCusker MP, Fanning S, Martins M. The ins and outs of RND efflux pumps in *Escherichia coli*. *Front Microbiol.* 2015;6(6):1-14.
  112. Piddock, LJV. Clinically relevant chromosomally encoded multidrug resistance efflux pumps in bacteria. *Clin Microbiol Rev.* 2006;19:382-402.
  113. Raymond MJ, Wöhrle RD, Call DR. Assessment and promotion of judicious antibiotic use on dairy farms in Washington State. *J Dairy Sci.* 2006;89(8):3228-3240.
  114. Sawant AA, Sordillo LM, Jayarao BM. A survey on antibiotic usage in dairy herds in Pennsylvania. *J Dairy Sci.* 2005;88(8):2991-2999.
  115. Michael CA, Dominey-Howes D, Labbate M. The antimicrobial resistance crisis: Causes, consequences, and management. *Front Public Heal.* 2014;2(9):1-8.
  116. Davies J. Origins and evolution of antibiotic resistance. *Microbiologia.* 1996;12(1):9-16.
  117. Golkar Z, Bagasra O, Gene Pace D. Bacteriophage therapy: A potential solution for the antibiotic resistance crisis. *J Infect Dev Ctries.* 2014;8(2):129-136.
  118. Martinez JL. General principles of antibiotic resistance in bacteria. *Drug Discov Today Technol.* 2014;11(1):33-39.
  119. Cox G, Wright GD. Intrinsic antibiotic resistance: Mechanisms, origins, challenges and solutions. *Int J Med Microbiol.* 2013;303(6-7):287-292.
  120. Hawkey PM. The origins and molecular basis of antibiotic resistance. *Br Med J.* 1998;317(7159):657-660.
  121. Le Roux, F., Blokesch M. Eco-evolutionary dynamics linked to horizontal gene transfer in *Vibrios*. *Annu Rev Microbiol.* 2018;8(72):89-110.
  122. Salverda MLM, Koomen J, Koopmanschap B, Zwart MP, De Visser JAGM. Adaptive benefits from small mutation supplies in an antibiotic resistance enzyme. *Proc Natl Acad Sci U S A.* 2017;114(48):12773-12778.
  123. Galimand M, Courvalin P, Lambert T. Plasmid-mediated high-level resistance to aminoglycosides in Enterobacteriaceae due to 16S rRNA methylation. *Antimicrob Agents Chemother.* 2003;47(8):2565-2571.
  124. Bogaerts P, Galimand M, Bauraing C, Deplano A, Vanhoof R, De Mendonca R, et al. Emergence of ArmA and RmtB aminoglycoside resistance 16S rRNA methylases in Belgium. *J Antimicrob Chemother.* 2007;59(3):459-464.
  125. Chen L, Chen ZL, Liu JH, Zeng ZL, Ma JY, Jiang HX. Emergence of RmtB methylase-producing *Escherichia coli* and *Enterobacter cloacae* isolates from pigs in China. *J Antimicrob Chemother.* 2007;59(5):880-885.
  126. Xia J, Sun J, Li L, Fang LX, Deng H, Yang RS, et al. First report of the Inc11/ST898 conjugative plasmid carrying *rmtE2* 16S rRNA methyltransferase gene in *Escherichia coli*. *Antimicrob Agents Chemother.* 2015;59(12):7921-7932.
  127. Peirano G, Pitout JDD. Molecular epidemiology of *Escherichia coli* producing CTX-M  $\beta$ -lactamases: the worldwide emergence of clone ST131 O25:H4. *Int J Antimicrob Agents.* 2010;35(4):316-321.
  128. Hopkins KL, Davies RH, Threlfall EJ. Mechanisms of quinolone resistance in *Escherichia coli* and *Salmonella*: Recent developments. *Int J Antimicrob Agents.* 2005;25(5):358-373.
  129. de Jong A, Muggeo A, El Garch F, Moyaert H, de Champs

- C, Guillard T. Characterization of quinolone resistance mechanisms in Enterobacteriaceae isolated from companion animals in Europe (ComPath II study). *Vet Microbiol.* 2018;216:159–67.
130. Olowe, B. M. Okanlawon, R. A. Olowe A.B. Antimicrobial resistant pattern of *Escherichia coli* from human clinical samples in Osogbo, south western Nigeria. *African J Microbiol Res.* 2008;2(1):8–11.
131. Schwarz S, Kehrenberg C, Doublet B, Cloeckaert A. Molecular basis of bacterial resistance to chloramphenicol and florfenicol. *FEMS Microbiol Rev.* 2004;28(5):519–542.
132. Liu J, Keelan P, Bennett PM, Enne VI. Characterization of a novel macrolide efflux gene, *mef (B)*, found linked to *sul3* in porcine *Escherichia coli*. *J Antimicrob Chemother.* 2009;63(3):423–426.
133. Pattishall KH, Acar J, Burchall JJ, Goldstein FW, Harvey RJ. Two distinct types of trimethoprim resistant dihydrofolate reductase specified by R plasmids of different compatibility groups. *J Biol Chem.* 1977;252(7):2319–2323.
134. Delannoy S, Devendec L Le, Jouy E, Fach P, Drider D, Kempf I. Characterization of colistin-resistant *Escherichia coli* isolated from diseased pigs in France. *Front Microbiol.* 2017;8(11):1–12.

# A Review of Antimicrobial Resistance (AMR) of Escherichia coli on Livestock and Animal Products: Public Health Importance

---

## ORIGINALITY REPORT

---

19%

SIMILARITY INDEX

16%

INTERNET SOURCES

16%

PUBLICATIONS

1%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Daniel Eibach, Cristina Belmar Campos, Ralf Krumkamp, Hassan M. Al-Emran et al. "Extended spectrum beta-lactamase producing Enterobacteriaceae causing bloodstream infections in rural Ghana, 2007–2012", International Journal of Medical Microbiology, 2016  
Publication 1%

---
- 2** Nguyen Nhung, Nguyen Cuong, Guy Thwaites, Juan Carrique-Mas. "Antimicrobial Usage and Antimicrobial Resistance in Animal Production in Southeast Asia: A Review", Antibiotics, 2016  
Publication 1%

---
- 3** Submitted to Universitas Airlangga  
Student Paper 1%

---
- 4** [livrepository.liverpool.ac.uk](http://livrepository.liverpool.ac.uk)  
Internet Source 1%

---
- 5** [www.ema.europa.eu](http://www.ema.europa.eu)  
Internet Source 1%

---

6	<a href="http://rcastoragev2.blob.core.windows.net">rcastoragev2.blob.core.windows.net</a> Internet Source	<1 %
7	<a href="http://vital.seals.ac.za:8080">vital.seals.ac.za:8080</a> Internet Source	<1 %
8	<a href="http://www.intechopen.com">www.intechopen.com</a> Internet Source	<1 %
9	<a href="http://repository.uhamka.ac.id">repository.uhamka.ac.id</a> Internet Source	<1 %
10	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	<1 %
11	<a href="http://mdpi.com">mdpi.com</a> Internet Source	<1 %
12	"Advances in Veterinary Dermatology", Wiley, 2017 Publication	<1 %
13	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
14	<a href="http://repository.up.ac.za">repository.up.ac.za</a> Internet Source	<1 %
15	<a href="http://www.scribd.com">www.scribd.com</a> Internet Source	<1 %
16	<a href="http://www.jscimedcentral.com">www.jscimedcentral.com</a> Internet Source	<1 %
17	<a href="http://eprints.lib.hokudai.ac.jp">eprints.lib.hokudai.ac.jp</a>	

Internet Source

<1 %

18

[lume.ufrgs.br](http://lume.ufrgs.br)

Internet Source

<1 %

19

Farzaneh Firoozeh, Mohammad Zibaei, Farzad Badmasti, Azad Khaledi. "Virulence factors, antimicrobial resistance and the relationship between these characteristics in uropathogenic Escherichia coli", Gene Reports, 2022

Publication

<1 %

20

Sabah A.A. Jassim, Richard G. Limoges. "Bacteriophages: Practical Applications for Nature's Biocontrol", Springer Science and Business Media LLC, 2017

Publication

<1 %

21

[openresearch-repository.anu.edu.au](http://openresearch-repository.anu.edu.au)

Internet Source

<1 %

22

[www.revistas.unitru.edu.pe](http://www.revistas.unitru.edu.pe)

Internet Source

<1 %

23

[mafiadoc.com](http://mafiadoc.com)

Internet Source

<1 %

24

[opus.uleth.ca](http://opus.uleth.ca)

Internet Source

<1 %

25

[www.pnas.org](http://www.pnas.org)

Internet Source

<1 %

26

ophrp.org

Internet Source

&lt;1 %

27

repozitorij.unizg.hr

Internet Source

&lt;1 %

28

Isabel Carvalho, Nadia Safia Chenouf, Rita Cunha, Carla Martins et al. "Antimicrobial Resistance Genes and Diversity of Clones among ESBL- and Acquired AmpC-Producing Escherichia coli Isolated from Fecal Samples of Healthy and Sick Cats in Portugal", Antibiotics, 2021

Publication

&lt;1 %

29

Jorge Rivera-Gomis, Pedro Marín, Julio Ota, Juan Sebastián Galecio, Cristina Martínez-Conesa, María José Cubero. "Resistance patterns to C and D antibiotic categories for veterinary use of Campylobacter spp., Escherichia coli and Enterococcus spp. commensal isolates from laying hen farms in Spain during 2018", Preventive Veterinary Medicine, 2021

Publication

&lt;1 %

30

Khameneh, Bahman, Roudayna Diab, Kiarash Ghazvini, and Bibi Sedigheh Fazly Bazzaz. "Breakthroughs in bacterial resistance mechanisms and the potential ways to combat them", Microbial Pathogenesis, 2016.

Publication

&lt;1 %

31 Michelle Mahony, Brendan McMullan, Jeremy Brown, Sean E Kennedy. "Multidrug-resistant organisms in urinary tract infections in children", *Pediatric Nephrology*, 2019  
Publication

---

32 Shumei Huang, Shuang Wang, Yan Li, Ming Fang et al. "Prevalence and transmission of mobilized colistin resistance (mcr-1) gene positive *Escherichia coli* in healthy rural residents in Shandong province, China", *Microbiological Research*, 2021  
Publication

---

33 [d.lib.msu.edu](http://d.lib.msu.edu)  
Internet Source

---

34 [e-space.mmu.ac.uk](http://e-space.mmu.ac.uk)  
Internet Source

---

35 [ebin.pub](http://ebin.pub)  
Internet Source

---

36 [repo.poltekkesdepkes-sby.ac.id](http://repo.poltekkesdepkes-sby.ac.id)  
Internet Source

---

37 [tel.archives-ouvertes.fr](http://tel.archives-ouvertes.fr)  
Internet Source

---

38 [www.formatex.info](http://www.formatex.info)  
Internet Source

---

39 [www.spandidos-publications.com](http://www.spandidos-publications.com)  
Internet Source

---

40

"Drug Resistance in Bacteria, Fungi, Malaria, and Cancer", Springer Science and Business Media LLC, 2017

Publication

<1 %

41

Kevin Alby, Melissa B. Miller. "Mechanisms and Detection of Antimicrobial Resistance", Elsevier BV, 2023

Publication

<1 %

42

Leonardo Pagani, Giada Fasani, Richard Aschbacher. "Chapter 22 What Healthcare Workers Should Know About the "One Health Approach" and the Global Impact of Antimicrobial Resistance", Springer Science and Business Media LLC, 2021

Publication

<1 %

43

Mabel Kamweli Aworh, Jacob Kwaga, Emmanuel Okolocha, Nwando Mba, Siddhartha Thakur. "Prevalence and risk factors for multi-drug resistant Escherichia coli among poultry workers in the Federal Capital Territory, Abuja, Nigeria", PLOS ONE, 2019

Publication

<1 %

44

Rosine Manishimwe, Martin Buhire, Alexie Uyisunze, Jean Bosco Turikumwenayo, Michael Tukei. "Caractérisation d'Escherichia coli résistant aux antibiotiques dans différents systèmes avicoles de la province de l'Est et de

<1 %



la ville de Kigali au Rwanda", Revue d'élevage  
et de médecine vétérinaire des pays  
tropicaux, 2017

Publication

45

Sabine Delannoy, Laetitia Le Devendec, Eric Jouy, Patrick Fach, Djamel Drider, Isabelle Kempf. "Characterization of Colistin-Resistant Escherichia coli Isolated from Diseased Pigs in France", *Frontiers in Microbiology*, 2017

Publication

<1 %

46

Sukanta Chowdhury, Sumon Ghosh, Mohammad Abdul Aleem, Shahana Parveen et al. "Antibiotic Usage and Resistance in Food Animal Production: What Have We Learned from Bangladesh?", *Antibiotics*, 2021

Publication

<1 %

47

Teillant, Aude, Charles H. Brower, and Ramanan Laxminarayan. "Economics of Antibiotic Growth Promoters in Livestock", *Annual Review of Resource Economics*, 2015.

Publication

<1 %

48

[bibalex.org](http://bibalex.org)

Internet Source

<1 %

49

[eksakta.ppj.unp.ac.id](http://eksakta.ppj.unp.ac.id)

Internet Source

<1 %

50

[ir.unilag.edu.ng](http://ir.unilag.edu.ng)

Internet Source

<1 %

51	<a href="http://journal.uad.ac.id">journal.uad.ac.id</a> Internet Source	<1 %
52	<a href="http://mdpi-res.com">mdpi-res.com</a> Internet Source	<1 %
53	<a href="http://opus.lib.uts.edu.au">opus.lib.uts.edu.au</a> Internet Source	<1 %
54	<a href="http://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a> Internet Source	<1 %
55	<a href="http://toniau.ac.ir">toniau.ac.ir</a> Internet Source	<1 %
56	<a href="http://ujcontent.uj.ac.za">ujcontent.uj.ac.za</a> Internet Source	<1 %
57	<a href="http://vtechworks.lib.vt.edu">vtechworks.lib.vt.edu</a> Internet Source	<1 %
58	<a href="http://www.alliedacademies.org">www.alliedacademies.org</a> Internet Source	<1 %
59	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	<1 %
60	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	<1 %
61	<a href="http://www.youscribe.com">www.youscribe.com</a> Internet Source	<1 %
62	Débora Cristina Sampaio de Assis, Guilherme Resende da Silva, Isabela Pereira Lanza, Ana	<1 %

Cláudia dos Santos Rossi Ribeiro, Ângela Maria Quintão Lana, Leonardo José Camargos Lara, Tadeu Chaves de Figueiredo, Silvana de Vasconcelos Caçado. "Evaluation of the Presence and Levels of Enrofloxacin, Ciprofloxacin, Sulfaquinoxaline and Oxytetracycline in Broiler Chickens after Drug Administration", PLOS ONE, 2016

Publication

63

Onah Otor Paul, Siyaka Abdulateef. "Bacterial Resistance to Commonly Prescribed Antibiotics in a Tertiary Care Hospital: A Retrospective Review of Evidence", Journal of Advances in Microbiology, 2019

Publication

<1 %

64

[repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

<1 %

65

[www.msjonline.org](http://www.msjonline.org)

Internet Source

<1 %

66

"Antimicrobial Drug Resistance", Springer Science and Business Media LLC, 2017

Publication

<1 %

67

Antibiotic Discovery and Development, 2012.

Publication

<1 %

68

Feiyang Ma, Shixin Xu, Zhaoxin Tang, Zekun Li, Lu Zhang. "Use of antimicrobials in food animals and impact of transmission of

<1 %

antimicrobial resistance on humans",  
Biosafety and Health, 2020

Publication

69

Muriel Masi, Mathias Winterhalter, Jean-Marie Pagès. "Chapter 4 Outer Membrane Porins", Springer Science and Business Media LLC, 2019

Publication

<1 %

70

Wesam Elsayed Gawad, Omneya Mohamed Helmy, Wael Mostafa Tawakkol, Abdelgawad Mohamed Hashem. "Antimicrobial Resistance, Biofilm Formation, and Phylogenetic Grouping of Uropathogenic Escherichia coli Isolates in Egypt: The Role of Efflux Pump-Mediated Resistance", Jundishapur Journal of Microbiology, 2018

Publication

<1 %

71

[scholarworks.gsu.edu](https://scholarworks.gsu.edu)

Internet Source

<1 %

72

[www.efsa.europa.eu](http://www.efsa.europa.eu)

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# A Review of Antimicrobial Resistance (AMR) of Escherichia coli on Livestock and Animal Products: Public Health Importance

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---