

Source details

Ethiopian journal of health sciences

Open Access ⓘ

Scopus coverage years: from 2013 to Present

Publisher: Research and publications office, Jimma University

ISSN: 1029-1857

Subject area: Medicine: General Medicine

Source type: Journal

CiteScore 2022

1.8 ⓘ

SJR 2022

0.344 ⓘ

[View all documents >](#) [Set document alert](#) [Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology ⓘ

CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022 ▼

1.8 = $\frac{901 \text{ Citations 2019 - 2022}}{514 \text{ Documents 2019 - 2022}}$

Calculated on 05 May, 2023

CiteScoreTracker 2023 ⓘ

1.3 = $\frac{563 \text{ Citations to date}}{440 \text{ Documents to date}}$

Last updated on 05 May, 2023 • Updated monthly

CiteScore rank 2022 ⓘ

Category	Rank	Percentile
Medicine		
General Medicine	#427/830	48th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)






SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Ethiopian journal of health sciences

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
<p>Ethiopia</p> <div style="background-color: #333; color: white; padding: 5px; margin-bottom: 5px;">  Universities and research institutions in Ethiopia </div> <div style="background-color: #333; color: white; padding: 5px;">  Media Ranking in Ethiopia </div>	<p>Medicine</p> <ul style="list-style-type: none"> ↳ Medicine (miscellaneous) 	<p>Research and publications office, Jimma University</p> <div style="background-color: #333; color: white; padding: 5px; margin-top: 10px;">  Jimma University in Scimago Institutions Rankings </div>	<h1>29</h1>
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	10291857	2013-2022	<p>Homepage</p> <p>How to publish in this journal</p> <p>asratab@yahoo.com</p>







SCOPE


The first issue of the Ethiopian Journal of Health Sciences, published by the College of Health Science of Jimma University, appeared in July 1990 with the title 'Bulletin of Jimma Institute of Health Sciences'. The journal used to appear twice a year until July 2007. Thereafter, it used to be published trice a year until it became a quarterly publication in 2014. Since January 2016, it is appearing every other month. It is an open access journal available at <http://ejhs.ju.edu.et>. The journal publishes peer-reviewed articles related to Public Health and Medicine. The objectives of the journal are: To provide current scientific and technological information on health and related fields for informed planning and decision making. To contribute to the scientific knowledge and practices in medicine, public health and related fields by providing a formal means for researchers to share their scientific research works, observations and experiences.




Join the conversation about this journal

Quartiles





● Cited documents ● Uncited documents



Ethiopian journal of health sciences

Q3 Medicine (miscellaneous) best quartile

SJR 2022 0.34

powered by scimagojr.com


← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool**.



Metrics based on Scopus® data as of April 2023



Ririh Yudhastuti 2 weeks ago

Thank you

← reply



Ririh Yudhastuti 2 weeks ago

I have sent an article for this journal with the title Integration of Climatic Factors , Distribution of Dengue Fever, Topography, n Holiday season in the Gayo Highlands region of Aceh province , February 2023, how are you doing, please provide info.

← reply



HOME

ABOUT US

EDITORIAL BOARD

INDEXING

MEMBERSHIP

CONTACT US

EJHS

- ▶ Issues

- ▶ Information to authors

- ▶ Information for reviewers

- ▶ Partnerships

- ▶ Links

Home

EDITORIAL BOARD

1. EDITORIAL BOARD

1.1. Editor-in-Chief

Prof. Abraham Haileamlak, MD, Pediatric Cardiologist, Jimma University
(asratab@yahoo.com, kasechab@gmail.com, Abraham.hamlak@ju.edu.et, ejhs@ju.edu.et)

1.2. Associate Editors

Prof. Tefera Belachew, MD, MPH, PhD, Nutritionist, Jimma University (teferabelachew@gmail.com)
Prof. Tsinuel Girma, MD, PhD, Pediatrician, Nutritionist, Jimma University (tsinuel@yahoo.com)
Prof. Gemed Abebe, M.Sc, PhD, Molecular Biologist, Jimma University
(gemedabebe@ju.edu.et, biftuus@yahoo.com)

1.3. Editorial Board Members

Prof. Kifle Woldemichael, MD, MPH, Epidemiologist, Jimma University (bethy_kifle@yahoo.com)
Mr. Fasil Tessema, M.Sc, Biostatistician, Jimma University (alazarfasil@yahoo.com ,
fasil.tessema@ju.edu.et)
Prof. Argaw Ambelu, M.Sc, PhD, Environmental Health Specialist, Jimma University
(aambelu@yahoo.com, argaw.ambelu@ju.edu.et, argawambelu@gmail.com)
Prof. Mirkuzie Woldie, MD, MPH, MOH, Ethiopia (mirkuzie@yahoo.com, mirkuzie.woldie@ju.edu.et)
Dr. Muluembet Abera, MPH, PhD, Reproductive Health Specialist, Jimma University
mulu_abera.ts2009@yahoo.com)
Prof. Yeshigeta Gelaw, MD, Ophthalmologist, Bahir-Dar University (dryeshi@yahoo.com)
Prof. Markos Tesfaye, MD, Psychiatrist, Oslo, Norway (tesmarkos@yahoo.com)
Dr. Wondwosen Kassahun, M.Sc, PhD, Biostatistician, USA (wondkyt@yahoo.com)
Prof. Esayas Kebede, MD, PhD, Internist, Jimma University (esakgd@gmail.com)
Dr. Tsedeke Asaminew, MD, Ophthalmologist, SPHMMC (tsatsedeke@gmail.com)
Dr. Bosena Tebeje, M.Sc, Reproductive Health Nursing, Jimma University (bosenatebeje@yahoo.co.uk)
Prof. Delenasaw Yewhalaw, M.Sc, PhD, Entomologist, Jimma University (delenasawye@yahoo.com,
delenasaw.yewhalaw@ju.edu.et)
Dr. Alemseged Abdissa, M.Sc, PhD, Microbiologist, AHRI (alemable@yahoo.com)
Mr. Henok Asefa, M.Sc, Biostatistician, Jimma University (menothan@gmail.com)
Dr. Seid Mussa, M.Sc, Social Pharmacist, Jimma University (plss1176@yahoo.com)

2. EDITORIAL MANAGER

Dr. Tekle Ferede, MA, PhD, Linguist, Jimma University (tekle.ferede2014@gmail.com)

3. INTERN EDITORS

Sr. Enatfenta Sewmehon, M.Sc Nurse, Jimma University (enatfenta@gmail.com)
Mr. Yibeltal Siraneh, MPH, Health Service Management Specialist, Jimma University
(yibeltal_siraneh@yahoo.com)

4. EDITORIAL CONSULTANTS

Prof. EHO Parry (UK) (eldryd@thet.org)
Prof. Jeanette Magnus (Norway) (j.h.magnus@gmail.com, j.h.magnus@medisin.uio.no)
Prof. Ib Christian Bygbjerg (Denmark) (iby@sund.ku.dk)
Prof. Yemane Berhane (Ethiopia) (yemaneberhane12@gmail.com)
Prof. Tsige Gebremariam (Ethiopia) (tsigegm@yahoo.com, tsige.gebremariam@aau.edu.et)
Prof. Eyasu Makonnen (Ethiopia) (eyasu.makonnen@aau.edu.et)
Prof. William Bazeyo (Uganda) (wbazeyo@gmail.com, wbazeyo@musph.ac.ug)
Prof. David Serwadda (Uganda) (dserwada@imul.com)
Prof. Lulu Muhe (Ethiopia) (muhe1952@gmail.com, muhel@who.int)
Dr. Christine Laine (USA) (claine@acponline.org)
Dr. Assaye Kassie (USA) (assaye.nigussie@gatesfoundation.org)
Prof. Atalay Alem (Ethiopia) (atalayalem@yahoo.com, atalay.alem@gmail.com)

[Home](#)[Current](#)[Archives](#)[About](#)[Home](#) / [Archives](#) / [Vol. 32 No. 3 \(2022\)](#)

Articles

Table of content

yibeltal Siraneh



Editorial message

Abraham Haileamlak



Prevalence of COVID-19 Vaccine Side Effects among Early-Vaccinated Healthcare Workers in Eastern Ethiopia

Godana Jarso, Wassie Gebi, Meyrema Abdo, Misgana Lemma, Endashaw Abebe, Bekana Lemessa, Biniyam Tefera Deressa



Clinical Manifestations, Imaging Procedures and Laboratory Parameters among Hospitalized Patients with COVID-19 in Ilam Province, Western Iran

Mohammad Reza Kaffashian, Maryam Shirani, Maryam Koupaei, Nourkhoda Sadeghifard, Iraj Ahmadi, Aliashraf Mozafari, Ali Nazari, Mohsen Heidary, Saeed Khoshnood



Admission Pattern and Treatment Outcome in Pediatric Intensive Care Unit, Tertiary Hospital, Addis Ababa, Ethiopia

Gemechu Edae, Atnafu Mekonnen Tekleab, Melaku Getachew, Tigist Bacha



Factors Associated with ICU Mortality at Hawassa University Comprehensive Specialized Hospital (HUCSH)

Abdi Bati Wotiye, Emnet Tesfaye Shimber, Biniyam A. Ayele



Risk Factors of Birth Asphyxia among Newborns at Debre Markos Comprehensive Specialized Referral Hospital, Northwest Ethiopia: Unmatched Case-Control Study

Yoseph Merkeb Alamneh, Ayenew Negesse, Yared Asmare Aynalem, Wondimeneh Shibabaw Shiferaw, Mihretie Gedefew, Melkamu Tilahun, Yidersal Hune, Abtie Abebaw, Yalemgeta Biazin, Tadesse Yirga Akalu



Bacterial Sepsis among Children with Congenital Heart Disease in Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia

Edomgenet Tesfaye, Henok Tadele



Establishing a Self-sustaining Emergency Medicine Point-of-Care Ultrasound Curriculum in an Academic Teaching Hospital in Ethiopia

Anne Aspler, Fraser Kegel, Temesgen Beyene, Tigist Zewdu, Berhanu Tesfaye, Alexandra McKnight, Eileen Cheung, Jennifer Bryan, Claire Acton



Factors Associated with the Extent of Coronary Artery Disease and the Attained Outcome of Percutaneous Coronary Intervention at Gesund Cardiac and Medical Center, Addis Ababa, Ethiopia

Bekele Alemayehu Shashu, Ararso Baru



Electrocardiographic Changes and Serum Troponin Levels in Patients with Acute Stroke: A Prospective Cross-Sectional Study

Etedal Ahmed A. Ibrahim, Salma Mohamed Taha, Khabab Abbasher Hussien Mohamed Ahmed, Mazin S. Haroun, Mohammed Eltahier Abdalla Omer



Modeling Time to Cure of Deep Vein Thrombosis Using Cox Proportional Model in Southwest of Ethiopia

Gurmessa Nugussu Gelcho, Mosisa Girma Bekele



Mass Hysteria among Beneficiary Students of the School-Feeding Program in Addis Ababa, Ethiopia

Solomie Jebessa, Handsome Deksiso, Muluwork Tefera, Yonas Bahretibeb



Impact of Stunting on Development of Children between 1-3 Years of Age

Muhammad R. D. Mustakim, Irwanto, Roedi Irawan, Mira Irmawati, Bagus Setyofoedi



Early Postnatal Care Utilization among Rural Women in Horo Guduru Wollega Zone, Ethiopia

Lalisa Ayele Woldasemayat, Abiru Neme Negawo, Chaluma Kumela Mengesha, Tilahun Fufa Debela



Knowledge of the Therapeutic Goals of Diabetes Care among Patients with Type 2 Diabetes at a Tertiary Hospital in Ethiopia

Melaku Taye, Kehabtimer Shiferaw, Paulos Efreem, Getahun Tarekegn



Correlation between Radiological Images and Histopathological Type of Meningioma: A Cohort Study

Andi Ihwan, Rauf Rafika, Muhammad Husni Cangara, Kevin Jonathan Sjukur, Muhammad Faruk



Assessing the Complications of Monopolar Transurethral Resection of the Prostate (M-TURP) Using Clavien-Dindo Complications Grading System

Lijalem Mekonnen Geremew, Samuel Amare Gelaw, Andualem Deneke Beyene



Clinical Profiles and Surgical Outcome of Hypospadias Repair at a Teaching Hospital in Ethiopia

Maru Gama, Birhan Abitew, Kirubel Abebe



Asymptomatic Malaria in Households and Neighbors of Laboratory Confirmed Cases in Raya Kobo District, Northeast Ethiopia

Yimer Melese, Megbaru Alemu, Mulat Yimer, Banchamlak Tegegne, Tigist Tadele



Efficacy of Praziquantel Treatment and Schistosoma Mansonii Infection among Primary School Children in Kemisse Town, Northeast Ethiopia

Meslo Sema Berhanu, Seyfe Asrade Atnafie, Tahir Eyayu Ali, Aderaw Adamu Chekol, Habtamu Biazin Kebede



The Effect of Teff Seed on Hematological Findings and Anthropometric Measurements

Eftal Geççil Demir, Nadide Gizem Tarakçı, Ruken Aslınur Samancı, Merve Cambaz, Şeymanur Bilici, Dilara Tuygan, Büşra Çalık, Ayşe Sümeyye Çiftçi



Antibacterial Effects of Artemisia afra Leaf Crude Extract Against Some Selected Multi-Antibiotic Resistant Clinical Pathogens

Alemtsehay Beyene Haile, Tamene Milkessa Jiru



Imaging Findings of Concomitant Pulmonary and Central Nervous System Sarcoidosis: A Case Report

Tewodros Endale Balcha, Abebe Mekonnen Woldeyohannes, Azmera Gissila Aboye



Pyogenic Liver Abscess Presenting as an Initial Manifestation of Underlying Hepatocellular Cancer: A Case Report in Ethiopia

Hailemichael Desalegn, Ayantu Tesfaye, Paulos Shume



COVID-19 Vaccination and Disability in Ethiopia: Ensuring No One is Left Behind

Abubeker Alebachew Seid



Journal Identifiers

eISSN: 2413-7170
print ISSN: 1029-1857



ORIGINAL ARTICLE**Impact of Stunting on Development of Children between 1-3 Years of Age****Muhammad R. D. Mustakim¹, Irwanto^{1*}, Roedi Irawan¹, Mira Irmawati¹, Bagus Setyo-boedi¹****OPEN ACCESS**

Citation: Muhammad R. D. Mustakim, Irwanto, Roedi Irawan, Mira Irmawati, Bagus Setyo-boedi. Impact of Stunting on Development of Children between 1-3 Years of Age. *Ethiop J Health Sci.* 2022;32 (3):569. doi:<http://dx.doi.org/10.4314/ejhs.v32i3.13>

Received: October 28, 2021

Accepted: December 30, 2021

Published: May 1, 2022

Copyright: © 2022 Muhammad R.D., et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Nil

Competing Interests: The authors declare that this manuscript was approved by all authors in its form and that no competing interest exists.

Affiliation and Correspondence:

¹Department of Child Health, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia
Email: irwanto.idris@gmail.com

ABSTRACT

BACKGROUND: Stunting occurs due to chronic malnutrition and is a major problem for children in developing countries. It is important to evaluate the impact of stunting on the development of children. This study aimed to investigate the impact of stunting on the development of children between 1-3 years of age.

METHODS: This cross-sectional study was conducted from July 2020 to March 2021 in Surabaya, Indonesia. A questionnaire and growth assessment were done, following the development measurement to stunted and non-stunted children who met the inclusion and exclusion criteria. Development was measured by the Denver Developmental Screening Test II (DDST-II), and Cognitive Adaptive Test/Clinical Linguistic & Auditory Milestone (CAT/CLAMS) scales.

Results: Three hundred children are included in this study, consisting of 150 stunted and 150 non-stunted children. Stunted children had a higher risk to be suspected of delayed development compared to non-stunted children. The Crude Odd Ratio was 2.98, 4.24, 4.75 with the p-value 0.006, 0.001, and 0.001 respectively. The Adjusted Odd Ratio was 0.34, 0.24, 0.21 with p-value of 0.008, 0.001, and 0.001 respectively.

CONCLUSION: Stunting is associated with suspected development delay among children 1-3 years of age. Initiatives related to prevention need to be established and nutrition advice needs to be provided.

KEYWORDS: Children, Development, Stunting

INTRODUCTION

Optimal nutritional care is one of the important factors needed to produce optimal growth and development, especially in children under five years of age. Nutritional deficiencies at that time will have an impact on growth and determine the further development of children (1). One form of malnutrition that is currently a global problem is stunting. Stunting is an impact that occurs due to chronic malnutrition and is a major problem for children in rural areas who experience developmental disabilities. The United Nations Children's Fund (UNICEF) states that the prevalence of stunting in the under-five population globally reaches

21.9% (2,3). Based on these data, the highest number of stunting cases is in Africa and Asia, including Indonesia. Indonesia's basic Health Research survey shows that more than a third of children (30.8%) of the population under five are stunted. This condition must be overcome.

The most frequent form of child malnutrition is stunting (4). Malnutrition in early life might cause inflammation, changes in leptin levels, and increased glucocorticoids resulting in epigenetic changes. These changes may cause neurodevelopmental disorders, changes in neurogenesis and cell apoptosis as well as dysfunction of synapses resulting in developmental delay (5). It was concluded that malnutrition affects areas of the brain involved in cognition, memory and locomotor skills (6-8). The relationship between stunting and cognitive function also has been demonstrated (9). Children who were persistently stunted had significant lower cognition than children who were not stunted (10). The effects of stunting on neurocognitive function are severe. Stunted children have stunted brains and live stunted lives, preventing entire communities from developing (4). Growth stalling often starts in utero and lasts for at least the first two years after birth. Human development is hampered by the severe irreparable physical and neurocognitive harm that comes with stunted growth. Children with stunting have a 3.6 times higher risk of cognitive impairment than children without stunting (11). One effort that can be done to prevent stunting is to improve the nutritional intake for pregnant women and toddlers. This practical approach can be applied to prevent stunting, with regards to reducing the prevalence of child developmental disorders. In connection with the negative impact of stunting on children's development, it is necessary to evaluate the development of children who are stunted. Indonesia has the fifth-highest level of stunting in children among countries (3). This condition was exacerbated by the Covid-19 pandemic which had an impact on the economy of the community, causing an increase in the number of cases of stunting due to a decrease in

the ability of parents to meet children's nutritional needs. The high incidence of stunting surely carries high burden for the country. Research conducted to assess the effect of stunting on children's development has not been done much. This condition triggers the researchers to further observe the impact of stunting on the development of children, especially in children between 1-3 years of age.

METHODS

We conducted a cross-sectional study from July 2020 to March 2021. The population in this study were children aged 1 to 3 years old who were in Surabaya, Indonesia. We used consecutive sampling by collecting the participants with stratified random sampling. The participants of this study were stunted and non-stunted children who met the inclusion and exclusion criteria. We collected the participants from Primary Health Care which are located in Surabaya, Indonesia. We categorized the participants of the study into two groups, stunted and non-stunted children. The inclusion criteria of stunted children are 1) children aged from 1 to 3 years old who diagnosed stunted in Primary Health Care 2) proportionate short stature 3) has a history of chronic malnutrition, and 4) parents are willing to sign an informed consent to participate in the study. The definition of stunted children was based on WHO criteria with a length/height below - 2 Standard Deviation who have experienced linear growth faltering (12). The inclusion criteria of non-stunted children are as follows: 1) healthy children aged from 1 to 3 years, 2) parents are willing to sign an informed consent to participate in the study. The exclusion criteria of the study are: 1) children have a history of congenital anomaly, 2) has a history of other chronic diseases 3) has severe neurological deficits, and 4) children with short stature due to genetic or hormonal disorders. The ethical clearance of this study was approved and issued by the Health Research Ethics Committee, Faculty of Medicine, Universitas Airlangga.

Data collection was carried out by visiting and measuring the patients one by one after obtaining permission from the Surabaya Public Health Department. The researcher has been

swabbed PCR-COVID19 and declared negative. The subject and their family living in the same house have been swabbed Antigen-COVID19 and declared non-reactive. After the parents are willing to sign the informed consent. We deliver a brief explanation regarding the study and we gave a questionnaire for parents to be filled in. The questionnaire contains the data on the characteristics of participants including chronological age, gender, history of breastfeeding, socio-economic, and parental education. A questionnaire was also conducted to ensure that no exclusion criteria were found in the study participants. Following the questionnaire, we started to measure the growth in children. After the growth measurement was done, the children will be assessed for development using 2 measuring instruments. The first measuring instrument was the Denver Developmental Screening Test (DDST-II) to measure fine motor, gross motor, and personal-social development (13). The participants are categorized as suspects if they have one or more delays and/or two or more cautions, while considered as normal if they have no delay in any domain and no more than one caution. The second instrument was the Cognitive Adaptive Test/Clinical Linguistic & Auditory Milestone (CAT/CLAMS) scales to measure cognitive function (14). We categorized the participants as suspect if they have scored below 85, whereas the normal participant was classified if they have scored above or equal to 85.

Descriptive analysis was performed using statistical measures of mean \pm standard deviation (SD), number (percentage), median, and interquartile range (IQR). Mann-Whitney U

tests are used for the data that are not normally distributed. The bivariate analysis was tested to seek the correlation between stunting and the children's development. The Crude Odd Ratio (COR) was obtained using the Chi-Square test and Adjusted Odd Ratio (AOR) was acquired using Binary Logistic Regression to perceive the determinant factors of stunting in children's development. The analysis of the data was done with IBM SPSS Statistic Version 23. A p-value < 0.05 was considered statistically significant.

RESULTS

A total of 300 participants were included in this study. We divided the participants into two groups, 150 stunted children and 150 non-stunted children. The mean age of the children is 23.12 months ranging from 12 to 35 months. The data of socio-demographic and characteristics of all participants are presented in Table 1. The mean difference of actual height in stunted children and non-stunted children is 6.78 cm. There are significant differences between stunted and non-stunted children especially from the history of exclusive breastfeeding and physical appearance such as body weight, height, and head length ($p = 0.001$) (Table 1). In Table 1, we also compared the risk factors that might be the causes as to why children are stunted. The children who are categorized as the suspect in delayed development are found mostly in stunted children. Significant differences, mostly on motor function (both in fine and gross motor function), are found in development of stunted and not stunted children (Table 2).

Table 1: Socio-demographic and characteristics of stunted and non-stunted children.

	Non-stunted N = 150	Stunted N = 150	<i>p</i>
Characteristic of children			
Child's age (months)	22.91 ± 6.26	23.33 ± 6.54	0.502 ^a
Sex			0.562 ^b
Female	64 (42.7)	70 (46.7)	
Male	86 (57.3)	80 (53.3)	
Gestational age (weeks)	38 [36 – 29]	38 [25 – 42]	0.052 ^b
Term	147 (98)	139 (92.7)	
Preterm	3 (2)	11 (7.3)	
Type of delivery			
Spontaneous	120 (80)	88 (58.7)	
Caesarian section	30 (20)	62 (41.3)	
Birth weight (gram)	3030 ± 230.11	2953.27 ± 463.62	0.001 ^{b*}
≥ 2500 gram	149 (99.3)	134 (89.3)	
< 2500 gram	1 (0.7)	11 (10.7)	
Mother's age when pregnant	26.53 ± 6.60	26.94 ± 5.38	0.913 ^a
Exclusive breastfeeding			0.001 ^{b*}
Yes	139 (92.7)	102 (68)	
No	11 (46.3)	48 (32)	
Actual weight (kg) ^a	10.59 ± 1.96	9.67 ± 1.86	0.001 ^{a*}
Actual height (cm) ^a	85.49 ± 8.07	78.71 ± 6.02	0.001 ^{a*}
Actual head length (cm) ^a	47.76 ± 2.78	46.88 ± 3.04	0.001 ^{a*}
Socio-economic of family			
Father's education			0.100 ^b
Primary School	0 (0)	6 (2)	
Middle School	18 (12)	16 (10.7)	
Secondary School	66 (44)	66 (44)	
Tertiary education	66 (44)	62 (41.3)	
Mother's education			0.183 ^b
Primary School	2 (1.3)	8 (5.3)	
Middle School	16 (10.7)	16 (11.3)	
Secondary School	94 (62.7)	82 (54.7)	
Tertiary education	38 (25.3)	44 (29.3)	
Family income			0.491 ^b
< regional minimum wage	113 (75.3)	119 (79.3)	
≥ regional minimum wage	37 (24.7)	31 (20.7)	
Parenting			0.376 ^b
Parents	135 (90)	129 (86)	
Other family member	13 (8.7)	20 (13.3)	
Babysitter	2 (1.3)	1 (0.7)	

Data are presented as N (%), Mean ± Standard Deviation, Median [Interquartile range];

The data are not normally distributed; ^a Mann-Whitney U test & ^b Chi-Square are used; *a p-value < 0.05 was statistically significant.

Table 2. The results of development on stunted and non-stunted children between 1-3 years of age.

	Non-stunted N = 150	Stunted N = 150	<i>p</i>
DENVER II Developmental Status			
Normal	141 (94)	126 (84)	0.006*
Suspect	9 (6)	24 (16)	
CAT/CLAMS			
CAT			
Normal	122 (81.3)	76 (50.7)	0.001*
Suspect	28 (18.7)	74 (49.3)	
CLAMS			
Normal	114 (76)	60 (40)	0.001*
Suspect	36 (24)	90 (60)	

*a p-value < 0.05 was statistically significant; Chi-square analysis was used for the analysis.

Table 3 summarizes the COR and AOR of developmental function. The table shows that stunted children had a higher risk for children to experience the development delay compared to non-stunted children COR: 2.984, $p = 0.006$; AOR: 0.335, $p = 0.008$. We also found the same situation in the CAT/CLAMS instrument measuring the development of stunted children with $p = 0.001$ (Table 3). Stunted children can have the risk of experiencing delayed non-

language visual-motor skills (CAT) by 4.242 times compared to non-stunted children (AOR 0.236; $p = 0.001$). This case was also the same with language skill (CLAMS), stunting for children have the risk of at least 4.750 times experiencing delayed language skills compared to not stunted children (AOR 0.211; $p = 0.001$). Table 4 also present determinant factors which contributed in suspect delayed development.

Table 3: Bivariate and multivariate results between stunted and delayed development.

	Crude OR	95% CI (Lower – Upper)	<i>p</i>	Adjusted OR	95% CI (Lower – Upper)	<i>p</i>
DENVER II						
Stunted	2.984	1.337 – 6.660	0.006*	0.335	0.150 – 0.748	0.008*
Yes						
No						
CAT						
Stunted	4.242	2.520 – 7.141	0.001*	0.236	0.140 – 0.397	0.001*
Yes						
No						
CLAMS						
Stunted	4.750	2.889 – 7.809	0.001*	0.211	0.128 – 0.346	0.001*
Yes						
No						

Crude OR used Chi-Square Test; Adjusted OR used Binary Logistic Regression;

*a p-value < 0.05 was statistically significant.

Table 4: The Clinical risk factors of children between 1-3 years old compared with development Adjusted OR used Binary Logistic Regression; *a p-value < 0.05 was statistically significant.

Variable	Adjusted OR	95% CI (Lower – Upper)	p
DENVER II			
Birth weight (gram)	0.793	0.202 – 3.118	0.740
Gestational age (weeks)	1.145	0.229 – 5.720	0.869
Father's education	0.820	0.318 – 2.113	0.681
Mother's education	0.465	0.200 – 1.082	0.075
CAT			
Birth weight (gram)	0.592	0.202 – 1.743	0.342
Gestational age (weeks)	0.715	0.227 – 2.246	0.565
Father's education	1.491	0.800 – 2.779	0.209
Mother's education	0.431	0.227 – 0.816	0.010*
CLAMS			
Birth weight (gram)	0.510	0.175 – 1.482	0.216
Gestational age (weeks)	0.858	0.272 – 2.709	0.794
Father's education	0.990	0.548 – 1.788	0.973
Mother's education	0.497	0.270 – 0.915	0.025*

DISCUSSION

This study found a significant relationship between stunting with development of children aged 1-3 years. Delayed development in stunted children is caused by nutritional deficiencies that occur early in life. The intrauterine period is the first stage of the critical period of child development. In that period, nutritional factors play an important role in the maturation process of the central nervous system (15). A study conducted on animals and humans showed that the chronicity, time, and severity of nutritional deficiencies have an effect on brain development that will have an impact on subsequent developmental abilities (16). We found that children who were stunted had a greater risk of experiencing delays in motor development, both gross and fine motor. If the muscle mechanism has not developed properly, motor movement will not be perfect. This eventuality occurs in children with stunted development abnormalities, striped muscle or striated muscle that regulates subconscious actions that develop at a slower rate, and there can be no coordinated voluntary action before the child is in normal condition (17).

These results are coherent with a study that shows that acute malnutrition that occurs in children aged 1 year is associated with developmental delays in gross motor, fine motor, social, and language interactions (18). Similar results were also obtained from a study on stunted children aged 9-24 months, which showed that the fine motor skills of stunted children were worse than not stunted children (19). Another study in Bangladesh showed that stunted children had lower motor scores than not stunted children. They also reported that stunting at <2 years of age was a predictor of delays in motor development (20).

As previously described, several studies stated that the delays in motor development experienced by stunted children were mainly related to the maturation process of the central nervous system. However, low height in stunted children was also a factor in the delay (15). A study in 2013 showed that there were significant associations between gross motor development and anthropometry measures at 12 months. Increment of one height z-score in terms of the decrease in the average time taken to 'lie to sit' of 1.80 months (p = 0.03) (21). A meta-analysis study conducted in 2015 shows a significant

correlation between height and motor skills in children in low-middle income Countries with a correlation value of 0.24. The meta-analysis also shows that height measures significantly affect children's walking ability earlier ($p < 0.05$) (22).

We also found that stunted children also had significantly lower cognitive scores than those who were not stunted. The results of the present study are similar to a Peruvian study that showed that when compared to their non-stunted peers, children who were stunted from the age of 6 months to 6 years or stunted during childhood had considerably worse scores on cognitive capabilities (verbal vocabulary and quantitative test scores) at the age of 4.5 to 6 years (22). A study in Ethiopia found that stunting harms the cognitive development of children at the age of five and eight years old (11). The quantitative assessment for cognitive function score was higher for non-stunted children with 8.60 percentage points of difference (in favour of non-stunted children) with a significant statistical value ($p < 0.01$) (11). The link between linear growth and cognitive development in the first two years of life was discovered in a meta-analysis of 29 LMICs, but the meta-analysis was unable to integrate educational, follow-up, and environmental data to describe the link between stunting and cognitive impairment in children (22). However, whether these changes are permanent or reversible is still a point of contention in the literature. A recent study showed that children who experienced catch-up growth had no difference in verbal vocabulary and quantitative test scores compared to non-stunted children. The authors concluded that children can recover from early nutritional deficiencies and improve their cognitive performance (23).

Our analysis showed that gestational week was positively associated with gross motor skills. This result is in line with a study in 2014 that stated that small for gestational age infants are more susceptible to motor developmental delay but may decrease with increasing age when compared to term infants (24). This is in contrast with the study done by Vungarala and Rajeswari (2018) that found no significant correlation between gestational age and motor

development at eight months of corrected age (25). This study confirms that other factors such as the father's education are also important determinants of cognitive function. A father's education was positively associated with a child's cognitive function (26). This phenomenon is associated with better verbal activity skills in fathers who have higher education (i.e., college-level) (27). A systematic review regarding the impact of father involvement on children's cognitive skills showed that children whose fathers are content with their parenting and financially supportive of their families have superior linguistic and cognitive skills, according to research. In this study, the father's education is one of the determinants that affect children's cognitive function. Another study in Ecuador stated that parents' education has important implications for child cognitive development. The schooling and vocabulary levels of mothers were strong predictors of the cognitive development of young children (28).

In this study, we also found that fewer children in the stunting group received exclusive breastfeeding than the not stunted group. Breastfeeding supplies the newborn with colostrum, which is high in nutrients, as well as antibodies, which are essential for the formation of the gut microbiota and immune system. Any additional drink or food introduced before the age of four months is linked to an increased risk of gastrointestinal disorders, which can lead to growth retardation, nutritional deficiencies, and predisposition to infectious diseases in the first two years of life. The link between exclusive breastfeeding and length-for-age and weight-for-age was verified in a study by Kuchenbecker et al. ($p = 0.001$) (29). Exclusively breastfed infants under the age of six months were taller, heavier, and less likely to be stunted than non-exclusively breastfed infants.

We also found that stunted children also had lower birth weight than those who were not stunted. History of birth weight less than 2500 grams were found to be more in the stunting group than the control group. These results are also consistent with a study, which stated that birth weight less than 2500 grams was a risk

factor for stunting in children aged 6-24 months (AOR = 5.3) (30). Similar studies were also found in studies conducted in Ethiopia which stated that birth weight is a strong predictor in determining the anthropometric status of children in the future, this is because most babies with low birth weight are not able to catch up with normal growth in childhood (11).

This study is the first that analyzes the Impact of Stunting on the Development of Children between 1-3 Years of Age while in the COVID-19 pandemic. However, we admit that this study has limitations concerning other factors that are not properly discussed. Certain characteristics, which may be major predictors of poor motor and cognitive skills in children throughout their early years of life, were not evaluated. For example, maternal health, growth hormone, nutritional deficiency, lack of kid stimulation, violent exposure, and some environmental factors.

Stunting is associated with delayed development among children 1-3 years of age, mostly in motor function. Initiatives related to prevention need to be established and nutrition advice needs to be provided. Our study also confirms that other factors such as the gestational week and the father's education are also important determinants of development. Periodic screening is necessary, especially in children aged 1-3 years. Stunting prevention programs must be one of the top priorities for health workers in services close to the community, especially primary health care. If the child shows symptoms of stunting, they are immediately referred to a paediatrician to be able to improve their nutritional status.

ACKNOWLEDGMENTS

We would like to thank the Surabaya Health Office and Primary Health Care in Surabaya, Indonesia for the permission so that we could collect the data research accordingly. We also would like to express our sincere gratitude to the health personnel from doctors, nurses, and other people in charge involved in this study for the help and support we've received, we can have the courage to complete the research.

REFERENCES

1. Mohseni M, Aryankhesal A, Kalantari N. Prevention of malnutrition among children under 5 years old in Iran: A policy analysis. *PLoS One*. 2019; 14(3): e0213136.
2. UNICEF. Malnutrition prevalence remains alarming: stunting is declining too slowly while wasting still impacts the lives of far too many young children. <https://data.unicef.org/topic/nutrition/malnutrition/> Accessed on April 15th 2021.
3. UNICEF. Improving child nutrition: the achievable imperative for global progress. New York, United Nations Publications Sales No.: E.13.XX.4. 2013: 1–114.
4. de Onis M, Branca F. Childhood stunting: a global perspective. *Matern Child Nutr*. 2016; 12: 12.
5. Martins VJB, Florêncio TMMT, Grillo LP, et al. Long-Lasting Effects of Undernutrition. *Int J Environ Res Public Health*. 2011; 8(6): 1817–46.
6. Prendergast AJ, Humphrey JG. The stunting syndrome in developing countries. *Paediatr Int Child Health*. 2014; 34(4): 250–65.
7. Ranade SC, Rose A, Rao M, Gallego J, Gressens P, Mani S. Different types of nutritional deficiencies affect different domains of spatial memory function checked in a radial arm maze. *Neuroscience*. 2008; 152(4): 859–66.
8. Soliman AT, De Sanctis V, Alaraj NM, et al. Early and Long-term Consequences of Nutritional Stunting: From Childhood to Adulthood. *Acta Biomed*. 2021; 92(1): e2021168.
9. Ocansey ME, Adu-Afarwaa S, Kumordzie SM, et al. The association of early linear growth and haemoglobin concentration with later cognitive, motor, and social-emotional development at preschool age in Ghana. *Matern Child Nutr*. 2019; 15(4): e12834.

10. Alam MA, Richard SA, Fahim SM, et al. Impact of early-onset persistent stunting on cognitive development at 5 years of age: Results from a multi-country cohort study. *PLoS One*. 2020; 15(1): e0227839.
11. Woldehanna T, Behrman JR, Araya MW. The effect of early childhood stunting on children's cognitive achievements: Evidence from young lives Ethiopia. *Ethiop J Health Dev*. 2017; 31(2): 75–84.
12. World Health Organization. Nurturing care for early childhood development: A framework for helping children survive and thrive to transform health and human potential. Geneva: 2018.
13. Frankenburg WK, Dodds JB. The Denver developmental screening test. *J Pediatr*. 1967; 71(2): 181–91.
14. Wachtel RC, Shapiro BK, Palmer FB, Allen MC, Capute AJ. CAT/CLAMS. A tool for the pediatric evaluation of infants and young children with developmental delay. Clinical Adaptive Test/Clinical Linguistic and Auditory Milestone Scale. *Clin Pediatr (Phila)*. 1994; 33(7): 410–5.
15. Muhoozi GKM, Atukunda P, Mwadime R, Iversen PO, Westerberg AC. Nutritional and developmental status among 6- to 8-month-old children in southwestern Uganda: a cross-sectional study. *Food Nutr Res*. 2016; 60: 30270.
16. Black MM. Impact of Nutrition on Growth , Brain , and Cognition. *Nestle Nutr Inst Workshop Ser*. 2018; 89: 185–95.
17. Komaini A, Mardela R. Differences of fundamental motor skills stunting and non stunting preschool children in Kindergarten in North Padang. *IOP Conf Ser: Mater Sci Eng* 2018;335 012131. <https://iopscience.iop.org/article/10.1088/1757-899X/335/1/012131/pdf>
18. Abessa TG, Bruckers L, Kolsteren P, Granitzer M. Developmental performance of hospitalized severely acutely malnourished under-six children in low- income setting. *BMC Pediatr*. 2017; 17(1): 197.
19. Chang SM, Walker SP, Christine SG. Early childhood stunting and later fine motor abilities. *Dev Med Child Neurol*. 2010; 52(9): 831-6.
20. Dwivedi D, Singh S, Singh J, Bajaj N, Singh H. Neurodevelopmental Status of Children aged 6-30 months with Severe Acute Malnutrition. *Indian Pediatr*. 2018; 55(2): 131–3.
21. Ireland PJ, Ware RS, Donaghey S, et al. The effect of height, weight and head circumference on gross motor development in achondroplasia. *J Paediatr Child Health*. 2013; 49(2): E122-7.
22. Sudfeld CR, McCoy DC, Danaei G, et al. Linear growth and child development in low- and middle-income countries: a meta-analysis. *Pediatrics*. 2015; 135(5): e1266.
23. Crookston BT, Penny ME, Alder SC, et al. Children who recover from early stunting and children who are not stunted demonstrate similar levels of cognition. *J Nutr*. 2010; 140(11): 1996–2001.
24. Moreira R, Magalhães L, Alves C. Effect of preterm birth on motor development, behavior, and school performance of school-age children: a systematic review. *J Pediatr (Rio J)*. 2014; 90(2): 119–34.
25. Vungarala P, Rajeswari M. Correlation of birth weight, gestational age and muscle tone with motor development of preterm infants. *Int J Physiother*. 2018; 5(2): 63–8.
26. Rollè L, Gullotta G, Trombetta T, et al. Father Involvement and Cognitive Development in Early and Middle Childhood: A Systematic Review. *Front Psychol*. 2019; 10: 2405.

27. Cabrera NJ, Volling BL, Barr R. Fathers Are Parents, Too! Widening the Lens on Parenting for Children's Development. *Child Dev Perspect.* 2018; 12(3): 152–7.
28. Schady, Norbert. Parents' Education, Mothers' Vocabulary, and Cognitive Development in Early Childhood: Longitudinal Evidence From Ecuador. *Am J Public Health.* 2011; 101(12): 2299–2307.
29. Kuchenbecker J, Jordan I, Reinbott A, et al. Exclusive breastfeeding and its effect on growth of malawian infants: Results from a cross-sectional study. *Paediatr Int Child Health.* 2015; 35(1): 14–23.
30. Berhe K, Seid O, Gebremariam Y, Berhe A, Etsay N. Risk factors of stunting (chronic undernutrition) of children aged 6 to 24 months in Mekelle City, Tigray Region, North Ethiopia: An unmatched case-control study. *PlosOne.* 2019; 14(6): e0217736.