

Adiponectin, anthropometric measurements and insulin resistance in adolescence with obesity

by Nur Aisyah Widjaja

Submission date: 12-Jun-2023 09:21PM (UTC+0800)

Submission ID: 2114498441

File name: 04._Adiponectin,_anthropometric_measurements.pdf (238.88K)

Word count: 3717

Character count: 20867

Adiponectin, anthropometric measurements and insulin resistance in adolescence with obesity

Nur Aisiyah Widjaja^{1,2*}, Rendi Aji Prihaningtyas¹, Meta Herdiana Hanindita¹,
Roedi Irawan¹, IDG Ugrasena¹ & Retno Handajani³

¹Department of Child Health, Dr. Soetomo General Academic Hospital, Faculty of Medicine, Universitas Airlangga, Jl. Mayjen Prof. Dr. Moestopo no 6-8, Surabaya, Indonesia; ²Medical Doctoral Program Student, Faculty of Medicine, Universitas Airlangga, Jl. Mayjen Prof. Dr. Moestopo no 47, Surabaya, Indonesia; ³Department of Biochemistry, Faculty of Medicine, Universitas Airlangga, Jl. Mayjen Prof. Dr. Moestopo no 47, Surabaya, Indonesia

ABSTRACT

Introduction: Obesity in adolescents can cause metabolic syndrome. Insulin resistance increases the risk of metabolic syndrome, which then increases the risk of premature death. Studies about anthropometric measurements and adiponectin levels as early markers of insulin resistance in obese adolescents are still limited. **Methods:** A cross-sectional study was performed on 59 obese adolescents aged 13–16 years. Obesity was established on the basis of the Centers for Disease Control and Prevention (CDC) curve (2000). Insulin and blood glucose level measurements were carried out using an enzymatic kit. Adiponectin levels were assayed using enzyme-linked immunosorbent assay (ELISA). The relationships between variables were evaluated by correlation analysis using SPSS. **Results:** Statistical tests showed a positive correlation between waist circumference ($r=0.421$; $p=0.001$) and Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) ($r=0.396$; $p=0.002$). Waist-to-hip ratio (WHR) and waist-to-height ratio (WHtR) had a weak positive correlation with insulin ($r=0.343$; $p=0.008$ and $r=0.311$; $p=0.017$) and HOMA-IR ($r=0.306$; $p=0.018$). There was a weak negative correlation between adiponectin and insulin in obese adolescents ($r=-0.278$; $p=0.033$). **Conclusion:** Anthropometric measurements (waist circumference, WHR and WHtR) and adiponectin can be used for early detection of insulin resistance and hyperinsulinemia in obese adolescents.

Keywords: Adiponectin, insulin, HOMA-IR, adolescents, obesity

31

INTRODUCTION

Obesity is a global problem. The number of obese individuals is increasing in Asia (Mazidi *et al.*, 2018). Obesity is associated with various complications, including metabolic syndrome, cardiovascular disease and type 2 diabetes mellitus (Asghar & Sheikh,

2017). These complications have varied effects, ranging from an increased risk of premature death to a reduction in quality of life (Hirko *et al.*, 2015; Morrison *et al.*, 2015).

In obesity, an increase in the size of fat cells is associated with inflammatory conditions. Visceral fat plays an important

*Corresponding author: Nur Aisiyah Widjaja
Department of Child Health, Dr. Soetomo General Academic Hospital, Faculty of Medicine, Universitas Airlangga, Jl. Mayjen Prof. Dr. Moestopo no 6-8, Surabaya, Indonesia
Tel: (+62)8123073379, E-mail: nuril08@yahoo.com
doi: <https://doi.org/10.31246/mjn-2019-0100>

role in the production of adipocytokines and other pro-inflammatory cytokines to cause inflammation. Pro-inflammatory cytokines are associated with insulin insensitivity (Asghar & Sheikh, 2017). Insulin resistance plays a role in the pathogenesis of type 2 diabetes mellitus (Tangvarasittichai, 2015), metabolic syndrome (Banerji, Lam & Chaiken, 2017) and cardiovascular disease (Ormazabal *et al.*, 2018).

Waist circumference is one of the indices used to assess a person's risk of metabolic syndrome (Prakaschandra & Naidoo, 2017). The associations of waist circumference, thigh circumference, waist-to-hip ratio (WHR) and waist-to-height ratio (WHtR) with insulin resistance have been studied previously. However, most studies were conducted in adult subjects of normal weight (Benites-Zapata *et al.*, 2019) or those with type 2 diabetes mellitus (Yoon *et al.*, 2016).

Adiponectin is an adipocytokine known to have anti-inflammatory, anti-atherogenic and insulin-sensitising effects (Achari & Jain, 2017; Orlando *et al.*, 2019). Adiponectin levels are known to decrease in obese adolescents (Orlando *et al.*, 2019). Adiponectin has been studied as a protective factor against complications of obesity, such as diabetes mellitus, metabolic syndrome, hypertension, dyslipidaemia and cardiovascular disease (Sharma, McClung & Abraham, 2016; Orlando *et al.*, 2019). The protective mechanism of adiponectin has been explained through various adiponectin signalling pathways (Ruan & Dong, 2016). However, clinical research on adiponectin as a marker of insulin resistance has mostly been conducted in adults and patients with type 2 diabetes mellitus (Aleidi *et al.*, 2015).

Studies on adiponectin and anthropometric measurements as initial markers for detecting insulin

resistance in obese adolescents are still limited in developing countries. This study aims to analyse the association of anthropometric measurements and adiponectin levels with fasting glucose, insulin and Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) levels in obese adolescents.

15 MATERIALS AND METHODS

Study design

This was a cross-sectional study conducted on obese adolescents. Respondents were considered obese if their Body Mass Index (BMI) was above the 95th percentile on the Centers for Disease Control and Prevention (CDC) 2000 BMI curve according to age and sex. The inclusion criteria were age 13–16 years, obese, and consent obtained from a parent/guardian to participate in the study. Respondents suffering from infections, inflammation, autoimmune diseases, cancer, chronic diseases and endocrine disorders, those who were smokers and who consumed alcohol were excluded from this study. This study was conducted after obtaining ethical approval from the ethics committee of the Dr Soetomo General Academic Hospital, Surabaya (No. 0411/KEPK/VII/2018).

Physical examinations

Respondents were examined for weight, height, waist circumference and thigh circumference. Weight was measured using a digital scale (Seca, Germany) accurate to 0.1 kg. Height was measured with a portable stadiometer (Seca, Germany) accurate to 0.1 cm. Body weight was measured with respondents standing straight and not wearing footwear or other accessories. Height was measured from the vertex of the head to the heel with respondents in standing position and not wearing footwear or a hat. BMI was calculated with the

formula of body weight (kilogrammes) divided by height squared (meter squared) and plotted onto the BMI curve according to age and sex (CDC, 2000). Waist circumference was measured using a tape measure accurate to 0.1 cm at the midpoint between the lowest rib and the endpoint of the iliac crest upon expiration. Hip circumference was measured using a tape measure at the widest area of the hip at the point of the greatest gluteal protuberance. WHR was calculated as waist circumference (cm) divided by hip circumference (cm). WHtR was calculated as waist circumference (cm) divided by height (cm).

Biochemistry examinations

Blood measurements were performed, including adiponectin, fasting blood glucose levels, insulin and HOMA-IR. Blood collection was carried out in the morning after a 12-hour fast. Blood was drawn from the vena mediana cubiti. Blood was centrifuged, and the serum was removed and stored at -70°C until an adiponectin examination was performed. The adiponectin examination was carried out using enzyme-linked immunosorbent assay (ELISA) in accordance with manufacturer's instructions. The examination of insulin and fasting blood glucose levels was carried out with an enzymatic kit according to standard procedures.

HOMA-IR was used to describe insulin resistance.

Statistical analysis

Adiponectin, fasting blood glucose, insulin and HOMA-IR levels were expressed as medians and percentiles due to skewed distributions. A normality test was performed on each variable using the Kolmogorov–Smirnov test. The associations between fasting blood glucose, insulin and HOMA-IR with anthropometric measurements and adiponectin levels were analysed by Pearson's and Spearman's rho correlation. Data analysis was performed using SPSS statistics software version 21.0.

RESULTS

In this study, there were 59 obese adolescents aged 13–16 years. A total of 32 (54.2%) adolescents were males and 27 (45.8%) were females. The median fasting insulin level and HOMA-IR value were 16.09 mU/ml and 2.85, respectively, in obese adolescents (Table 1).

A weak positive correlation was obtained between waist circumference ($r=0.421$; $p=0.001$) and HOMA-IR ($r=0.396$; $p=0.002$). WHtR also had a weak positive correlation with HOMA-IR ($r=0.306$; $p=0.018$) (Table 2). WHR and

Table 1. Characteristics of obese adolescents

Variable	Median (25 th –75 th)
Body Mass Index (kg/m ²)	31.25 (29.20–33.70)
Waist circumference (cm)	97.20 (94.00–107.00)
Hip circumference (cm)	105.00 (99.00–110.00)
WHR	0.95 (0.91–0.99)
WHtR	0.62 (0.59–0.66)
Adiponectin (ng/ml)	6841.90 (5204.66–10044.31)
Fasting blood glucose (mg/dl)	79.00 (75.00–84.00)
Insulin (mU/ml)	16.09 (10.87–22.82)
HOMA-IR	2.85 (2.07–4.24)

25th: percentile 25; 75th: percentile 75

WHR: Waist-to-hip ratio; WHtR: Waist-to-height ratio; HOMA-IR: Homeostatic Model Assessment of Insulin Resistance

Table 2. Correlation between anthropometric measurements and insulin resistance


Variable	Fasting blood glucose		Insulin		HOMA-IR	
	r	p	r	p	r	p
Body Mass Index [†]	0.168	-0.182	0.111	0.401	0.068	0.608
Waist circumference [†]	0.029	0.830	0.421	0.001*	0.396	0.002*
Hip circumference [†]	-0.094	0.480	0.034	0.276	0.152	0.251
WHR [‡]	0.105	0.213	0.343	0.008*	0.159	0.230
WHtR [‡]	0.108	0.416	0.311	0.017*	0.306	0.018*

[†]Spearman's rho correlation

[‡]Pearson's correlation

*significant at $p < 0.05$

WHR: Waist-to-hip ratio; WHtR: Waist-to-height ratio

**Figure 1.** Correlation of insulin with WHR and WHtR

WHtR had a weak positive correlation with insulin ($r=0.343$; $p=0.008$; $r=0.311$; $p=0.017$) (Figure 1).

There was a weak negative correlation between adiponectin and insulin levels ($r=-0.278$; $p=0.033$). Statistical tests did not detect a relationship between adiponectin and HOMA-IR or fasting blood glucose levels ($p > 0.05$) (Table 3). In this study, a negative correlation was found between adiponectin and insulin levels (Figure 2).

DISCUSSION

Pro-inflammatory cytokines are associated with various metabolic complications, such as insulin resistance (Asghar & Sheikh, 2017). As chronic, low-level inflammation occurs in obesity, it is therefore associated with insulin resistance (Lim *et al.*, 2015). There is an increase in fasting insulin levels and HOMA-IR in obese adolescents. HOMA-IR is an index of peripheral insulin sensitivity, which is used to predict


Table 3. Correlation between adiponectin and insulin resistance

Variable	Adiponectin	
	<i>r</i>	<i>p</i>
Fasting blood glucose [†]	0.036	0.789
Insulin [†]	-0.278	0.033*
HOMA-IR [‡]	-0.205	0.119

[†]Spearman's rho correlation

[‡]Pearson's correlation

*significant at $p < 0.05$

**Figure 2.** Correlation between adiponectin and insulin

insulin resistance. Increased HOMA-IR is accompanied by increased levels of blood glucose, low-density lipoprotein cholesterol, triglycerides, insulin and glycated haemoglobin (HbA1c), which are risk factors for metabolic syndrome (González-Jiménez *et al.*, 2016). BMI is associated with insulin secretion and insulin sensitivity in obesity (Wang *et al.*, 2016). An increase in HOMA-IR is proportional to an increase in the BMI of adolescents (González-Jiménez *et al.*, 2016), although this increase can also be found in healthy adolescents (Telford *et al.*, 2012). In this study, there was no relationship between BMI and fasting blood glucose levels, insulin or insulin

resistance. The findings of this study differed from those of previous studies which showed BMI to be correlated with insulin resistance (Cheng *et al.*, 2017; González-Jiménez *et al.*, 2016; Lim *et al.*, 2015). However, the strength of correlation between BMI with insulin resistance shown by HOMA-IR was found to be weak in adolescents (Convit, Wedin & Diaz-Gimenez, 2012).

Body fat composition has a larger effect on insulin resistance than body fat percentage (Cheng *et al.*, 2017). In this study, waist circumference and WHtR were positively correlated with insulin and HOMA-IR. Previous studies have suggested that waist circumference is

correlated with increased insulin levels (Cempaka & Sidiartha, 2017) and HOMA-IR (da Silva *et al.*, 2018; Lim *et al.*, 2015). Waist circumference is more strongly correlated with HOMA-IR than with BMI (Convit *et al.*, 2012). In a similar study, adolescents stated that BMI, along with increased waist circumference and systolic blood pressure, were risk factors for insulin resistance (González-Jiménez *et al.*, 2016).

This study found that WHR has a positive correlation with insulin, similar to studies conducted in China. A study on respondents with normal weight showed that WHR was associated with hyperinsulinemia after tests of glucose tolerance and insulin resistance (Benites-Zapata *et al.*, 2019). Hyperinsulinemia in obesity occurs due to compensatory insulin secretion under conditions of insulin resistance (Wang *et al.*, 2016). WHR has also been shown to have a positive correlation with HOMA-IR (Lim *et al.*, 2015). However, in this study, there was no relationship between WHR and HOMA-IR. One possible explanation may be the similarities in pancreatic beta-cell function disorders between obese and non-obese respondents with impaired glucose tolerance (Takahara *et al.*, 2013).

Adiponectin is produced by fat tissue. The concentration of adiponectin in plasma is 2–30 µg/ml (Sharma *et al.*, 2016). Although adiponectin is produced by fat tissue, but its level is decreased in obesity. Adiponectin is negatively correlated with metabolic syndrome (Ntzouvani *et al.*, 2016), prevents cardiovascular disease and improves insulin sensitisation (Stern, Rutkowski & Scherer, 2016). In this study, low levels of adiponectin were obtained. Adiponectin had no correlation with blood glucose and HOMA-IR levels, but had a negative correlation with fasting insulin levels.

The differences between our results and those of previous studies may have been due to several factors. The relationship between anthropometric measurements and insulin resistance can vary between study populations, influenced by ethnicity (Yoon *et al.*, 2016) and age (Chandler-Laney *et al.*, 2010). Similarly, adiponectin levels are also influenced by ethnicity and gender (Sharma *et al.*, 2016; Aleidi *et al.*, 2015). However, combinations of anthropometric measurements can be used to predict cardiometabolic risk in adolescents (Samouda *et al.*, 2015). In addition, adiponectin and anthropometric measurements are also useful for early detection of insulin resistance and hyperinsulinemia in obese adolescents.

This study has several limitations. Firstly, the sample size was limited and there were no non-obese subjects with normal BMI included as a comparison group in this study. Secondly, no serial measurements of adiponectin, blood glucose and insulin levels were performed in this study. This could have affected the accuracy of the results as blood glucose can be affected by dietary consumption and physical activity.

CONCLUSION

There was a positive correlation in waist circumference, WHR and WHtR with insulin, while a positive correlation was observed between waist circumference and WHtR with HOMA-IR. There was no correlation between adiponectin and blood glucose levels or HOMA-IR. Adiponectin had a negative correlation with fasting insulin levels. Adiponectin, waist circumference, WHR and WHtR can be used for early detection of insulin resistance and hyperinsulinemia in obese adolescents. This can be one step towards preventing metabolic syndrome in adulthood.

Acknowledgement 25

The authors thank Universitas Airlangga and Dr. Soetomo Academic General Hospital, Surabaya, Indonesia for supporting this research.

Authors' contributions 29

NAW, principal investor, conceptualised and designed the study, prepared the draft of the manuscript and reviewed the manuscript; RAP, led the data collection, data analysis and wrote the manuscript; MHH, reviewed the manuscript; RI, led the data collection and reviewed the manuscript; IDGU, reviewed the manuscript; RH, reviewed the manuscript.

11

Conflict of interest

The authors declare no conflict of interest in this study.

References

- Aleidi S, Issa A, Bustanji H, Khalil M & Bustanji Y (2015). Adiponectin serum levels correlate with insulin resistance in type 2 diabetic patients. *Saudi Pharm J* 23(3):250-256. <https://doi.org/10.1016/j.jsps.2014.11.011>
- Achari AE & Jain SK (2017). Adiponectin, a Therapeutic Target for Obesity, Diabetes, and Endothelial Dysfunction. *Int J Mol Sci* 18(6):1321-1338. <https://doi.org/10.3390/ijms18061321>
- Asghar A & Sheikh N (2017). Role of immune cells in obesity induced low grade inflammation and insulin resistance. *Cell Immunol* 315:18-26. <https://doi.org/10.1016/j.cellimm.2017.03.001>
- Banerji MA, Lam ML & Chaiken R (2017). Insulin Resistance and the Metabolic Syndrome. In L Poretsky (Ed.), *Principles of Diabetes Mellitus* (pp. 1-25). https://doi.org/10.1007/978-3-319-20797-1_34-2
- Benites-Zapata VA, Toro-Huamanchumo CJ, Urrunaga-Pastor D, Guarnizo-Poma M, Lazaro-Alcantara H, Paico-Palacios S, Pantoja Torres B & Ranilla-Seguín V del C. (2019). High waist-to-hip ratio levels are associated with insulin resistance markers in normal-weight women. *Diabetes Metab Syndr: Clinical Research & Reviews* 13(1):636-642. <https://doi.org/10.1016/j.dsx.2018.11.043>
- Cempaka VP & Sidiartha IGL (2017). Waist circumference and insulin levels in obese children. *Paediatr Indones* 57(4):194-197. <https://doi.org/10.14238/pi57.4.2017.194-7>
- CDC (2000). *Clinical Growth Charts. Children 2-20 years: BMI-for-age*. Centers for Disease Control and Prevention (CDC) and the National Center for Health Statistics, Maryland. From https://www.cdc.gov/growthcharts/cdc_charts.htm. [Retrieved December 10 2019].
- Chandler-Laney PC, Phadke RP, Granger WM, Muñoz AJ, Dalla Man C, Cobelli C, Ovalle F, Fernández JR & Gower BA (2010). Adiposity and β -cell function: relationships differ with ethnicity and age. *Obesity (Silver Spring)* 18(11): 2086-2092. <https://doi.org/10.1038/oby.2010.44>.
- Cheng YH, Tsao YC, Tzeng IS, Chuang HH, Li WC, Tung TH & Chen JY (2017). Body mass index and waist circumference are better predictors of insulin resistance than total body fat percentage in middle-aged and elderly Taiwanese. *Medicine* 96(39): e8126. <https://doi.org/10.1097/MD.00000000000008126>
- Convit AJ, Wedin KW & Diaz-Gimenez L (2012). Prediction of insulin resistance with anthropometric measures: Lessons from a large adolescent population. *Diabetes Metab Syndr Obes: Targets and Therapy* 5:219-225. <https://doi.org/10.2147/DMSO.S33478>
- da Silva CC, Vasques ACJ, Zambon MP, Camilo DF, De Bernardi Rodrigues AM, Antonio MARGM & Geloneze B (2018). Sagittal abdominal diameter resembles waist circumference as a surrogate marker of insulin resistance in adolescents-Brazilian Metabolic Syndrome Study. *Pediatr Diabetes* 19(5):882-891. <https://doi.org/10.1111/pedi.12664>
- González-Jiménez E, Schmidt-RioValle J, Montero-Alonso MA, Padez C, García-García CJ & Perona JS (2016). Influence of Biochemical and Anthropometric Factors on the Presence of Insulin Resistance in Adolescents. *Biol Res Nurs* 18(5):541-548. <https://doi.org/10.1177/1099800416648207>
- Hirko KA, Kantor ED, Cohen SS, Blot WJ, Stampfer MJ & Signorello LB (2015). Body Mass Index in Young Adulthood, Obesity Trajectory, and Premature Mortality. *Am J Epidemiol* 182(5):441-450. <https://doi.org/10.1093/aje/kwv084>
- Lim SM, Choi DP, Rhee Y & Kim HC (2015). Association between Obesity Indices and Insulin Resistance among Healthy Korean Adolescents: The JS High School Study. *PLOS ONE* 10(5):e0125238. <https://doi.org/10.1371/journal.pone.0125238>

- Mazidi M, Banach M, Kengne AP & Meta-analysis Collaboration Group (2018). Prevalence of childhood and adolescent overweight and obesity in Asian countries: A systematic review and meta-analysis. *Arch Med Sci* 14(6):1185-1203.
- Morrison KM, Shin S, Tarnopolsky M & Taylor VH (2015). Association of depression & health related quality of life with body composition in children and youth with obesity. *J Affect Disord* 172:18-23. <https://doi.org/10.1016/j.jad.2014.09.014>
- Ntzouvani A, Fragopoulou E, Panagiotakos D, Pitsavos C & Antonopoulou S (2016). Reduced circulating adiponectin levels are associated with the metabolic syndrome independently of obesity, lipid indices and serum insulin levels: A cross-sectional study. *Lipids Health Dis* 15(1):1-14. <https://doi.org/10.1186/s12944-016-0311-7>
- Orlando A, Nava E, Giussani M & Genovesi S (2019). Adiponectin and Cardiovascular Risk. From Pathophysiology to Clinic: Focus on Children and Adolescents. *Int J Mol Sci* 20(13):1-14. <https://doi.org/10.3390/ijms20133228>
- Ormazabal V, Nair S, Elfeky O, Aguayo C, Salomon C & Zuriaga FA (2018). Association between insulin resistance and the development of cardiovascular disease. *Cardiovasc Diabetol* 17(1):1-14. <https://doi.org/10.1186/s12933-018-0762-4>
- Prakaschandra R & Naidoo DP (2017). Increased waist circumference is the main driver for the development of the metabolic syndrome in South African Asian Indians. *Diabetes Metab Syndr: Clinical Research & Reviews* 11:S81-S85. <https://doi.org/10.1016/j.dsx.2016.12.011>
- Ruan H & Dong LQ (2016). Adiponectin signaling and function in insulin target tissues. *J Mol Cell Biol* 8(2):101-109. <https://doi.org/10.1093/jmcb/mjw014>
- Samouda H, de Beaufort C, Stranges S, Guinhouya BC, Gilson G, Hirsch M, Jacobs J, Leite S, Vailant M & Dadoun F (2015). Adding anthropometric measures of regional adiposity to BMI improves prediction of cardiometabolic, inflammatory and adipokines profiles in youths: A cross-sectional study. *BMC Pediatr* 15(1):1-9. <https://doi.org/10.1186/s12887-015-0486-5>
- Sharma M, McClung JA & Abraham NG (2016). Chapter 4 - Adiponectin: A Mediator of Obesity, Insulin Resistance, Diabetes, and the Metabolic Syndrome. In *Translational Research in Coronary Artery Disease* (pp. 33-42). <https://doi.org/10.1016/B978-0-12-802385-3.00004-8>
- Stern JH, Rutkowski JM & Scherer PE (2016). Adiponectin, Leptin, and Fatty Acids in the Maintenance of Metabolic Homeostasis through Adipose Tissue Crosstalk. *Cell Metab* 23(5):770-784. <https://doi.org/10.1016/j.cmet.2016.04.011>
- Takahara M, Katakami N, Kaneto H, Noguchi M & Shimomura I (2013). Pancreatic beta cell function in lean and obese Japanese with various degrees of glucose tolerance. *Endocr J* 60(7):923-930. <https://doi.org/10.1507/endocrj.EJ13-0046>
- Tangvarasittichai S (2015). Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World J Diabetes* 6(3):456. <https://doi.org/10.4239/wjd.v6.i3.456>
- Telford RD, Cunningham RB, Telford RM, Kerrigan J, Hickman PE, Potter JM & Abhayaratna WP (2012). Effects of Changes in Adiposity and Physical Activity on Preadolescent Insulin Resistance: The Australian LOOK Longitudinal Study. *PLoS ONE* 7(10):e47438. <https://doi.org/10.1371/journal.pone.0047438>
- Wang T, Ma X, Tang T, Jin L, Peng D, Zhang R, Chen M, Yan J, Wang S, Yan D, He Z, Jiang F, Cheng X, Bao Y, Liu Z, Hu C & Jia W (2016). Overall and central obesity with insulin sensitivity and secretion in a Han Chinese population: A Mendelian randomization analysis. *Int J Obes* 40(11):1736-1741. <https://doi.org/10.1038/ijo.2016.155>
- Yoon YS, Choi HS, Kim JK, Kim YI & Oh SW (2016). Differences in the associations of anthropometric measures with insulin resistance and type 2 diabetes mellitus between Korean and US populations: Comparisons of representative nationwide sample data. *Obes Res Clin Pract* 10(6):642-651. <https://doi.org/10.1016/j.orcp.2015.11.001>

Adiponectin, anthropometric measurements and insulin resistance in adolescence with obesity

ORIGINALITY REPORT

19%

SIMILARITY INDEX

16%

INTERNET SOURCES

13%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	pesquisa.bvsalud.org Internet Source	1%
2	www.alliedacademies.org Internet Source	1%
3	link.springer.com Internet Source	1%
4	aoco2019.icomes.or.kr Internet Source	1%
5	Submitted to Bournemouth University Student Paper	1%
6	Submitted to University of California, Merced Student Paper	1%
7	Submitted to University of East London Student Paper	1%
8	Submitted to Nottingham Trent University Student Paper	1%
9	eprints.nottingham.ac.uk Internet Source	1%

10	www.physiology.org Internet Source	1 %
11	medcraveonline.com Internet Source	1 %
12	theses.gla.ac.uk Internet Source	1 %
13	mbali.unizulu.ac.za Internet Source	<1 %
14	openrepository.aut.ac.nz Internet Source	<1 %
15	www.gavinpublishers.com Internet Source	<1 %
16	www.medrxiv.org Internet Source	<1 %
17	www.researchgate.net Internet Source	<1 %
18	www.scielo.br Internet Source	<1 %
19	www.ncbi.nlm.nih.gov Internet Source	<1 %
20	Liong Boy Kurniawan, Bachtiar Syamsir, Ilham Akbar Rahman, Endy Adnan, Tenri Esa, Yuyun Widaningsih, Uleng Bahrnun, Mansyur Arif. "Anthropometric features in predicting insulin resistance among non-	<1 %

menopausal Indonesian adult females",
Romanian Journal of Internal Medicine, 2020

Publication

21

Shao-Yuan Chuang, Wen-Ling Liu, Hsing-Yi Chang, Chih-Cheng Hsu, Wen-Harn Pan. "Central obesity and elevated blood pressure in middle life are associated with physical and cognitive impairment in later life: A retrospective design with repeated measures", Experimental Gerontology, 2023

Publication

<1 %

22

Yudi Her Oktaviono, Suryo Ardi Hutomo, Makhyan Jibril Al-Farabi, Angliana Chouw, Ferry Sandra. "Human umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration", F1000Research, 2020

Publication

<1 %

23

www.patientcareonline.com

Internet Source

<1 %

24

J. A. Osuna C., R. Gómez-Pérez, G. Arata-Bellabarba, V. Villaroel. "RELATIONSHIP BETWEEN BMI, TOTAL TESTOSTERONE, SEX HORMONE-BINDING-GLOBULIN, LEPTIN, INSULIN AND INSULIN RESISTANCE IN OBESE MEN", Archives of Andrology, 2009

Publication

<1 %

25

media.neliti.com

Internet Source

<1 %

26	pdffox.com Internet Source	<1 %
27	repositories.lib.utexas.edu Internet Source	<1 %
28	www.journals.usp.br Internet Source	<1 %
29	www.nature.com Internet Source	<1 %
30	www.pjps.pk Internet Source	<1 %
31	Chen Chen, Yi Ye, Yanbo Zhang, Xiong-Fei Pan, An Pan. "Weight change across adulthood in relation to all cause and cause specific mortality: prospective cohort study", BMJ, 2019 Publication	<1 %
32	Mohd Rahimi Che Jusoh, Stephen R. Stannard, Toby Mündel. "Sago supplementation for recovery from cycling in a warm-humid environment and its influence on subsequent cycling physiology and performance", Temperature, 2016 Publication	<1 %
33	Mona Abo Baker Abd Ellatef, Ebtessam A. Elsayed, Khalil Abd Elmaksoud Abd Elhamed, Mohammed Almohaithef. "Quality of Life of Obese School-Age Children and	<1 %

their Mothers' Perspectives in Cairo, Egypt",
Egyptian Journal of Health Care, 2017

Publication

34

Siroon Bekkering, Christoph Saner, Niels P. Riksen, Mihai G. Netea et al. "Trained Immunity: Linking Obesity and Cardiovascular Disease across the Life-Course?", Trends in Endocrinology & Metabolism, 2020

Publication

<1 %

35

academic.oup.com

Internet Source

<1 %

36

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

37

researcherslinks.com

Internet Source

<1 %

38

www.elsevier.es

Internet Source

<1 %

39

www.karger.com

Internet Source

<1 %

40

www.nel.edu

Internet Source

<1 %

41

Amrita Banerjee, Jagdish Singh. "Remodeling adipose tissue inflammasome for type 2 diabetes mellitus treatment: Current perspective and translational strategies", Bioengineering & Translational Medicine, 2019

<1 %

42 Takahara, Mitsuyoshi, Naoto Katakami, Hideaki Kaneto, Midori Noguchi, and Ichihiro Shimomura. "Pancreatic beta cell function in lean and obese Japanese with various degrees of glucose tolerance", *Endocrine Journal*, 2013.

Publication

43 Bin Liu, Huan Zheng, Guanghui Liu, Zhiling Li. "Adiponectin is Inversely Associated with Insulin Resistance in Adolescents with Nonalcoholic Fatty Liver Disease", *Endocrine, Metabolic & Immune Disorders - Drug Targets*, 2022

Publication

44 Tuba ONAY, ASLI UÇAR. "Which anthropometric measurement/ratio is a better predictor of non-alcoholic fatty liver disease?", *Research Square Platform LLC*, 2023

Publication

45 www.ijrcog.org

Internet Source

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Adiponectin, anthropometric measurements and insulin resistance in adolescence with obesity

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
