



SURAT KETERANGAN
Nomor : 3318/UN3.1.6/KP/2023

Yang bertanda tangan di bawah ini :

Nama : Prof. Dr. Mirni Lamid, drh., MP
NIP : 196201161992032001
Pangkat / Golongan : Pembina Utama Madya / IVd
Jabatan : Dekan

Dengan ini menerangkan :

Nama : Dr. Hani Plumeriastuti, drh., M.Kes.
NIP : 195908081987012001
Pangkat / Golongan : Pembina (Gol. IV/a)
Jabatan : Lektor Kepala

Telah melaksanakan penelitian dengan judul sebagai berikut :

No.	Judul Karya Ilmiah	Tahun Pelaksanaan Penelitian
1.	Genetic Identification of Shiga Toxin Encoding Gene from Cases of Multidrug Resistance (MDR) Escherichia coli Isolated from Raw Milk.	2021
2.	Molecular identification of blaCTX-M and blaTEM genes encoding extended-spectrum β -lactamase (ESBL) producing Escherichia coli isolated from raw cow's milk in East Java, Indonesia	2021
3.	Incidence of Escherichia coli producing Extended-spectrum beta-lactamase in wastewater of dairy farms in East Java, Indonesia	2023
4.	Detection of Multidrug-Resistant (MDR) Escherichia coli Isolated from Raw Milk in East Java Province, Indonesia.	2020
5.	Presence of multidrug resistance (MDR) and extended-spectrum beta-lactamase (ESBL) of Escherichia coli isolated from cloacal swab of broilers in several wet markets in Surabaya.	2021
6.	A comprehensive description of the exoskeleton of six Lobster species (Genus Panulirus) in Aceh Province, Indonesia.	2023
7.	Analisis Filogenetik Gen Hemagglutinin dan Neuraminidase Avian Influenza H9N2 Asal Ayam Petelur di Jawa Timur	2020
8.	Identification of Ectoparasites in Pearl Catfish (<i>Clarias gariepinus</i>) with One and Three Months Age in Maclele Cultivation, Tuban District, Tuban Regency.	2022
9.	Cases of Multidrug Resistance (MDR) in <i>Klebsiella pneumoniae</i> Isolated from Healthy Pigs.	2021



UNIVERSITAS AIRLANGGA

FAKULTAS KEDOKTERAN HEWAN

Kampus C Mulyorejo Surabaya 60115 Telp. (031) 5992785, 5993016 Fax (031) 5993015

Laman: <http://www.fkh.unair.ac.id>, e-mail: info@fkh.unair.ac.id

Adapun penelitian tidak perlu Uji Ethical Clearance karena **menggunakan hewan coba yang tidak disakiti**.

Demikian Surat Keterangan ini kami buat untuk dapat dipergunakan sebagai persyaratan pengusulan Jabatan Fungsional **Guru Besar**

Surabaya, 12 Juni 2023

Dekan,



Prof. Dr. Mirni Lamid, drh., MP

NIP 196201161992032001



ASEAN
University
Network



JOURNAL

of Parasite Science

J. Parasite Sci.



PUBLISHER COLLABORATION

Journal of Parasite Science



Collaboration
Division of Veterinary Parasitology
Faculty of Veterinary Medicine, Universitas Airlangga
and
Indonesian Veterinary Parasitology Association



EDITOR'S ADDRESS

Editorial Office Journal of Parasite Science

Division of Veterinary Parasitology
Faculty of Veterinary Medicine, Universitas Airlangga
Kampus "C" Jl. Mulyorejo Surabaya 60115
Phone: (031) 5992785; 5993016; Fax. (031) 5993015
e-mail: jparasitol@gmail.com ; jps@fkh.unair.ac.id
Homepage : <https://e-journal.unair.ac.id/JoPS>

EDITORIAL BOARD

Chief Editor

Dr. Kusnoto, DVM., M.Si., Universitas Airlangga, Indonesia

Chief's Assistant Editor

Prof. Muchammad Yunus, DVM., M.Kes., Ph.D., Universitas Airlangga, Indonesia

Editorial Board Member

Prof. Dr. Lucia Tri Suwanti, DVM., MP. Universitas Airlangga, Indonesia

Prof. Dr. Makoto Matsubayashi, DVM., Ph.D., Osaka Prefecture University, Japan

Assoc. Prof. Dr. Mahmut Kabalak, Hacettepe University, Turkey

Dr. Attaur Rahman, Abdul Wali Khan University Mardan, Pakistan

Dr. Muhammad Hassan Mushtaq, University of Veterinary and Animal Sciences Lahore, Pakistan

Aondohemba Samuel Nege, B.Fish., M.Sc., National Taiwan University, Taiwan, Province of China

Muhammad Thohawi Elziyad Purnama, DVM., M.Si. PSDKU Universitas Airlangga, Banyuwangi

Administration Staff

Dyah Ayu Kurniawati, DVM., M.Sc, Universitas Airlangga, Indonesia

Moch Arifudin, Universitas Airlangga, Indonesia

REVIEWERS

Editors, Authors and Readers of Journal of Parasite Science give the highest appreciation and gratitude to the experts below, as reviewers who has reviewed all the articles, whether published or rejected, according to the recommendations submitted to the editor in Vol. 5 No. 2 September 2021

April Hari Wardhana, DVM, M.Sc, Ph.D., Balai Besar Penelitian Veteriner Bogor, Bogor, Indonesia

Prof. Dr. Sri Subekti, DVM., DEA., Perkumpulan Pemberantasan Penyakit Parasit Indonesia

Prof. Dr. Upiek Kesumawati Hadi, DVM., MS., Institut Pertanian Bogor, Bogor, Indonesia

Dr. Raden Wisnu Nurcahyo, DVM., Universitas Gadjah Mada, Yogyakarta, Indonesia

Dr. Dwi Priowidodo, DVM., MP., Universitas Gadjah Mada, Yogyakarta, Indonesia

Dr. Nyoman Adi Suratma, DVM., MP., Universitas Udayana, Bali, Indonesia

Prof. Hiroshi Sato, DVM., Ph.D., Yamaguchi University, Jepang

Prof. Graeme Martin, B.Sc(Agric)., Ph.D W.Aust., University of Western Australia School of Agriculture and Environment, Australia

Prof. Dr. Muhammad Salisu Abubakar., Usmanu Danfodiyo University Sakoto, Nigeria

Dr. Zainab Alwan Makawi., University of Baghdad, Iraq

VISION AND MISSION

Journal of Parasite Science (JoPS) with registered number [P-ISSN \(2599-0993\)](#) ; [E-ISSN \(2656-5331\)](#) published by the Division of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga.

Journal of Parasite Science (JoPS) presenting research articles, case reports, community development activities and literature studies of veterinary medicine. Published twice a year on March and September. Since 2002 the Faculty of Veterinary Medicine has had Veterinary Medicine Media, because considering that there is only one journal that must accommodate many articles written by lecturers and students, various journals have been developed according to the disciplines in each department, one of which is the JoPS. JoPS can publish articles from various faculties and institutions related to the parasitology. JoPS which has been published since 2017 is here for accommodate publication obligations for undergraduate and postgraduate students as well as other parties conducting research in the field of parasitology. Loading of articles in the JoPS through the Open Journal System (OJS). Complete information for article loading and article writing instructions are available on the website and every issue. Incoming articles will go through a selection process by editors and reviewers.

VISION

To become a leading and reputable journal at the national and international level in veterinary parasitology science.

MISSION

1. Provide scientific communication media of Parasitology to participate in advancing science and technology in related sector;
2. Organizing the publication of scientific journals of Parasitology with a high impact factor in the development of science and technology;
3. Making journals as terms for the advancement and intellectual development of the academic community in welcoming the Universitas Airlangga World Class University;
4. Organizing an accountable and quality journal management to increase the number of intellectual products in the form of scientific journals;
5. Become a leading reference for the academic community and researchers of veterinary parasitology and published as a scientific journal.

Journal Scope: Journal of Parasite Science (JoPS) publishes the results of original research in all aspects of basic and applied parasitology, and ranging from parasites biodiversity, parasites of all wildlife, invertebrate and vertebrate, as well as host-parasite relationships of intrinsic biological interest to those of social and economic importance predominately in veterinary, human medicine and agriculture aspect. Original research includes the development of novel and innovative concepts and ideas, as well as experimental and observational science that raises new theory.

Language:

Main: English


The articles accepted by the Journal of Parasite Science (JoPS) are:

1. Research articles;
2. Case reports;
3. Community development articles;
4. Literature study articles.

TABLE OF CONTENTS

	Page
1. Ectoparasite Detection in Culling Layer Hen That Sold at Wonokromo Market Surabaya City (Nashihatun Sholihah, Sherbina Bai, Sri Hidanah, Dhandy Koesomo Wardhana, Agus Sunarso, Poedji Hastutiek, M. Gandul Atik Yuliani).....	38 - 41
2. Ovicidal Effects of Pineapple Peel Juice (<i>Ananas comosus L.</i>) as a Control of Ascaridians in Chicken (<i>Gallus Domesticus</i>) In Vitro (Indira Vania Carisya, Setiawan Koesdarto, Tjuk Imam Restiadi, Nove Hidajati).....	42 - 45
3. Identification of Ectoparasites and Gastrointestinal Tract Endoparasites in Stray Cats at Traditional Market of Bojonegoro City (Karina Rachmawati, Setiawan Koesdarto, Iwan Sahrial Hamid, Dian Ayu Permatasari).....	46 - 49
4. Prevalences of Gastrointestinal Parasitic in Dogs and Cats at Kaki 4 and K-5 Clinics Kediri City (Taufan Ary Handoko, Poedji Hastutiek, Mufasirin, Herry Agoes Hermadi, Nusdianto Triakoso).....	50 - 53
5. Ultrastructural Morphology Comparison of <i>Raillietina echinobothrida</i> and <i>Raillietina tetragona</i> in Native Chickens at Sioarjo Regency Using Scanning Electron Microscope (SEM) (Afiyah Rahmadani, Arimbi, Endang Suprihati, Nunuk Dyah Retno Lastuti, Sunaryo Hadi Warsito, Maudika Pamela Fiorenza).....	54 - 59
6. Identification of Ectoparasites in Pearl Catfish (<i>Clarias gariepinus</i>) with One and Three Months Age in Maclele Cultivation, Tuban District, Tuban Regency (Azizatun Nimah, Galuh Swa Ardhya, Hani Plumeriastuti, Setiawan Koesdarto, Budiarto).....	60 - 66

Identification of Ectoparasites in Pearl Catfish (*Clarias gariepinus*) with One and Three Months Age in Maclele Cultivation, Tuban District, Tuban Regency

¹⁾Azizatun Nimah*, ¹⁾Galuh Swa Ardhy, ²⁾Hani Plumeriastuti, ³⁾Setiawan Koesdarto,
⁴⁾Budiarto 

¹⁾Student, Faculty of Veterinary Medicine, Universitas Airlangga

²⁾Division of Veterinary Basic Medicine, Faculty of Veterinary Medicine, Universitas Airlangga

³⁾Division of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga

⁴⁾Division of Veterinary Public Health, Faculty of Veterinary Medicine, Universitas Airlangga

*Corresponding author: azizatunnimah@gmail.com

Abstract

This study aims to identify ectoparasites that attack pearl catfish and the incidence of ectoparasite infestations in 100 samples of one- and three-month-old pearl catfish. This research was conducted from January to March 2021 in Makalele Cultivation, Tuban District, Tuban Regency. The method used in this research is skin scraping and gill installation. Data analysis using Chi-Square Test with SPSS version 24.0 for windows. The results showed that there were 3 types of ectoparasites found, namely *Trichodina nobilis*, *Trichodina pediculus*, *Chilodonella* sp., *Dactylogyrus* sp. and the highest incidence of ectoparasites occurred in young fish compared to old fish with a percentage of 35% and 25%, respectively, and based on the Chi-Square test there was no significant difference ($p \geq 0.05$).

Keywords: *Clarias gariepinus*, incidence rate, *T. Nobilis*, *T. Pediculus*, *Chilodonella* sp.

Introduction

Tuban Regency is one of the regencies in East Java that has a high contribution to the fisheries business field and shows an increase from 2011 to now (RPJMD, 2016). Through fish farming business activities can reduce pressure on the utilization of marine resources. Catfish is one type of fish that has economic value and one type of freshwater fish that is often cultivated in the Tuban Regency. The type of catfish that is often cultivated in this area is the type of pearl catfish.

Pearl catfish cultivation can be found in Maculele Cultivation, Tuban District, Tuban Regency. This cultivation was established in 2017, This is due to the conditions in the area which have a stable temperature range among other areas. Murugaian (2008) stated that temperature is one of the factors that can affect the growth process of fish, as well as affect the fish's immune system (Ghufran, 2010). Problems that are often experienced in catfish farming are the incidence of disease and fish death.

The disease is an abnormality in fish caused by an unbalanced interaction between three components in an aquatic ecosystem, namely a weak host, a vicious pathogen, and a deteriorating environmental quality. Factors that play a role in the emergence of disease are internal and

external. Internal factors are factors that come from the internal environment such as genetic disorders, immunity, and body metabolism. External factors that affect pathogenic diseases are parasites, viruses, fungi, and bacteria (Khotimah et al., 2018).

Ectoparasite attacks can cause acute high mortality, changes in fish morphology, decreased fish productivity, and fish death will often occur without showing symptoms and if not treated immediately will cause mass death because of the transmission and spread of ectoparasites is very fast. Ectoparasites that can infest catfish include *Trichodina* sp., *Ichthyophthirius* multiplies, *Vorticella* sp., *Oodinium* sp., *Chilodonella* sp., *Dactylogyrus* sp., and *Gyrodactylus* sp. (Hardiroseyani, 2006).

This study used pearl catfish aged one and three months. The age difference was chosen because the pearl catfish with the age of three months was still classified as mature age and ready for harvest, while the pearl catfish with the age of one month is included in the young age group.



Materials and Methods

Material

The material needed in this study was to use 100 live pearl catfish (*Clarias gariepinus*) samples, NaCl 0.9%, aquadest, formalin 0.1%, and oil immersion.

Tools

The research equipment used in the examination of the samples were gloves, masks, Styrofoam ice boxes, scrap tools (scalpel/wooden spatula), pipettes, surgical scissors, tweezers, Petri dishes, object-glass, cover glass, trays, toothpicks, light microscopes, and the Lucida microscope.

Sample and Sample Size

This research was conducted during January-March 2021. The sample size was calculated based on the sample size formula Solving (Husein, 2013):

$$\text{Formula } n = \frac{N}{1 + Ne^2}$$

n = Number of samples; N = Population size; e = Sampling error.

$$n = \frac{2000}{1 + 2000(0,1)^2}$$

$$n = \frac{2000}{21}$$

$$n = 95,24$$

From the calculation results above, the sample size is rounded up to 100 samples.

Research Procedure

The method used is the method *cross-sectional study* and taking samples at the location directly in the than catfish Cultivation, Tuban District, Tuban Regency and taken by non-random sampling with the purposive method. Identification of ectoparasites will be conducted at the Parasitology Laboratory, Division of Veterinary Parasitology, Faculty of Veterinary Medicine, Airlangga University, Surabaya. Examination of ectoparasites on pearl catfish is carried out in two ways, namely, scraping or mounting gills. Identification of ectoparasites by scraping or scrapping method is done by scraping the skin of the fish, head to tail using a scalpel to get mucus. Taking fish slime is done with a strong push and carefully so as not to injure the fish's body. The mucus resulting from the scrapping was immediately placed on an object glass and dripped with physiological sodium chloride (0.9% NaCl) and then dried. Put into formalin 0,1% for 2-3 minutes of preservation then lid with cover glass. Observe with microscope with 400x-1000x magnification.

Data Analysis

Data from the identification of ectoparasites that infest pearl catfish (*Clarias gariepinus*) in the cultivation of Makalele, Tuban District, Tuban Regency presented descriptively, pictures, and tables. The results of the data to determine the difference in the incidence of

ectoparasite infestation at the age of one and three months using the Chi-Square Test in the SPSS for Windows 24.0 program.

Results

Pool Temperature Check

Temperature checks in each pool were also carried out. Pools A1 and B1 have the same temperature and A2 and B2 have almost the same temperature. The water temperature in the range of 28 - 29°C is still in the optimal temperature range for pearl catfish farming activities. Temperature is very influential on various chemical reactions in water and one of them is on the solubility of oxygen in the water and the metabolism of the fish's body, so it will affect the growth of fish.



Ectoparasite Infestation Examination

Based on the results of research that have been carried out on pearl catfish in Makalele Cultivation, Tuban District, Tuban Regency which was carried out from January to March 2021 on 100 samples of pearl catfish in which each pond was divided into 50, it was found that 60 fish were infested with ectoparasites using the native method, namely skin scrapings. (scrapping) and gill examination. From 100 samples of pearl catfish examined for skin and gills, positive results were obtained from 35 fish at one month and 25 at three months. Sampling carried out on the left and right gills and skin. One-month-old catfish found ectoparasite infestation as many as 32 positive samples on the left gill and as many as 31 positive samples on the right gill, 24 positive samples found on the left skin, and as many as 26 positive samples on the right skin. And in catfish aged three months, 23 positive samples were found on the left gills and 21 positive samples on the right gills, 20 positive samples were found on the left skin, and 21 positive samples on the right skin, the incidence of ectoparasite infestation in catfish One and three-month-old pearls. After being analyzed using the Chi-Square test with a significant value of 5% alpha to determine differences in the incidence of ectoparasite infestation in one and three-month-old pearl catfish, obtained p-value 0.066 (p>0.05) non-significant. The results of this study showed that there was no difference in the level of ectoparasite infestation between the age of one and three months of pearl catfish.

Ectoparasite Microscopic Examination

The results of the examination of ectoparasites found through microscopic examination of the body of the pearl catfish, namely *Trichodina* sp., *Chilodonella* sp., and *Dactylogyrus* sp. On observation under a microscope, *Trichodina* sp. transparent and



shaped like a ring filled with vibrating hairs. The characteristics of *Trichodina* sp. found in this study were found on the skin, fins, and gills. The shape of this parasite is spherical with the oral surface looking slightly concave and the outside of the cells seen surrounded by many cilia. This is by the research of Munawwaroh and Rahayu (2017) which states that *Trichodina* sp. has a concave body shape and has attachments located on the anterior and posterior parts. The diameter of *Trichodina* sp. found is 79.81µm. Image results of *Trichodina* sp. and camera lucida can be seen in Figures 1a and 1b.



Figures 1 (a) *Trichodina* sp. (b) *Trichodina* sp. with with camera lucida with 400x magnification. adhesive disk(Ad); Blades (B); Border Membrane (Bm); Cilia (C); Ray (R); Radial pins (Rp). 100 m bar scale.

The type of *Trichodina* sp. What was found was *Trichodina nobilis* with a body diameter of 79.81µm. *Trichodina nobilis* was found to have a larger body size, sharp blade apex tip, slightly curved distal blade margin to the anterior blade margin, and visible long, straight, and pointed rays. Based on the calculation results, the body diameter of this parasite is 79.81 m. The adhesive disc is concave with a diameter of 65.86 m, surrounded by a border membrane with a width


of 12.67 m. The denticle blade is pointed with a length of 8.76 m and 23 - 25 in number. The diameter of the denticle ring is 40.44 m. The ray is pointed at the tip with a length of 9.89 m and a width of the denticle span of 9.05 m. Size *Trichodina nobilis* and the results from the camera lucida can be seen in Figures 2a and 2b.



Figures 2 (a) *Trichodina nobilis* (b) *Trichodina nobilis* with camera lucida. Body diameter (BD); Blade length (BI); Blade membrane width (Bmw); Diameter of adhesive disc (Da); Denticle diameter (Dd); Denticle span (Ds); Ray length (R1).

The type of ectoparasite *Trichodina* sp. what was found next was *Trichodina pediculus* where the morphology was almost the same as *Trichodina nobilis*. In the results that have been researched obtained calculations for *Trichodina pediculus* showed the body diameter of this parasite 68.05 m. The adhesive disc is concave with a diameter of 58.34 m, surrounded by a border membrane with a width of 9.32 m. The denticle blade is pointed with a length of 6.79 m and a total of 26. The diameter of the denticle ring is 41.32 m. Rayon the end with a length of 10.89 m


and a width of the denticle span of 6.08 m. Van and Basson (1986) in Woo (2006) stated that *Trichodina pediculus* has a wide distribution in fresh water and is generally found on the skin of adult fish and fish fry. Size *Trichodina pediculus* and the results from the camera lucida can be seen in Figures 3a and 3b.



Figures 3 (a) *Trichodina pediculus* (b) *Trichodina pediculus* with camera lucida. Blade length (Bl); Blade membrane width (Bmw); Diameter of adhesive disc (Da); Denticle diameter (Dd); Denticle span (Ds); Ray length (Rl).

Other types of protozoa were also found, namely *Chilodonella* sp. This parasite is very much found on the skin, fins, and gills of pearl catfish. *Chilodonella* sp. is a protozoan that can reproduce rapidly by mitotic division but also by conjugation (Mahendra and Nurbadriati, 2019). The shape of the parasite found was oval with a convex dorsal and a flattened ventral had cilia, the size of the parasite found was 20.34 m long and 14.83 m wide, and the method of infestation of this parasite was seen in groups or colonies. *Chilodonella* sp. breed at a temperature of 0.5 - 200C. Under bad conditions, these parasites will


form cysts. *Chilodonella* sp. cannot live without a host for more than 12 - 24 hours. The presence of ectoparasites *Chilodonella* sp. The high temperature can be caused because these ectoparasites have a low-temperature tolerance range so they can maintain their existence. Himage result of *Chilodonella* sp. and camera lucida can be seen in Figures 4a and 4b.



Figures 4 (a) *Chilodonella* sp. (b) *Chilodonella* sp. with lucida camera with 1000x magnification. Cilia (C); Contractile vacuole (Cv); Cytopharynx (Cy); Digested vacuole (Dv); Macronucleus (Ma); Micronucleus (Mi).

In addition to the protozoan parasites, this study also found monogenean parasites, namely *Dactylogyrus* sp. This parasite attaches to the gills of fish. The shape of this worm is dorsoventral flat and bilaterally symmetrical. On the dorsal side, there are prohaptor organs and eyespots, while on the ventral side there are opisthaptor organs. In the posterior part of the body, there is an opisthaptor with a pair of median hooks. The characteristics of the worms found are dark in color, with a body length of 0.23 mm and a width of 0.06 mm. On *Dactylogyrus* sp. The life cycle will start from an adult worm that lays eggs and hatches into a ciliated larva (onchomiracidium) which has more than one cilia and eye spot and

will swim until it attaches to the gills and then becomes an adult worm (Kabata, 1985). These worms are oviparous and have haptors, which are organs for attachment. Parasite *Dactylogyrus* sp. is capable of releasing 4-10 eggs every 24 hours and will increase in number with increasing temperature. Himage and organs from *Dactylogyrus* sp. and camera lucida can be seen in Figures 5a, 5b, and 5c.



Figures 5 (a) *Dactylogyrus* sp. (b) *Dactylogyrus* sp. with with camera lucida with 400x magnification. (c) Organs of *Dactylogyrus* sp. Anchor (A); Dorsal bar (Db); Head organs (Ho); Eyes (E); Marginal hooks (Mh); Ovaries (O); Pharynx (F); testes; Prostate gland (Pg); Stomach (S); Testis (T); Vitellaria (V); Ventral bar (Vb).

Discussion

The results of research conducted on 100 samples of pearl catfish obtained from Makalele

Cultivation, Tuban District, Tuban Regency from January to March 2021 were infested with ectoparasites where the age of one-month pearl catfish was 35% and the age of 3 months pearl catfish was 25%. To know differences in the incidence of ectoparasite infestation in one and three month old pearl catfish In this study, Chi Square analysis was carried out which showed no significant or non-significant difference ($p \geq 0.05$). This result is of course contrary to the opinion Irianto (2005) who stated that young fish are more susceptible to disease because the immune response in fish is fully formed when the fish are mature, in contrast to young fish, although young fish have an immune response but work less efficiently. In addition to age, several things affect the level of ectoparasite infestation in fish, including; species, temperature, nutritional conditions, physiological conditions, and population density (Hardi, 2015). Judging from the population of 1-month-old catfish ponds, there are more fish populations, but the size of 3-month-old catfish is much larger than 1-month-old catfish, resulting in a higher population density.

The high fish population density allows a decrease in the quality of aquaculture water (Shafrudin et al., 2006). This is by the statement of Sekar et al. (2016) that environmental conditions in the maintenance container and water quality can affect the increase in prevalence values and poor environmental conditions can cause fish conditions to become weak so that parasites will very easily attack the host. This will make the fish become stressed so that there is an unbalanced relationship between fish, the environment, and pathogens, and fish are easily infected with parasites. The water quality in Makalele Cultivation looks poor and there is still many fish floating on the surface of the water. The water condition of each pool looks a little cloudy. This of course can lead to easy ectoparasites that infest pearl catfish and can cause disease.

Based on the results of the identification found three types of ectoparasites infest pearl catfish in Makalele Cultivation, Tuban District, Tuban Regency. The infestation consisted of one and three months of age. The types of ectoparasites that have been found are *Trichodina* sp., *Chilodonella* sp., and *Dactylogyrus* sp. The percentage of pearl catfish in Makalele Cultivation, Tuban District, Tuban Regency aged one month consisted of 5% that were positive for *Trichodina* sp., 3% positive samples were infested with *Chilodonella* sp., 5% positive samples were infested with *Dactylogyrus* sp., 5% positive samples were infested with *Trichodina* sp. + *Chilodonella* sp., and 17% of infected positive samples *Trichodina*

sp. + *Dactylogyrus sp.* In a three-month-old pearl catfish consists of 6% positive infected with *Trichodina sp.*, 9% positive sample infested *Chilodonella sp.*, 1% positive samples were infested with *Dactylogyrus sp.*, 5% infested positive samples *Trichodina sp.* + *Dactylogyrus sp.*, and 4% of the positive samples that were infested with *Trichodina sp.* + *Chilodonella sp.*

Total *Trichodina sp.* quite a lot found on the surface of the fish body because this parasite contains a lot of mucus and epithelial tissue which is a good place to live and a place to find food. This parasite attacks a lot and irritates the gills, fins, and skin. The possibility that can affect the parasitic attack height *Trichodina sp.* in this study, namely the content of organic matter and the density of bacteria in the waters. This corresponds to the statement Nugroho et al., (2012) content organic matter and density are biological indicators that are used to describe the quality of fish rearing in general.

The form of the parasite These rings are ciliated and move rapidly from one organ to another (Bell and Burt 2008). Men massage by Pramono and Syakuri (2008), *Trichodina sp.* can stick to the surface of the body and will rotate 360° using cilia so that it will damage the surrounding cells and eat the destroyed epithelial cells to irritate the body surface.

The first *Trichodina nobilis* ectoparasite was identified by Chen (1963) found in Cyprin. fishidae. The results showed that *Trichodina nobilis* mostly infested the outer body of the fish and only a few infested the gills. This is following Vera et al. (2003) which states that *Trichodina nobilis* uses the host's body, especially on the skin for microhabitat.

This study also found *Trichodina pediculus* species, according to Dana et al. (2002). *Trichodina pediculus* attacks the skin and gills, irritating. The life cycle of *Trichodina nobilis* and *Trichodina pediculus* is quite fast and has only one definitive host and no intermediate hosts (Riko et al., 2012). *Trichodina nobilis* and *Trichodina pediculus* reproduce by dividing themselves or commonly called binary. When the denticle divides from the parent cell, it produces daughter cells. Transmission of *Trichodina nobilis* and *Trichodina pediculus* occurs through direct contact from infected fish to healthy fish.

Another type of protozoan ectoparasite found was *Chilodonella sp.* These ectoparasites mostly attack the skin, fins, and gills. The parasite found is in the form of oral and there are cilia on the entire surface of the body which is used as a means of locomotion. This is following the opinion of Mahasri and Kismiyati (2015) that the parasite *Chilodonella sp.* oval-shaped like a heart,

the shape of the macronucleus is oval and the shape of the micronucleus is circular. How to infest this parasite in groups or colonies and often these ectoparasites are found in large numbers, when there is a decrease in temperature the fish will be easily infected with *Chilodonella sp.* The phylum Protozoa has a very fast life cycle and reproduction because it can reproduce in approximately 24 hours and therefore rapidly divides and is transmitted to fish.others (Amirullah et al., 2012). According to Tobler et al. (2007) Reproduction of *Chilodonella sp.* by mitotic division but also by conjugation. *Chilodonella sp.*

The ectoparasite worm *Dactylogyrus sp.* was also found in this study and found only on the gills, so it is referred to as gill worms and is one of the largest monogenean genera. Ectoparasite *Dactylogyrus sp.* has an opisthaptor, which is an organ to attach to the target organ of its host, which is equipped with two pairs of penetrating organs, which are like anchors. In this study, *Dactylogyrus sp.* is seen to have 2 pairs of eyes that look like black dots and have 4 indentations on the head. This is by the statement of Yuli et al. (2017) there are 2 pairs of black eyes on the anterior. Schmidt et al., (2006) explained the mechanism of the direct life cycle of monogenean parasites, which generally start with a single host, starting from an egg, becoming a ciliated larva oncomiracidium, then growing up. On *Dactylogyrus sp.* The life cycle will start from an adult worm that lays eggs and hatches into a ciliated larva (oncomiracidium) which has more than one cilia and eyespot and will swim until it attaches to the gills and then becomes an adult worm (Kabata, 1985). These worms are oviparous and have haptors, which are attachment organs that are equipped with two pairs of anchors and 14 lateral hooks.

Conclusion

The conclusion of the research on the identification of ectoparasites in pearl catfish (*Carias gariepinus*) at the age of one and three months in Maclele Cultivation, Tuban District, Tuban Regency, is a type of ectoparasite that infests pearl catfish in Makalele Cultivation, Tuban District, Tuban Regency. The species are *Trichodina nobilis*, *Trichodina pediculus*, *Chilodonella sp.*, and *Dactylogyrus sp.* On the incidence of ectoparasite infestation among pearl catfish aged one month and three months. there was no significant difference (p 0.05).

References

- Amirullah, S., Y. Dhahiyat and I. Rustikawati. 2012. Intensity and Prevalence of Ectoparasites in Fish in the Upper Cimanuk

- River, Garut Regency, West Java. *Journal of Fisheries and Marine Affairs*, 3(4), 271-282.
- Bell, G. and A. Burt. 2008. The Comparative Biology of Parasite Species Diversity: Internal Helminths of Freshwater Fish. *The Journal of Animal Ecology* 60(3): 1047-1064.
- Chen, CL 1963. Studies on Ectoparasitic Trichodinids from Freshwater Fish, Tadpole, and Crustaceans in China. *Acta Hydrobiologica Sinica* 3: 99-111.
- Dana, D., I. Effendi, K. Sumawidjaja and Y. Hadiroseyani. 2008. Trichodina Parasites on Seeds of Betutu Fish (*Oxyeleotris marmorata*). *Indonesian Journal of Aquaculture*. 1(1) : 5-8.
- Ghufran, M. and Kordi 2004. Management of Fish Pests and Diseases. Jakarta: Rineka Cipta and Bina Adiaksara.
- Hardi, EH 2015. Parasites of Aquatic Biota. Mulawarman University Press. Samarinda.
- Hardiroseyani. 2006. Catfish Parasite Intarization *Clarias* sp. in the Bogor area. *Indonesian Journal of Aquaculture*, 5(2). Bogor: 167-177.
- Husein, U. 2013. Research Methods for Business Thesis and Thesis Second Edition. Jakarta: Rajawali Press.
- Irianto, A. 2005. Teleostei Fish Pathology. Gadjah Mada University Press. Yogyakarta. 144-152.
- Kabata, Z. 1985. Parasites and Diseases of Fish Cultured in the Tropics. London and Philadelphia: Taylor and France: 318.
- Khotimah, A., Rokhmani, and E. Riwidharso. 2018. Prevalence and abundance of *Vorticella* sp. on Bkaur Crab (*Scylla serrata*) Landed at the Sleko Fish Auction Place, Cilacap Regency, Central Java. *Pros Sem Nas Masy Biodiv Indonesia* 4(1): 87-91.
- Mahasri, G and Kismiyati. 2015. Textbook of Parasites and Fish Diseases I (Science of Protozoan Diseases in Fish and Shrimp). Faculty of Fisheries and Marine Affairs. Universitas Airlangga. Surabaya. 1(1): 21-29.
- Mahendra and Nurbadriati. 2019. The prevalence and intensity of ectoparasites in Tawes (*Puntius javanicus*) in Meunasah Krueng Village, Beutong District, Nagan Raya Regency. *AKUKULTURA Journal* Available online at Volume 3(1) P-ISSN: 2579-4752, E-ISSN: 2620-7397.
- Murugaian, P. 2008. Effect of Temperature on the Behavioral and Physiological Responses of Catfish, *Mystus gulio* (Hamitto). Department of Biotechnology, Vinyaka Missions University.
- Nugroho, RA, LT Pambudi, D. Chilmawati and AHC Haditomo. 2012. Application of Aquaponic Technology in Freshwater Fish Cultivation for Optimization of Production Capacity. *Fisheries scientific journal*. 8(1) : 46-51.
- Pramono, TB and H. Syakuri. 2008. Parasite Infection on the Body Surface of the Nile Fish (*Osteochilus hasselti*) Traded at PPI Purbalingga. *Fisheries Scientific Periodic FPIK, Jenderal Sudirman University, Purwokerto*. 3(2).
- Tuban Regency Medium Term Development Plan [RPJMD] 2016 to 2021. 2016. Tuban Regency. 1-10.
- Riko, YA, Rosidah and T. Herawati. 2012. Intensity and Prevalence of Ectoparasites in Milkfish (*Chanos chanos*) in KJA in Cirata Reservoir, Cianjur Regency, West Java. *Journal of Fisheries and Marine Affairs*. 3(4) 231-241.
- Schmidt, LJ, MP Gaikowski and WH Gingerich. 2006. Environmental assessment for the use of hydrogen peroxide in aquaculture for treating external fungal and bacterial diseases of cultured fish and fish eggs. La Crosse, Wisconsin: USGS Report. 180.
- Sekar, PM, AHC Haditomo and Desrina. 2016. Monogenea Infestation of Freshwater Consumed Fish in Aquaculture Ponds, Ngrajek Village, Magelang. *Journal of Aquaculture Management and Technology* Volume 5, Number 1. Semarang. 162-170.
- Shafrudin, D., Yuniarti and M. Setiawati. 2006. Effect of seed density of African catfish (*Clarias* sp.) on production in aquaculture systems with nitrogen control through the addition of wheat flour. *J Indonesian Aquaculture* 5(2):137-147.
- Tobler, M., I. Schlupp, J. Francisco, G. de Leon, M. Glaubrech and M. Plath. 2007. Extreme habitats as a refuge from parasite infections. Evidence from an extremophile fish. *parasitology*. 1-6.
- Vera, N., P. Simonovic and P. Vesna. 2003. Preferences of Trichodinids (Ciliata, Peritrichia) occurring on fish-pond carp for particular organs and some morphological implication. *Acta Veterinaria (Belgrade)* 53(1): 41- 46.
- Woo, JL 2006. Fish Disease and Disorder Parasites. The University of Guelph. CAB. International. Canada.
- Yuli, S., H. Harris, and IA Yusanti. 2017. Ectoparasite attack rate on catfish (*Pangasius hypophthalmus*) cultured in floating net cages on the Musi River, Palembang. *Journal of Fisheries and Aquaculture Sciences* 12 (2): 50-58.

AUTHOR INDEX

Afiyah Rahmadani	54	M. Gandul Atik Yuliani	38
Agus Sunarso	38	Maudika Pamela Fiorenza	54
Arimbi	54	Mufasirin	50
Azizatun Nimah	60	Nashihatun Sholihah	38
Budiarto	60	Nove Hidajati	42
Dian Ayu Permatasari	46	Nunuk Dyah Retno Lastuti	54
Dhandy Koesoemo Wardhana	38	Nusdianto Triakoso	50
Endang Suprihati	54	Poedji Hastutiek	38, 50
Galuh Swa Ardhya	60	Setiawan Koesdarto	42, 46, 60
Hani Plumeriastuti	60	Sherbina Bai	38
Herry Agoes Hermadi	50	Sri Hidanah	38
Indira Vania Carisya	42	Sunaryo Hadi Warsito	54
Iwan Sahrial Hamid	46	Taufan Ary Handoko	50
Karina Rachmawati	46	Tjuk Imam Restiadi	42

SUBJECT INDEX

- Ascariasis 42, 45
Ascaridia galli 42, 43, 44, 45
Culling Layer Hen 38, 40
Cat 50, 51, 52, 53
Clarias gariepinus 60, 61, 65, 66
Chilodonella sp. 60, 61, 63, 64, 65
Dog 50, 51, 52, 53
Ectoparasite 38, 39, 40, 41, 46, 47, 48, 49
Endoparasite 46, 47, 48, 49
Gastrointestinal 46
Incidence rate 60
Native chicken 54, 55, 56, 58
Parasite 50, 51, 52, 53
Prevalence 50, 51, 52, 53
Raillietina echinobothrida 54, 55, 56, 57, 58
Raillietina tetragona 54, 55, 56, 57, 58
Ovicidal Effects 42, 43, 45
Scanning Electron Microscope 54, 55, 56, 57, 58
Stray Cat 46, 47, 48, 49
Trichodina nobilis 60, 62, 65
Trichodina pediculus 60, 62, 65
Wonokromo Market 38, 39, 40, 41

SUBSCRIBED FORM

AREA	Issue* (give sign √)		
	1 year	2 year	3 year
Surabaya	<input type="checkbox"/> Rp. 200.000,-	<input type="checkbox"/> Rp. 350.000,-	<input type="checkbox"/> Rp. 450.000,-
East Java	<input type="checkbox"/> Rp. 250.000,-	<input type="checkbox"/> Rp. 400.000,-	<input type="checkbox"/> Rp. 500.000,-
Other Java	<input type="checkbox"/> Rp. 300.000,-	<input type="checkbox"/> Rp. 450.000,-	<input type="checkbox"/> Rp. 550.000,-
Other Countries	<input type="checkbox"/> USD 50	<input type="checkbox"/> USD 75	<input type="checkbox"/> USD 100

*(published 2 times a year)

I propose to subscribe to Journal of Parasite Science

Name:

Institution:

Office address and Phone No:

.....

Home address and Phone No:

.....

.....

Country:

Fax:

Email:

Postal code:

Date:

Signature:

I pay for this journal subscription using (give sign √):

Western Union Cheque with MTCN No:

Bank transfer / online to:

Bank Mandiri

Account no 142 00 1670625 0

Dr. Kusnoto, drh., M.Si

Send proof of payment with the subscription form to email jps@fkh.unair.ac.id

AUTHOR GUIDELINES

Journal of Parasite Science (JoPS) with registered number [P-ISSN \(2599-0993\)](#) ; [E-ISSN \(2656-5331\)](#) published by the Division of Veterinary Parasitology, Faculty of Veterinary Medicine, Universitas Airlangga.

Journal of Parasite Science (JoPS) presenting research articles, case reports, community development activities and literature studies of veterinary medicine. Published twice a year on March and September. Since 2002 the Faculty of Veterinary Medicine has had Veterinary Medicine Media, because considering that there is only one journal that must accommodate many articles written by lecturers and students, various journals have been developed according to the disciplines in each department, one of which is the JoPS. JoPS can publish articles from various faculties and institutions related to the parasitology. JoPS which has been published since 2017 is here for accommodate publication obligations for undergraduate and postgraduate students as well as other parties conducting research in the field of parasitology. Loading of articles in the JoPS through the Open Journal System (OJS). Complete information for article loading and article writing instructions are available on the website and every issue. Incoming articles will go through a selection process by editors and reviewers. Manuscripts should be typed in double spaced, except for titles, abstracts, titles of tables and tables, headings of figures, reference, and attachments, typed according to separate provisions. Paragraph formatting: new paragraph begins (first line) 0.3 inches. The standard font for writing is Times New Roman 12. Using A4 paper size (8.27 x 11.69)". Using English. Tables / Illustrations / Images must be of high contrast, also include separate scanning files (photos) with papers in JPG file format. Table captions, pictures or other explanations in the attachment should be typed 1 (one) space.

Title should be concise but informative, as far as possible in no more than 12 words.

Abstract written in English, do not exceed 250 words, containing elements of background, material and methods, results and conclusions.

Keywords maximum of 6 (six) words or phrases, written after the abstract in each languages, alphabetically ordered. As far as possible avoid using keywords from the title.

Introduction consists of the background, research objectives, problem formulation and benefits. The background includes four main ideas, namely: problem statement, problem scale, problem chronology and research approach. Contains a description of the facts and expectations for the completion of the research. Described based on concepts built from references in the literature. The research question and/or hypothesis (if any) in the introduction should not write in the form of points or should not be numbered.

Method in details so that other author could do the similar research with the written method in the article. Materials and methods are brief descriptions of the research design, how to conduct research and mention the materials/ objects and tools used.

Results and Discussion compared with other research that ever conducted or discuss based on the theoretical framework that is used in the research. Use up-to-date references such as books that published in the last ten years and journal articles that published in the last five years (except the reference that being recommended by the reviewer in this field).

Tables are made without vertical lines (use only lines at the top and bottom of the table as well as for separating heading from the main table), with table title placed before the table, numbered in Arabic numerals (Table X), and have to be referred in the text. The description of the table is placed after the table; it must be concise but clear enough so that the table separately can be understood without referring to the text. The table along with the title and description are placed after the References.

Figures presented are only those that support the findings of the study, and not restatement of data from tables in the form of figures. When resulted data in the form of figure is more informative, interesting or significant,

presentation of data in table form is not required. Figure title is placed after the figure, numbered in Arabic numerals (Figure Y), and have to be referred in the text. The description of the figure is placed under the title of the figure; it must be concise but clear enough so that the figure separately can be understood without referring to the text. Image (in JPEG format) is sent in separate file. The title and description of the figure are placed after the References.

Conclusion in one until two paragraphs without using the bullet points or number.

Approval of Ethical Commission have to be stated (number and institution) if the manuscript is constructed based on a research using live animals.

Acknowledgements are delivered to the research funders (state the name, number and recipient name of the grant, if applicable), and to those who have helped carry out the research.

References should be written according to the APA guidelines, that uses the family name of the author at the beginning of the sequence. Example: Family Name, Abbreviated the First Name and the Middle Name. Do not use commas or point for the abbreviated name. For the year of the references give bracket without points. The reference from a journal article, anthology books or book chapter must include the page. References from the internet not journals are not recommended.

The example of the references as below:

Online journal article

Johnson, K. D., Winkelman, C., Burant, C. J., Dolansky, M., & Totten, V. (2014). The factors that affect the frequency of vital sign monitoring in the emergency department. *JEN: Journal of Emergency Nursing*, 12(2), 27-35.
<https://doi.org/10.1016/j.jen.2012.07.023>

Book (authored or entire book) *

Judd, D. M., Sitzman, K., & Davis, M. (2010). *A history of American nursing: Trends and eras*. Jones and Bartlett Publishers.

Chapter in an edited book *

Lynch, S. (2010). Pain management. In S. R. Hardin and R. Kaplow (Eds.), *Cardiac surgery essentials for critical care nursing* (3rd ed., pp. 287-309). Jones & Bartlett.

PowerPoint slides (from classroom website)

Nicholson, A. (n.d.). RN to BSN program orientation [PowerPoint slides]. Canvas @UIowa: <https://uiowa.instructure.com/>

Webpage

Nelson, L. (2019). Nurses say violent assaults against healthcare workers are a silent epidemic. <https://nurse.org/articles/-/workplace-violence-in-nursing-and-hospitals/>

Report

Institute of Medicine (2010). The future of nursing: Leading change, advancing health [report brief]. <https://www.nap.edu/resource/12956/Future-of-Nursing-2010-Report-Brief.pdf>

Review Article

Journal of Parasite Science received review articles on topics included in the scope of the journal which are of current interest. Reviews are invited by the editor. The review must be a comprehensive analysis and perspective on the state of the field and where it is heading. Reviews will be subject to the same peer review process as is applied for original papers. The manuscript is arranged in the same way as the original article with an unstructured abstract (maximum of 250 words). Manuscripts should be typed in double spaced, except for titles, abstracts, titles of tables and tables, headings of figures, reference, and attachments, typed according to separate provisions. Paragraph formatting: new paragraph begins (first line) 0.3 inches. The standard font for writing is Times New Roman 12. Using A4 paper size (8.27 x 11.69)". Using English. Tables / Illustrations / Images must be of high contrast, also include separate scanning files (photos) with papers in JPG file format. Table captions, pictures or other explanations in the attachment should be typed 1 (one) space.

Case reports:

Journal of Parasite Science receives quality and interesting case reports with topics included in the scope of this journal. All reports are peer reviewed as is applied for original articles. All instructions are the same as for the original articles.

Indexing By :



Plagiarism Checker



Management Reference Tools



Division of Veterinary Parasitology
Faculty of Veterinary Medicine, Universitas Airlangga
Kampus "C" Jl. Mulyorejo Surabaya 60115
Phone: (031) 5992785; 5993016; Fax. (031) 5993015
e-mail: jparasitol@gmail.com ; jps@fkh.unair.ac.id
Website : <https://e-journal.unair.ac.id/JoPS>

