

MEDIA

p-ISSN : 0215-8930; e-ISSN : 2775-975X

MEDIA

KEDOKTERAN HEWAN

VETERINARY MEDICINE JOURNAL

MKH (VET. MED. J) VOL. 33, NO. 3, AUGUST 2022




Figure source: Ridwan et al. 2021. Media Kedokteran Hewan, 32 (3): 105-113

Media Kedokteran Hewan (Veterinary Medicine Journal)

Volume 33, No.3, September 2022

Media Kedokteran Hewan (p-ISSN: 0215-8930) (e-ISSN: 2775-975X) (established 1985) publishes all aspects of veterinary science and its related subjects. Media Kedokteran Hewan publishes periodically three times a year (January, May, and September). Media Kedokteran Hewan publishes original articles, review articles, and case studies in Indonesian or English, with an emphasis on novel information of excellent scientific and/or clinical quality, relevant to domestic animal species and biotechnology of veterinary medicine from researchers, lecturers, students, and other practitioners around Indonesia and worldwide.

Editor in Chief

Yulianna Puspitasari, drh., M.VSc., Ph.D. Universitas Airlangga, Indonesia

Associate Editor

Dian Ayu Permatasari, drh., M.Vet. Universitas Airlangga, Indonesia
Dhandy Koesoemo Wardhana, drh., M.Vet. Universitas Airlangga, Indonesia

Assistant Editor

Rosa Kartika Al Jihadi, K5 Perawatan Hewan Kota Kediri

Editorial Board Members

Prof. Dr. Pudji Srianto, drh. M.Kes. Universitas Airlangga, Indonesia
Prof. Dr. Widya Paramita Lokapirnasari, drh., MP. Universitas Airlangga, Indonesia
Dr. Attaur Rahman, College of Animal Husbandry & Veterinary Sciences, Abdul Wali Khan University Mardan, Pakistan
Dr. Shekhar Chhetri, DVM, M.Sc. Royal University of Bhutan, Bhutan
Dr. Ugbo Emmanuel Nnabuike, Ebonyi State University, Nigeria
Dr. Abzal Abdramanov, M.Sc. Kazakh National Agrarian Research University, Kazakhstan
Dr. Jagdish Gudewar, B.VSc., M.VSc. College of Veterinary & Animal Sciences, Parbhani, India
Abdul Rehman, DVM., M.Phil., Ph.D. University of Veterinary and Animal Sciences, Lahore, Pakistan
Dr. Freshinta Jellia Wibisono, drh., M.Vet. Universitas Wijaya Kusuma, Surabaya, Indonesia
Roza Azizah Primatika, S.Si, M.Si. Universitas Gadjah Mada, Indonesia
Fidi Noraini, drh., M.Si. Universitas Brawijaya, Indonesia
Dr. Nanik Hidayatik, drh., Universitas Airlangga, Indonesia



Media Kedokteran Hewan

Veterinary Medicine Journal
Volume 33 No.3 2022



Media Kedokteran Hewan by [Unair](https://unair.ac.id) is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/)

Mohammad Thohawi Elziyad P, drh., M.Si. Universitas Airlangga, Indonesia
Ragil Angga Prastiya, drh., M.Si. Universitas Airlangga, Indonesia
Mirza Atikah Madarina Hisyam, drh., M.Si. Universitas Airlangga, Indonesia

Peer Reviewer:

Prof. Dr. Imam Mustofa, drh, M.Kes. Universitas Airlangga, Indonesia
Prof. Dr. Fedik Abdul Rantam, drh. Universitas Airlangga, Indonesia
Prof. Dr. Ir. Sri Hidanah, MS. Universitas Airlangga, Indonesia
Prof. Ooi Hong-Kean, DVM.,Ph.D. Japan Science and Technology Agency, Japan
Sattar Mengal, DVM., M.Phil., Ph.D. Lasbela University of Agriculture, Water and
Marine Sciences, Pakistan
Innocent Damudu Peter, DVM., M.VSc., Ph.D. University of Maiduguri, Nigeria
Mohd Farhan Hanif bin Reduan, DVM., Ph.D. Universiti Malaysia Kelantan, Malaysia
Nurrul Shaqinah Nasruddin, DVM, Ph.D. National University of Malaysia, Malaysia
Norfitriah Mohamed Sohaimi, DVM., Ph.D. Universiti Putra Malaysia, Malaysia
Dr. Mark Hiew Wen Han, Universiti Putra Malaysia, Malaysia
Taira Kensuke, DVM., Ph.D. Azabu University, Japan
Shin Yoon-Joo, Seoul National University, South Korea
Rey Oronan, University of the Philippines, Los Banos
I Ketut Mudite Adnyane, drh., M.Si., Ph.D., PAVet. Institut Pertanian Bogor
University, Indonesia
Eddy Sukmawinata, DVM., M.Sc., Ph.D. Department of Infectious Diseases, Faculty
of Medicine, University of Miyazaki, Japan
Dede Setiawan, DVM, Charles River Laboratories Montreal, Quebec, Canada
Dr. Sri Rahmatul Laila, Institut Pertanian Bogor University, Indonesia
Dr. Chaerul Basri, Institut Pertanian Bogor University, Indonesia
Wredha Sandhi Ardha Prakoso, drh., M.Sc. Karantina Hewan, Badan Karantina
Pertanian
Dilasdita Kartika Pradana, drh., M.Si. Balai Besar Veteriner Kelas I Denpasar, Bali,
Indonesia
Dr. Sarasati Windria, drh. Universitas Padjajaran, Bandung, Indonesia
Drh. Mudhita Zikrullah Ritonga, S.KH.,M.Vet, Universitas Syiah Kuala, Indonesia
Shelly Kusumarini R, drh., M.Si. Universitas Brawijaya, Indonesia
Syaiful Rizal, drh., M.Si. Badan Riset dan Inovasi Nasional, Indonesia
Suryo Kuncorojakti, drh., M.Vet., Ph.D. Universitas Airlangga, Indonesia
Adiana Mutamsari Witaningrum, drh., M.Vet. Universitas Airlangga, Indonesia
Oky Setyo Widodo, drh., M.Si. Universitas Airlangga, Indonesia
Martia Rani Tacharina, drh., M.Si. Universitas Airlangga, Indonesia

Editorial Office : Faculty of Veterinary Medicine, Universitas Airlangga
Kampus C Universitas Airlangga Mulyorejo, Surabaya 60115
Telp. (031)5992785, 5993016 Fax. (031) 5993015
e-mail : medvetj@journal.unair.ac.id



Author Guidelines

Veterinary Medicine Journal is a journal that contains scientific articles on veterinary medicine and animal husbandry biotechnology published by Faculty of Veterinary Medicine Airlangga University 3 times per year on January, May and September. Veterinary Medicine Journal received manuscripts in the form of original research articles, review articles and case reports in Indonesian and English. Manuscripts received must be original, current and have never been published or are being planned to be published in other scientific journals.

Manuscripts must be submitted online through the Open Journal System (OJS) in Word format. The entire text is typed in Book Antiqua 11pt double spaced. The title is written with a Title case (bold, 14pt, align center). The full length of the manuscripts is a maximum of 12 pages of HVS paper. Italicize only for species names or terms that have not been standardized as Indonesian. Define abbreviations upon first appearance in the text. Do not use non-standard abbreviations unless they appear at least three times in the text. Keep abbreviations to a minimum. Avoid unnecessary duplication of text.

The first page contains titles in Indonesian and English, followed by full names of all authors without titles and initials (bold, center), followed by the name and complete address of the respective institution (marked with numeric superscripts) and e-mail of corresponding authors (marked with *superscript).

The second page forward contain:

ABSTRACT in English and Indonesian, **INTRODUCTION, MATERIALS AND METHODS, RESULTS, DISCUSSION, CONCLUSIONS, APPROVAL OF ETHICAL COMMISSION, ACKNOWLEDGEMENTS** and **REFERENCES**.

Title should be concise but informative, as far as possible in no more than 12 words.

Abstract written in Indonesian and English, do not exceed 200 words, containing elements of background, material and methods, results and conclusions.

Keywords maximum of 5 (five) words or phrases, written after the abstract in each languages, alphabetically ordered. As far as possible avoid using keywords from the title.

Introduction should be brief, containing elements of background, problems, objectives and reference sources that support.

Research materials (materials and equipment) do not need to be mentioned separately, but rather integrated in the method used, complete with the brand and catalog number if applicable. Do not include common supplies, such as test tubes, pipette tips, beakers, etc. or standard laboratory equipment.

Method must be concise but sufficiently detailed (with reference or modification) so research can be repeated by other researchers.

Results are displayed in a concise but clear narrative with/without tables or figures.

Tables are made without vertical lines (use only lines at the top and bottom of the table as well as for separating heading from the main table), with table title placed before the table, numbered in Arabic numerals (**Table X**), and have to be referred in the text. The description of the table is placed after the table; it must be concise but clear enough so that the table separately can be understood without referring to the text. The table along with the title and description are placed after the References.

Figures presented are only those that support the findings of the study, and not restatement of data from tables in the form of figures. When resulted data in the form of figure is more informative, interesting or significant, presentation of data in table form is not required. Figure title is placed after the figure, numbered in Arabic numerals (**Figure Y**), and have to be referred in the text. The description of the figure is placed under the title of the figure; it must be concise but clear enough so that the figure separately can be understood without referring to the text. Image (in JPEG format) is sent in separate file. The title and description of the figure are placed after the References.

Discussion contains explanation of what are found related to **the importance of your study** and how it may be able to answer the research question, comparison of findings (internally, between research data, and externally, compared with findings from other studies) and cause-effect analysis.

Conclusion does not only repeat the results of the study, but summarize the findings into a narrative that impacts on the development of science and/or practitioners in the field of veterinary reproduction.

Approval of Ethical Commission have to be stated (number and institution) if the manuscript is constructed based on a research using live animals.

Acknowledgements are delivered to the research funders (state the name, number and recipient name of the grant, if applicable), and to those who have helped carry out the research.

References are sorted alphabetically based on the author's last name. The *titles of journals* should be *abbreviated* according to the style *used* by each journal. Sixty (60%) scientific journal/article and (40%) text book.

Roitt, I., J. Brostoff, and D. Male. 2005. Immunology. 6th Ed. Black Well Scientific Pub. Oxford.

Staropoli, I., J.M. Clement, M.P. Frenkiel, M. Hofiung and V. Deuble. 2006. Dengue-1 virus envelope glycoprotein gene expressed in recombinant baculovirus elicits virus neutralization antibody in mice and protects them from virus challenge. Am.J. Trop. Med. Hygi; 45: 159-167.

Review article



Veterinary Medicine Journal received review articles on topics included in the scope of the journal which are of current interest. Reviews are invited by the editor. The review must be a comprehensive analysis and perspective on the state of the field and where it is heading. Reviews will be subject to the same peer review process as is applied for original papers. The manuscript is arranged in the same way as the original article with an unstructured abstract (maximum of 200 words). The number of references is limited to 50-70, with 75% of them have to be from publications in the past 10 years. The total length should not exceed 12 pages (HVS paper, double spaced 11pt Book Antiqua).

Case reports:

Veterinary Medicine Journal receives quality and interesting case reports with topics included in the scope of this journal. All reports are peer reviewed as is applied for original articles. All instructions are the same as for the original articles.



**Media Kedokteran Hewan
(Veterinary Medicine Journal)**

Vol. 33 No. 3 September 2022

LIST OF CONTENT

	Page
1. Ergothioneine Modulates Biomarkers of Oxidative Stress, Biochemical Profiles and Rectal Temperatures Arabian Stallions Following Exercise in A Hot-Humid Environment Adakole S. Adah¹, Joseph O. Ayo, Peter I. Rekwot, Tagang Aluwong, Deborah A. Adah	163-176
2. Pathology and Characterization of Fowl Pox Virus Infection in a Turkey-Chicken Backyard Flock, Nigeria Alfiana Laili Dwi Agustin, Novarina Sulsia Ista'In Ningtyas	177-187
3. <i>Escherichia coli</i> Infections, and Antimicrobial Resistance in Poultry Flocks, in North Central Nigeria Olatunde B. Akanbi, Isaac D. Olorunshola, Peter Osilojo, Eunice Ademola, G.O.A. Agada, Julius O. Aiyedun, Christiana Ibrinke Odita and Shola David Ola-Fadunsin	188-207
4. The correlation between antioxidant levels of superoxide dimustase (SOD) with the abnormalities in different diluent levels and shelf life of Muscovy Duck semen stored at cold temperatures Fitriani, Erna Yuniati, M. Irwan Zakir	208-213
5. Phylogenetic Study of CyHV-3 Virus Glycoprotein Encoding Genes Koi Fish in Several Regions of East Java Indra Sukma Putra, Mohammad Fathoni, Suwarno, Jola Rahmahani	214-232



6. Cytotoxicity of Kembang Bulan (*Tithonia diversifolia*) Ethanolic Leaf Extract on Rabbit Limbal Mesenchymal Stem Cells 233-239

Hani Plumeriastuti, Vida Seanita Zahro, Nusdianto Triakoso, Fedik Abdul Rantam, Iwan Sahrial Hamid, Mustofa Helmi Effendi, Annise Proboningrat

7. Case report of *Spinturnix* spp. infestation on Fruit Bats (*Rousettus* spp.) in Banyuwangi 240-250

Fransiska Okta Zania, Audina Putri Geraldine, Citra Kurnia Putri, Ryanka Edila, Aditya Yudhana



Cytotoxicity of Kembang Bulan (*Tithonia diversifolia*) Ethanolic Leaf Extract on Rabbit Limbal Mesenchymal Stem Cells

Hani Plumeriastuti^{1*}, Vida Seanita Zahro², Nusdianto Triakoso³, Fedik Abdul Rantam⁴, Iwan Sahrial Hamid⁵, Mustofa Helmi Effendi⁶, Annise Proboningrat¹

¹Division of Veterinary Pathology, Department of Veterinary Medicine, Faculty of Veterinary Medicine, Universitas Airlangga

²Bachelor of Veterinary Medicine Study Program, Department of Veterinary Medicine, Faculty of Veterinary Medicine, Universitas Airlangga

³Division of Veterinary Clinic, Department of Veterinary Medicine, Faculty of Veterinary Medicine, Universitas Airlangga

⁴Division of Veterinary Microbiology, Department of Veterinary Medicine, Faculty of Veterinary Medicine, Universitas Airlangga

⁵Division of Veterinary Basic Medicine, Department of Veterinary Medicine, Faculty of Veterinary Medicine, Universitas Airlangga

⁶Division of Veterinary Public Health, Department of Veterinary Medicine, Faculty of Veterinary Medicine, Universitas Airlangga

*E-mail: hani-p@fkh.unair.ac.id

ABSTRACT

Kembang Bulan (*Tithonia diversifolia*) is known as an herbal plant that has many useful properties to cure disease, such as diabetes, malaria, and other infections. Instead of much usefulness, *T. diversifolia* has main biological properties that may induce toxicity in this plant known as sesquiterpene lactones. The aim of this study was to identify the cytotoxicity effect of *T. diversifolia* leaf extract on limbal mesenchymal stem cells of rabbits using the MTT assay. *T. diversifolia* leaf extract was divided into 4 concentrations (0.5%, 0.25%, 0.125%, and 0.0625%) and given to limbal mesenchymal stem cells of rabbits in 96-well microplates, which would be incubated for 24 hours. The cytotoxicity result was obtained using the MTT assay method. Every well would be observed to see the cell life level of each concentration. The optical density absorbance was calculated using a microplate reader. *T. diversifolia* leaf extract with a concentration of 0.0625% has the biggest viability with a level of existence of limbal mesenchymal stem cells of 50%, and the other *T. diversifolia* leaf extract concentrations of 0.5%, 0.25%, and 0.125% have less than 50% level of existence of limbal mesenchymal stem cells.

Keyword: LC50, live cell, Mexican sunflower, MTT assay



INTRODUCTION

The usage of traditional medicine is safer compared to synthetic medicine (Yuan et al., 2016). One of the most common traditional medicines was made from *Tithonia diversifolia* leaves, which belong to the family Asteraceae and are commonly used by people from South Asia to cure many kinds of illnesses (Ajao and Moteetee, 2017). In Indonesia, this plant is usually called "Daun Insulin" (Ammar et al., 2021). *T. diversifolia* can be used as medicine for diabetic, malaria, and many other infectious diseases (Di Giacomo et al., 2015).

T. diversifolia has the potential to be developed as a standardized herbal medicine that is safe for health purposes. The first step in drug development is the pre-clinical test, which is a toxicity test conducted both in vitro and in vivo (Fischer et al., 2020). A cytotoxicity test was conducted for evaluation and screening using culture in vitro. The advantages of this test include its simplicity, speed, sensitivity, and lack of the need for an animal experiment (Li et al., 2015).

The MTT (methyl thiazolyl diphenyl-tetrazolium) assay is an easy and low-cost technique commonly used to evaluate the cytotoxicity of many drug candidates. The idea behind this assay was that the amount of formazan crystal formation was proportional to the number of living cells, including their metabolic activities (Li et al., 2012).

The MTT assay employs 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide via a redox reaction inside the cell. It is determined by the cell's ability to convert MTT salt into formazan, as indicated by the color changing from yellow to a blue-purple like form. The MTT assay works on the premise that only living cells have the ability to change the tetrazolium ring via mitochondrial activity. The result of this process can be examined spectrophotometrically using a microplate reader (Buranaamnuay, 2021; Proboningrat et al., 2021). The intensity of the purple color correlates with the number of living cells. The percentage of death cells is then calculated using the number of living cells (Proboningrat et al., 2021).

The standard for measuring toxicity is LC50 (lethal concentration at 50%). It depends on the incubation period (Zhang et al., 2007). A toxicity test on primary cells used lethal concentration (LC50) as the standard, where the sensitivity of the drug was measured according to the rising number of death cells after drug treatment compared to the decreasing number of life cells in the untreated control. While the cell line inhibitory concentration (IC50) is suitable to measure the toxicity of the drug (He et al., 2016).

Limbal mesenchymal stem cells have recently been investigated as a cell-based alternative therapy for treating degenerative diseases. The ability of stem cells to self-renew and



differentiate is intriguing for cytotoxicity assay applications. Stem cells are an excellent model for predicting toxicity because they can be used to assess cytotoxic effects on cell viability and the cell differentiation process. Besides, they may be more relevant for predicting human or animal toxicity. Furthermore, they are a well-established in vitro model for human or animal multipotent cell populations, they do not require as many supplements as NHK, and they are commercially available or can be self-isolated in our laboratory (Abud et al., 2015).

The aim of this study was to observe the effect of the serial concentration of *Tithonia diversifolia* leaf extract on rabbit limbal mesenchymal stem cells.

MATERIALS AND METHODS

Materials

Materials used in this study were *Tithonia diversifolia* leaves, ethanol 70%, aerosol carboxyl, sterilized aquadest, limbal mesenchymal stem cell isolated from rabbit, phosphate buffer saline (PBS) 10%, fetal bovine serum (FBS), 3-(4,5-dimethylthiazol-2-yl) 2,5-diphenyl tetrazolium bromide (MTT), dimethyl sulfoxide (DMSO), modified minimum essential medium eagle-alpha (MEM- α), and trypsin EDTA.

Experimental Design

This study consists of two control groups and four treatment groups. The control groups were cell control (limbal mesenchymal stem cells

cultured in MEM- α medium) and media control (MEM- α medium only). Treatment groups consisted of limbal mesenchymal stem cells treated with *T. diversifolia* leaf extract at concentrations of 0.5%, 0.25%, 0.125%, and 0.0625%.

Preparation of *Tithonia diversifolia* Leaf Extract

2.5 kg of *T. diversifolia* leaf powder was macerated in 25 L of 70% ethanol (60 °C, 3 hours). The liquid extract was then evaporated at 60 °C to leave 4.5 liters of liquid. Then, it was mixed with 58 grams of carboxyl aerosol for 5 minutes until it became homogeneous. The resulting thick extract was dried in an oven at 45 °C. Prior to the in vitro assay, the dried powder of *T. diversifolia* leaf extract was dissolved in MEM- α medium with concentrations of 0.5% (5,000 $\mu\text{g}/\text{mL}$), 0.25% (2,500 $\mu\text{g}/\text{mL}$), 0.125% (1,250 $\mu\text{g}/\text{mL}$), and 0.0625% (625 $\mu\text{g}/\text{mL}$).

MTT Assay

The cytotoxicity of *T. diversifolia* leaf extract was evaluated using 3-(4,5-dimethylthiazol-2-yl) 2,5-diphenyl tetrazolium bromide (MTT) assay. Rabbit limbal mesenchymal stem cells were cultured in 96 well microplate with density of 3.5×10^3 cells/well for 24 hours (CO₂ 5%, 37 °C). First, the medium from all wells was removed. Cells were then treated with four concentrations of *T. diversifolia* leaf extract (0.5%, 0.25%, 0.125%, and 0.0625%) for 24 h. The medium was removed, and the cells were incubated with 25 μL MTT (5 mg/mL) for 4 hours (37 °C). The cells were observed using an inverted microscope to see the presence of formazan crystals. Next, 50-



100 μ L of DMSO was added to each well to dissolve the formazan crystals. The absorbance of dissolved formazan was measured spectrophotometrically using a microplate reader with a wavelength of 595 nm. The darker the color, the higher the optical density (OD), which indicates the number of living cells. The percentage of live cells is calculated using the formula:

$$\text{Live cells (\%)} = \left(\frac{\text{OD of treatment cells} + \text{OD of media control}}{\text{OD of cell control} + \text{OD of media control}} \right) \times 100\%$$

Data Analysis

The regression relationship between the concentration of *T. diversifolia* leaf extract and the percentage of viable limbal mesenchymal stem cells was analyzed using probit analysis to obtain a lethal concentration (LC50) value.

RESULTS AND DISCUSSION

The result showed that the rise in *Tithonia diversifolia* leaf extract concentration caused the death of limbal mesenchymal stem cells as well, due to the cell's inability to reduce MTT into formazon crystal. It is observed that the blue-purple form cannot be observed while the color of the medium containing phenol red is dominant (Figures 1b, 1c, and 1d). The positive control produced a different result. It can be observed that there was a transformation from MTT into formazon crystal marked by a blue-purple color with high density (Figure 1a). Most living cells were found at the lowest concentration of *T. diversifolia* leaf extract (Figure 1e).

A material that will be developed into a new drug candidate must pay attention to its toxic properties. The drug substance used must be non-toxic, non-irritating, have biocompatibility properties, and not have adverse effects on the biological environment, both locally and systemically (Meizarini, 2005).

According to Passoni et al. (2013), the toxic compounds in *T. diversifolia* leaf extract that are thought to be responsible for causing toxic effects are sesquiterpene lactones and chlorogenic acid derivatives, which cause damage to the liver and kidneys. Leaf extracts containing higher concentrations of chlorogenic acid derivatives were given to rats for 90 days and caused liver damage. Meanwhile, extracts rich in sesquiterpene lactones cause kidney damage. The reported safe dose of *T. diversifolia* leaf extract is less than 100 mg/kg body weight.

The mechanism of toxicity of *T. diversifolia* leaves related to sesquiterpene lactones includes alkylating and non-alkylating mechanisms. Alkylation toxicity by sesquiterpene lactones is known to induce ROS accumulation. Meanwhile, the non-alkylating mechanism of sesquiterpene lactones is known to inhibit a class of CA²⁺ transport enzymes, which causes cytosolic calcium levels to increase, thus leading to oxidative DNA damage (Amorim et al., 2013).






Figure 1. Microscopy of limbal mesenchymal stem cells at magnifications of 40-100x. (a) Cell control; (b) extract treatment with a concentration 0,5%; (c) extract treatment with a concentration 0,25%; (d) extract treatment with a concentration 0,125%; and (e) extract treatment with a concentration 0,0625%.

The parameter commonly used in the cytotoxicity test of an herbal substance is the LC50 value. LC50 indicates a concentration value that results in 50% inhibition of cell proliferation and indicates the potential toxicity of a compound to cells. This value is a benchmark for conducting cell kinetics observation tests. The LC50 value can indicate the potential of a

compound as a cytotoxic agent; the greater the LC50 value, the less toxic the compound will be (Djajanegara et al., 2009).

Observations on the cytotoxicity test of *T. diversifolia* leaf extract with concentrations of 0.5%, 0.25%, 0.125%, and 0.0625% in rabbit limbal mesenchymal stem cells for 24 hours showed a different percentage of living

cells in each treatment. The percentage of limbal mesenchymal stem cells that survived the treatment with *T. diversifolia* leaf extract at 0.0625% was 50%. *T. diversifolia* leaf extract with concentrations of 0.5%, 0.25%, and 0.125% yielded a value of less than 50%. It was noted that at concentrations of 0.5%, 0.25%, and 0.125%, the number of living cells that could be observed was 14%, 29%, and 33%, respectively.

The result of this study revealed that the higher the concentration of *T. diversifolia* leaf extract, the lower the percentage of living cells from limbal mesenchymal stem cells, indicating a higher toxic effect of *T. diversifolia* leaf extract.

CONCLUSIONS

It can be concluded that the greater the concentration of *T. diversifolia* leaf extract, the lower the number of viable rabbit limbal mesenchymal stem cells. The concentration of *T. diversifolia* leaf extract required to kill 50% of rabbit limbal mesenchymal stem cells (LC50) within 24 hours is 0.059%.

ACKNOWLEDGEMENTS

The authors express their gratitude to the Stem Cell Research and Development Center, Universitas Airlangga, Surabaya, for providing the rabbit limbal mesenchymal stem cells for this study.

REFERENCES

- Abud, A.P.R., J. Zych, T.L Reus, C. Kuligovski, E. de Moraes, B. Dallagiovanna, and A.M. de Aguiar. 2015. The use of human adipose-derived stem cells based cytotoxicity assay for acute toxicity test. *Regulato. Toxicol. Pharmacol*; 73(3): 992-998.
- Ajao, A.A. and A.N. Moteetee. 2017. *Tithonia diversifolia* (Hemsl) A. Gray. (Asteraceae: Heliantheae), an invasive plant of significant ethnopharmacological importance: A review. *South African J. Botany*; 113: 396-403.
- Ammar, L.A., B. Kurniawati, D. Anggorowati, A.P. Cahyaningsih, and A.D. Setyawan. 2021. Ethnobotanical study of the medicinal plant used by local communities in karst area of Pacitan District, East Java, Indonesia. *Int. J. Trop. Drylands*; 5(2): 84-93.
- Amorim, M.H.R., R.M.G da Costa, C. Lopes, and M.M Bastos. 2013. Sesquiterpene lactones: adverse health effects and toxicity mechanisms. *Crit. Rev. Toxicol*; 43(7): 559-579.
- Buranaamnuy, K. 2021. The MTT assay application to measure the viability of spermatozoa: A variety of the assay protocols. *Open Vet. J*; 11(2): 251-269.
- Di Giacomo, C., L. Vanella, V. Sorrenti, R. Santangelo, I. Barbagallo, G. Calabrese, C. Genovese, S. Mastrojeni, S. Ragusa, and R. Acquaviva. 2015. Effects of *Tithonia diversifolia* (Hemsl.) A. Gray extract on adipocyte differentiation of human mesenchymal stem cells. *PLoS One*; 10(4): e0122320.
- Djajanegara, I. and P. Wahyudi. 2009. *Pemakaian sel HeLa dalam uji*



- sitotoksisitas fraksi kloroform dan etanol ekstrak daun *Annona squamosa*. *J. Ilmu Kefarmasian Indonesia*; 7(1): 7-11.
- Fischer, I., C. Milton, and H. Wallace. 2020. Toxicity testing is evolving! *Toxicol. Res*; 9(2): 67-80.
- He, Y., Q. Zhu, M. Chen, Q. Huang, W. Wang, Q. Li, Y. Huang, and Y. Di. 2016. The changing 50% inhibitory concentration (IC50) of cisplatin: a pilot study on the artifacts of the MTT assay and the precise measurement of density-dependent chemoresistance in ovarian cancer. *Oncotarget*. 7(43): 70803-70821.
- Li, W., J. Zhou, and Y. Xu. 2015. Study of the in vitro cytotoxicity testing of medical devices. *Biomed. Rep*; 3(5): 617-620.
- Li, Y., W. Huang, S. Huang, J. Du, and C. Huang. 2012. Screening of anti-cancer agent using zebrafish: comparison with the MTT assay. *Biochem. Biophys. Res. Comm*; 422(1): 85-90.
- Meizarini, A., E. Munadzirroh, and P. Rachmadi. 2005. Sitotoksisitas bahan restorasi cyanoacrylate dengan variasi perbandingan powder dan liquid menggunakan MTT assay. *J. Penelit. Med. Eksak*; 6(1): 16-25.
- Passoni, F.D., R.B. Oliveira, D.A. Chagas-Paula, L. Gobbo-Neto, and F.B. Da Costa. 2013. Repeated-dose toxicological studies of *Tithonia diversifolia* (Hemsl.) A. gray and identification of the toxic compounds. *J. Ethnopharm*; 147(2): 389-394.
- Proboningrat, A., A.N.M. Ansori, A. Fadholly, N. Putri, M.K.J. Kusala, and A.B. Achmad. 2021. First Report on the Cytotoxicity of Pinus merkusii Bark Extract in WiDr, A Human Colon Carcinoma Cell Line. *Res. J. Pharm. Tech*; 14(3): 1685-1688.
- Proboningrat, A, S. Jayanti, A. Fadholly, A.N.M. Ansori, N. Putri, M.K.J. Kusala, S.A. Sudjarwo, F.A. Rantam, and A.B. Achmad. 2021. The Cytotoxicity of Ethanolic Extract of *Allium cepa* L. on Hela Cell Lines. *Res. J. Pharm. Tech*; 14(9): 4969-4972.
- Yuan, H., Q. Ma, L. Ye, and G. Piao. 2016. The Traditional Medicine and Modern Medicine from Natural Products. *Mol*; 21(5): 559.
- Zhang, M., D. Aguilera, C. Das, H. Vasquez, P. Zage, V. Gopalakrishnan, and J. Wolff. 2007. Measuring cytotoxicity: a new perspective on LC50. *Anticancer Res*; 27(1A): 35-38.