The background of the cover is a close-up photograph of a sea urchin. The urchin's body is covered in numerous sharp, white spines that radiate outwards. The central part of the urchin, where the mouthparts are located, shows a pattern of brown and white spots. The surrounding environment appears to be a rocky seabed with some reddish and purple hues, possibly from algae or other marine life.

ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 24 - Number 3 - March 2023

Front cover: Mauritia arabica (Linnaeus, 1758)
(PHOTO: JEAN-MARIE GRADOT)

Published monthly

PRINTED IN INDONESIA

ISSN: 1412-033X

E-ISSN: 2085-4722



9 771412 033962



9 772085 472966

BIODIVERSITAS

Journal of Biological Diversity
Volume 24 – Number 3 – March 2023

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Ajar Nath Yadav** (India), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaee** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Soendjoto** (Indonesia), **Mohamed M.M. Najim** (Srilanka), **Mohib Shah** (Pakistan), **Nurhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Seyed Aliakbar Hedayati** (Iran), **Seyed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Shubhi Avasthi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Wawan Sujarwo** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:

S u t a r n o

EDITORIAL MEMBERS:

English Editors: **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:

biodiversitas.mipa.uns.ac.id; smujo.id/biodiv



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or *Biodiversitas* encourages submission of manuscripts dealing with all aspects of biodiversity, including plants, animals, and microbes at the level of the gene, species, and ecosystem. Ethnobiology papers are also considered.

Article types The journal seeks original full-length: (i) **Research papers**, (ii) **Reviews**, and (iii) **Short communications**. Original research manuscripts are limited to 8,000 words (including tables and figures) or proportional to articles in this publication number. Review articles are also limited to 8,000 words, while Short communications should be less than 2,500 words, except for pre-study.

Submission: The journal only accepts online submissions through the open journal system (<https://smujo.id/biodiv/about/submissions>) or email the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). Please ensure that the manuscript is submitted using the *Biodiversitas* template, which can be found at (<https://biodiversitas.mipa.uns.ac.id/D/guidance.htm>). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, and a paragraph describing the claimed novelty of the findings versus current knowledge. Please also provide a list of five potential reviewers in your cover letter. Submission of a manuscript implies the submitted work has not been published (except as part of a thesis or report, or abstract) and is not being considered for publication elsewhere. When a group writes a manuscript, all authors should read and approve the final version of the submitted manuscript and its revision; and agree on the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis, drafting the manuscript, and correcting the revision. All authors must be responsible for the work's quality, accuracy, and ethics.

Ethics Author(s) must be obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors can reproduce articles as long as they are not used for commercial purposes. For the new invention, authors must manage its patent before publication.

Open Access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance Only articles written in US English are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double-blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. Manuscripts will be rejected if the content does not align with the journal scope, does not meet the standard quality, is in an inappropriate format, or contains complicated grammar, dishonesty (i.e., plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biological significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting typographical errors. The corrected proofs should be returned in 7 days to avoid publication delays. The accepted papers will be published online in chronological order at any time but printed at the end of each month.

A charge Authors are charged USD 320 (IDR 4,500,000). Additional charges may be billed for language improvement, USD 75-150 (IDR 1,000,000-2,000,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by email and sending back the uncorrected proofs. **Manuscript preparation** Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering sizes can be applied in presenting tables and figures (9 pt). Word processing program or additional software can be used; however, it must be PC compatible, use the *Biodiversitas* template, and be Microsoft Word based (.doc or .rtf, not .docx).

Scientific names of species (incl. subspecies, variety, etc.) should be written in italics, except in italicized sentences. Scientific names (genus, species, author) and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. The Genus name can be shortened after the first mention, except where this may generate confusion. Name of the author can be eliminated after the first mention. For example, *Rhizopus oryzae* L. UICC 524 can be written hereinafter as *R. oryzae* UICC 524. Using trivial names should be avoided.

Biochemical and chemical nomenclature should follow the order of the IUPAC - IUB. For DNA sequences, it is better to use Courier New font. Standard chemical abbreviations can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurements** should use IS denominations, and other systems should use equivalent values with the denomination of IS mentioned first. A dot should not follow abbreviations like g, mg, mL, etc. Minus index (m⁻², L⁻¹, h⁻¹) suggested being used, except in things like "per-plant" or "per-plot." **Mathematical equations** can be written down in one column with text;

in that case, they can be written separately. **Numbers** one to ten are written in words, except if it relates to measurement, while values above them are written in number, except in early sentences. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent." Avoid expressing ideas with complicated sentences and verbiage/phrasing, and use efficient and effective sentences.

The **article's title** should be written in a compact, clear, and informative sentence, preferably not more than 20 words. Author name(s) should be completely written. Name and institution address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscripts written by a group, author for correspondence, and address are required. The first page of the manuscript is used for writing the above information.

The **Abstract** should not be more than 200 words. Include between five and eight **Keywords**, using scientific and local names (if any), research themes, and special methods used, and sorted from A to Z. All important **abbreviations** must be defined at their first mention. The running title is about five words. The **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences; however, manuscripts with long discussions should be divided into subtitles. Thorough discussion represents the causal effect that mainly explains why and how the research results occurred and did not only re-express the mentioned results in sentences. A **Conclusion** should be given at the end of the Discussion. Finally, **acknowledgments** are expressed in brief; all sources of institutional, private, and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest must be noted.

Figures and Tables of three pages maximum should be clearly presented. Include a label below each figure and above each table (see example). Colored figures can only be accepted if the manuscript's information can be lost without those images; the chart is preferred to use black and white images. The author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned as sources. Although there is no **Appendix**, all data or data analysis is incorporated into Results and Discussions. For comprehensive/broad data, supplementary information can be provided on the website.

References In the text, give the author names followed by the year of publication and arrange from oldest to newest and from A to Z; in citing an article written by two authors, both of them should be mentioned; however, for three and more authors only the first author is mentioned followed by et al. For example, Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation should be avoided, as shown with the word "cit." Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in alphabetical order. Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). Please include DOI links for journal papers. The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning: a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158. DOI: 10.13057/biodiv/d070213

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in the book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds.). *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.). *Toward Mount Lawu National Park: Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from the internet: Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4:187. www.molecularsystembiology.com

THIS PAGE INTENTIONALLY LEFT BLANK

Home (<https://smujo.id/biodiv/index>)
/ Archives (<https://smujo.id/biodiv/issue/archive>)
/ Vol. 24 No. 3 (2023)



(<https://smujo.id/biodiv/issue/view/371>)

Vol. 24 No. 3 (2023)

Information

For Readers

(<https://smujo.id/biodiv/information/readers>)

For Authors

(<https://smujo.id/biodiv/information/authors>)

For Librarians

(<https://smujo.id/biodiv/information/librarians>)

Journals List

Biodiversitas Journal of
Biological Diversity
(<https://smujo.id/biodiv>)

Nusantara Bioscience
(<https://smujo.id/nb>)

Prosiding Seminar
Nasional Masyarakat
Biodiversitas Indonesia
(<https://smujo.id/psnmbi>)

Asian Journal of
Agriculture
(<https://smujo.id/aja>)

Full Issue

Front Cover (<https://smujo.id/biodiv/issue/view/371/329>)

Articles

The diversity of butterflies (Lepidoptera: Rhopalocera) at Tanahbala Island, North Sumatra, Indonesia
(<https://smujo.id/biodiv/article/view/9212>)

MAIRAWITA, STUDY DUHA, HENNY HERWINA

PDF (<https://smujo.id/biodiv/article/view/9212/6675>)

Habitat use of proboscis monkey (*Nasalis larvatus*) in production forests of West Kalimantan, Indonesia
(<https://smujo.id/biodiv/article/view/13359>)

WARDATUTTHOYYIBAH, SATYAWAN PUDYATMOKO, MUHAMMAD ALI IMRON, SENA ADI SUBRATA

PDF (<https://smujo.id/biodiv/article/view/13359/6679>)

Short Communication: Diversity, biomass, and carbon stock of seagrass community in three coastal waters of Minahasa Peninsula, North Sulawesi, Indonesia

Asian Journal of
Ethnobiology
(<https://smujo.id/aje>)

Asian Journal of Forestry
(<https://smujo.id/ajf>)

Asian Journal of Natural
Product Biochemistry
(<https://smujo.id/jnpb>)

Asian Journal of Tropical
Biotechnology
(<https://smujo.id/bbs>)

International Journal of
Bonorowo Wetlands
(<https://smujo.id/bw>)

Cell Biology and
Development
(<https://smujo.id/cbd>)

Indo-Pacific Journal of
Ocean Life
(<https://smujo.id/ol>)

International Journal of
Tropical Drylands
(<https://smujo.id/td>)

(<https://smujo.id/biodiv/article/view/13666>)

BILLY T. WAGEY, CAROLUS PARUNTU, RIDWAN
LASABUDA, ALEX KAMBEY

PDF (<https://smujo.id/biodiv/article/view/13666/6677>)

Diversity of bat ectoparasites from the caves of
selected Key Biodiversity Areas in Central
Visayas, Philippines

(<https://smujo.id/biodiv/article/view/13263>)

GILBERT A. BEJEC, LILIBETH A. BUCOL, AGUSTIN B.
ANCOG, ALBERT C. PAGENTE, JAZZEL JANE M. PANERIO,
ANNA LEE N. BEJEC, JACINTO D. BELANIZO, DENNY JANE
S. TUASTOMBAN, REIZL P. JOSE

PDF (<https://smujo.id/biodiv/article/view/13263/6671>)

Biosurfactant production of entomopathogenic
Bacillus subtilis BK7.1, as potential biocontrol
bacteria, isolated from Baluran National Park,
East Java, Indonesia

(<https://smujo.id/biodiv/article/view/13658>)

SALAMUN, RIZKY DANANG SUSETYO, NI'MATUZHROH,
FATIMAH, ALMANDO GERALDI, AGUS SUPRIYANTO, TRI
NURHARIYATI, FARAH AISYAH NAFIDIASTRI, NABILATUN
NISA', ENDARTO

PDF (<https://smujo.id/biodiv/article/view/13658/6676>)

Reviewers List

Reviewers

(<https://smujo.id/biodiv/reviewers/index>)

Visitor Statistics

Statistics

(<https://smujo.id/info/stats>)



(<https://info.flagcounter.com/JKAr>)

Characterization of silver nanoparticles (AgNPs) synthesized from Piper ornatum leaf extract and its activity against food borne pathogen Staphylococcus aureus
(<https://smujo.id/biodiv/article/view/13586>)

FIRLI RAHMAH PRIMULA DEWI, VUANGHAO LIM, A'LIYATUR ROSYIDAH, FATIMAH, SRI PUJI ASTUTI WAHYUNINGSIH, UMMI ZUBAIDAH

PDF (<https://smujo.id/biodiv/article/view/13586/6669>)

Spontaneously growing plants on revegetation sites of former coal mine in South Kalimantan Province, Indonesia
(<https://smujo.id/biodiv/article/view/13534>)

MOCHAMAD ARIEF SOENDJOTO, MAULANA KHALID RIEFANI, DIDIK TRIWIBOWO, FAZLUL WAHYUDI, DODIK CHOIRUN, YUDHA PAHING PERDANA

PDF (<https://smujo.id/biodiv/article/view/13534/6670>)

Assessing soil conservation techniques on sloping lands in the humid tropics area of Indonesia in the context of maize cultivation
(<https://smujo.id/biodiv/article/view/13423>)

RISMA NESWATI, SOFYAN ABDULLAH, YUNUS MUSA, NASARUDDIN

PDF (<https://smujo.id/biodiv/article/view/13423/6678>)

Biosurfactant production of entomopathogenic *Bacillus subtilis* BK7.1, as potential biocontrol bacteria, isolated from Baluran National Park, East Java, Indonesia

SALAMUN^{1,2,3,4,*}, RIZKY DANANG SUSETYO^{1,5}, NI'MATUZHROH^{1,2,3}, FATIMAH^{1,2,3}, ALMANDO GERALDI^{1,2,3}, AGUS SUPRIYANTO^{1,2}, TRI NURHARIYATI^{1,2}, FARAH AISYAH NAFIDIASTRI⁶, NABILATUN NISA⁷, ENDARTO⁸

¹Laboratory of Microbiology, Department of Biology, Faculty of Science and Technology, Universitas Airlangga. Kampus C, Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115, East Java, Indonesia. Tel.: +62-315-936501, ✉email: salamun@fst.unair.ac.id

²Research Group for Applied Microbiology and Bioresource Technology, Universitas Airlangga. Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115, East Java, Indonesia

³University of Co-E-Research Center for Bio-Molecule Engineering, Universitas Airlangga. Jl. Mulyorejo Surabaya, Surabaya 60115, East Java, Indonesia

⁴Laboratory of Entomology, Institute of Tropical Diseases, Universitas Airlangga. Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115, East Java, Indonesia

⁵Institute of Life Science, Technology and Engineering, Universitas Airlangga. Kampus C, Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115, Surabaya 60115, East Java, Indonesia

⁶Laboratory of Microbiology, Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Surabaya. Jl. Ketintang, Surabaya 60213, East Java, Indonesia

⁷Laboratory of Molecular Genetics, Department of Biology, Faculty of Science and Technology, Universitas Airlangga. Jl. Dr. Ir. H. Soekarno, Mulyorejo, Surabaya 60115, East Java, Indonesia

⁸Research Development, Baluran National Park. Jl. Raya Banyuwangi-Situbondo Km 35, Situbondo 68374, East Java, Indonesia

Manuscript received: 27 January 2023. Revision accepted: 25 March 2023.

Abstract. Salamun, Susetyo RD, Ni'matuzhroh, Fatimah, Geraldi A, Supriyanto A, Nurhariyati T, Nafidiastri FA, Nisa' N, Endarto. 2023. Biosurfactant production of entomopathogenic *Bacillus subtilis* BK7.1, as potential biocontrol bacteria, isolated from Baluran National Park, East Java, Indonesia. *Biodiversitas* 24: 1785-1792. Biosurfactants as biocontrol agents have received much attention for pest control and disease vectors. The research aimed to identify the species and genetic relationship, hemolytic activity, detect coding genes, and trial production of biosurfactants on various substrates of entomopathogenic *Bacillus* sp. BK7.1 isolated from natural soil in Baluran National Park, East Java, Indonesia. Biosurfactant screening was carried out by testing hemolytic activity, surface tension, and emulsification activities, detecting coding genes of biosurfactant biosynthesis, and testing biosurfactant production in various substrates. The results of the molecular identification by amplifying the 16S rRNA gene using the Polymerase Chain Reaction (PCR) method for *Bacillus* sp. BK7.1 has a genetic similarity of 98.68% with *B. subtilis* subsp. *inaquosorum* strain BGSC 3A28. Screening showed positive hemolytic activity results, reduced surface tension, increased emulsification activities, and the production of biosurfactant in glucose, glycerol, and molasses substrates. The PCR results showed that *Bacillus* sp. BK7.1 had *urfAA* and *urfAD* genes encoding surfactin biosynthesis, giving it the potential to produce bioinsecticide compounds. Based on these studies, the indigenous entomopathogenic *B. subtilis* BK7.1 can be developed as environmentally friendly microbial bioinsecticides for pest control and disease vectors.

Keywords: *Bacillus subtilis* BK7.1, biosurfactant production, crop protection, entomopathogenic, hemolytic activity, *urfAA-urfAD* gene

INTRODUCTION

Controlling insect pests and insect vectors with chemical insecticides is used widely all around the globe (Safni et al. 2018). However, the chemical insecticides have a negative impact on control of disease vector and pest because they cause insect resistance (Şengül et al. 2022). There are a number of biocontrol methods available to resolve these problems. Entomopathogens are natural enemies that can produce toxic metabolites against insect pests and plant pathogens. Biocontrol methods can be used as an alternative in fighting diseases transmitted by vector mosquitoes, plant pathogens, and insect pests. These methods do not cause pollution and are environmentally friendly (Thomas 2017).

Biocontrol agents using *Bacillus* strains are methods that have been widely developed because they are proven to be environmentally friendly (Bergamasco et al. 2013; Syaharuddin et al. 2018; Abdel-Aziz et al. 2020; Qureshi et al. 2021). A group of bacteria, fungi, and yeasts have produced biosurfactants are capable of producing biosurfactants with different surface activities and molecular structures (Santos et al. 2018). Several groups of microbes can synthesize biosurfactants, which can be used to replace non-biodegradable and non-environmental friendly synthetic surfactants (Moro et al. 2018). The biosurfactant produced by *Bacillus* is one of the entomopathogenic mechanisms that have caused the death of insects. Biosurfactants are unique microbial metabolites that appear in biological action against plant pathogens and insect pests.

Biosurfactants have many interesting features including high levels of biodegradability and optimal activity under extreme conditions (Khedher et al. 2017). Following previous studies, *Bacillus subtilis*, *Bacillus amyloliquefaciens*, and *Bacillus velezensis* produce biosurfactant and are efficient biocontrol agents against different targets (Revathi et al. 2013; Nafidiastri et al. 2021). *Bacillus* sp. is able to synthesize lipopeptide biosurfactants, such as surfactin, fengicin, and iturin (Th  atre et al. 2021). Surfactin consists of 7 amino acids bonded to a carboxyl group and a fatty acid hydroxyl group at carbon atoms number 12-16, synthesized by a complex mechanism, catalyzed by Nonribosomal Peptide Synthetase (NRPS) and encoded by the *srfA* operon. Surfactin can suppress plant diseases through strong biosurfactant activity (Cawoy et al. 2014) by inhibiting bacterial growth, lysing cell membranes or destroying them through physicochemical interactions (Deleu et al. 2013), suppressing fungi by promoting colonization of beneficial bacteria (Jia et al. 2015), and triggering systemic resistance (Cawoy et al. 2014). Biosurfactants have been applied in various industrial and petroleum fields (Nwaguma et al. 2016; Pele et al. 2019; Gomaa et al. 2019). Biosurfactants are lower in toxicity, more biodegradable and environmentally friendly, harmless, and work more specifically (De Almeida et al. 2016; Martins and Martins et al. 2018; Gayathiri et al. 2022). Biosurfactants are stable and efficient under unfavorable salinity, pH and temperature conditions often encountered in the petroleum industry (Silva et al. 2014). Biosurfactants can reduce surface and interfacial tension, as well as suitable emulsifiers and dispersing agents and are widely used in the industrial sector (Mulligan et al. 2014).

Perspective studies to find entomopathogenic *Bacillus* spp. are still being carried out to find the safest way to control disease vectors transmitted by mosquitoes. The results of screening tests for potential initial toxicity against *Aedes aegypti* Linnaeus, 1762 larvae have reported that 68 entomopathogenic *Bacillus* sp. which have been isolated from 30 natural soil samples with potential status variations from low to very high. In the affirmation test, there were three isolates coded BK7.1, BK7.2, and BK5.2, with the highest entomopathogenic potential status, larval mortality rates at 48 hours of exposure were 93, 87 and 70%, respectively (Salamun et al. 2020). *Bacillus* sp. BK5.2, after molecular identification, has been identified as *Bacillus thuringiensis* BK5.2 which produces an entomopathogenic cry toxin (Salamun et al. 2021). The identification of *Bacillus* sp. BK7.1 has been carried out through morphological and physiological characterization (Salamun et al. 2020). It is necessary to carry out molecular identification and mechanism of action of *Bacillus* sp. BK7.1 as an entomopathogenic bacteria. Genetic characteristics were used in this study to determine the species and their relationships in the phylogenetic tree, as well as the detection of biosurfactant coding genes and the screening of biosurfactant activities such as hemolytic activity, surface tension, emulsification activity, and production on various substrates.

MATERIALS AND METHODS

Isolation and identification of bacteria

Bacillus sp. BK7.1 was isolated from Baluran National Park soil samples. This isolate was identified conventionally such as macroscopic, microscopic, and physiological characters first and then at molecular level through 16S rRNA (Salamun et al. 2020). This bacteria was maintained aerobically on NB agar plates and was regularly transferred into fresh NB medium slant for short-term storage.

Molecular identification using 16 S rRNA gene

The DNA genome of *Bacillus* sp. BK7.1 was isolated according to the Thermo Scientific GeneJet Genomic DNA Purification Kit, visualized under ultraviolet by electrophoresis, purity and concentration with a Thermo Scientific Multiskan GO Microdroplet Spectrophotometer; purity was calculated by the ratio between the values of 260 nm and 280 nm in the DNA samples (Meena et al. 2020). Amplifying genomic DNA of *Bacillus* sp. BK7.1 utilized 16S rRNA primers (27f and 1492r) was examined by electrophoresis on 1% agarose gel followed by ethidium bromide (EtBr) dye and visualized under ultraviolet light, then purified and sequenced. Amplicon results were then aligned and contigs were developed from the sequences using the BioEdit Sequence Alignment Editor software for Windows. The 16S rRNA nucleotide sequence was aligned with 16S rRNA gene sequences from other microorganisms published in GenBank. Genetic similarity was determined to contig alignment and phylogenetic tree construction using the Program of Mega 7. The phylogenetic tree was designed by inputting FASTAs from BLAST species (Kumar et al. 2016).

Screening biosurfactant activities

Screening of biosurfactants was carried out by three methods, hemolytic activity, surface tension value, and emulsification activity. Hemolytic activity using blood agar media inoculated with *Bacillus* sp. BK7.1 by spots method and incubated for two days at room temperature and zone of inhibition observed around the colony. Surface tension was measured with *du nouy* tensiometer, with 50% tween 20 as a positive control and nutrient broth as a negative control. The decrease in the surface tension value (10 mN/m) indicated the potential to produce biosurfactants. The emulsification activity was measured by inserting a 2 mL supernatant fraction and kerosene in a test tube. This mixture was stirred on vortex mixer for 1 minute, incubated for 24 hours at room temperature, and measured after the emulsion height was stable. The percentage (%) of the emulsion layer height (cm) divided by the total solution height was calculated as the emulsion index value (E24).

Detection *srfAA* and *srfAD* surfactin gene

Amplification of the *srfAA* and *srfAD* surfactin genes of *Bacillus* sp. BK7.1 was carried out by using primers selected according to the literature. Electrophoresis and visualization were performed under UV Transluminator. Forward primer F-5' TCGGGACAGGAAGACATCAT 3'

and reverse primer R-5' CCACTCAAACGGATAATCCTGA 3' for *urfAA* gene (Mora et al. 2020; Kim et al. 2020). Forward primer F-5' ATGAGCCAACCTTCAAATCATTG 3' and reverse primer R-5' TCACGATTGAATGATT GGATGCT 3' for *urfAD* gene. The amplicons were aligned and developed from the sequences by the BioEdit Sequence Alignment Editor for Windows software. The nucleotide sequences are translated into a protein to be formed. The translation of the nucleotide sequence aligned with BLASTp from the other *Bacillus*, which has been published on GenBank.

Biosurfactant production

The biosurfactant production activity begins by providing synthetic mineral water (SMW), by dissolving one by one, 3 g (NH₄)₂SO₄, 10 g NaCl, 0.2 g MgSO₄·7H₂O, 0.01 g CaCl₂, 0.001 g MnSO₄·H₂O, 0.001 g H₃BO₃, 0.001 g ZnSO₄·7H₂O, 0.001 g CuSO₄·5H₂O, 0.005 g CoCl₂·6H₂O, and 0.001 g NaMoO₄·2H₂O into 900 mL distilled water, respectively. The elements phosphate and iron were made separately. The phosphate elements dissolved 5 g of KH₂PO₄ and 2 g of K₂HPO₄ into 50 mL of distilled water, while the iron element dissolved 0.0006 g of FeSO₄·7H₂O into 50 mL of distilled water, respectively. The phosphate and iron elements were sterilized using an autoclave for 15-20 minutes at 121°C with 1 atm.

A 250 mL culture bottle was prepared to be filled with 86.4 mL of SMW and added 2% substrates (glucose, glycerol, molasses) solution, was homogenized and ensured that the pH was 7.0. The culture vial was sterilized by autoclave for 15-20 minutes at 121°C 1 atm. After sterilization, the culture was cooled at room temperature, then 4.8 mL of phosphate and iron elements were added. Then each added 4% (4 mL) of bacterial culture with an absorbance value of 0.5 Optical Density in 650 nm. The culture solution was incubated at room temperature for 96 hrs with an agitation of 130 rpm. Every 24 hrs, bacterial biomass, surface tension value, and emulsification activity against diesel and kerosene were measured until 96 hrs incubation.

RESULTS AND DISCUSSION

Identification of 16S rRNA gene

Purity and concentration of DNA genome of *Bacillus* sp. BK 7.1 obtained a 1.782 and a 31 ng/μL and after being confirmed with agarose gel electrophoresis 1% in Figure 1. *Bacillus* sp. BK7.1 has a size of 1449 bp of the 16S rRNA nucleotide sequence, which similarity to *Bacillus subtilis*

subsp. *inaquosorum* strain BGSC 3A28, homology level of 98.68% (Table 1).

The phylogenetic analysis results where *Bacillus* sp. BK7.1 and some strain of known *Bacillus* are presented in Figure 2. The closest relative of *Bacillus* sp. BK7.1 is a strain of *Bacillus subtilis* strain SBMP4, and this grouping only shows the closeness of the strains based on the similarity of the 16S rRNA sequence, and does not describe the ability to produce biosurfactants, especially surfactin.

Screening of biosurfactant activity

The hemolytic activity of *B. subtilis* BK7.1 can be seen in Figure 3. The surface tension value of the supernatant fraction of *B. subtilis* BK7.1 of 49.17 mN/m can be seen in Table 2. When compared with the surface tension value of the control in the form of distilled water and the control media of Nutrient Broth (NB), the value of the culture supernatant of *B. subtilis* BK7.1 experienced a decrease in surface tension value of 15.21 mN/m from the NB media control and 22.83 mN/m from the distilled water control. The emulsification index value of the supernatant *B. subtilis* BK7.1 of 18.02%, which was left for one hour while after 24 hours the emulsification index value becomes 25.53%, where it was decreased by 21.92% (Figure 4). The emulsification index value indicates the stability of the emulsion and lines that produce values above 50%.







Figure 1. The electrophoresis results of DNA genome (A) and 16S rRNA gene (B) of *Bacillus* sp. BK7.1 on 1% agarose gel. Descriptions: M 100 bp DNA marker, S sample, S1 sample of DNA genome, S2 sample of 16SrRNA gene

Table 1. The species of *Bacillus* sp. BK7.1 based on approach 16S rRNA gene with Basic Local Alignment Search Tools (BLAST) program

| Species | Accession no. | E value | %ID | Query cover (%) |
|---|---------------|---------|-------|-----------------|
| <i>Bacillus subtilis</i> subsp. <i>inaquosorum</i> strain BGSC 3A28 | NR_104873.1 | 0.0 | 98.68 | 99 |
| <i>Bacillus subtilis</i> strain DSM 10 | NR_027552.1 | 0.0 | 98.61 | 99 |
| <i>Bacillus subtilis</i> strain JCM 1465 | NR_113265.1 | 0.0 | 98.61 | 99 |

Table 2. Value of surface tension (mN/m) of supernatant fraction of *Bacillus subtilis* BK7.1 on treatment variation

| Treatment | Surface Tension |
|--|-----------------|
| Control of sterile water | 72 |
| Control of Nutrient Broth (NB) medium, room temperature, pH = 7 | 64.38 |
| Control of Tween 20 at 50% solution | 37.11 |
| Supernatant of <i>Bacillus subtilis</i> BK7.1 (24 hours), room temperature, pH = 8 | 49.17 |
| Δ Surface tension of supernatant of <i>Bacillus subtilis</i> BK7.1 against sterile water | 22.83 |
| Δ Surface tension of supernatant of <i>Bacillus subtilis</i> BK7.1 against NB medium | 15.21 |

**Figure 2.** Analysis of the phylogenetic tree of *Bacillus* sp. BK7.1 based on cladograms of other species and strains, and *Escherichia coli* as an outgroup species**Figure 3.** Screening biosurfactant using hemolytic activity in *Bacillus subtilis* BK7.1 on blood agar plate media. A. Isolate, B. Clear zone around the colony, C. Blood agar plate, R. Replicates**Figure 4.** The emulsification activity of supernatant *Bacillus subtilis* BK7.1 on the kerosene substrate. A. 1 hour of exposure, B. 24 hours of exposure. a. kerosene, b. emulsion, c. isolate

Detection *srfAA* and *srfAD* surfactin gene

The encoding gene of surfactin discovered sizes scale 201 bp, expected as *srfAA* gene, and 723 bp expected as *srfAD* gene (Figure 5). In Table 3 showed that the similarity results, which have a value of 91.04% because there are several differences in amino acids possessed by *B. subtilis* BK 7.1 and other strains of *B. subtilis*. This can be caused by the presence of gene diversity even in the same *B. subtilis* group.

Biosurfactant production



Biosurfactant productions of *B. subtilis* BK7.1 on glucose, glycerol, and molasses substrates can be detected through a bacterial growth curve, surface tension value, and emulsification activity (Figure 6). The growth activity of *B.*

subtilis BK7.1 showed on various substrates in Figure 6a. The isolates had grown well on SMW media with the addition of glucose, glycerol, and molasses as substrates. On glucose substrate with up to 72 hrs incubation, isolates still showed an exponential phase, and 96 hrs incubation entered the stationary phase, as well as on glycerol substrate. On molasses substrate, it still showed an exponential phase until 96 hrs incubation.

The results showed that the emulsification activity of *B. subtilis* BK7.1 on the three substrates tended to increase up to 96 hrs of incubation (Figures 6.B, 6.C), which proved that the isolate produced surfactin. On glucose substrate, the highest emulsification activity occurred at 96 hrs of incubation. Decreased in surface tension values are shown in Figure 6.D.

Table 3. The results of Basic Local Alignment Search Tools (BLAST) analysis of *srfAA* and *srfAD* protein isolates of *Bacillus subtilis* BK 7.1

| Protein | Species | Accession no. | E value | %ID | Query cover (%) |
|---|--------------------------------------|----------------|---------|-------|-----------------|
| Surfactin non-ribosomal peptide synthetase <i>srfAA</i> | <i>Bacillus subtilis inaquosorum</i> | WP_060397903.1 | 9e-34 | 91.04 | 100 |
| Surfactin biosynthesis thioesterase <i>SrfAD</i> | <i>Bacillus subtilis</i> group | WP_075750164.1 | 5e-178 | 99.17 | 99 |

**Figure 5.** The electrophoresis results of *srfAD* (a) and *srfAA* (b) surfactin gene amplification of *Bacillus subtilis* BK7.1. Description: M 100 bp DNA marker, SA sample of *srfAA* surfactin gene 201 bp, SD sample of *srfAD* surfactin gene 729 bp**Figure 6.** Biosurfactant productions of *Bacillus subtilis* BK7.1 on glucose, glycerol, and molasses substrates, incubation period 0-, 24-, 48-, 72-, 96-hours. Descriptions: A. Cells density, B. Emulsification activity (1 hour), C. Emulsification activity (24 hours), D. Surface tension value

Discussions

Conventional identification of *Bacillus* sp. BK7.1 has been carried out. Based on the macroscopic, microscopic, and physiological characteristics of *Bacillus* sp. BK7.1 has similarities with *Bacillus sphaericus* (Salamun et al. 2020). Researchers suggest further research to confirm the species name, by identifying the 16S rRNA gene. The electrophoresis results from 16S rRNA gene amplification of *Bacillus* sp. BK7.1 showed a band over 1500 bp in size (Figure 1). *Bacillus* sp. BK7.1 had a 98.68% similarity to *Bacillus subtilis* subsp. *inaquosorum* strain BGSC 3A28 based on molecular identification. The gene of 16S rRNA could be used for the identification of microorganisms because it is one of the genes with specific characteristics (Pearson 2013). The 16S RNA gene sequencing is was a fast and accurate method for bacterial identification. Bacteria represented the same genus if they have a similarity index above 95% and the same species above 97% (Johnson et al. 2019; Srinivasan et al. 2015). The similarity was less than 100% because there were variations in amino acid sequences that affect the genotypic character but do not affect the phenotypic character (Johnson et al. 2019).

Research has shown that *B. subtilis* strain SBMP4 could control pathogenic fungi such as *Aspergillus* and *Fusarium* in early *Arachis hypogea* plants (Syed et al. 2020). *Bacillus* has adapted to and grown in extreme environmental conditions, forms endospores that are resistant to stress, and secretes various secondary metabolites such as surfactin (Shafi et al. 2017). Another essential characteristic was the abundance of secondary metabolites and moderate dietary requirements with a fast growth rate (Yadav et al. 2016; Mishra and Arora 2018). Biosurfactant lipopeptides from entomopathogenic microbes could act as biocontrol, especially antimicrobials and anti-biofilms (Abdel-Aziz et al. 2020; Qureshi et al. 2021). Surfactin produced by *B. subtilis* was one of the most effective biosurfactants. Surfactin reduced the surface tension of water up to 27 mN/m, with a critical micelle concentration of 0.01 g/L and high emulsification activity and has shown antimicrobial, antiviral, and antitumor activity (Gudina et al. 2013, 2015).

Controlling insects can use biosurfactants introduced as an alternative to synthetic chemicals. Many reports that biosurfactant activity produced by the *Bacillus* strain could kill adult mosquitoes. The hemolytic activity of *B. subtilis* BK7.1 could be seen in Figure 3. The clear zone on the hemolytic activity test by biosurfactants has caused lysis of the red blood cell membrane, and the cells secrete hemoglobin. The hemolytic activity occurred through two different mechanisms, at a high concentration occur, cell membrane lysis, and at low concentrations increase, membrane permeability to solutes and cause osmotic lysis

(Zaragosa et al. 2010). The inhibition zone formed in the observation of hemolytic activity indicates a biosurfactant production process; the larger the lysis diameter of blood agar, the higher the biosurfactant concentration (Singh 2012).

Bacteria could produce biosurfactants if they can reduce surface tension values by ≥ 10 mN/m (Oliveira et al. 2021). The surface and interfacial tension decrease is caused by the presence of hydrophobic and hydrophilic groups in the biosurfactants, where these compounds can accumulate between the liquid phases (Kapadia and Yagnik 2013). The entomopathogenic activity of biosurfactants against *A. aegypti* was caused by surfactin produced by *B. subtilis*. Surfactin triggers the surface tension of the water, causing a lack of oxygen underwater. The concentration of O₂ caused the larval spiracles of *A. aegypti* to open so that it can cause the insect death. In addition, surfactin could be very active against pH, temperature around 25-42°C, and UV stability, making it enjoyable to develop as a larvicidal agent (Guimarães et al. 2019).

The emulsification index value of *B. subtilis* BK7.1 was a low category. Lipopeptides such as surfactin consist of cycloheptapeptides with amino acids attached to fatty acids of a different chain. This chemical structure caused surfactin to be amphiphilic and able to mix in both polar and non-polar solvents, while this amphiphilic structure allows surfactin to form emulsions. The characteristics of surfactin were involved in cell attachment and cause membrane disruption (Chen et al. 2022). The ability of surfactin to bind Ca²⁺ caused a conformational change in the peptide cycle and allows it to be incorporated into the phospholipid bilayer (Khedher et al. 2015, Khedher et al. 2017).

The emulsification activity of *B. subtilis* BK7.1 in 1-hour observation tended to decrease compared to 24 hrs observation. This difference has shown that the emulsion was unstable because the isolate produces biosurfactants which act as active surface molecules only in decreasing surface tension. Based on molecular weight, biosurfactants have been classified into low and high molecular weight biosurfactants. Low molecular weight biosurfactants, including glycolipids, phospholipids, and lipopeptides, were efficient in reducing surface tension. Meanwhile, high molecular weight biosurfactants, such as proteins, lipopolysaccharides, lipoproteins or complex mixtures of these biopolymers, were more effective in stabilizing oil-in-water emulsions as emulsifiers (Uzoigwe et al. 2015). This result was supported by the fact that the biosurfactant produced by *B. subtilis* 21332 has shown high emulsification activity values on glucose substrates up to 55.2% (Zhu et al. 2016). In contrast to the reported that value of the emulsification activity of *B. subtilis* 573 to 27.1%, with the addition of 1% bacterial culture concentration (Pereira et al. 2013), while in this study, the addition of 4%. Differences in the addition of culture affected the activity of biosurfactants produced by bacteria. The higher the concentration of bacterial culture added to the media, the density of bacteria in the substrate also increases and affects the speed of using the available substrate to produce biosurfactants.

Bacillus species had *srfAA* gene, which encodes phosphopantetheinyl transferase and contributes to the nonribosomal biosynthesis of surfactin (Plaza et al. 2015). The nonribosomal peptide synthetase complex was coded by *srfAA* and *srfAD* gene known as surfactin synthetase. The *srfAA* and *srfAD* genes have contributed to the control of surfactin biosynthesis gene expression. The 4-phosphopantetheinyl transferase was an activating enzyme for the *srfA* multienzyme complex. The *srfAA*, *srfAB*, *srfAC*, and *srfAD* genes were involved in the assembly of heptamodular non-ribosomal peptide (NPRS) synthesis in which the modular enzyme contains a typical N-terminal in the CLP-BGCs domain and acylates the first amino acid, glutamine with various 3-OH fatty acids derived from of primary metabolism (Théâtre et al. 2021). The surfactin gene transformed surfactin synthetase into an active form. The production of biosurfactants especially surfactin, that have *Bacillus* influenced by *srfAA* and *srfAD* gene (Plaza et al. 2015). Table 3 showed that the similarity results have a value of 91.04%, because there were several differences in amino acids possessed by *B. subtilis* BK 7.1 and other strains of *B. subtilis*. The presence of gene diversity could cause this even in the same *B. subtilis* group.

The results of this study have also reported that there are differences in the production of biosurfactants. The higher emulsification activity from *B. subtilis* 573 to 48.4% (Pereira et al. 2013), *B. subtilis* 21332 up to 55.2% (Zhu et al. 2016), and *B. subtilis* N3-4P up to 38.3% (Zhu et al. 2016) on mineral salt media containing using different carbon sources than glycerol. The production of biosurfactant by *Bacillus nealsonii* S2M in glycerol substrate has been able to emulsify various hydrocarbons in 55% (Phulpoto et al. 2020).

Biosurfactant production of *B. subtilis* BK7.1 observed through surface tension values is shown in Fig. 6d. Glucose and sucrose substrates have been reported as the best carbon sources for the biosurfactant production process by the *Bacillus* group (Abdel-Mawgoud et al. 2008). *B. subtilis* BK7.1 reduced the surface tension up to 51.47 mN/m at 48 h incubation. *B. subtilis* B30, in 2% glucose substrate has the lowest surface tension value (25.56 mN/m) (Al-Wahaibi et al. 2014). The difference in surface tension reduction was caused by different species and strains of bacteria, as well as the level of their ability to utilize various substrates. Variations in nucleotide sequences between bacteria species affected the formation of biosurfactant biosynthetic genes.

On the glycerol substrate, *B. subtilis* BK7.1 has reduced the surface tension to 53.67 mN/m at 48 h, 42.01 at 96 h, and 54.36 at 72 h incubation, respectively. *B. subtilis* N3-4P has grown better on glycerol substrate than glucose, hexadecane, and diesel. This *B. subtilis* N3-4P decreased the surface tension to 27.8 mN/m on glycerol substrate (Zhu et al. 2016). The same has been reported that the difference in the value of the decrease in surface tension by *B. subtilis* 309, *B. subtilis* 311, and *B. subtilis* 573 on glycerol and glucose substrates, with the value of the decrease in surface tension on glycerol substrates 29.7, 30.1, and 29.9 mN/m, but on glucose substrates 29.2, 29.0, and 29.5 mN/m, respectively (Pereira et al. 2013).

The value of the surface tension of *B. subtilis* BK7.1, on molasses substrate, was 45.91 mN/m. *B. subtilis* SNW3 on molasses substrate was able to reduce the surface tension up to 41 mN/m (Umar et al. 2021), *B. subtilis* ATCC 6633 up to 30.48 mN/m (Kashkouli et al. 2011), and *B. subtilis* RSL-2 up to 24.09 mN/m (Verma et al. 2020). This difference has been due to the influence of various concentrations of molasses substrate. *B. subtilis* BK7.1 used 2% molasses, *B. subtilis* ATCC 6633 used 3% molasses (Kashkouli et al. 2011), and *B. subtilis* RSL-2 used 5% molasses (Verma et al. 2020). In addition, the efficiency of biosurfactant production by *B. subtilis* 3KP with molasses substrate was influenced by the instability of the biosurfactant product. Differences in composition and nutrient content in molasses, suspected related to the processing of sugar from the molasses (Ni'matuzahroh et al. 2017). The difference in sugar content of molasses as the main carbon source for the growth of *B. subtilis* 3KP bacteria has affected the productivity of biosurfactant production (Ni'matuzahroh et al. 2017).

Indigenous entomopathogenic *B. subtilis* BK7.1 isolated from Baluran National Park, East Java, Indonesia, 98.68% similarity to *B. subtilis* subsp. *inaquosorum* strain BGSC 3A28. The results of screening for biosurfactant activity showed positive hemolytic activity, decreased surface tension, and increased emulsification activity. The *urfAA* and *urfAD* genes were detected encoding surfactin, which has the capacity for biosurfactant production on various glucose, glycerol, and molasses substrates. *B. subtilis* BK7.1 produced biosurfactant, the potential to develop for environmentally friendly biocontrol agent for biopesticides in agriculture and disease vector control in public health. Therefore, this research needs to be followed up to detect the chemical components of biosurfactants produced by these bacteria.

ACKNOWLEDGEMENTS

The authors would like to express appreciation for the financial support from the internal funding Featured Faculty Research (*Penelitian Unggulan Fakultas*) 2022-2023 and Featured Basic Research (*Penelitian Dasar Unggulan*) 2023 Universitas Airlangga. The authors grate full to the Dean of the Faculty of Science and Technology (*Fakultas Sains dan Teknologi*) and Head of Research and Community Services Institutions (*Lembaga Penelitian dan Pengabdian Masyarakat*), and Chancellor of Universitas Airlangga, Surabaya, Indonesia.

REFERENCES

- Abdel-Aziz MM, Al-Omar MS, Mohammed HA, Emam TM. 2020. In vitro and ex vivo antibiofilm activity of a lipopeptide biosurfactant produced by the entomopathogenic *Beauveria bassiana* strain against *Microsporum canis*. *Microorganisms* 8 (2): 232. DOI: 10.3390/microorganisms8020232.
- Abdel-Mawgoud AM, Aboulwafa MM, Hassouna NA. 2008. Optimization of surfactin production by *Bacillus subtilis* isolate BS5. *Appl Biochem Biotechnol* 150 (3): 305-325. DOI: 10.1007/s12010-008-8155-x.
- Al-Wahaibi Y, Joshi S, Al-Bahry S, Elshafie A, Al-Bemani A, Shibulal B. 2014. Biosurfactant production by *Bacillus subtilis* B30 and its application in enhancing oil recovery. *Colloids Surf B* 114: 324-333. DOI: 10.1016/j.colsurfb.2013.09.022.
- Bergamasco VB, Mendes DRP, Fernandes OA, Desidério JA, Lemos MVF. 2013. *Bacillus thuringiensis* Cry11a10 and Vip3Aa protein interactions and their toxicity in *Spodoptera* spp. (Lepidoptera). *J Invert Pathol* 112 (2): 152-158. DOI: 10.1016/j.jip.2012.11.011.
- Cawoy H, Mariutto M, Henry G, Fisher C, Vasilyeva N, Thonart P. 2014. Plant defence stimulation by natural isolates of *Bacillus* depends on efficient surfactin production. *Mol Plant Microbe Interact* 27 (2): 87-100. DOI: 10.1094/MPMI-09-13-0262-R.
- Chen X, Lu Y, Shan M, Zhao H, Lu Z, Lu Y. 2022. A mini-review: mechanism of antimicrobial action and application of surfactin. *World J Microbiol Biotechnol* 38 (8): 143. DOI: 10.1007/s11274-022-03323-3.
- De Almeida DG, Soares SRCF, Luna JM, Rufino, RD, Santos VA, Banat IM, Sarubbo LA. 2016. Biosurfactants: Promising molecules for petroleum biotechnology advances. *Front Microbiol* 7: 1718. DOI: 10.3389/fmicb.2016.01718.
- Deleu M, Lorent J, Lins L, Brasseur R, Braun N, El Kirat K, Nylander T, Dufrene YF, Mingeot-Leclercq MP. 2013. Effects of surfactin on membrane models displaying lipid phase separation. *Biochim Biophys Acta Biomembr* 1828: 801-815. DOI: 10.1016/j.bbmem.2012.11.007.
- Gayathiri E, Prakash P, Karmegam N, Varjani S, Awasthi MK, Ravindran B. 2022. Biosurfactants: Potential and eco-friendly material for sustainable agriculture and environmental safety—a review. *Agronomy* 12 (3): 662. DOI: 10.3390/agronomy12030662.
- Gomaa EZ, El-Meihy RM. 2019. Bacterial biosurfactant from *Citrobacter freundii* MG812314.1 as a bioremoval tool of heavy metals from wastewater. *Bull Natl Res Cent* 43 (1): 1-14. DOI: 10.1186/s42269-019-0088-8.
- Gudina EJ, Fernandes EC, Rodrigues AI, Teixeira JA, Rodrigues LR. 2015. Biosurfactant production by *Bacillus subtilis* using corn steep liquor as culture medium. *Front Microb* 6: 59. DOI: 10.1016/j.biortech.2014.11.069.
- Gudiña EJ, Rangarajan V, Sen R, Rodrigues LR. 2013. Potential therapeutic applications of biosurfactants. *Trends Pharmacol Sci* 34 (12): 667-675. DOI: 10.1016/j.tips.2013.10.002.
- Guimarães CR., Pasqualino IP, da Mota FF, de Godoy MG, Seldin L, de Castilho LVA, Freire DMG. 2019. Surfactin from *Bacillus velezensis* H₂O-1: Production and physicochemical characterization for postsalt applications. *J Surfactants Detergents* 22: 451-462. DOI: 10.1002/jsde.12250.
- Jia K, Gao YH, Huang XQ, Guo RJ, Li SD. 2015. Rhizosphere inhibition of cucumber *Fusarium* wilt by different surfactin excreting strains of *Bacillus subtilis*. *Plant Pathol J* 31: 140-151. DOI: 10.5423/ppj.oa.10.2014.0113.
- Johnson JS, Spakowicz DJ, Hong BY, Petersen LM, Demkowicz P, Chen L, Leopold SR, Hanson BM, Agresta HO, Gerstein M, Sodergren E, Weinstock GM. 2019. Evaluation of 16S rRNA gene sequencing for species and strain-level microbiome analysis. *Nat Commun* 10 (1): 5029. DOI: 10.1038/s41467-019-13036-1.
- Kapadia SG, Yagnik BN. 2013. Current Trend and Potential for Microbial Biosurfactants. *Asian J Exp Biol Sci* 4: 1-8.
- Kashkouli RY, Mogharei A, Mousavian S, Vahabzadeh F. 2011. Performance of artificial neural network for predicting fermentation characteristics in biosurfactant production by *Bacillus subtilis* ATCC 6633 using Sugar Cane Molasses. *Intl J Food Eng* 7 (6). DOI: 10.2202/1556-3758.1939.
- Khedher SB, Boukedi H, Dammak M, Feki OK, Boudawara TS, Mesrati LA, Tounsi S. 2017. Combinatorial effect of *Bacillus amyloliquefaciens* AG1 biosurfactant and *Bacillus thuringiensis* Vip3Aa16 toxin on *Spodoptera littoralis* larvae. *J Invert Pathol* 144: 11-17. DOI: 10.1016/j.jip.2017.01.006.
- Khedher SB, Boukedi H, Kilani-Feki O, Chaib K, Laarif A, Mesrati LA, Tounsi S. 2015. *Bacillus amyloliquefaciens* AG1 biosurfactant: histopathological effects and diversity of its putative receptors on *Tuta absoluta* midgut. *J Invertebr Pathol* 132: 42-47. DOI: 10.1016/j.jip.2015.08.010.
- Kim YT, Kim SE, Lee WJ, Fumei Z, Cho MS, Moon JS, Oh HW, Park HY, Kim SU. 2020. Isolation and characterization of a high iturin yielding *Bacillus velezensis* UV mutant with improved antifungal activity. *PLoS ONE* 15 (12): e0234177. DOI: 10.1371/journal.pone.0234177.

- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 33: 1870-1874. DOI: 10.1093/molbev/msw054.
- Martins PC, Martins VG. 2018. Biosurfactant production from industrial wastes with potential remove of insoluble paint. *Intl Biodeterior Biodegrad* 127: 10-16. DOI: 10.1016/j.ibiod.2017.11.005.
- Meena KR, Sharma A, Kanwar SS. 2020. Antitumoral and Antimicrobial Activity of Surfactin Extracted from *Bacillus subtilis* KLP2015. *Intl J Pept Res Ther* 26 (1): 423-433. DOI: 10.1007/s10989-019-09848-w.
- Mishra J, Arora NK. 2018. Secondary metabolites of fluorescent pseudomonads in biocontrol of phytopathogens for sustainable agriculture. *Appl Soil Ecol* 125: 35-45. DOI: 10.1016/j.apsoil.2017.12.004.
- Mora I, Cabrefiga J, Montesinos E. 2015. Cyclic lipopeptide biosynthetic genes and products, and inhibitory activity of plant-associated *Bacillus* against phytopathogenic bacteria. *PLoS ONE* 10 (5): e0127738. DOI: 10.1371/journal.pone.0127738.
- Moro GV, Almeida RTR, Napp AP, Porto C, Pilau EJ, Ludtke DS, Moro AV, Vainstein MH. 2018. Identification and ultra-high-performance-liquid chromatography coupled with high-resolution mass spectrometry characterization of biosurfactants including a new surfactin, isolated from oil-contaminated environments. *Microb Biotechnol* 11: 759-769. DOI: 10.1111/1751-7915.13276.
- Mulligan CN, Sharma SK, Mudhoo A. 2014. *Biosurfactants: Research Trends and Applications*. CRC Press, Florida. DOI: 10.1201/b16383.
- Nafidistri FA, Susetyo RD, Nurhariyati T, Supriyanto A, Geraldi A, Ni'matuzahroh, Fatimah, Salamun. 2021. Biosurfactant activity of indigenous *Bacillus* sp. ES4.3 isolated from endemic breeding sites of dengue hemorrhagic fever vector in Surabaya, East Java, Indonesia. *Biodiversitas* 2 (12): 5375-5381. DOI: 10.13057/biodiv/d221219.
- Ni'matuzahroh, Trikurniadewi N, Pramadita ARA, Pratiwi IA, Salamun, Fatimah, Sumarsh S. 2017. Biodegradation of naphthalene and phenanthren by *Bacillus subtilis* 3KP. *AIP Conf Proc* 1854: 020026. DOI: 10.1063/1.4985417.
- Nwaguma LV, Chikere CB, Okpokwasili GC. 2016. Isolation characterization, and application of biosurfactant by *Klebsiella pneumoniae* strain IVN51 isolated from hydrocarbon-polluted soil in Ogoniland, Nigeria. *Bioresour Bioproc* 3 (1): 1-13. DOI: 10.1186/s40643-016-0118-4.
- Oliveira EM, Sales VH, Andrade MS, Zilli JÉ, Borges WL, de Souza TM. 2021. Isolation and characterization of biosurfactant-producing bacteria from Amapaense Amazon soils. *Intl J Microbiol* 2021: 9959550. DOI: 10.1155/2021/9959550.
- Pearson WR. 2013. An Introduction to Sequence Similarity ("Homology") Searching. *Curr Protoc Bioinform* 42 (1): 1-9. DOI: 10.1002/0471250953.bi0301842.
- Pele MA, Ribeaux DR, Vieira ER, Souza AF, Luna MAC, Rodriguez DM, Andrade RFS, Alviano DS, Alviano CS, Barreto-Berger E, Santiago ALCMA, Campos-Takaki GM. 2019. Conversion of renewable substrates for biosurfactant production by *Rhizopus arrhizus* UCP 1607 and enhancing the removal of diesel oil from marine soil. *Electron J Biotechnol* 38: 40-48. DOI: 10.1016/J.EJBT.2018.12.003.
- Pereira JFB, Gudina EJ, Costa R, Vitorino R, Teixeira JA, Coutinho JAP, Rodrigues LR. 2013. Optimization and characterization of biosurfactant production by *Bacillus subtilis* isolates towards microbial enhanced oil recovery applications. *Fuel* 111: 259-268. DOI: 10.1016/j.fuel.2013.04.040.
- Phulpoto IA, Yu Z, Hu B, Wang Y, Ndayisenga F, Li J, Liang H, Qazi MA. 2020. Production and characterization of surfactin-like biosurfactant produced by novel strain *Bacillus nealsonii* S2MT and its potential for oil contaminated soil remediation. *Microb Cell Fact* 19: 145. DOI: 10.1186/s12934-020-01402-4.
- Plaza G, Chojniak J, Rudnicka K, Paraszkiwicz K, Bernat P. 2015. Detection of biosurfactants in *Bacillus* species: genes and products identification. *J Appl Microb* 119: 1023-1034. DOI: 10.1111/jam.12893.
- Qureshi KA, Bholay AD, Rai PK, Mohammed HA, Khan RA, Azam F, Jaremko M, Emwas AH, Stefanowicz P, Waliczek M, Kijewska M, Ragab EA, Rehan M, Elhassan GO, Anwar MJ, Prajapati DK. 2021. Isolation, characterization, anti-MRSA evaluation, and in-silico multi-target anti-microbial validations of actinomycin X₂ and actinomycin D produced by novel *Streptomyces smyrnaeus* UKAQ_23. *Sci Rep* 11: 14539. DOI: 10.1038/s41598-021-93285-7.
- Revathi K, Chandrasekaran R, Thanigaivel A, Kirubakaran SA, Sathish-Narayanan S, Senthil-Nathan S. 2013. Effects of *Bacillus subtilis* metabolites on larval *Aedes aegypti* L. *Pestic Biochem Physiol* 107: 369-376. DOI: 10.1016/j.pestbp.2013.10.005.
- Safni I, Lisnawita, Lubis K, Tantawi AR, Murthi S. 2018. Isolation and characterization of rhizobacteria for biological control of root-knot nematodes in Indonesia. *J Intl Soc Southeast Asian Agric Sci* 24 (1): 67-81.
- Salamun, Fatimah, Fauzi A, Praduwana SN, Ni'matuzahroh. 2021. Larvicidal toxicity and parasporal inclusion of native *Bacillus thuringiensis* BK5.2 against *Aedes aegypti*. *J Basic Clin Physiol Pharmacol* 32 (4): 379-384. DOI: 10.1515/jbcp-2020-0472.
- Salamun, Ni'matuzahroh, Fatimah, Findawati V, Susetyo RD, Al-Batati N, Nurhariyati T, Supriyanto A. 2020. Prospect of native entomopathogenic *Bacilli* from Baluran National Park as biological control of dengue fever vector. *Ann Biol* 36 (2): 232-237.
- Santos DKF, Rufino RD, Luna JM, Santos VA, Sarubbo LA. 2016. Biosurfactants: Multifunctional biomolecules of the 21st century. *Intl J Mol Sci* 17 (3): 401. DOI: 10.3390/ijms17030401.
- Şengül DMŞ, Canpolat E. 2022. Plant-Based bioinsecticides for mosquito control: impact on insecticide resistance and disease transmission. *Insects* 13 (2): 162. DOI: 10.3390/insects13020162.
- Shafi J, Tian H, Ji M. 2017. *Bacillus* species as versatile weapons for plant pathogens: a review. *Biotechnol Biotechnol Equipment* 31 (3): 446-459. DOI: 10.1080/13102818.2017.1286950.
- Silva RCF, de Almeida DG, Rufino RD, Luna JM, Santos VA, Sarubbo LA. 2014. Application of biosurfactants in the petroleum industry and the remediation of oil spills. *Intl J Mol Sci* 15: 12523-12542. DOI: 10.3390/ijms150712523.
- Singh V. 2012. Biosurfactant - isolation, production, purification, and significance. *Intl J Sci Res Public* 2 (7): 1-4. DOI: 10.4172/2157-7463.1000124.
- Srinivasan R, Karaoz U, Volegova M, MacKichan J, Kato-Maeda M, Miller S, Nadarajan R, Brodie EL, Lynch SV. 2015. Use of 16S rRNA gene for identification of a broad range of clinically relevant bacterial pathogens. *PLOS ONE* 10 (2): e0117617. DOI: 10.1371/journal.pone.0117617.
- Syahrudin A, Marzuki, Sumarheni, Evary YM. 2018. Isolation and identification of *Bacillus alcalophilus* from *Kappaphycosalvarezii* and their antibacterial activity against human pathogens. *Asian J Microbiol Biotechnol Environ Sci* 20 (1): 94-99.
- Syed S, Tollamadugu NVKVP, Lian B. 2020. *Aspergillus* and *Fusarium* control in the early stages of *Arachis hypogaea* (groundnut crop) by plant growth-promoting rhizobacteria (PGPR) consortium. *Microbiol Res* 240: 126562. DOI: 10.1016/j.micres.2020.126562.
- Théâtre A, Cano-Prieto C, Bartolini M, Laurin Y, Deleu M, Niehren J, Fida T, Gerbinet S, Alanjary M, Medema MH, Léonard A, Lins L, Arabolaza A, Gramajo H, Gross H, Jacques P. 2021. The Surfactin-like lipopeptides from *Bacillus* spp.: Natural biodiversity and synthetic biology for a broader application range. *Front Bioeng Biotechnol* 9: 623701. DOI: 10.3389/fbioe.2021.623701.
- Thomas MB. 2017. Biological control of human disease vectors: a perspective on challenges and opportunities. *BioControl* 63 (1): 61-69. DOI: 10.1007/s10526-017-9815-y.
- Umar A, Zafar A, Wali H, Siddique MP, Malik ZA, Ahmed S. 2021. Surfactin-like biosurfactant production and optimization by *Bacillus subtilis* SNW3: Product characterization and its influence on seed development and plant growth. *Res Square* 2021: 1-18. DOI: 10.21203/RS.3.RS-550205%2FV1.
- Uzoigwe C, Burgess JG, Ennis CJ, Rahman PK. 2015. Bioemulsifiers are not biosurfactants and require different screening approaches. *Front Microbiol* 6: 245. DOI: 10.3389/fmicb.2015.00245.
- Verma R, Sharma S, Kundu LM, Pandey LM. 2020. Experimental investigation of molasses as a sole nutrient for the production of an alternative metabolite biosurfactant. *J Water Proc Eng* 38: 101632. DOI: 10.1016/J.JWPE.2020.101632.
- Yadav AN, Sachan SG, Verma P, Saxena AK. 2016. Bioprospecting of plant growth promoting psychrotrophic Bacilli from the cold desert of north western Indian Himalayas. *Indian J Exp Biol* 54 (2): 142-150.
- Zaragosa A, Aranda FJ, Espuny MJ, Teruel JA, Marques A, Manresa A, Ortiz A. 2010. Hemolytic activity of a bacterial trehalose lipid biosurfactant produced by *Rhodococcus* sp.: evidence of colloid-osmotic mechanism. *Langmuir* 26 (11): 8567-8572. DOI: 10.1021/la904637k.
- Zhu Z, Zhang B, Chen B, Cai Q, Lin W. 2016. Biosurfactant production by marine-originated bacteria *Bacillus subtilis* and Its application for crude oil removal. *Water Air Soil Pollutant* 227: 328. DOI: 10.1007/s11270-016-3012-y.



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Biodiversitas

COUNTRY

[Indonesia](#)Universities and research
institutions in Indonesia

Media Ranking in Indonesia

SUBJECT AREA AND CATEGORY

[Agricultural and Biological Sciences](#)
[Animal Science and Zoology](#)
[Plant Science](#)[Biochemistry, Genetics and Molecular
Biology](#)
[Molecular Biology](#)

PUBLISHER

[Biology department, Sebelas Maret
University Surakarta](#)Universitas Negeri Sebelas
Maret in Scimago Institutions
Rankings

H-INDEX

19

PUBLICATION TYPE

[Journals](#)

ISSN

1412033X, 20854722

COVERAGE

2014-2021

INFORMATION

[Homepage](#)[How to publish in this journal](#)grahameagleton@gmail.com

SCOPE

"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of gene, species, and ecosystem.

Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS ?

options

| | | | | |
|--|--|--|--|--|
| <p>1 Diversity CHE</p> <p>41% similarity</p> | <p>2 PeerJ USA</p> <p>39% similarity</p> | <p>3 Biologia (Poland) DEU</p> <p>37% similarity</p> | <p>4 Tropical Life Sciences Research MYS</p> <p>33% similarity</p> | <p>5 Proceedings of the National Academy of Sciences India IND</p> <p>32% similarity</p> |
|--|--|--|--|--|

SJR








Total Documents



Citations per document





Biodiversitas

Q3 Animal Science and Zoology best quartile

SJR 2021 0.29

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç"
```

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization tool.**

