


Indonesian J. Pharm.  
Volume 27 Issue 3 (2016)  
July-September



ISSN : 2338-9427  
Formerly ISSN : 0126-1037

# Indonesian Journal of Pharmacy (Indonesian J. Pharm.)

Accredited by DGHE (DIKTI) No. 58/DIKTI/Rep/2013



STT NO. 1652/SK/DITJEN PPG/SST/1990

Faculty of Pharmacy  
Universitas Gadjah Mada





Home > Archives > **Vol 30 No 4, 2019**

## Vol 30 No 4, 2019

### Table of Contents

#### Articles

**The effect of Phyllanthus niruri L extracts on human leukemic cell proliferation and apoptosis induction**

*Nanda Ayu Puspita, Hasen Alhebshi*

DOI:10.14499/indonesianjpharm30iss4pp241 | Abstract view:0 | PDF download:0

**The Effect of Ursolic Acid from Plantago Lanceolata Leaves on Leukocytes Migration and Chemokines Level**

*Nanang Fakhruddin, Yuvianti Dwi Franyoto, Eny Dwi Astuti, Arief Nurrochmad, Subagus Wahyuono*

DOI:10.14499/indonesianjpharm30iss4pp252 | Abstract view:5 | PDF download:1

**Effects of Peel Extract from Citrus reticulata and Hesperidin, A Citrus Flavonoid, on Macrophage Cell Line**

*Muthi' Ikawati, Inna Armandari, Annisa Khumaira, Yogi Ertanto*

DOI:10.14499/indonesianjpharm30iss4pp260 | Abstract view:0 | PDF download:0

**Potential Deleterious Effects of L-Citrulline Supplementation in Isoproterenol-Induced Myocardial Infarction: Focus on Nitrosative Stress**

*Gregorius Bhaskara Wikanendra, Wawaimuli Arozal, Kusmardi Kusmardi, Vetrizah Juniantito, Andrea Laurentius*

DOI:10.14499/indonesianjpharm30iss4pp269 | Abstract view:0 | PDF download:0

**Response Surface Methodology used in the Optimization of RP-HPLC Condition for Quantitative Analysis of Carmine and Rhodamine B**

*Reyna Nuvitasari, Abdul Rohman, Sudibyo Martono*

DOI:10.14499/indonesianjpharm30iss4pp276 | Abstract view:99 | PDF download:18

**Pharmacoeconomic Rationale of Zinc Supplementation In The Management of Acute Diarrhea In Children With Rotavirus Infection In Indonesia**

*Mohamad Saifudin Hakim, Serhii Soloviov, Yati Soenarto, Iryna Dzyublyk, Maryia Leleka, Viktor Trokhymchuk*

DOI:10.14499/indonesianjpharm30iss4pp285 | Abstract view:0 | PDF download:0

**The capability of Several Population-based Approach Software to Analyze Sparse Drug Plasma Concentration Data after Intra-Venous Bolus Injection**

*Akhmad Kharis Nugroho, Lukman Hakim*

DOI:10.14499/indonesianjpharm30iss4pp293 | Abstract view:90 | PDF download:15

**Quantitative Analysis of Multi-components in Curcuma xanthorrhiza by Single Marker**

*Badrunanto Badrunanto, Wulan Tri Wahyuni, Hera Nirwati, Mohamad Rafi*

DOI:10.14499/indonesianjpharm30iss4pp301 | Abstract view:153 | PDF download:49

**Quality of Life of Patient with Hypertension in Primary Health Care in Bandar Lampung**

*Sudewi Mukaromah Khoirunnisa, Atika Dalili Akhmad*

DOI:10.14499/indonesianjpharm30iss4pp309 | Abstract view:0 | PDF download:0

**In Vitro Study: Effect of Cobalt(II) Chloride Against Dengue Virus Type 1 in Vero Cells**

*Teguh Hari Sucipto, Yovilianda Maulitva Untoro, Harsasi Setyawati, Anisa Maharani, Nova Faridatus Sholihah, Siti Churrotin, Ilham Harlan Amarullah, Soegeng Soegijanto*

DOI:10.14499/indonesianjpharm30iss4pp316 | Abstract view:0 | PDF download:0

Indonesian J Pharm indexed by:



Microsoft Academic

Focus & Scope

Author Guideline

Author Fees

Online Submission

Editorial Board

Peer Reviewer

Subscription Form

Screening for Plagiarism

Visitor Statistics

241

252

260

269

276

285

293

301

309

316

This journal has been published by faculty of pharmacy Universitas Gadjah Mada in collaboration with IAI



CITATION ANALYSIS

► SCOPUS

► GOOGLE SCOLAR

TEMPLATE



TOOLS



NOTIFICATIONS

► View

► Subscribe

USER

Username

Password

Remember me

Login



[000001501646](#) View My Stats

**INFORMATION**

- ▶ For Readers
- ▶ For Authors
- ▶ For Librarians

**JOURNAL CONTENT**

Search

Search Scope

- Browse
- ▶ By Issue
  - ▶ By Author
  - ▶ By Title



[000001501647](#)

**ISSN BARCODE**

**ISSN (PRINT)**



**KEYWORDS**

1,2,4-triazoles, LC MS, forced degradation **Anticancer**  
 Antituberculosis, Benzoylsalicylhydrazide, Methyl Salicylate, InHA, Microwave synthesis, Molecular docking **Apoptosis**  
 Atherosclerosis, curcumin, SNEDDS, nanosuspension, nanoemulsion, solid lipid nano, VCAM-1, IL-6, preventive measure AutoDock Vina Biofilm, Intravenous catheter, colonization, Staphylococcus epidermidis LCMS/MS PRP, diabetic rat testis, caspase-3 mRNA expression, IGF-1, mRNA in expression THPGV-0, THGPV-1, curcumin analog, cytotoxic, anticancer Thymol, Eugenol, polymicrobial biofilm, MBIC50, MBEC50 Turbinaria decurrens Bory. anti-collagenase anti-elastase anti-tyrosinase **antibacterial**  
**antioxidant** fucoxanthin patient medication counseling service vavain derivative  $\alpha$ -glucosidase



[Home](#) > [About the Journal](#) > **Editorial Team**

## Editorial Team

### Editor in Chief

Prof. Dr. Abdul Rohman, Faculty of Pharmacy, Gadjah Mada University 55281 Yogyakarta, Indonesia.

### Editorial Board

Prof. Dr. Shufeng Zhou, Department of Pharmaceutical Sciences, University of South Florida Tampa, United States  
 Prof. Dr. Kazutaka Maeyama, Ehime University, Department of Pharmacology, Japan  
 Prof. Dr. Masashi Kawaichi, Nara Institute of Science and Technology, Division of Gene Function in Animals, Japan  
 Prof. Dr. Gunawan Indrayanto, Universitas Airlangga, Faculty of Pharmacy, Indonesia  
 Prof. Dr. Veeresh P. Veerapur, Sree Siddaganga College of Pharmacy, Pharmaceutical Chemistry Department, India  
 Prof. Dr. Agung Endro Nugroho, Universitas Gadjah Mada, Faculty of Pharmacy, Department of Pharmacology and Clinical Pharmacy, Indonesia  
 Prof. Dr. Lee E. Kirsch, University of Iowa, Division of Pharmaceutics and Translational Therapeutics, United States  
 Prof. Dr. Henk Timmerman, Vrije Universiteit Amsterdam, Division of Medicinal Chemistry, Netherlands  
 Prof. Dr. Jeroen Kool, Vrije Universiteit Amsterdam, Division of BioAnalytical Chemistry, Netherlands  
 Dr. Saikat Kumar Basu, University of Lethbridge, Department of Biological Sciences, Canada  
 Dr. Joseph David Francis Tucci, La Trobe University, School of Pharmacy and Applied Science, Australia  
 Dr. Chuda Chittasupho, Srinakharinwirot University, Department of Pharmaceutical Technology, Thailand  
 Dr. Supang Khonde, University of Phayao, School of Pharmaceutical Sciences, Thailand  
 Dr. Pudjono Pudjono, Universitas Gadjah Mada, Faculty of Pharmacy, Department of Pharmacology and Clinical Pharmacy, Indonesia  
 Dr. Montarat Thavorncharoensap, Faculty of Pharmacy, Department of Pharmacy, Mahidol University, Thailand  
 Dr. Karuna Shanker, Central Institute of Medicinal and Aromatic Plants India, Department of Analytical Chemistry, India  
 Dr. Jun An, Sun Yat-Sen University, Department of Cardiothoracic Surgery, China  
 Dr. Mohammed Emamussalehin Choudhury, Department of Pharmacology, Bangladesh Agriculture University, Bangladesh  
 Dr. Abdul Wahab, Department of Pharmacy, Kohat University of Science and Technology (KUST), Pakistan  
 Dr. Tony Hadibarata, Curtin University Sarawak Malaysia, Department of Environmental Engineering, Malaysia  
 Dr. Shahin Gavanji, Department of Biotechnology, Faculty of Advanced Sciences and Technologies, University of Isfahan, Isfahan, Iran, Islamic Republic of  
 Dr. Kesavanarayanan K S, Department of Pharmacology and Toxicology, College of Pharmacy, University of Hail, Hail, Kingdom of Saudi Arabia., Saudi Arabia

### Indonesian J Pharm indexed by:



000001501648 [View My Stats](#)

[Focus & Scope](#)

[Author Guideline](#)

[Author Fees](#)

[Online Submission](#)

[Editorial Board](#)

[Peer Reviewer](#)

[Subscription Form](#)

[Screening for Plagiarism](#)

[Visitor Statistics](#)

This journal has been published by faculty of pharmacy Universitas Gadjah Mada in collaboration with IAI



[CITATION ANALYSIS](#)

[▶ SCOPUS](#)

[▶ GOOGLE SCOLAR](#)

[TEMPLATE](#)



[TOOLS](#)

[MENDELEY](#)

[grammarly](#)

[EndNote®](#)  
...Bibliographies Made Easy™

[zotero](#)

[NOTIFICATIONS](#)

[▶ View](#)

[▶ Subscribe](#)

[USER](#)

Username

Password

Remember me

**INFORMATION**

- ▶ For Readers
- ▶ For Authors
- ▶ For Librarians

**JOURNAL CONTENT**

Search

Search Scope

Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title

**Visitors**

000001501649


**ISSN BARCODE****ISSN (PRINT)****KEYWORDS**

1,2,4-triazoles, LC MS, forced degradation **Anticancer**  
 Antituberculosis,  
 Benzoylsalicylhydrazide, Methyl Salicylate, InHA, Microwave synthesis,  
 Molecular docking **Apoptosis**  
 Atherosclerosis, curcumin, SNEDDS, nanosuspension, nanoemulsion, solid lipid nano, VCAM-1, IL-6, preventive measure AutoDock Vina Biofilm, Intravenous catheter, colonization, Staphylococcus epidermidis LCMS/MS PRP, diabetic rat testis, caspase-3 mRNA expression, IGF-1, mRNA in expression THPGV-0, THGPV-1, curcumin analog, cytotoxic, anticancer Thymol, Eugenol, polymicrobial biofilm, MBIC50, MBEC50 Turbinaria decurrens Bory. anti-collagenase anti-elastase anti-tyrosinase  
**antibacterial**  
**antioxidant** fucoxanthin patient medication counseling service vavain derivative  $\alpha$ -glucosidase

# BUKTI SCOPUS

## Indonesian Journal of Pharmacy

<b>COUNTRY</b> Indonesia 	<b>SUBJECT AREA AND CATEGORY</b> Health Professions ↳ Pharmacy Medicine ↳ Pharmacology (medical) Pharmacology, Toxicology and Pharmaceutics ↳ Pharmaceutical Science	<b>PUBLISHER</b> Universitas Gadjah Mada - Faculty of Pharmacy	<b>H-INDEX</b> <h1 style="font-size: 2em;">7</h1>
<b>PUBLICATION TYPE</b> Journals	<b>ISSN</b> 23389427, 23389486	<b>COVERAGE</b> 2016-2021	<b>INFORMATION</b> <a href="#">Homepage</a> <a href="#">How to publish in this journal</a> <a href="mailto:mfi@ugm.ac.id">mfi@ugm.ac.id</a>



## *In Vitro* Study: Effect of Cobalt(II) Chloride Against Dengue Virus Type 1 in Vero Cells

Yovilianda Maulitiva Untoro<sup>1</sup>, Teguh Hari Sucipto<sup>2\*</sup>, Harsasi Setyawati<sup>1</sup>, Anisa Maharani<sup>1</sup>, Novia Faridatus Sholihah<sup>1</sup>, Siti Churrotin<sup>2</sup>, Ilham Harlan Amarullah<sup>2</sup> and Soegeng Soegijanto<sup>2</sup>

1. Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Campus C, Surabaya Indonesia 60115

2. Dengue Study Group, Institute of Tropical Disease, Universitas Airlangga, Surabaya Indonesia 60115

### Info Article

Submitted: 05-07-2019

Revised: 23-10-2019

Accepted: 27-11-2019

\*Corresponding author  
Teguh Hari Sucipto

Email:  
Teguhharisucipto  
@staf.unair.ac.id

### ABSTRACT

Dengue virus (DENV) serotypes possess various types from DENV-1 to DENV-4 that are enveloped viruses belong to the genus Flavivirus of the Flaviviridae. Dengue vaccine or antiviral has not yet been clinically approved for humans, even though there have been significant efforts toward this end. Antiviral activity against DENV is needed to develop to be an alternative drug for the DENV virus. Cobalt(II) chloride has been used in the treatment and prevention of diseases of humans since ancient times. This study aimed to investigate the antiviral effects and Cytotoxicity of Cobalt(II) chloride. This compound was further investigated for its inhibitory effect on the replication of DENV-1 in Vero cells. The antiviral activity and cytotoxic was measured by WST-1 assay. The IC<sub>50</sub> value of the Cobalt(II) chloride for DENV-1 was 0.38 µg/ml. The cytotoxicity of Cobalt(II) chloride to Vero cell suggests that the CC<sub>50</sub> value was 2.91 µg/ml. The results of this study demonstrate the anti-dengue serotype 1 inhibitory activity of Cobalt(II) chloride was highly toxic.

**Keywords:** Cobalt(II) chloride, DENV-1, Antiviral activity

### INTRODUCTION

Dengue virus (DENV) is a positive-sense RNA virus replicating in the membranous compartments of cytoplasm. Viral infection induces the dengue virus with significant human viral pathogens transmitted by *Aedes aegypti*. Moreover, about 50 million infections occur each year, and over 2.5 billion people at risk (Gubler, 2002). There are four serotypes of the dengue virus. Four serotypes of dengue virus: DENV-1, DENV-2, DENV-3, and DENV-4) are transmitted by the vector mosquitoes such as *Aedes aegypti* and *Aedes albopictus* (Halstead, 2015). Vaccines have been developed for DENV infection (Woodland, 2015). Recently, WHO immunization group (Strategic Advisory Group of Experts; SAGE) has recommended the use of partially effective dengue vaccine (a live attenuated tetravalent dengue vaccine developed by Sanofi Pasteur; CYD-TDV, named Dengvaxia) that has been licensed and used in 11 countries including Brazil, Mexico, Singapore, Thailand and Indonesia (Aguiar *et al.*, 2016). Nevertheless, control of the dengue virus through the use of vaccination has proved to be elusive

(Burke *et al.*, 2001). A new approach to controlling DENV infection is needed. Antiviral activity of DENV is required to develop an alternative drug for the dengue virus. The reason caused by the alternative drug is not an effective antiviral treatment for DENV, and the patient is not supportively-treated with any specific treatment measures (Zandi *et al.*, 2011).

Cobalt(II) chloride is an inorganic compound of cobalt and chlorine, with the formula CoCl<sub>2</sub>. It is usually supplied as the hexahydrate CoCl<sub>2</sub>·6H<sub>2</sub>O, which is one of the most commonly used cobalt compounds in the laboratory (Greenwood *et al.*, 1997). Cobalt is mainly found in the corrin ring of vitamin B12 (also known as cobalamin) and also supports metabolic and red blood processes (Chang *et al.*, 2010). Cobalt(II) chloride has been used in the treatment and prevention of diseases of humans since ancient times. It has been known for 45 years that small doses of cobalt produce polycythemia in many types of animals, including humans (Devlin *et al.*, 1967). In 1976, Duckham was investigated about the treatment of refractory anemia of chronic renal



failure with cobalt chloride. The research has shown that cobalt is a useful addition to the available therapy in the management of the refractory anemia of chronic renal failure (Duckham *et al.*, 1975). Enteric-coated cobalt chloride 25 mg twice daily is recommended for 12 weeks. The anemia will respond in the majority of patients during this time. It appears that within this period of treatment, the serum cobalt levels will reach a peak and level off. They suggest that either the maintenance dose of cobalt chloride is reduced to 25mg daily or that courses of three months be given intermittently.

In a previous study, Dutta *et al.* investigated the coordination of different ligands to copper (II), and cobalt(III) metal centers enhance the Zika virus and dengue virus loads in both arthropod cells and human keratinocytes (Dutta *et al.*, 2017). These findings suggest that the use of Cu(II) or Co(III) conjugation to organic compounds, in insect repellents and/or food additives could enhance DENV-2/ZIKV loads in human cells and perhaps induce pathogenesis in infected individuals or individuals pre-exposed to such conjugated complexes. However, there are just published data about anti-DENV type 2 activities of cobalt compounds. Therefore, the research aimed to determine the antiviral activity of Cobalt(II) chloride against dengue virus type 1 and the cytotoxicity of Cobalt(II) chloride to Vero cell.

## MATERIALS AND METHODS

The chemical reagents used in this research were the Cobalt(II)-chloride (Sigma-Aldrich, Germany), dimethyl sulfoxide (Merck 99.98%, Germany), Minimum Essential Eagle Medium (Sigma-Aldrich, Germany), Dengue virus serotype 1 Surabaya Isolate (AB915377), Methanol (Merck 99.98%, Germany), Vero cell (African green monkey kidney), Viral ToxGlo Assay (Promega, USA), and Cell Proliferation Reagent WST-1 (Roche Applied Science).

### Antiviral activity assay

Vero Cells with concentration  $1 \times 10^6$  cells/10 mL were seeded into a 96 well plate and incubated plates containing cells at 37°C in a humidified CO<sub>2</sub> incubator for at least 4h (and up to 24h) to facilitate attachment and allow cells to recover from seeding stresses. One hundred microliters of Dengue virus with concentration  $4 \times 10^5$  FFU/mL stock were combined with various concentrations of CoCl<sub>2</sub>. The concentrations of CoCl<sub>2</sub> used in

this study are 50 µg/mL, 25 µg/mL, 12.5 µg/mL, 6.25 µg/mL, 3.13 µg/mL, 1.56 µg/mL (Zandi, 2011). After one hour of exposure to the dengue virus at room temperature, 100 µL of the CoCl<sub>2</sub> treated virus was added to Vero cells in individual wells. Cells were infected for 24h. The unabsorbed virus was removed and replaced with MEM. Twenty-four hours post-infection 100 µL of Viral ToxGlo. 100 µL of ATP Detection Reagent was added to each well of a 96-well plate (25 µL to each well of a 384-well plate) and wait at least 10min before measuring luminescence. Calculate IC<sub>50</sub> values by plotting net RLU values (subtracting the average of blank wells) versus compound concentration. The IC<sub>50</sub> value is the compound concentration that produced a 50% increase in ATP levels compared to virus and no-virus controls.

### In vitro cytotoxicity assay

Cytotoxicity used WST-1 cell proliferation reagent by Roche Applied Science, Mannheim, Germany. The assay is very sensitive: it can detect 1,000 cells/well of a 96-well plate reader. Vero cells ( $1 \times 10^5$  cells/mL), 500 µL of serial dilution compound, and a total of 10 µL of Cell Proliferation Reagent was added to each well of a 96-well plate and incubated under 5% CO<sub>2</sub> at 37°C for 1h. The various concentration of CoCl<sub>2</sub> are 50 µg/mL, 25 µg/mL, 12.5 µg/mL, 6.25 µg/mL, 3.13 µg/mL, 1.56 µg/mL (Zandi, 2011). The plate was read at 450nm (main filter) and 655 nm (reference filter) using an iMark™ Microplate Absorbance Reader.

## RESULT AND DISCUSSION

### Antiviral activity

Antiviral activity is a homogeneous, add-mix-measure method intended that used in research studies to identify cytopathic effect (CPE) induced by a viral infection. Viral ToxGlo assay a more accurate quantitative measure of viral inhibition. Cells were either incubated with MEM, infected with dengue virus, or infected with CoCl<sub>2</sub> treated dengue virus. A significant inhibitory activity to that of the complex Cobalt (II) chloride was displayed against the tested pathogenic DENV-1 in Vero cells. In the inhibitory activity test, we studied the ability of the compound to produce a direct virus-inactivating effect. The basis of this test is the measurement of the ability of living cells based on mitochondrial activity from cell culture. This method is based on the ATP generated by active cells on a live-cell. Microplate reader calculated the live-cell after incubation for 24h.



The percentage inhibition of the development of dengue virus type-1 by the test sample of Co (II) chloride are presented on the figure (Figure 1).




Figure 1. Inactivation of DENV-1 at variation concentrations of Cobalt(II) chloride

The mechanism analogy of inhibition DENV-2 from the previous study, it is also possible that an intact compound can diffuse across the cell membrane or virus capsid, or that Co(II) solute can enter cells through the transport and ion/voltage-gated channels. While Co(II) themselves can interact with oxidative organelles or redox-active protein to induce reactive oxygen species (ROS) in cells, Co(II) produced can also induce ROS by various chemical reactions, and ROS can break DNA strands and alter gene expression. Another possible mechanism is that Co(II) can chelate with biomolecules or dislodge the metal ions in some metalloproteins, leading to dysfunctional proteins and further cell inactivation (Broglie *et al.*, 2015).

The  $IC_{50}$  value was determined from the concentration-response curve (Figure 1); the  $IC_{50}$  value was  $0.38\mu\text{g/mL}$ ,  $R^2$  was 0.9406. Based on the value of the  $IC_{50}$  Cobalt(II)-Chloride was a highly toxic compound. The antiviral activity of Cobalt has been described for distinct viruses. Delehanty *et al.* (2008) investigated the antiviral activity of Co(III)-hexamine to Sindbis virus replication. Co(III)-hexamine, significantly inhibited Sindbis virus replication in baby hamster kidney (BHK) cells in a dose and time-dependent manner. In plaque assays, the incubation of Co(III)-hexamine with Sindbis virus resulted in a dose-dependent decrease in virus replication when measured at both 24 and 48h post-infection. Over the concentration range of 0–5mM Co(III)-hexamine, the  $IC_{50}$  for the inhibition of viral replication was determined to be  $0.10\pm 0.04\text{mM}$  at 48h. Analysis by flow cytometry confirmed that Co(III) hexamine mediated a concomitant dose-dependent increase in BHK cell viability and a decrease in the percentage of Sindbis virus-infected cells

( $IC_{50}=0.13\pm 0.04\text{mM}$ ). These research findings demonstrate for the first time that Co(III) hexamine possesses potent antiviral activity (Delehanty *et al.*, 2008).

In the previous study, the majority of the inorganic ions tested enhanced the anti-herpes simplex virus type 1 (HSV-1) activity,  $EC_{50}$  (effective concentration) of  $ZnCl_2$ ,  $CuCl_2$ , and  $FeCl_3$  were  $1.430\mu\text{M}$ ,  $121\mu\text{M}$ , and  $35.8\mu\text{M}$ . All cations tested showed higher antiviral activity except for zinc. The mechanism of action was not determined during study, but evidence suggests that these chelates may target the extracellular attachment between the virion glycoprotein B and the heparin sulfate proteoglycans on the cells' surface (Langland *et al.*, 2018). The result of Co(III) complexes containing N, O donor ligands activity for HSV-1 has inhibited replication virus with as little as  $5\mu\text{g/mL}$  required for active antiviral (Chang *et al.*, 2010). Many different metal ions are required to support the metabolism of cells. The suggestion has been made that all nucleotidyl transferases are metalloenzymes, and the involvement of zinc ions in both DNA and RNA polymerases is well documented. The attachment of a copper complex to an mRNA that is synthesized late in the infective cycle of viruses would affect the function of the mRNA, and this may account for the antiviral activity of these compounds (Hutchinson, 1985).




Figure 2 Cytotoxicity of Cobalt (II) chloride for Vero Cells at variation concentrations

### Cytotoxicity of Cobalt(II) chloride to Vero Cell

Cobalt (II) chloride was screened for its cytotoxicity against Vero cells at different concentrations to determine the  $CC_{50}$  by WST-1 assay. The  $CC_{50}$  value was found to increase with an increasing concentration of the test compound (Figure 2). The  $CC_{50}$  of Cobalt (II) chloride for Vero cells was  $2.91\mu\text{g/mL}$ . In this study, we have examined the relationship between the

concentration of Vero cells in the culture medium and the cytotoxic potency of Cobalt(II)-chloride.

In the previous study, Sucipto *et al.* investigated the antiviral activity of Copper(II) chloride dehydrate against dengue virus type-2 in Vero cell. It has been revealed that Co(II) is more toxic than Cu(II) with a CC<sub>50</sub> value of 5.03 µg/mL (Sucipto *et al.*, 2017). Copper(II) was non-toxic to human erythrocyte cells even at a concentration of 500 µg/mL (Lv *et al.*, 2006). The effects of metal solutions containing Al(III), Cd(II), Hg(II), and Pb(II) to Vero cells were 6.25, 0.83, 3.7, 2.9, and 3.4 µg/mL respectively (Traoré *et al.*, 1999). The CoCl<sub>2</sub>·6H<sub>2</sub>O being the most effective antiproliferative agent, hence it was further tested against cancer cells and induced maximum cell death in IMR-32 followed by PC-3 and A549 with values of 7.12, 21.91, and 29.81 µg/mL respectively (Mahey *et al.*, 2016). Thus, apoptosis was found to execute in cells by over generation of ROS (reactive oxygen species) and concomitant damage to the mitochondrial membrane.

For Vero cells incubated with polyoxometalate-stabilized gold nanoparticles for 24h was from 20-100 µg/mL, approximately 93-95% was shown to be alive, 0.1-0.3% apoptotic. This data is commensurate with little or no toxicity when compared to the control sample (Gabas *et al.*, 2016). The synthesis of metal silver nanoparticles effect to Vero cell lines was found to be 18.15 µg/mL. The cytotoxic effects had indicated the occurrence of active physicochemical interaction of silver atoms with the functional groups of intracellular proteins, as well as with the nitrogen bases and phosphate groups in DNA (Prasannaraj *et al.*, 2017).

## CONCLUSION

Cobalt(II) chloride was shown to have antiviral activity against DENV-1, a member of the *Flavivirus* genus, *in vitro*. Antiviral activity results suggest that the IC<sub>50</sub> value was 0.38 µg/ml. Cytotoxicity of Cobalt(II) chloride to Vero cell suggests that the CC<sub>50</sub> value was 2.91 µg/ml. Based on the value of the IC<sub>50</sub> and CC<sub>50</sub> value, Cobalt(II) chloride was a highly toxic compound.

## ACKNOWLEDGMENTS

This study was funded by the Student Creativity Program (Program Kreativitas Mahasiswa, PKM) grant from the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia and Mandatory Research Grant (Hibah Riset Mandat) of Universitas

Airlangga 2019. The authors thank the Institute of Tropical Disease Universitas Airlangga for a research internship opportunity in the Dengue Laboratory; Japan Initiative for Global Research Network on Infectious Disease (J-GRID); and for the Center of Excellence (COE) program by the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia.

## REFERENCES

- Aguiar M., Stollenwerk N., Halstead SB., 2016. The impact of the newly licensed dengue vaccine in endemic countries. *PLoS Negl. Trop. Dis.* 10: 1-23.
- Brogie JJ., Alston B., Yang C., Ma L., Adcock AF., Chen W., Yang L., 2015. Antiviral activity of gold/copper sulfide core/shell nanoparticles against human norovirus virus-like particles. *PloS ONE.* 10: 1-14
- Burke DS., Monath TP., 2001. *Flaviviruses.* DM. Knipe, PM. Howley (Eds), *Fields-Virology* (4<sup>th</sup> Edition), Lippincott Williams & Wilkins, Philadelphia. pp. 852-921.
- Chang EL., Simmers C., Knight DA., 2010. Cobalt Complexes as Antiviral and Antibacterial Agents. *Pharmaceuticals.* 3: 1711-1728.
- Delehanty JB., Bongard JE., Thach DC., Knight DA., Hickey TE., Chang EL., 2008. Antiviral properties of cobalt(III)-complexes. *Bioorg. Med. Chem.* 16: 830-837.
- Devlin JG., Brien T., Stephenson N., 1967. Case of Cobalt Poisoning. *Brit. Med. J.* 1: 544-545
- Duckham JM., Lee HA., 1976. The Treatment of Refractory Anaemia of Chronic Renal Failure with Cobalt Chloride. *Q. J. Med.* 178: 277-294.
- Dutta S., Celestine MJ., Khanal S., Huddleston A., Simms C., Arca JF., Mitra A., Heller L., Kraj PJ., Ledizet M., Anderson JF., Neelakanta G., Holder AA., Sultana H., 2018. Coordination of different ligands to copper(II) and cobalt(III) metal centers enhances Zika virus and dengue virus loads in both arthropod cells and human keratinocytes. *Biochim. Biophys. Acta Gen. Subj.* 1862: 40-50
- Gabas IM., Stepien G., Moros M., Mitchell SG., de la Fuente JM., 2016. *In vitro* cell cytotoxicity profile and morphological response to polyoxometalate-stabilised gold nanoparticles. *New J. Chem.* 40: 1039-1047
- Greenwood NN., and A. Earnshaw. 1997. *Cobalt, Rhodium and Iridium, in Chemistry of the Elements.* Butterworth-Heinemann, Oxford, England. pp. 1113-1143.

- Gubler DJ., 2002. Epidemic dengue/dengue hemorrhagic fever as a public health, social and economic problem in the 21st century. *Trends Microbiol.* 10: 100–103.
- Halstead SB., 2008. Dengue Virus-mosquito Interactions. *Annu. Rev. Entomol.* 53: 273–291.
- Hutchinson DW., 1985. Metal chelators as potential antiviral agents. *Antiviral Res.* 5: 193-205
- Langland J., Jacobs B., Wagner C., Ruiz G., Cahill TM., 2018. Antiviral activity of metal chelates of caffeic acid and similar compounds towards herpes simplex, VSV-Ebola pseudotyped and vaccinia viruses. *Antiviral Res.* 160: 143-150
- Lv J., Liu T., Cai S., Wang X., Liu L., Wang Y., 2006. Synthesis, structure and biological activity of cobalt(II) and copper(II) complexes of valine-derived schiff bases. *J. Inorg. Biochem.* 100: 1888–1896.
- Mahey S., Kumar R., Arora R., Mahajan J., Arora S., Bhardwaj R., Thukal AK., 2016. Effect of cobalt(II) chloride hexahydrate on some human cancer cell lines. *Springer Plus.* 5:1-14
- Prasannaraj G., Venkatachalam P., 2017. Green engineering of biomolecule-coated metallic silver nanoparticles and their potential cytotoxic activity against cancer cell lines. *Adv. Nat. Sci.: Nanosci. Nanotechnol.* 8: 1-11
- Sucipto TH., Churrotin S., Setyawati H., Kotaki T., Martak F., Soegijanto S., 2017. Antiviral Activity of Copper(I)Chloride Dihydrate Against Dengue Virus type-2 In Vero Cell. *Indonesian J. Trop. Infect. Dis.* 6: 84-87
- Traoré A., Bonini M., Dano SD., Creppy EE., 1999. Synergistic effects of some metals contaminating mussels on the cytotoxicity of the marine toxin okadaic acid. *Arch. Toxicol.* 73: 289-295
- Woodland DL., 2015. Vaccines Against Dengue Virus. *Viral Immunol.* 28: 75.
- Zandi K., Teoh BT., Sam SS., Wong PF., Mustafa MR., Abubakar S., 2011. Antiviral Activity of Four Types of Bioflavonoid Against Dengue Virus Type-2. *Virol. J.* 8: 1–11.