

A new modified logistic growth model for empirical use

by Windarto Windarto

Submission date: 22-Feb-2023 10:32AM (UTC+0800)

Submission ID: 2020093382

File name: 2018_A_new_modified_logistic_growth_model_for_empirical_use.pdf (256.97K)

Word count: 4658

Character count: 22564

A new modified logistic growth model for empirical use

12 Windarto*, Eridani and Utami Dyah Purwati

Department of Mathematics, Faculty of Science and Technology,
Universitas Airlangga, Indonesia
*Email: windarto@fst.unair.ac.id

Abstract

Richards model, Gompertz model, and logistic model are widely used to describe growth model of a population. The Richards growth model is a modification of the logistic growth model. In this paper, we present a new modified logistic growth model. The proposed model was derived from a modification of the classical logistic differential equation. From the solution of the differential equation, we present a new mathematical growth model so called a WEP-modified logistic growth model for describing growth function of a living organism. We also extend the proposed model into couple WEP-modified logistic growth model. We further simulated and verified the proposed model by using chicken weight data cited from the literature. It was found that the proposed model gave more accurate predicted results compared to Richard, Gompertz, and logistic model. Therefore the proposed model could be used as an alternative model to describe individual growth.

Keywords: mathematical model, growth function, modified logistic growth.

1. INTRODUCTION

Optimum food utilization strategy is one of the important efforts to increase meat production of livestock. The dynamics of livestock growth over time is needed to obtain an optimal growth strategy of animal feeds. Mathematical models of the growth curve could be used to determine the selection of suitable feeding materials for livestock development [1]. In addition, the growth curve could also be used to determine the age of livestock slaughter to be optimal. Moreover, the growth curve model could be used as a parameter in pre-harvest methods in large livestock such as cattle, buffalo, goats, and sheep. The mathematical model of livestock growth could also be used to analyze the efficiency of livestock production over the lifetime (lifetime production efficiency) [2].

The growth process of livestock, including poultry, could be measured from mass (weight) profile of the livestock versus time [3], [4]. Livestock and poultry growth generally follows a sigmoidal pattern. Poultry growth usually starts with an accelerating growth phase from hatching. Then, poultry attains the maximum growth rate at a certain time (the inflection time). After that, poultry growth is decelerating. At final phase, poultry weight generally tends to a limiting value (asymptote) mature weight [1], [5].

Many nonlinear growth curves have proposed to describe and fit the sigmoid relationship between poultry weight and time. Logistic model, Gompertz model, and Richards model are commonly for describing a relationship between poultry weight and time [1], [3], [5]. Richards and Gompertz models gave good descriptions of weight growth in many animal species including chicken, cattle, elks, ostrich, turkey, and emus. Gompertz growth model has applied as the growth model for chicken. Gompertz model fitted many actual data and its parameters had biological interpretations [6], [7], [8]. Moreover, the Gompertz model has good fitting for weight information whose inflection points occur, when approximately 35 - 40% of growth have been achieved [5].

Accurate and straightforward growth models are useful for describing life individual growth. In this paper, we present a new mathematical model for the growth function of a living organism. The model was derived from the modified logistic differential equation. Then, the model was implemented to describe body weight growth of chicken (rooster and hen), where the growth data cite from literature. The accuracy of predicted results from the model was compared to the standard logistic model, Gompertz model and Richards model.

This paper is organized as follows. Section 2 presents some modified logistic growth models. The proposed model and its main property is discussed in section 3. Implementations of the proposed model, logistic model,

16
Gompertz model and Richard model on chicken (rooster and hen) data cited from the literature are presented in Section 4. Conclusions are written in the last section.

2. MODIFIED LOGISTIC GROWTH MODEL

The first mathematical model describing population growth is the Malthus model or exponential model [9]. Let $y(t)$ is population size at time t . In the exponential model, the growth rate $\frac{dy}{dt}$ is assumed proportional the size of existing population $y(t)$. The exponential model could be represented by the following differential equation

$$y(t) = Y_0 \exp(rt). \quad (1)$$

20 Here r is the proportional growth rate parameter. The exponential growth model in Eq. (1) is rarely used to describe population growth, since it produces an unbounded population growth.

The exponential growth model was improved by logistic growth model. In the logistic model, a population grows until it attains a maximum capacity [9]. The logistic model is based on the assumption that the growth rate $\frac{dy}{dt}$ is proportional to the existing population and the remaining resources available to the existing population. The logistic growth model be expressed as

$$y(t) = \frac{K}{1 + \exp(-rt) \left(\frac{K}{Y_0} - 1 \right)} = \frac{K}{1 + \exp[-r(t - t_{\text{inf}})]}, \quad (2)$$

where

$$t_{\text{inf}} = \frac{1}{r} \ln \left(\frac{K}{Y_0} - 1 \right). \quad (3)$$

When $y(t)$ represents body weight of a livestock at time t , then parameter K in Eq. (2) could be considered as the mature weight (the maximum weight that could be attained by livestock). Here t_{inf} is the inflection time (the optimal time of a population growth).

The logistic growth model has various modifications. One of the modified version is the shifted logistic function. The first version of the shifted logistic function could be presented in the following form [10]

$$y(t) = K \left(\frac{1}{1 + \exp(-r(t - t_{\text{inf}}))} - \frac{1}{1 + \exp(rt_{\text{inf}})} \right). \quad (4)$$

The second version and the third version of the shifted logistic function could be expressed as [11]

$$y(t) = \frac{K}{1 + \exp(-r(t - t_{\text{inf}}))} + L \quad (5)$$

and

$$y(t) = \frac{K + Mt}{1 + \exp(-r(t - t_{\text{inf}}))} + L \quad (6)$$

respectively. Here, L and M are additional parameters.

Modification of the logistic growth model also occurred in the differential equations model. The logistic differential equation has been modified into von Bertalanffy, Richards, Gompertz, Blumberg, Turner et al. and Tsoularis differential equations. The von Bertalanffy differential equation has the following form [12], [13]

$$\frac{dy}{dt} = ry^{\frac{2}{3}} \left(1 - \left(\frac{y}{K} \right)^{\frac{1}{3}} \right), \quad y(0) = Y_0. \quad (7)$$

Richards (1959) proposed a modified logistic differential equation so-called Richards differential equation. The Richards differential equation has the following form [13], [14]

$$\frac{dy}{dt} = ry \left(1 - \left(\frac{y}{K} \right)^{\beta} \right), \quad y(0) = Y_0. \quad (8)$$

Gompertz differential equation is a limiting case of a modified logistic differential equation. The Gompertz differential equation is derived from

$$\frac{dy}{dt} = \lim_{\beta \rightarrow 0} \frac{ry \left(1 - \left(\frac{y}{K}\right)^\beta\right)}{\beta} = ry \ln\left(\frac{K}{y}\right), \quad y(0) = Y_0. \quad (9)$$

Blumberg (1968) also introduced a modification of logistic differential equation so called the hyper logistic function, accordingly [13], [15]

$$\frac{dy}{dt} = ry^\alpha \left(1 - \frac{y}{K}\right)^\gamma, \quad y(0) = Y_0. \quad (10)$$

Turner et al. (1976) proposed a modified logistic differential equation which they named the generic growth function. The modification has the following form [13], [16]

$$\frac{dy}{dt} = ry^{1+\beta(1-\gamma)} \left(1 - \left(\frac{y}{K}\right)^\beta\right)^\gamma, \quad y(0) = Y_0. \quad (11)$$

Tsoularis (2001) proposed a more general modification of logistic differential equation. The Tsoularis differential equation has the form [13]

$$\frac{dy}{dt} = ry^\alpha \left(1 - \left(\frac{y}{K}\right)^\beta\right)^\gamma, \quad y(0) = Y_0. \quad (12)$$

In the next section, we propose another version of a modified logistic differential equation.

3. THE PROPOSED MODEL

The logistic growth model and the modified logistic growth model presented in the previous section could be represented in the Kolmogorov form

$$\frac{dy}{dt} = yP(y) \quad (13)$$

for some continuous function P . For classical (standard) logistic differential equation, the function P is $P(y) = r\left(1 - \frac{y}{K}\right)$. In the logistic growth model, it is assumed that the growth rate of a population is proportional to the population number at the current time. Here, we modify the model in Eq. (13) in more general form, namely

$$\frac{dy}{dt} = F(y) \quad (14)$$

for some continuous function F . A simple growth model satisfies Eq. (14) but it does not satisfy the Kolmogorov form in Eq. (13), is the monomolecular model. The monomolecular model satisfy the following differential equation [17]

$$\frac{dy}{dt} = q - sy, \quad y(0) = Y_0. \quad (15)$$

Here, q could be considered as constant growth rate while s could be considered as the death rate of a population.

In this section, we propose a generalized model of the monomolecular model and the standard logistic growth model. We extend the monomolecular model and the logistic differential equation model into the following differential equation.

$$\frac{dy}{dt} = (q + ry) \left(1 - \frac{y}{K}\right), \quad y > 0 \quad (16)$$

and the initial condition $y(0) = Y_0 > 0$. Note that region of biological interest of the model in Eq. (16) is $\mathbf{R}_+ := \{x \in \mathbf{R} : x > 0\}$, since a life organism could not grow from nothing. Here, q and r could be considered as constant growth rate and proportional growth rate respectively.

The modified logistic growth model in Eq. (16) has one equilibrium, namely $y = K$. Global stability of the equilibrium is presented in the following theorem.

Theorem 3.1. *The equilibrium $y = K$ is globally asymptotically stable.*

Proof: We define a Lyapunov function $V: \mathbf{R} \rightarrow \mathbf{R}$ by $V(y) = (y - K)^2$. The function V is a $C^\infty(\mathbf{R})$ function. In addition, the equilibrium $y = K$ is the global minimum of V . Moreover, V is a definite positive function around the equilibrium where every $y \in \mathbf{R} \setminus \{K\}, V(y) > 0$. The time derivative of V computed along solutions of the mathematical model in Eq. (16) is given by the expression

$$\frac{dV}{dt} = \frac{-2}{K} (q + ry)(y - K)^2.$$

Since all parameters in the model are positive and the variable y is positive, it follows that $\frac{dV}{dt} \leq 0$ for $y > 0$. In addition $\frac{dV}{dt} = 0$ if and only if $y = K$. Therefore the greatest compact invariant set in $\{y \in \mathbf{R}_+ : \frac{dV}{dt} = 0\}$ is the singleton $\{K\}$. By LaSalle's invariance principle [18], the equilibrium $y = K$ is globally asymptotically stable in \mathbf{R}_+ .

The population weight at the inflection time (t_{inf}) could be determined as follows. By differentiating both sides of Eq. (16) and setting $\frac{d^2y}{dt^2}(t_{\text{inf}}) = 0$, we find

$$y(t_{\text{inf}}) = \frac{K}{2} - \frac{q}{2r}. \tag{17}$$

Hence, the population weight at the inflection time for this model is smaller than the values obtained from the logistic growth model. Exact values of the inflection time could be obtained whenever the analytical solution of the model in Eq. (16) could be found.

The differential equation in Eq. (16) could be written as

$$\left(\frac{r}{q + ry} + \frac{1}{K - y} \right) dy = \left(\frac{q}{K} + r \right) dt.$$

By integrating the left side with respect to y and the right side with respect to t gives

$$\ln \left(\frac{q + ry}{K - y} \right) = \left(\frac{q}{K} + r \right) t + c_0 \tag{18}$$

for some constant c_0 . The mathematical expression in the Eq. (18) could be written as

$$\frac{q + ry}{K - y} = c_1 \exp \left(\left(\frac{q}{K} + r \right) t \right), c_1 = \exp(c_0). \tag{19}$$

By solving Eq. (19) for y , it could be obtained explicit solution of the modified logistic differential equation as

$$y(t) = \frac{c_1 K \exp \left(\frac{qt}{K} + rt \right) - q}{r + c_1 \exp \left(\frac{qt}{K} + rt \right)}. \tag{20}$$

By substituting the initial condition $y(0) = Y_0$, then $c_1 = \frac{qY_0 + q}{K - Y_0}$. Hence, the explicit solution in Eq. (20) could be written as

$$y(t) = \frac{K - q \left(\frac{K - Y_0}{rY_0 + q} \right) \exp \left(\frac{-qt}{K} - rt \right)}{1 + r \left(\frac{K - Y_0}{rY_0 + q} \right) \exp \left(\frac{-qt}{K} - rt \right)}. \tag{21}$$

By defining the following parameters

$$\alpha = \frac{q}{K} + r, \quad A = K - q \left(\frac{K - Y_0}{rY_0 + q} \right), \quad B = r \left(\frac{K - Y_0}{rY_0 + q} \right), \tag{22}$$

then the modified logistic growth model in Eq. (21) could be written as

$$y(t) = \frac{K - (K - A) \exp(-\alpha t)}{1 + B \exp(-\alpha t)}. \quad (23)$$

Here α, A, B, K are positive parameters and $A \leq K$. The parameter α is effective growth rate, K is the maximum capacity (mature weight), while the parameter A, B are corresponding to initial weight and inflection time. The inflection time (t_{inf}) of the model in Eq. (23) is

$$t_{inf} = \frac{\ln B}{\alpha} = \frac{K}{q + rK} \ln \left(r \left(\frac{K - Y_0}{rY_0 + q} \right) \right). \quad (24)$$

The inflection time in (24) could be determined by evaluating the second derivative of y in Eq. (23) and setting $\frac{d^2y}{dt^2}(t_{inf}) = 0$. If the constant growth rate parameter (q) is zero, then the inflection time in Eq. (24) could be simplified into Eq. (3). From Eq. (24), the modified logistic growth model in Eq. (23) could be presented in the following form

$$y(t) = \frac{K - (K - A) \exp(-\alpha t)}{1 + \exp(-\alpha(t - t_{inf}))}. \quad (25)$$

Since there are some well-known modified logistic growth model, then the presented growth model presented in Eq. (29) could be called by a WEP-modified logistic growth model. Here WEP comes from Windarto-Eridani-Purwati.

4. EXTENSION OF THE PROPOSED MODEL

It is well known that the length and weight of fish species will grow until they attain some maximum values. By applying the presented model in the previous section, the dynamics of fish weight and fish length could be modeled by following differential equations

$$\frac{dW}{dt} = (q_w + r_w W) \left(1 - \frac{W}{K_w} \right), \quad W(0) = w_0, \quad (26)$$

and

$$\frac{dL}{dt} = (q_l + r_l L) \left(1 - \frac{L}{K_l} \right), \quad L(0) = l_0, \quad (27)$$

respectively. Here, $W(t)$ and $L(t)$ are fish weight and fish length at time t respectively. In Eq. (26)-(27), q_w, q_l are constant growth rate of fish weight and fish length, while r_w, r_l are proportional growth rate of fish weight and fish length respectively. By applying the analytical solution of the previous section, we found dynamic of fish weight and fish length could be described by

$$W(t) = \frac{K_w - q_w \left(\frac{K_w - w_0}{r_w w_0 + q_w} \right) \exp \left(\frac{-q_w t}{K_w} - r_w t \right)}{1 + r_w \left(\frac{K_w - w_0}{r_w w_0 + q_w} \right) \exp \left(\frac{-q_w t}{K_w} - r_w t \right)} \quad (28)$$

and

$$L(t) = \frac{K_l - q_l \left(\frac{K_l - l_0}{r_l l_0 + q_l} \right) \exp \left(\frac{-q_l t}{K_l} - r_l t \right)}{1 + r_l \left(\frac{K_l - l_0}{r_l l_0 + q_l} \right) \exp \left(\frac{-q_l t}{K_l} - r_l t \right)} \quad (29)$$

respectively.

It is also well known that there is length-weight relationship (LWR) of fish species. A mathematical equation was used to show relationships between the average weight of fish at a given length [19], [20]. The length-weight relationship is given by

$$W(t) = aL(t)^b. \quad (30)$$

Here a and b are empirical parameters. Typically, the b parameters range from 2 to 4. Fish can attain either isometric or allometric growth. Isometric growth indicates that both fish length and fish weight are increasing at the same rate [20]. Here, we estimated parameters empirical parameters a and b for the threadfin bream (*Nemipterus marginatus*) from the South China Sea, where the data cited from literature [21]. The length-weight relationship data is shown in Table I.

TABLE I: Length-weight relationship data for the threadfin bream (*Nemipterus marginatus*)

L (cm)	W (grams)	L (cm)	W (grams)
8.1	6.3	16.6	65.6
9.1	9.60	17.7	69.4
10.2	11.6	18.7	76.4
11.9	18.5	19	82.5
12.2	26.2	20.6	106.6
13.8	36.1	21.9	119.8
14.8	40.1	22.9	169.2
15.7	47.3	23.5	173.3

By taking natural logarithm of eq. (30), the relationship could be converted into

$$\ln W = \ln a + b \ln L. \quad (31)$$

Then by using linear regression method, we find the parameter values of $\ln a = -4.53786$ and $b = 3.05735$. Hence the parameter value of a is 0.01070. Since the parameter value of b does not significantly differ from 3, then the species attain isometric growth [20]. In order to estimate parameters in Eq. (28) and (29), we need fish weight and fish length data over time. In the next section, we apply the proposed model (WEP-modified logistic growth model) to some secondary data cited from the literature.

5. APPLICATION OF THE PROPOSED MODEL

In this section, the proposed model is implemented to describe chicken body weight (rooster and hen) growth, where the data are cited from literature [3], [22]. Rooster (x) and hen (y) body weight at different age (t) are presented in Table 2. In addition, accuracy result of the proposed model will be compared to the logistic model, Gompertz model, and Richards model. The logistic model was presented in Eq. (2), while Richards and Gompertz differential equations were presented in Eq. (8) and (9) respectively. Analytical solution of the Richards differential equation in Eq. (8) was given by

$$y_R(t) = \frac{K}{\left[1 + \beta \exp(-r\beta(t - t_{\text{inf}}))\right]^{\frac{1}{\beta}}}, \quad (32)$$

where the inflection time $t_{\text{inf}} = \frac{1}{r\beta} \ln\left(\frac{\left(\frac{K}{y_0}\right)^\beta - 1}{\beta}\right)$. By defining $m = \beta + 1$, $r^* = r\beta$, then the Richards growth model in Eq. (32) could be expressed as

$$y_R = K \left[1 - (1 - m) \exp(-r^*(t - t_{\text{inf}}))\right]^{\frac{1}{1-m}}. \quad (33)$$

Exact solution of the Gompertz differential equation in Eq. (9) was given by

$$y_G(t) = \frac{K}{\exp\left(\exp(-r(t - t_{\text{inf}}))\right)} \quad (34)$$

where $t_{inf} = \frac{1}{r} \ln \left(\ln \left(\frac{K}{y_0} \right) \right)$. Some authors used the following Gompertz-Laid growth model [3]

$$y_G(t) = W_0 \exp \left(\exp(rt_{inf}) \left(1 - \exp(-rt) \right) \right). \quad (35)$$

Here, W_0 is initial chicken weight in the Gompertz model and m is the shape parameter in Richards model. For $m = 2$, then the Richards model could be simplified into logistic model. For m tends to one, then the Richards model could be simplified into the Gompertz model.

TABLE II: Means of rooster and hen chicken weight data

t (days)	x (grams)	y(grams)	t (days)	x (grams)	y (grams)
0	37	36.68	42	519.72	436.51
3	41.74	40.8	45	577.27	480.31
6	59.19	57.33	48	633.59	522.91
9	79.94	77.24	51	667.18	547.23
12	102.96	97.96	54	717.17	583.56
15	132.13	121.92	57	786.35	631.77
18	170.18	155.08	71	1069.28	832.57
21	206.56	184.24	85	1326.49	1009.48
24	250.71	218.37	99	1589.71	1183.8
27	285.27	247.12	113	1859.26	1440.18
30	324.92	279.58	127	2015.44	1561.89
33	372.83	319.55	141	2142.31	1619.34
36	417.41	355.13	155	2220.54	1680.29
39	469.13	396.32	170	2262.63	1717.78

There are four parameters in the model should be estimated, namely parameter α (effective growth rate), K (maximum weight/ mature weight of chicken), the inflection time t_{inf} and parameter A (correspond to the initial chicken weight). Since growth function of the model is explicitly presented in the Eq. (25), then nonlinear regression procedures could be applied to estimate the parameters.

The parameters α , K , t_{inf} and A are estimated such that the normalized residual sum of squares (NRSS)

$$NRSS = \sum_i \frac{(z_i - \hat{z}_i)^2}{(z_i - \bar{z})^2}, z = x \vee z = y \quad (36)$$

is minimum. In Eq. (36), \bar{z} is the average of z and \hat{z}_i is chicken weight at time i predicted from the model. The normalized residual sum of square corresponds to the determination coefficient via the following relation

$$R^2 = 1 - NRSS. \quad (37)$$

Parameters in the logistics, Gompertz, and Richards model also be estimated with a similar manner. The accuracy of the predicted results could also be measured by evaluation of Mean Absolute Percentage Error (MAPE), which is given by the following formula

$$MAPE = \sum_i \frac{1}{n} \left| \frac{z_i - \hat{z}_i}{z_i} \right| 100\%. \quad (38)$$

Here n is the number of observational data. The nonlinear least square (nls) procedure of R open source software is used to estimate parameters of the proposed model, logistic, Gompertz, and Richards model. R open source software was built by the R Foundation for Statistical Computing. Estimation results of the proposed model, logistic, Gompertz, and Richards model for rooster and hen weight, the determination coefficient (R^2) and Mean Absolute Percentage Error (MAPE) for the models are presented in Table III, while the dynamics of rooster weight and hen weight are shown in Fig. 1 and Fig. 2 respectively.




Fig. 1: Dynamic of rooster weight.




Fig. 2: Dynamic of hen weight.

TABLE III: Estimated parameters for the proposed model, logistic, Richards and Gompertz growth model

Model	Parameters	Rooster	Hen
The proposed model (WEP-modified logistic growth model)	Mature weight (K)	2399.749	1847.162
	Effective growth rate (α)	0.031	0.029
	Inflection time (t_{inf})	71.584	69.015
	A	166.323	183.061
	NRSS	0.00031	0.00119
	R^2	0.99969	0.99881
Logistic model	MAPE	0.04754	0.04267
	Mature weight (K)	2279.904	1739.652
	Growth rate (r)	0.040	0.039
	Inflection time (t_{inf})	74.677	73.331
	NRSS	0.00357	0.00501
	R^2	0.99643	0.99499
Richards model	MAPE	0.299927	0.25398
	Mature weight (K)	2512.972	1945.342
	Growth rate (r^*)	0.023	0.021
	Inflection time (t_{inf})	64.307	61.344
	Shape parameter (m)	1.054	0.978
	NRSS	0.00071	0.00175
Gompertz model	R^2	0.99929	0.99825
	MAPE	0.07373	0.06552
	Growth rate (r)	0.022	0.021
	Inflection time (t_{inf})	63.498	61.704
	Mature weight (K)	2539.651	1936.385
	NRSS	0.00073	0.00175
	R^2	0.99927	0.99825
	MAPE	0.06007	0.07031

It could be seen from Fig. 1 and Fig. 2 that rooster growth and hen growth follow sigmoidal patterns. Rooster growth and hen growth starts by an accelerating growth phase from hatching. Then, the chicken attains a maximum growth rate at the inflection time. At final phase, the chicken weight tends to a mature weight. Qualitatively, all of the models, describe the chicken growth well, as seen in the figures. But, if we compare its MAPE, as seen in Table III, we see that the logistic model have the biggest MAPE, and it mean that its accuracy is poorer than the other models. This apparently due to the logistic model is not accurate in predicting the dynamics of rooster and hen weight at the early times (Fig. 1 and Fig. 2). By adding one additional parameter (q) to the presented model, the dynamics of rooster and hen weight could be better estimated by using the presented model.

From the Table III, it was found that the growth rate (the effective growth rate) or the maturation rate (α in the proposed model, r in the logistic and Gompertz model and r^* in the Richards model) was higher in rooster than in hen. This result is consistent with the result from Aggrey (2002) [3]. It also could be found that inflection time of the proposed model is relatively close to inflection time of the logistic model. In addition, inflection time of the Richards model is relatively close to the Gompertz model. It is apparently due to the shape parameter m in the Richards model is close to one. Moreover, it was found that the proposed model, logistic model, Richards model, and Gompertz model produced a high determination coefficient (R^2 is greater than 0.99). Although the determination coefficients of the four models did not differ significantly, Mean Absolute Percentage Error (MAPE) of the models considerably varied. It was found the proposed model has the smallest MAPE, which is 4.754% in rooster and 4.267% in hen. This indicates that the proposed model could be used as an alternative model to describe poultry growth curve or individual growth.

6. CONCLUSION

A new growth model was presented in this paper. The model was derived from the modification of logistic differential equation. The proposed model also was simulated and verified using rooster and hen weight data cited from the literature. The estimation results from the model were compared to the logistic model, Richards, and Gompertz growth model. It was found that the model gave better results compared to the logistic model, Richards, and Gompertz growth model. It indicates that model could be used as an alternative model to describe poultry growth curve or individual growth.

ACKNOWLEDGEMENT

Part of this research was funded by General Directorate of Higher Education (Ditjen DIKTI), Ministry of Research, Technology and Higher Education Republic Indonesia through "Penelitian Unggulan Perguruan Tinggi (PUPT)" project.

14
CONFLICT OF INTEREST

The authors do not have conflict of interest in regard to this research or its funding.

REFERENCES

- [1] M. Selvaggi, V. Laudadio, C. Dario and V. Tufarelli. Modelling Growth Curves in a Nondescript Italian Chicken Breed: an Opportunity to Improve Genetic and Feeding Strategies. *J. Poult. Sci.*, 52: 288–294, 2015.
- [2] I. Inounu, D. Mauluddin, R.R. Noor and Subandriyo. Analisis Kurva Pertumbuhan Domba Garut dan Persilangannya. *Jurnal Ilmu Ternak dan Veteriner*, 12(4): 286–299 (Text in Indonesian), 2007.
- [3] S.E. Aggrey. Comparison of Three Nonlinear and Spline Regression Models for Describing Chicken Growth Curves. *Poultry Science*, 81:1782–1788, 2002.
- [4] H. Nešetřilová. Multiphasic growth models for cattle. *Czech J. Anim. Sci.*, 50 (8): 347–354, 2005.
- [5] A.O. Raji, S.T. Mbap, and J. Aliyu. Comparison of different models to describe growth of the japanese quail (*coturnix japonica*). *Trakia Journal of Sciences*, 2:182–188, 2014.
- [6] N.B. Anthony, D.A. Emmerson, K.E. Nestor, W.L. Bacon, P.B. Siegel and E.A. Dunnington. Comparison of growth curves of weight selected populations of turkeys, quail and chickens. *Poultry Science*, 70:13–19, 1991.
- [7] R.E. Ricklefs. Modification of growth and development of muscles of poultry. *Poultry Science*, 64:1563–1576, 1985.
- [8] S. Mignon-Grasteau, C. Beaumont, E. Le Bihan-Duval, J.P. Poivey, H. de Rochambeau and F.H. Richard. Genetic parameters of growth curve parameters in male and female chickens. *British Poultry Science*, 40:44–51, 1999.
- [9] J. Stewart. Calculus Early Transcendental Seventh Edition. *Brooks/Cole Cengage Learning*, 2012.
- [10] M.A.J.S. van Boekel. Kinetic Modeling of Reactions in Foods, *CRC Press*, 2009.
- [11] P.J. Moatea, L. Dougherty, M.D. Schnall, R.J. Landis, R.C. Boston. A modified logistic model to describe gadolinium kinetics in breast tumors. *Magnetic Resonance Imaging*, 22:467–473, 2004.
- [12] L. von Bertalanffy. A quantitative theory of organic growth. *Human Biology*, 10(2): 181–213, 1938.
- [13] A. Tsoularis. Analysis of logistic growth models. *Res. Lett. Inf. Math. Sci.*, 2: 23–46, 2001.
- [14] F.J. Richards. A flexible growth function for empirical use. *Journal of Experimental Botany*, 10(29): 290–300, 1959.
- [15] A.A. Blumberg. Logistic growth rate functions. *Journal of Theoretical Biology*, 21: 42–44, 1968.
- [16] M.E. Turner, E. Bradley, K. Kirk and K. Pruitt. A Theory of Growth. *Mathematical Biosciences*, 29: 367–373, 1976.
- [17] J. France, J. Dijkstra, Ms. Dhanoa. Growth functions and their application in animal science. *Annales de zootechnie*, 45 (Suppl1): 165–174, 1996.
- [18] J.P. LaSalle. The stability of dynamical systems. *SIAM*, Philadelphia, 1976.
- [19] J.E. Beyer. On length-weight relationship computing the mean weight of the fish of a given length class. *Fish Bytes*, 5(10):11-13, 1987.
- [20] O.S. Ogunola, O.A. Onada, A.E. Falaye. Preliminary evaluation of some aspects of the ecology (growth pattern, condition factor and reproductive biology) of African pike, *Hepsetus odoe* (Bloch 1794), in Lake Eleiyele, Ibadan, Nigeria. *Fisheries and Aquatic Sciences*, 21:12, 2018.
- [21] D. Pauly. Some simple methods for the assessment of tropical fish stocks. *FAO Fisheries Technical Paper*, 234: 52, 1983.
- [22] Windarto, S.W. Indratno, N. Nuraini, and E. Soewono. A comparison of binary and continuous genetic algorithm in parameter estimation of a logistic growth model. *AIP Conference Proceedings*, 1587, 2014.

A new modified logistic growth model for empirical use

ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

15%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	repositorio.unesp.br Internet Source	1%
2	scik.org Internet Source	1%
3	tru.uni-sz.bg Internet Source	1%
4	Onyinebifun Emmanuel Biu, Maureen Tobechukwu Nwakuya, Gamage Tubona. "Assessment of Asymptotic and Logistics Growth Models on A Chemist Data", European Journal of Mathematics and Statistics, 2022 Publication	1%
5	Aggrey, S.. "Comparison of three nonlinear and spline regression models for describing chicken growth curves", Poultry Science, 2002. Publication	1%
6	Daisuke Satoh. "Discrete Gompertz equation and model selection between Gompertz and logistic models", International Journal of Forecasting, 2021 Publication	1%

7	Submitted to Universiti Teknologi Malaysia Student Paper	1 %
8	pdfs.semanticscholar.org Internet Source	1 %
9	Undergraduate Texts in Mathematics, 2012. Publication	1 %
10	Windarto, Fatmawati, Kamara Mustiko Putri. "A Maize Foliar Disease Mathematical Model with Standard Incidence Rate", IOP Conference Series: Materials Science and Engineering, 2019 Publication	1 %
11	O. Bamgbose. "Galeoides Decadactylus: A bio- indicator for chlorina' hydrocarbons in the Nigerian marine environment", Journal of Environmental Science and Health Part A, 02/1993 Publication	1 %
12	Windarto, Fatmawati, Nadiyah Nurlaily Nuzulia. "On mathematical model approach to competition dynamic of shipping companies in Surabaya", AIP Publishing, 2021 Publication	<1 %
13	Youness Mir, François Dubeau. "A Piecewise Model for Two-phase Growth Phenomena", Communications in Statistics - Simulation and Computation, 2019	<1 %

-
- 14 www.foodandnutritionjournal.org <1 %
Internet Source
-
- 15 fas.biomedcentral.com <1 %
Internet Source
-
- 16 Submitted to iGroup <1 %
Student Paper
-
- 17 www.coursehero.com <1 %
Internet Source
-
- 18 Gerardo Chowell, Richard Rothenberg, Kimberlyn Roosa, Amna Tariq, James M. Hyman, Ruiyan Luo. "Chapter 5 Sub-epidemic Model Forecasts During the First Wave of the COVID-19 Pandemic in the USA and European Hotspots", Springer Science and Business Media LLC, 2022 <1 %
Publication
-
- 19 Submitted to University of Witwatersrand <1 %
Student Paper
-
- 20 nozdr.ru <1 %
Internet Source
-
- 21 Lidan Jiang, Fang Zou, Yali Qiao, Ying Huang. "Patent analysis for generating the technology landscape and competition situation of renewable energy", Journal of Cleaner Production, 2022 <1 %

22

mafiadoc.com

Internet Source

<1 %

23

pure.rug.nl

Internet Source

<1 %

24

Edem, Edem Thomas, Ofunama, Preye, Eyo, Victor `Oscar. "Length-weight Relationship, Growth Pattern and Condition Factor of the Silver Catfish *Chrysichthys nigrodigitatus* (Lacepède, 1803) from Lower River Benue, Nigeria", *Asian Journal of Fisheries and Aquatic Research*, 2018

Publication

<1 %

25

eudl.eu

Internet Source

<1 %

26

Li, Yu Qi, Huan Zhang, and Yi Ran Liu. "Regression Analysis of Observed Foundation Settlement by Modified Logistic Growth Model", *Advanced Materials Research*, 2011.

Publication

<1 %

27

Windarto, Utami Dyah Purwati, Nadiyah Amalia. "A Mathematical Model of Social Media Popularity with Standard Incidence Rate", *IOP Conference Series: Materials Science and Engineering*, 2019

Publication

<1 %

28

"Applications of Evolutionary Computation",
Springer Science and Business Media LLC,
2016

Publication

<1 %

29

Alberto Oliveri, Martina Maselli, Matteo Lodi,
Marco Storace, Matteo Cianchetti. "Model-
Based Compensation of Rate-Dependent
Hysteresis in a Piezoresistive Strain Sensor",
IEEE Transactions on Industrial Electronics,
2019

Publication

<1 %

30

Joseph Junior Aduba, Behrooz Asgari.
"Analysing and forecasting the diffusion of
electronic payments system in Nigeria",
Technology Analysis & Strategic Management,
2021

Publication

<1 %

31

Witzel, F., R. Fritsche-Guenther, N. Lehmann,
A. Sieber, and N. Bluthgen. "Analysis of
impedance-based cellular growth assays",
Bioinformatics, 2015.

Publication

<1 %

32

advancesindifferenceequations.springeropen.com

Internet Source

<1 %

33

dergipark.org.tr

Internet Source

<1 %

34 Poudel, Diwakar, Leif K. Sandal, and Sturla F. Kvamsdal. "Stochastically Induced Critical Depensation and Risk of Stock Collapse", *Marine Resource Economics*, 2015. <1 %

Publication

35 Qiu Zhai, Sihan Yan, Huakun Wang, Yu Zhang. "Experimental study on ship's mooring force of arched longitudinal beam wharf", 2021 7th International Conference on Hydraulic and Civil Engineering & Smart Water Conservancy and Intelligent Disaster Reduction Forum (ICHCE & SWIDR), 2021 <1 %

Publication

36 U Haberlandt, B Klöcking, V Krysanova, A Becker. "Regionalisation of the base flow index from dynamically simulated flow components — a case study in the Elbe River Basin", *Journal of Hydrology*, 2001 <1 %

Publication

37 eprints.undip.ac.id <1 %

Internet Source

38 www.hindawi.com <1 %

Internet Source

39 www.ncbi.nlm.nih.gov <1 %

Internet Source

40 MONDHER BELLALAH, THEMA, CEREG. "Valuing Lease Contracts Under Incomplete <1 %

Information: A Real-Options Approach", The Engineering Economist, 2002

Publication

41

S. H. Saker. "Oscillation and attractivity of discrete nonlinear delay population model", Journal of Applied Mathematics and Computing, 09/2007

Publication

<1 %

42

W.B. Roush, S.L. Branton. "A comparison of fitting growth models with a genetic algorithm and nonlinear regression", Poultry Science, 2005

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

A new modified logistic growth model for empirical use

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
