

INTERNATIONAL JOURNAL OF SCIENTIFIC ADVANCES

”



VOLUME 3 ISSUE 5 : SEP-OCT 2022

ISSN: 2708 - 7972

WWW.IJSCIA.COM



VOLUME 3 | ISSUE 5 : SEP - OCT 2022

Eye Pupil Localisation and Labeling Using a Small Size Database and YOLOv4 Object Detection Algorithm

🕒 30 October 2022 • 📖 Volume3 Issue 5

Eye Pupil Localisation and Labeling Using a Small Size Database and YOLOv4 Object Detection Algorithm
Varuzhan H. Baghdasaryan Abstract Eye-related

Read More »

🔗 <https://doi.org/10.51542/ijscia.v3i5.22>

Central Neuroaxial Block Anesthesia: A Safe Choice for Caesarean Section in Patients with Mitral Stenosis

🕒 28 October 2022 • 📖 Volume3 Issue 5

Central Neuroaxial Block Anesthesia: A Safe Choice for Caesarean Section in Patients with Mitral Stenosis
Dewa Ayu Mas Shintya Dewi1*

Read More »

🔗 <https://doi.org/10.51542/ijscia.v3i5.21>

Magnetic Monopoles and Lenz's Law in Hyperbolic Medicine





“Hyperbolic Medicine” the study of the

[Read More »](#)

<https://doi.org/10.51542/ijscia.v3i5.20>

Efficacy Assessment of Decentralized Wastewater Treatment Systems in treating Faecal Sludge within Dar es Salaam City

🕒 27 October 2022 • 📖 Volume3 Issue 5

Efficacy Assessment of Decentralized Wastewater Treatment Systems in treating Faecal Sludge within Dar es Salaam City Mwingira Catherine* & Alexander

[Read More »](#)

<https://doi.org/10.51542/ijscia.v3i5.19>

Effectiveness of Health Education about Stunting Toward Improvement of Community Knowledge

🕒 18 October 2022 • 📖 Volume3 Issue 5

Effectiveness of Health Education about Stunting Toward Improvement of Community Knowledge Muhammad Faizi1, Nur Rochmah1*, Yuni Hisbiyah1, Rayi Kurnia Perwitasari1,

[Read More »](#)

<https://doi.org/10.51542/ijscia.v3i5.18>

To Explore the Division of Responsibilities Between General Practitioners and Specialists

🕒 17 October 2022 • 📖 Volume3 Issue 5

To Explore the Division of Responsibilities Between General Practitioners and Specialists Jian Gao Abstract This paper introduces research about the

[Read More »](#)

<https://doi.org/10.51542/ijscia.v3i5.17>

A Multimodal Hate Speech Classification Process Using Dual Feature Extraction Techniques

🕒 13 October 2022 • 📖 Volume3 Issue 5

A Multimodal Hate Speech Classification Process Using Dual Feature Extraction Techniques Chibuike Onuoha1, Ikerionwu Charles2 and Obi Nwokonkwo1 Abstract Racist

[Read More »](#)

<https://doi.org/10.51542/ijscia.v3i5.16>

Physical Examination and Clinical Course of Poisoning and Drug Overdose

🕒 13 October 2022 • 📖 Volume3 Issue 5

Physical Examination and Clinical Course of Poisoning and Drug Overdose Elendu Chukwuka1*, Chiagozie Perpetua Ayabazu2, Akinbayo A. Akintunde2, Ibukunoluwa V.

[Read More »](#)

<https://doi.org/10.51542/ijscia.v3i5.15>





Non-Cirrhotic Hepatocellular Carcinoma with Duodenal Metastasis in

Chronic Hepatitis B: Case Report

11 October 2022 • Volume3 Issue 5

Non-Cirrhotic Hepatocellular Carcinoma with Duodenal Metastasis in Chronic Hepatitis B: Case Report I Gusti Lanang Rama Dwi Suputra* & I

Read More »

<https://doi.org/10.51542/ijscia.v3i5.14>

Distribution of Body Mass Index and Random Plasma Glucose as Early Detection of Diabetes Mellitus in Indonesian Students

11 October 2022 • Volume3 Issue 5

Distribution of Body Mass Index and Random Plasma Glucose as Early Detection of Diabetes Mellitus in Indonesian Students Nur Rochmah1,

Read More »

<https://doi.org/10.51542/ijscia.v3i5.13>

Assessment and Analysis of Undergraduate Fall Enrollment Trends of The Selected College of Agriculture at the 1890 Land-Grant Universities: 1996 To 2018

11 October 2022 • Volume3 Issue 5

Assessment and Analysis of Undergraduate Fall Enrollment Trends of The Selected College of Agriculture at the 1890 Land-Grant Universities: 1996

Read More »

<https://doi.org/10.51542/ijscia.v3i5.12>

A Review Article on Etiology, Clinical Presentation, Laboratory Evaluation and Treatment of Autoimmune Hemolytic Anemia

28 September 2022 • Volume3 Issue 5

A Review Article on Etiology, Clinical Presentation, Laboratory Evaluation and Treatment of Autoimmune Hemolytic Anemia Jamil Ahmad1*, Faryal Gohar2, Zeyd

Read More »

<https://doi.org/10.51542/ijscia.v3i5.11>

An Assessment of E-Learning Strategies in Enhancing Quality Education in Cameroon During the COVID-19 Pandemic in Cameroon: Case of Government Bilingual High School Yaoundé

24 September 2022 • Volume3 Issue 5

An Assessment of E-Learning Strategies in Enhancing Quality Education in Cameroon During the COVID-19 Pandemic in Cameroon: Case of Government

Read More »

<https://doi.org/10.51542/ijscia.v3i5.10>

Cardiac Shock in Patients Hospitalized for Acute Myocardial Infarction

23 September 2022 • Volume3 Issue 5





CALL FOR PAPERS
VOLUME 4 ISSUE 2 [Click Here](#)  **Mar - Apr 2023**



EDITORIAL BOARD

Editor-in-Chief

Dr. T. J. Ashaolu

Email: editor@ijscia.com, ijsciaeditor@gmail.com

Assistant Editor-in-Chief

Joseph O. Ashaolu

National Yang-Ming University, Taipei, Taiwan

Rp 199,0 JT	Rp 109,0 JT	Rp 80,0 JT	Rp 108,0 JT	

Beli Mobil Pasti Dapat Garansi
OLX Indonesia

EDITORIAL MEMBERS

Prof. Emeritus MNO Sadiku
Prairie View A&M University
Texas, USA

Professor Ramesh C. Gupta
Former Vice Chancellor, Radha Govind University
Jharkhand, India

Professor Massoud Kaykhali
University of Sistan & Baluchestan
Zahedan, Iran

Professor Elzbieta Macioszek
Silesian University of Technology
Gliwice, Poland

Professor Dariusz Jacek Jakobczak
Koszalin University of Technology
Koszalin, Poland

Asst. Prof. Esam AL Lawati
A' Sharqiyah University
Ibra, Sultanate of Oman

Dr. S. A. O. Adeyeye
Ten Duc Thang University
Ho Chi Minh City, Vietnam

Asst. Prof. Ibrahim Khalifa
Benha University
Moshtohar, Egypt

Professor Dr. Ho Soon Min
INTI International University
Nilai, Malaysia

Dr. B. J. Alegbeleye
St Elizabeth Catholic General Hospital
Kumbo-Nso, Cameroon

Dr. Ridwan Babatunde Ibrahim
Academia Sinica
Taipei, Taiwan

Editorial board -

Prof. Dr. Eng. Ahmed kadhim Hussein
Babylon University
Babylon City, Hillia, Iraq

Dr. Murat Basar
IVF Laboratories Bahçeci Sağlık Grubu
Istanbul, Turkey

Dr. Bassam Saeed
Farah Association for Child with Kidney Disease
Damascus, Syria

Asst. Prof. Selezneva Irina Stanislavovna
Ural Federal University
Yekaterinburg, Russia

Professor Michael Gr. Voskoglou
GTES Institute of Western Greece
Patras, Greece

Assoc. Prof. Thoudam Paraskumar Singh
MGM College of Engineering and Technology Kamothe,
Navi Mumbai, India

Dr. Isaac Delali Kottoh
Ghana Atomic Energy Commission
Accra, Ghana

Assoc. Prof. Godfred A Menezes
RAK Medical & Health Sciences University
Ras Al Khaimah, United Arab Emirate

Assoc. Prof. Syed Kamran Sami
Balochistan University of Information Technology
Quetta, Pakistan

Asst. Prof. Muhammad Amin
Balochistan University of Information Technology
Quetta, Pakistan

Ma. Gladys B. Aquino
De La Salle Medical and Health Sciences Institute
Dasmariñas, Philippines

Assoc. Prof. Vahe Davtyan
Brusov State University
Yerevan, Armenia

Dr. Sylvain Somé Marie
National Yang-Ming University
Taipei, Taiwan

Professor Teresa Orbera
University of Oriente
Santiago de Cuba, Cuba

Dr. Magdaline Joseph Kwaji
Bioresources Development Center
Odi, Bayelsa, Nigeria

Dr. Sampson Tsumori Akoko

Editorial board -

Dr. Sampson Kwame-Akpan
Kwame Nkrumah University of Science and Technology
Kumasi, Ghana

Asst. Prof. Nisarg G. Gandhewar
SB Jain Institute of Technology
Nagpur, India

Dr. Manoj Khandelwal
Federation University
Ballarat, Victoria, Australia

Dr. Taimur Sharif
Newman University
Birmingham, UK

Dr. Tinni Dutta
University of Calcutta
Kolkata, India

Professor Dr. Jacob Oluwoye
Alabama A&M University
Normal, Alabama, USA

Dr. Benjamin D. Sookhoo
University of South Florida
Tampa, Florida, USA

Victor Eyo Assi
University of Uyo
Uyo, Akwa Ibom, Nigeria

Dr. Bid D. Dhruvasasad
Sarvajanic College of Physiotherapy
Gujarat, India

Dr. Ankit Bhargava
Consultant Physio & Fitness Expert-Wendt India Ltd.
India

Dr. Manoranjan Tripathy
Dev Sanskriti Vishwavidyalaya
Uttarakhand, India

Asst. Prof. Abbas Ar. Mohamed
PMBAH National Guard Health Affairs
Al Madinah Al Munawwarah, Kingdom of Saudi Arabia

Asst. Prof. Nurul Mohammad Zayed
Daffodil International University
Dhaka, Bangladesh

Prof. Dr. Hasibun Naher
Brac University
Dhaka, Bangladesh

Davis Ojima J.P. Bursar
Ignatius Ajuru University of Education
Port Harcourt, Rivers, Nigeria

Dr. Okogwu Antonia
University of Port Harcourt

Editorial board -

University of Port Harcourt
Port Harcourt, River, Nigeria

Dr. Garyfalia Charitaki
Hellenic Open University (University of Thessaly)
Volos, Greece

Dr. Munish Kumar Sharma
University of Alberta
Edmonton, Canada

Dr. Niyonzima N. Francois
Deputy Vice Chancellor Academic and Research INES
Ruhengeri, Rwanda

Dr. Srishti Sharma
M. Patel College of Physiotherapy
Gandhi Nagar, India

Professor R. K. Mathukia
Junagadh Agricultural University
Junagadh, Gujarat, India

Professor Hamid Ali Abed Al-Asadi
Basra University
Basra, Iraq

Dr. Firas Abdullah Thweny Al-Saedi
Nahrain University
Baghdad, Iraq and Private Researcher, New Jersey, USA

Dr. George Fuh Kum
University of Yaoundé I
Yaoundé, Cameroon

Professor Mohamed El Houseny El Sebeay Shams
Mansoura University
Mansoura, Egypt

Dr. Peter Ashlame Agu
Nasarawa State University
Keffi, Nigeria

Dr. Asadullah Baloch
Mir Chakar Khan Rind University
Quetta, Pakistan

Professor Ignatius Isaac Dambudzo
Zimbabwe Open University,
Harare, Zimbabwe

Professor Hamid Ali Abed Alasadi
Iraq University College
Basra, Iraq

Dr. Ammar Daher Bashatweh
Independent Scholar
Irbid, Jordan

Subrat Kumar Mahapatra
Central University
Cuttack, West Bengal

Dr. Xiong Chen
Hubei University of Medicine
Shiyan City, China



ISSN: 2708-7972



This work is licensed under a Creative Commons Attribution 4.0 (International) Licence.(CC BY-NC 4.0).

Navigations

- > [About IJSCIA](#)
- > [Editorial Board](#)
- > [Achives](#)
- > [Research Areas](#)
- > [Ethical Guidelines](#)
- > [Article Processing Charges](#)
- > [Submit Paper](#)
- > [Payment Link](#)

Download

- > [Sample Paper Format](#)
- > [Copyright Form](#)

Support

- > [Contact Us](#)
- > [F.A.Q](#)

Distribution of Body Mass Index and Random Plasma Glucose as Early Detection of Diabetes Mellitus in Indonesian Students

Nur Rochmah¹, Muhammad Faizi^{1*}, Yuni Hisbiyah¹, Rayi Kurnia Perwitasari¹, Jihan Kalishah², Qorri 'Aina², Tyas Maslakhathien Nuzula²

¹Department of Child Health, Faculty of Medicine, Dr. Soetomo General Hospital Universitas Airlangga, Surabaya, East Java, Indonesia

²Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia

*Corresponding author details: Muhammad Faizi; muhammad.faizi@fk.unair.ac.id

ABSTRACT

Background: Diabetes mellitus (DM) is a chronic metabolic disorder characterized by abnormal elevated blood glucose levels resulting from defects in insulin secretion, insulin action, or both. The prevalence of DM in children and adolescents is increasing worldwide. Until recently, early detection of DM was still not widely done in Indonesia. Therefore, this study aimed to measure BMI and random plasma glucose in Indonesian students. **Methods:** This descriptive study involving 8th and 9th grade students at Alif Laam Miim Islamic Boarding School Surabaya aged 13 - 18 years old. Height and weight were measured using standard anthropometry. Capillary blood samples were taken on the index or middle finger of the student using a single-use lancet. Statistical analysis was conducted with SPSS using the T-test, Mann-Whitney, and Chi square test. **Results:** This study included 95 subjects with the mean age of 14.65 ± 1.83 years old. There is no significant difference of BMI ($p = 0.63$) and random plasma glucose ($p = 0.09$) between males and females. The distribution of BMI for underweight, healthy weight, overweight, and obesity were 6.32%; 64.21%; 14.74%; 14.74% respectively. None of them had a high range of random plasma glucose. **Conclusion:** Indonesian students in Surabaya had a healthy weight and a normal range of plasma glucose.

Keywords: diabetes mellitus; body mass index; random plasma glucose; children; adolescence

INTRODUCTION

Diabetes mellitus (DM) is a long-term metabolic condition marked by abnormally high blood sugar levels brought on by deficiencies in insulin secretion, insulin action, or both [1]. Based on their etiopathogenetic, there are two groups of DM types, including type 1 and type 2 DM. Type 1 Diabetes Mellitus (T1DM) is defined by an inability to produce insulin due to the destruction of pancreatic-cells, which is typically the outcome of T-cell-mediated autoimmunity. In contrast, Type 2 Diabetes Mellitus (T2DM) is characterized by insulin resistance and insufficient insulin secretory compensation [2].

T1DM was the most prevalent form of diabetes in children, whereas T2DM was more prevalent in adults and obese individuals [3]. Recent research indicates that the prevalence of T2DM in children and adolescents has increased over the past several decades [2, 4, 5]. The prevalence of T2DM in children and adolescents has increased due to the lifestyle factors, which also contribute to the global rise in obesity [4]. In the United States, the prevalence among adolescents with T2DM is increasing from 9 to 12.5 cases per 100.000 youth [5]. The incidence of T2DM in children increases with age and the peak age onset of T2DM occurs at puberty [6]. Childhood onset of T2DM is associated with greater risk of microvascular and macrovascular complication [5]. The Indonesian Pediatric Society (IPS) had only recorded 1,249 Indonesian children with T1DM from 2017–2019.

The prevalence of T1DM in Indonesia raises over 10 years, from 3.88 per 100 million in 2000 to 28.19 per 100 million in 2010. The diagnosis and treatment of T1DM in children is often delayed. In Indonesia, the number of T1DM with Diabetic Ketoacidosis (DKA) at diagnosis remained high (71%) [7]. Consequently, screening for DM in children is essential in order to provide an early diagnosis and intervention to prevent future problems. The screening for diabetes mellitus consisted of measuring glycemia and doing a risk assessment for DM [8].

In 2010, the proportion of adolescents in Indonesia was around 43.5 million people. Badan Pusat Statistik (BPS) Indonesia said that there are 426,786 adolescents aged 10-19 in Surabaya [9]. Approximately, 87% of adolescents have a sedentary lifestyle and a poor diet. It can result in obesity and raise the risk of metabolic disorders, including diabetes in old age [10]. The World Health Organization (WHO) has suggested Body Mass Index (BMI) assessment as the simplest method for classifying obesity [11]. BMI measurement is a simple and inexpensive method for screening for obesity and other weight disorders [12]. Until recently, early detection of diabetes mellitus was still not widely done in Indonesia, especially in the student population. Therefore, we aimed to measure BMI and random plasma glucose in Indonesian students.

METHODS

This descriptive study involving 8th and 9th grade students at Alif Laam Miim Islamic Boarding School Surabaya aged 13 - 18 years old. This study was approved by the Faculty of Medicine Health Research and Ethics Committee with ethical clearance number 139/EC/KEPK/FKUA/2022. Informed consent was already obtained from the subject's guardians. Height and weight were measured using standard anthropometry done by healthcare professionals. Height was measured to the nearest 0.1 cm using a wall-mounted tape measure (Onemed) while body weight was measured to the nearest 0.1 kg (Onemed). Capillary blood samples were taken on the index or middle finger of the student using a single-use lancet to measure random plasma glucose (ACCU CHECK). Students who refused to have their height, weight, and random plasma glucose measured were excluded from this study.

Body Mass Index (BMI) was calculated by dividing weight in kilograms with the square of height in meters. BMI was categorized for underweight (<5th percentile), healthy weight (5th percentile - <85th percentile), overweight (>85th percentile - <95th percentile), and obesity (≥95th percentile) based on the BMI-for-age percentile growth charts [12].

Random plasma Glucose was categorized by normal (<200 mg/dL) and high (≥200 mg/dL) [13]. Data was entered in and statistical analysis was performed using SPSS version 17 with p-value < 0.05 was considered as statistically significant. The variables were expressed as mean ± standard deviation and frequency (percentage). Difference variable in numeric data between male and female was analyzed using T-test or Mann-Whitney U test based on the Kolmogorov-Smirnov normality test. Meanwhile, categorical data was analyzed with Chi-square or Fisher-exact to compare frequencies.

RESULT

This study enrolled 95 students in Surabaya. The demographic characteristics of subjects are shown in Table 1. The mean age of the subject is 14.65 ± 1.83 years old. There is no significant difference of BMI (p = 0.63) and random plasma glucose (p = 0.09) between males and females. BMI was further classified into four categories based on the CDC criteria [12]. The distribution of the BMI based on the categories can be seen in Figure 1. From 95 students, none of them had a high range of random plasma glucose that can be seen in Table 3.

TABLE 1: Demographic Characteristics of Samples

Characteristic	All (n = 95)	Males (n = 42)	Females (n = 53)	p-value
Age (years)	14.65 ± 1.83	14.80 ± 1.06	14.79 ± 1.04	0.96
Anthropometry				
Height (m)	1.57 ± 0.08	1.64 ± 0.05	1.52 ± 0.07	0.00*
Weight (kg)	56.03 ± 13.06	60.80 ± 14.78	52.33 ± 10.03	0.00*
Body mass index (kg/m ²)	22.60 ± 4.57	22.48 ± 4.78	22.68 ± 4.47	0.63
Random plasma glucose (mg/dL)	95.10 ± 9.25	93.07 ± 10.86	96.32 ± 7.68	0.09
Parental Ethnicity				
Javanese	79 (83.2)	37 (88.1)	42 (79.2)	0.58
Madura	13 (13.7)	4 (9.5)	9 (17.0)	
Other	3 (3.2)	1 (2.4)	2 (3.8)	
History of family with DM*				
Yes	16 (16.8)	6 (14.3)	10 (18.9)	0.55
No	79 (83.2)	36 (85.7)	43 (81.1)	

DM, Diabetes Melitus; *, statistically significant




FIGURE 1: Distribution of Body Mass Index (BMI)

TABLE 2: Distribution of Random Plasma Glucose

Category	Total (n)	Percent (%)
Normal (<200 mg/dL)	95	100%
High (>200 mg/dL)	0	0

DISCUSSION

This study reveals that the majority of students had a healthy BMI (64.21%) and that the proportion of students with overweight and obesity is the same (14.74%). The prevalence of overweight is approximately 18% and obesity is approximately 30.7%, which is quite low compared to previous studies conducted among university students in Indonesia [14]. Higher prevalence of obesity and overweight compared to this study also found in other countries like Boston (34% of females and 32% of males), India (27%), Qatar (46% of males and 44% of females), and Morocco (42%) [15-18]. However, Norwegian has a lower number of children and adolescents with obesity (5.2% of males and 8.0% of females) [19]. A significantly lower prevalence of overweight and obesity in our study compared to other nations may be attributable to the fact that the study population resides in a boarding school that provides daily meals. The school also has a tight policy prohibiting kids from purchasing food from outside the school. It restricts kids' access to junk food and other unhealthy foods that may contribute to obesity. Another study with boarding school students as the population found a reduced prevalence of overweight and obesity among children and teenagers [20].

The screening program will be influenced by a number of variables, such as: 1) disease (severity, prevalence, potential pre-clinical detection, extended latency period, improved outcomes with early identification); 2) screening test (validity, reliability, ease of use, ability to detect disease pre-clinically); and 3) screening program [21]. The ADA suggests evaluating children and teenagers who are overweight and have two risk factors, such as acanthosis nigricans, hypertension, hyperlipidemia, or a first- or second-degree relative with a history of type 2 diabetes. The ADA and the American Academy of Pediatrics advise screening at-risk patients every two years beginning at age 10 or the start of puberty before age ten [22]. Elevated blood glucose levels lead to the development of problems during the lengthy pre-clinical phase of diabetes. So many healthcare guidelines now advise screening for the disease in order to detect diabetes early. The objective is to identify asymptomatic patients so that earlier treatment can have a positive effect on the disease's progression. The most probable culprits in the development of T1DM are genetic susceptibility, environmental factors, vitamin D processing, and viruses. There is no substantial evidence indicating the root cause as of yet [21].

None of the students in this study had a high range of RPG based on ADA criteria (>200 mg/dL). In our study, around 26 children had RPG 100 mg/dL and the maximum value of RPG was 123 mg/dL in one child. It contrasts with a study from India who found that 5.3% of the obese children population in their study had an elevated RPG (>130 mg/dL) [23]. Study from Qatar also found 6% of boys and 2% of girls with elevated blood sugar (RPG \geq 140 mg/dL and/or Fasting Plasma Glucose \geq 100 mg/dL) [17]. Nevertheless, in a study with samples of the US adult population without diagnosed diabetes, a single RBG \geq 100 mg/dL was found strongly associated with undiagnosed diabetes and was the single strongest predictor of undiagnosed diabetes outperforming other risk factors [24].

International Diabetes Federation (IDF) also recommends that individuals with RPG value 100 - 199 mg/dL should undergo a formal diabetes testing [8]. Until now, the recommended screening and diagnostic tools for prediabetes and diabetes are Fasting Plasma Glucose (FPG), Oral Glucose Tolerance Test (OGTT) and HbA1c [13]. In actual practice, however, most practitioners employ RPG as a screening tool because it may be performed and prescribed on the same day as the patient's visit [25, 26]. To the best of our knowledge, this is the first study in Indonesia that shows a distribution of BMI and RPG as early detection of DM in children and adolescents. Nevertheless, this study had several limitations: first, a small amount of population size was used in this study; second, even though BMI was the easiest way to use for screening obesity, it can't measure total amount of body fat in the body directly; third, this study uses RPG as a screening test of dysglycemia instead of FPG, OGTT, or HbA1c.

CONCLUSION

Our study showed that most adolescents in a boarding school in Surabaya had a healthy weight and a normal range of plasma glucose. However, it's important to continuously do BMI and glucose screening as an early detection of DM, especially in overweight and obesity adolescents. Further studies with more complete and bigger data are needed to make a better analysis between BMI and RPG as an early detection of DM.

REFERENCES

- [1] American Diabetes Association (ADA). (2012). Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes. *Diabetes care*, 43(1), S14-S31. doi: 10.2337/dc20-S002.
- [2] Craig, M. E., Jefferies, C., Dabalea, D., Balde, N., Seth, A., Donaghue K.C. (2014). Definition, epidemiology, and classification of diabetes in children and adolescents. *Pediatric Diabetes*, 15(S20), 4-17. doi:10.1111/PEDI.12186.
- [3] Tan S.Y., Mei Wong J.L., Sim Y.J., Wong S.S., Mohamed Elhassan S.A., Tan S.H., et al. (2018). Type 1 and 2 diabetes mellitus: A review on current treatment approach and gene therapy as potential intervention. *Diabetes & Metab Syndr*, 13(1), 364-372. doi: 10.1016/j.dsx.2018.10.008.
- [4] Botero, D., & Wolfsdorf, J.I. (2005). Diabetes mellitus in children and adolescents. *Archives of Medical Research*, 36(3), 281-290. doi: 10.1016/J.ARCMED.2004.12.002.
- [5] Koren, D., & Evotsky, L.L. (2021). Type 2 Diabetes Mellitus in Childhood and Adolescence. *Pediatrics in Review*, 42(4), 167-179. doi: 10.1542/pir.2019-0236
- [6] Dileepan, K., & Max Feldt, M. (2013) Type 2 Diabetes Mellitus in Children and Adolescents, *Pediatrics in Review*, 34(12), 541-548. doi: 10.1542/PIR.34-12-541.

- [7] Pulungan, A.B., Fadiana, G., & Annisa, D. (2021). Type 1 Diabetes Mellitus in Children: Experience in Indonesia. *Clinical Pediatric Endocrinology*, 30(1), 11-18. doi: 10.1297/cpe.30.11
- [8] International Diabetes Federation (IDF). (2012). Global Guideline for Type 2 Diabetes. Belgium: IDF.
- [9] Badan Pusat Statistik (BPS) Indonesia. (2013). Statistik Indonesia 2011. Jakarta: BPS
- [10] Pramono, A., et al. (2014). Kontribusi Makanan Jajan dan Aktivitas Fisik terhadap Kejadian Obesitas pada Remaja di Kota Semarang. *Gizi Indonesia*, 37(2), 129-136. doi: 10.36457/GIZINDO.V37I2.158.
- [11] Behera, A.A., Behera B.K, Dash, S., & Mishra, S. (2014). Effect of body mass index on gender difference in lung functions in Indian population. *Int J Clin Exp Physiol*, 1(3),229-231.
- [12] Centre for Disease Control (CDC) (2021). About Child and Teen BMI. accessed on: https://www.cdc.gov/healthyweight/assessing/bmi/childrens_bmi/about_childrens_bmi.html
- [13] American Diabetes Association (ADA). (2020). Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes-2020. *Diabetes Care*, 43(Suppl1), 14-31. Doi: 10.2337/dc20-S002
- [14] Oktariza, R.T., Kalanjati, V.P., & Tirthaningsing, N.W. (2021). Body Mass Index, Waist-Hip Ratio and Fasting Glucose amongst the University Students. *Fol Med Indones*, 57(1), 53-57. Doi: 10.20473/fmi.v57i1.14661
- [15] Field, A.E., Cook, N. R., Gilman, M.W. (2005). Weight Status in Childhood as a Predictor of Becoming Overweight or Hypertensive in Early Adulthood. *Obes Res*, 13(1), 163-169. Doi: 10.1038/oby.2005.21
- [16] Khadilkar, V.V., Khadilkar, A.V., Cole, T.J., Chiplonkar, S.A., & Pandit, D. (2011). Overweight and Obesity Prevalence and Body Mass Index Trends in Indian Children. *Int J Pediatr Obes*, 6(2-2), 2016-224. Doi: 10.3109/17477166.2010.541463.
- [17] Mamtani, R., Lowenfels, A.B., Sheikh, J., Cheema D., Al-Hamaq A., Matthis, S.A., et al. (2014). Adolescent prediabetes in a high-risk Middle East Country: a cross-sectional study. *JRSM Open*, 5(8), 1-5. Doi: 10.1177/2054270414536550
- [18] Mehdad, S., Hamrani, A., Kari, K.E., Hamdouchi, A.E., Barakat A., Mzibri M.E., et al. (2012). Body Mass Index, Waist Ciurcumferenece, Body Fat, Fasting Blood Glucose in a Sample of Moroccan Adolescents Aged 11-17 Years. *J Nutr Metab*, 510458. Doi: 10.1155/2012/510458
- [19] Bjorge, T., Engeland, A., Tversal, A., Smith, G.D. (2008). Body Mass Index in Adolescence in Relation to Cause-specific Mortality: A Follow Up of 230.000 Norwegian Adolescents. *American Journal of Epidemiology*, 168(1), 30-37. Doi: 10.1093/aje/kwn096
- [20] Kola-Raji, B.A., Balogun, M.R., & Odugbemi, T.O. (2017). A comparative study of nutritional status of adolescents from selected private and public boarding secondary schools in Ibadan, South Western Nigeria. *Journal of Medicine in the Tropics*, 19 (1), 49-55. Doi: 19. 49. 10.4103/jomt.jomt_43_16.
- [21] Peer, N., Balakrishna, Y., & Durao, S. (2020). Screening for type 2 diabetes mellitus. *Cochrane Database of Systematic Reviews*, 5. <https://doi.org/10.1002/14651858.CD005266.pub2>
- [22] Pippit, K., Li, M., & Gurgle, H.E. (2016). Diabetes Mellitus: Screening and Diagnosis. *American Family Physician*, 93 (2), 103-109
- [23] Khadilkar, A.V., Lohiya, N., Mistry, S., Chiplonkar, S., Khadilkar, V., Kajale, N., et al. (2019). Random Blood Glucose Concentrations and their Association with Body Mass Index in Indian School Children. *Indian Journal of Endocrinology and Metabolism*, 23(5), 529-535. Doi: 10.4103/ijem.IJEM_536_19
- [24] Bowen, M.E., Xuan, L., Lingvay, I., Halm, E.A. (2015). Random Blood Glucose: A Robust Risk Factor for Type 2 Diabetes. *J Clin Endocrinol Metab*, 100(4), 1503-1510. Doi: 10.1210/jc.2014-4116
- [25] Lee, J.M., Gebremariam, A., Wu, E.L., LaRose, J., Gurney, J.G. (2011). Evaluation of Nonfasting Tests to Screen for Childhood and Adolescent Dysglycemia. *Diabetes Care*, 34(12), 2597-2602. Doi: 10.2337/dc11-0827
- [26] Saudek, C.D., Herman, W.H., Sacks, D.B., Bergenstal, R.M., Edelman, D.M., Davidson, M.B. (2008). A New Look at Screening and Diagnosing Diabetes Mellitus. *J Clin Endocrinol Metab*, 93(7):2447-2453. Doi: 10.1210/jc.2007-2174