

Diabetes duration and thyroid stimulating hormone levels in children with type 1 diabetes mellitus

by Nur Rochmah

Submission date: 15-Mar-2023 01:10PM (UTC+0800)

Submission ID: 2037577094

File name: ing_hormone_levels_in_children_with_type_1_diabetes_mellitus.pdf (294.73K)

Word count: 2308

Character count: 12370

Diabetes duration and thyroid stimulating hormone levels in children with type 1 diabetes mellitus

Nur Rochmah, Muhammad Faizi

Abstract

Background Children with type 1 diabetes mellitus (T1DM) are at risk of thyroid dysfunction. An association between diabetes duration and thyroid stimulating hormone level remains inconclusive.

Objective To assess for a possible association between diabetes duration and thyroid stimulating hormone levels in children with T1DM.

Methods We conducted a cross-sectional study from January to June 2017 in the Pediatric Endocrine Outpatient Clinic at Dr. Soetomo Hospital. Subjects were children with T1DM aged 7 to <18 years. Exclusion criteria were children with diabetic ketoacidosis, previously diagnosed thyroid problems, and hospitalization in the pediatric intensive care unit (PICU).

Results From the 55 regular patients in our outpatient clinic, 34 patients were included in the study. Nineteen (54.3%) subjects were male, and the overall mean age was 11.3 years. Subjects' mean duration of diabetes was 3 years and their mean thyroid stimulating hormone concentration was 3.76 mIU/L. Pearson's correlation test revealed no significant association between duration of diabetes and thyroid stimulating hormone level ($r_s = -0.068$; $P = 0.703$).

Conclusion There is no significant association between duration of diabetes and thyroid stimulating hormone levels in children with T1DM. [Paediatr Indones. 2018;58:80-3; doi: <http://dx.doi.org/10.14238/pi58.1.2018.80-3>].

Keywords: diabetes duration; thyroid stimulating hormone; type 1 diabetes mellitus children

According to the World Health Organization (WHO) and the International Diabetes Federation (IDF), the prevalence of diabetes is on the rise.^{1,2} Data from the Pediatric Endocrine Working Group, Indonesian Pediatric Society, showed that 1,153 patients suffered from T1DM until April 2016.³ Thyroid dysfunction is reportedly higher among T1DM patients.⁴⁻⁹ Among diabetic adult populations, 15-30% reported autoimmune thyroiditis compared to 5-22% in children.⁸ In the non-diabetic population, 2-10% adults and 1-4% children reportedly have the condition.^{10,11} To date, serum thyroid stimulating hormone levels in T1DM patients have rarely been studied in Indonesia. Therefore, the objective of this study was to assess for a possible association between diabetes duration and thyroid stimulating hormone concentration in children with type 1 diabetes mellitus.

Methods

This cross-sectional study was carried out from January to June 2017 in the Pediatric Endocrine Outpatient

From the Department of Child Health, Airlangga University Medical School/Dr. Soetomo Hospital, Surabaya, East Java, Indonesia.

Reprint requests to: Nur Rochmah, Department of Child Health, Airlangga University Medical School/Dr. Soetomo Hospital, Jl Prof Dr Moestopo 6-8, Surabaya. Tel. +6231-5501748. E-mail: drnurrochmah@gmail.com.

Clinic (OPC) at Dr Soetomo Hospital, Surabaya, East Java. Subjects were children with T1DM aged 7 to <18 years. Exclusion criteria were diabetic ketoacidosis, previously diagnosed thyroid problems, and hospitalization in the PICU. Blood specimens were processed by an ADVIA Centaur immunoassay system, using an electrochemilluminescence immunoassay (ECLIA) method to measure TSH levels. Statistical analysis was done with Pearson's correlation test. Results with P values <0.05 were considered to be statistically significant. FT4 levels were obtained for subjects with abnormal TSH levels.

Table 1. Baseline characteristics of subjects

Characteristics	(N=34)
Male sex, n (%)	19 (55.9)
Mean body weight (SD), kg	33.34 (12.6)
Mean height (SD), cm	136.6 (16.63)
Mean body mass index (SD), Z-score	-0.68 (1.58)
Mean age (SD), years	11.3 (3.5)
Mean duration of diabetes mellitus (SD), years	3 (1.98)
Mean TSH (SD), mIU/L	3.76 (8.48)

The normal limits for FT4 were 1.0-2.1 ng/dL in 2 to 7-year-olds and 0.8-1.9 ng/dL in 8 to 20-year-olds. The normal TSH limits were 0.7-5.7 mIU/L in 2 to 7-year-olds and 0.7-5.7 mIU/L in 8 to 20-year-olds. The diagnosis of primary hypothyroidism was made in those with low FT4 and high TSH. Subclinical hypothyroidism was diagnosed in those with high TSH and normal FT4; hyperthyroidism in those with low TSH and high T3 and T4; and subclinical hyperthyroidism in those with normal TSH and high T3 and T4.¹²

Results

There were 55 T1DM patients who regularly visited our OPC. Thirty-four patients met the inclusion criteria and were included in the study.

There were 2 male patients with high TSH levels, 25 and 45 mIU/L, respectively. These patients were subsequently found to have normal FT4 levels, hence, they were diagnosed with subclinical hypothyroidism. Mean TSH was 3.76 (SD 8.48) mIU/L, ranging from




Figure 1. Association between duration of diabetes and TSH levels in children with T1DM

0.033 to 45 mIU/L. Pearson's correlation test revealed no significant association between duration of diabetes and thyroid stimulating hormone concentration ($r_s = -0.068$; $P = 0.703$).

There were 2 patients with high TSH, one with duration of illness 1 year with TSH 25 mIU/L and one patient 2 years 45 mIU/L (Figure 1).

Discussion

The mean TSH concentration in our subjects was 3.76 (SD 8.48) mIU/L. The International Society for Pediatrics and Adolescent Diabetes Mellitus recommends that screening of thyroid function by measuring thyroid stimulating hormone (TSH) and anti-thyroid peroxidase antibodies at the time of diabetes diagnosis and, thereafter, every second year in asymptomatic individuals without goiter, or in the absence of thyroid autoantibodies. More frequent assessment is indicated otherwise.¹³ Kabelitz et al. and Loviselli et al. reported the prevalence of autoimmune thyroiditis in population was 2.9-3.4%,^{14,15} while Kalaoumenou et al. reported 4.6% in Greek population.¹⁶ Subclinical hypothyroidism was found in 7-20%^{5,7} compared to adults with rates of 1-10%,^{7,8} and 2-6% in the general population of children.⁹

In our study, subjects' mean duration of diabetes was 3 years and mean age was 11.3 years, in which in the periode of puberty. Thyroid dysfunction apparent at diabetes onset^{10,12} or years thereafter.^{10,14,15} Peak of autoimmune thyroiditis incidence in the early until mid puberty.^{17,18}

Two of our male patients were diagnosed with

subclinical hypothyroidism. Females have been reported to be at risk for autoimmune thyroiditis.^{19,20} Sharifi *et al.* and Araujo *et al.* stated that gender predisposition of patients suffered from thyroid dysfunction was varied.^{20,21} Subclinical hypothyroidism is frequently observed in T1DM.^{13,21-27}

We found no significant association between duration of diabetes and TSH levels in children with T1DM. Past studies have shown that the longer the duration of diabetes, the higher the prevalence of autoimmune thyroiditis.²²⁻²⁶ A previous study reported that prevalence of autoimmune thyroiditis in T1DM patients increased post-puberty.²⁵ Another study stated that the peak prevalence of thyroid antibody was observed after the age of 15 years or a duration of diabetes of 3.5 years.²⁶

Thyroid stimulating hormone is a sensitive method to detect thyroid dysfunction. Normal TSH has a high negative predictive value to exclude thyroid disease and TSH changes can be detected earlier than FT4 changes. Ramasamy *et al.* stated that TSH > 2.2 mIU/L was predictor of hypothyroidism in T1DM, with 83% sensitivity and 72% specificity.²⁸ However, TSH is of limited value for diagnosing hypothyroidism in central hypothyroidism and acute illness. The TSH needs to be rechecked after the acute illness to distinguish between non-thyroidal illness syndrome and actual hypothyroidism.²⁹ In addition, the TSH examination is less expensive than the thyroid antibody test. Screening once every 2 years is safe, effective, cost-efficient, as well as useful for avoiding the trauma of unnecessarily frequent blood sampling.¹³

In conclusion, there is no significant association between duration of diabetes and thyroid stimulating hormone in children with T1DM.

Acknowledgement

We would like to thank all the medical staff of the Pediatric Endocrine Outpatient Clinic at Dr. Soetomo Hospital for their assistance in our study.

Conflict of Interest

None declared.

References

1. World Health Organization. Global report on diabetes. Geneva: WHO; 2016. p.21
2. International Diabetes Federation (IDF). Diabetes atlas. 7th ed. Belgium: International Diabetes Federation; 2015. [cited: 2017, June 2]. Available from: <http://www.idf.org/diabetesatlas>.
3. UKK Endokrin Anak. Unpublished data. 2016.
4. Krzewska A, Ben-Skowronek I. Effect of associated autoimmune diseases on type 1 diabetes mellitus incidence and metabolic control in children and adolescents. *BioMed Res Int*. 2016;2016:6219730.
5. Jonsdottir B, Anderson C, Carlson A, Delli A, Forsander G, Ludvigsson J, *et al.* Thyroid autoimmunity in relation to islet autoantibodies and HLA-DQ genotype in newly diagnosed type 1 diabetes in children and adolescents. *Diabetologia*. 2013;56:1735-42.
6. Denzer C, Karges B, Näke A, Rosenbauer J, Schober E, Schwab KO, *et al.* Subclinical hypothyroidism and dyslipidemia in children and adolescents with type 1 diabetes mellitus. *Eur J Endocrinol*. 2013;168:601-8.
7. Ghawil M, Tonutti E, Abusrewil S, Visentini D, Hadeed I, Miotti V, *et al.* Autoimmune thyroid disease in Libyan children and young adults with type 1 diabetes mellitus. *Eur J Pediatr*. 2011;170:983-7.
8. Van den Driessche A, Eenkhoorn V, Van Gaal L, De Block C. Type 1 diabetes and autoimmune polyglandular syndrome: a clinical review. *Neth J Med*. 2009 67:376-87.
9. Canaris GJ, Manowitz NR, Mayor G, Ridgway EC. The Colorado thyroid disease prevalence study. *Arch Intern Med*. 2000;160:526-34.
10. Wu T, Flowers JW, Tudiver F, Wilson JL, Punyasavatsut N. Subclinical thyroid disorders and cognitive performance among adolescents in the United States. *BMC Pediatr*. 2006;6:12.
11. Kordonouri O, Hartmann R, Deiss D, Wilms M, Grüters-Kieslich A. Natural course of autoimmune thyroiditis in type 1 diabetes: association with gender, age, diabetes duration, and puberty. *Arch Dis Child*. 2005;90:411-4.
12. De Boer, Lafranchi SH, 2007. Pediatric thyroid testing issues. *Pediatr Endocrinol Rev*, 5;1:570-7. In: Kappy MS, Allen DB, Geffner ME. *Pediatric practice endocrinology*. 2010: 107-30.
13. Kordonouri O, Klingensmith G, Knip M, Holl RW, Menon PSN, Aanstoot HJ, *et al.* Other complications and diabetes-associated conditions in children and adolescents. *Pediatric Diabetes* 2014; 15 (Suppl. 20): 270-8.

14. Umpierrez GE, Latif KA, Murphy MB, Lambeth HC, Stentz F, Bush A, *et al.* Thyroid dysfunction in patients with type 1 diabetes: A longitudinal study. *Diabetes Care.* 2003;26:1181-5.
15. Severinski S, Banac S, Severinski NS, Ahel V, Cvijovic K. Epidemiology and clinical characteristics of thyroid dysfunction in children and adolescents with type 1 diabetes. *Coll Antropol.* 2009;33:273-9.
16. Glastras SJ, Craig ME, Verge CF, Chan AK, Cusumano JM, Donaghue KC. The role of autoimmunity at diagnosis of type 1 diabetes in the development of thyroid and celiac disease and microvascular complications. *Diabetes Care.* 2005;28:2170-5.
17. Kim EY, Shin HC, Yang SW. Polymorphisms of HLA class II predispose children and adolescent with type 1 diabetes mellitus to autoimmune thyroid disease. *Autoimmunity.* 2003;36:177-81.
18. Warncke K, Frohlich-Reiterer EE, Thon A, Hofer SE, Wiemann D, Holl RW, *et al.* Polyendocrinopathy in children, adolescents, and young adults with type 1 diabetes: a multicenter analysis of 28,671 patients from the German/Austrian DPV-Wiss database. *Diabetes Care.* 2010;33:201-7.
19. Barker JM, Yu J, Yu L, Wang J, Miao D, Bao F, *et al.* Autoantibody "subspecificity" in type 1 diabetes: risk for organ-specific autoimmunity clusters in distinct groups. *Diabetes Care.* 2005; 28:850-5.
20. Sharifi F, Ghasemi L, Mousavinasab N. Thyroid function and anti-thyroid antibodies in Iranian patients with type 1 diabetes mellitus: influences of age and sex. *Iran J Allergy Asthma Immunol.* 2008;7:31-6.
21. Araujo J, Brandão LAC, Guimarães RL, Santos S, Falcao EA, Milanese M, *et al.* Prevalence of autoimmune thyroid disease and thyroid dysfunction in young Brazilian patients with type 1 diabetes. *Pediatr Diabetes.* 2008;9:272-6.
22. Shun CB, Donaghue KC, Phelan H, Twigg SM, Craig ME. Thyroid autoimmunity in type 1 diabetes: systematic review and meta-analysis. *Diabet Med.* 2014;31:126-35.
23. Lee YJ, Jung SY, Jung HW, Kim SY, Lee YA, Shin CH, *et al.* Unfavorable course of subclinical hypothyroidism in children with Hashimoto's thyroiditis compared to those with isolated non-autoimmune hyperthyrotropinemia. *J Korean Med Sci.* 2017;32:124-9.
24. Kakleas K, Paschali E, Kefalas N, Fotinou A, Kanariou M, Karayianni C, *et al.* Factors for thyroid autoimmunity in children and adolescents with type 1 diabetes mellitus. *J Med Sci.* 2009;114:214-20.
25. Kordonouri O, Klinghammer A, Lang EB, Gruters-Kleslich A, Grabert M, Holl RW. Thyroid autoimmunity in children and adolescents with type 1 diabetes: a multicenter survey. *Diabetes Care.* 2002;25:8:1346-50.
26. Kakleas K, Soldatou A, Karachaliou F, Karavanaki K. Associated autoimmune diseases in children and adolescents with type 1 diabetes mellitus (T1DM). *Autoimmun Rev.* 2015;14:781-97.
27. Muhame RM, Mworzi EA, Ma Assey K. Thyroid autoimmunity and function among Ugandan children and adolescents with type-1 diabetes mellitus. *Pan Afr Med J.* 2014;19:137.
28. Ramasamy V, Kadiyala R, Fayyaz F, Mallipedhi A, Baglioni P, Okosieme OE. Value of baseline serum thyrotropin as a predictor of hypothyroidism in patients with diabetes mellitus. *Endocr Pract.* 2011;17:26-32.
29. Stockigt J. Case finding and screening strategies for thyroid dysfunction. *Clin Chim Acta.* 2002;315:111-24.

Diabetes duration and thyroid stimulating hormone levels in children with type 1 diabetes mellitus

ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

17%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 www.degruyter.com 1%

Internet Source
- 2 R. Kadiyala. "Thyroid dysfunction in patients with diabetes: clinical implications and screening strategies : Thyroid dysfunction and diabetes", International Journal of Clinical Practice, 04/30/2010 1%

Publication
- 3 Shuang Wang, Wen Wang, Guojing Yu, Lin Wan, Yuying Fan, Hongjie Wang, Tong Liu, Taoyun Ji, Qingzhu Liu, Lixin Cai, Xiaoyan Liu. "Safety and efficacy of rapid withdrawal of anti - seizure medications during long - term video - electroencephalogram monitoring in children with drug resistant epilepsy: A retrospective study", Epilepsia Open, 2023 1%

Publication
- 4 Kostas Kakleas, Evangelia Paschali, Nikos Kefalas, Aspasia Fotinou, Maria Kanariou, Christina Karayianni, Kyriaki Karavanaki. 1%

"Factors for thyroid autoimmunity in children and adolescents with type 1 diabetes mellitus", Upsala Journal of Medical Sciences, 2009

Publication

5

www.mdpi.com

Internet Source

1 %

6

repository-tnmgrmu.ac.in

Internet Source

1 %

7

C. R. Prieto, H. Cardenas, H. B. Croxatto.

"Variability of breast sucking, associated milk transfer and the duration of lactational amenorrhoea", Reproduction, 1999

Publication

1 %

8

repository.unair.ac.id

Internet Source

1 %

9

Gretchen M. Oakley, Kristine A. Smith, Shaelene Ashby, Richard R. Orlandi, Jeremiah A. Alt. "Sleep Dysfunction is an Independent Predictor of Productivity Losses in Patients with Chronic Rhinosinusitis", Annals of Otolaryngology, Rhinology & Laryngology, 2021

Publication

1 %

10

Nikhil Shah, Anuradha Khadilkar, Ketan Gondhalekar, Vaman Khadilkar. "Prevalence of Dyslipidemia in Indian Children with Poorly

1 %

Controlled Type 1 Diabetes Mellitus", Pediatric Diabetes, 2020

Publication

11

ris.utwente.nl

Internet Source

1 %

12

pure.rug.nl

Internet Source

1 %

13

sciencescholar.us

Internet Source

1 %

14

"Correlation of heart failure severity and N-terminal pro-brain natriuretic peptide level in children", 'Paediatrica Indonesiana - Indonesian Pediatric Society'

Internet Source

1 %

15

Yin, Yan-Wei, Qian-Qian Sun, Bei-Bei Zhang, Ai-Min Hu, Qi Wang, Hong-Li Liu, Zhi-Zhen Hou, Yi-Hua Zeng, Rui-Jia Xu, and Long-Bao Shi.

"The lack of association between interleukin-6 gene -174 G/C polymorphism and the risk of type 1 diabetes mellitus: A meta-analysis of 18,152 subjects", Gene, 2013.

Publication

1 %

16

Lidiya Daniel, Demoz Haile, Gudina Egata.

"Disordered Eating Behaviours and Body Shape Dissatisfaction among Diabetic Adolescents: a Cross sectional study", Research Square Platform LLC, 2022

Publication

1 %

17 Ramli Ramli, Marselina Sattu, Aulia Muliathi S. Ismail, Erni Yusnita Lalusu et al. "Factors Influencing the Incidence of Stunting in Jaya Bakti Village, Pagimana District, Banggai Regency", Open Access Macedonian Journal of Medical Sciences, 2022

Publication

1 %

18 Sharath Kumar. "Wolfram syndrome: important implications for pediatricians and pediatric endocrinologists", Pediatric Diabetes, 2010

Publication

1 %

19 www.smj.org.sa

Internet Source

1 %

20 "Behavioral Diabetes", Springer Science and Business Media LLC, 2020

Publication

<1 %

21 "Thyroid Function Testing", Springer Nature, 2010

Publication

<1 %

22 G. de Sousa, S.R. Tittel, M. Häusler, P. M. Holterhus et al. "Type 1 Diabetes and Epilepsy in childhood and adolescence: Do Glutamic Acid Decarboxylase autoantibodies play a role? Data from the German/Austrian/Swiss/Luxembourgian DPV Registry", Pediatric Diabetes, 2020

Publication

<1 %

23

Konstantinos Kakleas, Lydia Kossyva, Anastasia Korona, Nikolitsa Kafassi, Spyridon Karanasios, Kyriaki Karavanaki. "Predictors of associated and multiple autoimmunity in children and adolescents with type 1 diabetes mellitus", *Annals of Pediatric Endocrinology & Metabolism*, 2022

Publication

<1 %

24

Pediatric Endocrinology, 2013.

Publication

<1 %

25

RF Afolabi, ME Palamuleni. "Exploring determinants of under-5 stunting in Malawi using a generalised linear mixed model", *South African Journal of Child Health*, 2021

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Diabetes duration and thyroid stimulating hormone levels in children with type 1 diabetes mellitus

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
