

Correlation between Fibrinogen

by Rika Ainun Tikha

Submission date: 25-Jun-2023 10:22AM (UTC+0800)

Submission ID: 2122054424

File name: Correlation_between_Fibrinogen.pdf (337.21K)

Word count: 1991

Character count: 10536

Correlation between Fibrinogen Level with Middle Cerebral Artery Pulsatility Index in Acute Ischemic Stroke

Rika Ainun Tikha¹, Sita Setyowatie², Muhammad Hamdan³, Achmad Firdaus Sani²,
Mohammad Saiful Ardhi²

¹Resident Student, ²Lecturer, Division of Neurovascular, Department of Neurology, ³Lecturer Division of Neurobehaviour, Department of Neurology, Dr. Soetomo Hospital/Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

Abstract

Background: High levels of fibrinogen will increase the blood viscosity. Pulsatility index parameter in transcranial doppler can measure blood vessel resistance which influence by cerebral blood flow velocity and blood viscosity. This study has aim to find correlation between fibrinogen with PI in acute thrombotic stroke patient.

Method: A cross sectional design was used in this study. The samples were patient with acute ischemic stroke in neurological ward Dr. Soetomo Hospital Surabaya from January 2019 to December 2020. Fibrinogen level was measured using *claus* method. PI was calculated by peak systolic velocity, end diastolic velocity and mean vessel velocity, which were measured in middle cerebral artery.

Result: This study involved 32 samples consisted 22 males and 10 females. Mean age of subjects was 55.37 ± 10.55 . The mean fibrinogen level is 356.95 ± 139.57 . Mean MCA PI is 1.03 ± 0.34 . There is positive correlation with weak correlation strength between fibrinogen level with MCA PI in acute thrombotic stroke patients and statistically significant ($p=0.024$, $r=0.398$).

Conclusion: There is a positive correlation between fibrinogen level with PI MCA in acute thrombotic stroke.

Keywords: acute ischemic stroke, fibrinogen, MCA PI

Introduction

Stroke is the second leading cause of death (11.8% of all deaths) worldwide after ischemic heart disease and the leading cause of Disability Adjusted Life Years (DALYs).¹ Globally, there were 6.5 million deaths due to stroke in 2013. Around 87% of every stroke incident were ischemic stroke.² Stroke affects all the quality of life and become a serious problem in Asia, which has more than 60% of the world population with the majority of countries are lower middle income countries.^{3,4}

Ischemic stroke occurs due to impaired cellular perfusion and metabolism which is closely related to the condition of blood flow, blood vessels and blood viscosity. High blood viscosity increases the risk of

cerebrovascular disease.⁵

Fibrinogen plays an important role for the interaction of erythrocytes and platelets in acute local inflammation condition, so that if fibrinogen levels increase, there will be an increase in erythrocyte aggregation and platelet aggregation which causes often occurs in acute stroke.^{6,7} Aside from that, fibrinogen in itself has a large molecular weight compared to its plasma proteins counterparts. These characteristics causes high levels of fibrinogen will increase blood viscosity.⁸

One of the Transcranial Doppler (TCD) parameters is the Pulsatility Index (PI) which evaluates the peripheral resistance of blood vessels.⁹ The most important factors

influencing the PI are cerebral blood flow velocity and blood viscosity.¹⁰

The purpose of this study is to determine whether fibrinogen level is correlated with the PI as both are positively correlated with the blood viscosity level concurrently.

7 Method

This study is an observational analytic study with a cross sectional design. Inclusion criteria were acute thrombotic stroke patient in acute phase, ages over than 18 years, National Institutes of Health Stroke Scale (NIHSS) < 15, and has complete medical record data. The exclusion criteria were: patients who suffers from sepsis, brain tumor, acute myocardial infarction, dysfunction of coagulation factor. Based on the formula for calculating the sample size of correlation study, the sample amount is 32.

All subjects underwent the same laboratory and clinical examinations. The fibrinogen level were measured using *claus* method with Sysmex CS2500 when patient were admitted in neurological ward. TCD examination was performed by one neurologist certified in neurosonology with SONARA TCD system (Natus,

Neurology USA). Doppler signals from the MCA were obtained at depths of 45-65 mm. PI was calculated as the difference between the peak systolic and end diastolic velocities divided by the mean flow velocity. Data were analyzed using Statistical Package for the Social Sciences (SPSS) software version 25.0. The normality of the data distribution was checked by the Kolmogorov Smirnov test. The Spearman statistical test was used to determine the correlation between the two variables with an abnormal distribution.

The research committee at the Dr. Soetomo Hospital Surabaya, approved this study ethical approval number 0180/KEPK/IV/2021.

Result

This study involved a total sample of 32 subjects. The total number of male patients were 22 (68.8%) and female patients were 10 (31.3%). The mean age was 55.37±10.55. There were no differences between with clinical manifestation among patients. Of these, right hemiparesis occurred in 17 (53.1%) and left hemiparesis in 15 (46.9%) patients. General characteristics are shown in table 1

Table 1: General Characteristics

Variable	n (%)	Mean ± SD
Sex		
Male	22 (68.8%)	
Female	10 (31.3%)	
Age		55.37 ± 10.55
Hemiparesis		
Right hemiparesis	17 (53.1%)	
Left hemiparesis	15 (46.9%)	

The mean level fibrinogen in this study was 356.96±139.58mg/dl with minimum level was 99.50 mg/dl and maximum level 723.40mg/dl. This data can be seen in table 2

Table 2 Mean value of fibrinogen level

Variable	Minimum	Maximum	Mean ± SD
Fibrinogen	99.50	723.40	356.96 ± 139.58

The result of MCA PI examination showed the mean PI value was 1.03 ± 0.34 , with minimum PI value was 0.56 and maximum PI value 2.17. This data can be seen in table 3

Table 3 Mean value of MCA PI

Variable	Minimum	Maximum	Mean ± SD
Pulsatility index	0.56	2.17	1.03 ± 0.34

CORRELATION BETWEEN FIBRINOGEN LEVEL WITH MCA PI

15 There was a positive correlation with weak correlation strength between fibrinogen level with MCA PI and statistically significant with $p < 0.05$. The correlation coefficient is 0.398, as seen in Table 4.

Table 4 Correlation between Fibrinogen Level with MCA PI

Variable	Correlation coefficient	p-value
Fibrinogen with PI MCA	0.398	0.024

Discussion

Since January 2019 until December 2020, we found 32 patients met the research criteria. Of these subject, 22 (68.8%) were males and 10 (31.3%) were females. This study shows that male suffered more acute thrombotic stroke than female. This is in accordance with several studies which state that male is more affected by stroke than woman. Unmodifiable characteristics of sex as a risk factor is making this demographic knowledge becomes an important feature. Other risk factor that also unmodifiable are age and race.¹¹

The mean age of patients was 55.37 ± 10.55 .¹⁰ This is consistent with data from other study that risk of stroke doubles every decade above 55 age.¹² While in developing countries most stroke cases occur in people less than 60 years and low awareness about the treatment approaches.¹³⁻¹⁵

The baseline data that has been collected are test for normality with Kolmogorov -Smirnov test (KS test). This test aims to determine the distribution of normal or abnormal data. Analysis using the KS test found that the distribution data was abnormal. Therefore, the analysis to determine the correlation between fibrinogen level with MCA PI in acute ischemic stroke was performed using Spearman's correlation analysis.

In this study, researchers found a significant relationship between fibrinogen level with MCA PI in acute ischemic stroke patients. The result of the p value is 0.024 (< 0.05) which is significant. This means that the higher fibrinogen level, the greater PI MCA value.

The results of this study are supported by other research which states that increased blood viscosity associated with acute ischemic stroke.¹⁶ In addition, there are other studies that state fibrinogen is a factor that

independently affects blood viscosity in a population of acute ischemic stroke patient.⁸ Other studies state that the higher the blood viscosity, the greater vascular resistance, resulting in an increase PI value.¹⁷

Conclusions and Suggestions

There was positive correlation with correlation coefficient ($r=0.398$) between fibrinogen level with MCA PI in acute ischemic stroke which was statistically significant ($p<0.05$). Further research need to be done, especially research that analyze confounding variables in blood viscosity and cerebral blood vessel imaging.

Ethical Clearance: This study received an ethical approval number.

Source of Funding: This research was carried out through individual funding.

Conflict of Interest: There was no conflict of interest from this study.

References

1. Feigin VL, Norrving B, Mensah GA. Global Burden of Stroke. *Circ Res*. 2017;120(3):439-448. doi:10.1161/CIRCRESAHA.116.308413
2. Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart Disease and Stroke Statistics'2017 Update: A Report from the American Heart Association. *Circulation*. 2017;135(10):e146-e603. doi:10.1161/CIR.0000000000000485
3. Venkatasubramanian N, Yoon BW, Pandian J, Navarro JC. Stroke epidemiology in south, east, and south-east asia: A review. *J Stroke*. 2017;19(3):286-294. doi:10.5853/jos.2017.00234
4. Ali DKA. Quality of Life of Patients with Ischemic Stroke Versus Hemorrhagic Stroke. *Medico-Legal Updat*. 2020;20(2):576-581. doi:https://doi.org/10.37506/mlu.v20i2.1172
5. Song SH, Kim JH, Lee JH, Yun YM, Choi DH, Kim HY. Elevated blood viscosity is associated with cerebral small vessel disease in patients with acute ischemic stroke. *BMC Neurol*. 2017;17(1):1-10. doi:10.1186/s12883-017-0808-3
6. Rasyid A. Peran Viskositas Darah Pada Stroke Iskemik Akut. *Maj Kedokt Neuro Sains Perhimpun Dr*. 2014;31(3). <http://www.neurona.web.id/paper/864.pdf>.
7. Saiful M. Hubungan Kadar hs-CRP dengan Keluaran Klinis Berdasarkan Indeks Barthel pada Pasien Stroke Iskemia Akut. *Aksona*. 2020;3:127-134.
8. Rasyid A, Harris S, Kurniawan M, Mesiano T, Hidayat R. Fibrinogen and LDL influence BV and Outcome of Acute Ischemic Stroke. pdf. *Ann Neurosci*. 2019;26(3-4) 30-34. doi:10.1177/0972753119900630
9. Kassab MY, Majid A, Farooq MU, et al. Transcranial doppler: An introduction for primary care physicians. *J Am Board Fam Med*. 2007;20(1):65-71. doi:10.3122/jabfm.2007.01.060128
10. Han SW, Lee SS, Kim SH, et al. Effect of cilostazol in acute lacunar infarction based on pulsatility index of transcranial doppler (ECLIPse): A multicenter, randomized, double-blind, placebo-controlled trial. *Eur Neurol*. 2013;69(1):33-40. doi:10.1159/000338247
11. A. Boehme, C. Esenwa ME. Stroke: Risk factors and prevention. *J Pak Med Assoc*. 2018;60(5):412. doi:10.1161/CIRCRESAHA.116.308398. Stroke
12. Donkor ES. Stroke in the 21st Century: A Snapshot of the Burden, Epidemiology, and Quality of Life. *Stroke Res Treat*. 2018;2018. doi:10.1155/2018/3238165
13. Connor MD, Walker R, Modi G, Warlow CP. Burden of stroke in black populations in sub-Saharan Africa. *Lancet Neurol*. 2007;6(3):269-278. doi:10.1016/S1474-4422(07)70002-9
14. Johnston SC, Mendis S, Mathers CD. Global variation in stroke burden and mortality: estimates from monitoring, surveillance, and modelling. *Lancet Neurol*. 2009;8(4):345-354. doi:10.1016/S1474-4422(09)70023-7
15. S.Kadam P, B.Kanase S. Awareness of Various

- Forms of Treatment Approaches in Stroke amongst Rural Population. *Med Leg Updat.* 2020;20(3):148-154. doi:<https://doi.org/10.37506/mlu.v20i3.1386>
16. Furukawa K, Abumiya T, Sakai K, et al. Increased Blood Viscosity in Ischemic Stroke Patients with Small Artery Occlusion Measured by an Electromagnetic Spinning Sphere Viscometer. *J Stroke Cerebrovasc Dis.* 2016;25(11):2762-2769. doi:10.1016/j.jstrokecerebrovasdis.2016.07.031
17. De Riva N, Budohoski KP, Smielewski P, et al. Transcranial doppler pulsatility index: What it is and what it isn't. *Neurocrit Care.* 2012;17(1):58-66. doi:10.1007/s12028-012-9672-6

Correlation between Fibrinogen

ORIGINALITY REPORT

15%

SIMILARITY INDEX

14%

INTERNET SOURCES

9%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	repository.sustech.edu Internet Source	1%
2	www.karger.com Internet Source	1%
3	academicjournals.org Internet Source	1%
4	thieme-connect.com Internet Source	1%
5	www.e-sciencecentral.org Internet Source	1%
6	mail.phcogj.com Internet Source	1%
7	garuda.kemdikbud.go.id Internet Source	1%
8	www.journalpulmonology.org Internet Source	1%
9	neura.edu.au Internet Source	1%

10	www.scielo.org.za Internet Source	1 %
11	mohw.go.kr Internet Source	1 %
12	www.cureus.com Internet Source	1 %
13	e-journal.unair.ac.id Internet Source	1 %
14	journals.viamedica.pl Internet Source	1 %
15	openres.ersjournals.com Internet Source	1 %
16	bmcneurol.biomedcentral.com Internet Source	1 %
17	www.bmj.com Internet Source	1 %
18	Ramesh Raj Acharya, Nikunja Yogi, Susmita Khadka, Dipak Kunwar, Rajyashree Kunwar, Rakshya Shrestha, Adhyashree Karki. "Clinical Profile and Risk Factors of Patients in Stroke: A Cross-Sectional Study", Eastern Green Neurosurgery, 2020 Publication	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Correlation between Fibrinogen

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
