

Source details

Sri Lanka Journal of Child Health Open Access ①	CiteScore 2021 0.3	Û
Scopus coverage years: from 2011 to 2022		
Publisher: Sri Lanka College of Paediatricians	SJR 2021	(j)
ISSN: 1391-5452 E-ISSN: 2386-110X	0.136	
Subject area:		
Source type: Journal	SNIP 2021	(i)
	0.239	Ŭ
View all documents > Set document alert Save to source list Source Homepage		

CiteScore CiteScore rank & trend Scopus content coverage

Г			×
	i	Improved CiteScore methodology	
		CiteScore 2023 counts the citations received in 2020-2023 to articles, reviews, conference papers, book chapters and data	
		papers published in 2020-2023, and divides this by the number of publications published in 2020-2023. Learn more $>$	
L			



CiteScore rank ①

Category

Rank Percentile

View CiteScore methodology \succ CiteScore FAQ \succ Add CiteScore to your site ${\rm g}^{\rm O}$

Q




Sri Lanka Journal of Child Health 8

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Sri Lanka	Medicine Pediatrics, Perinatology and Child Health	Sri Lanka College of Paediatricians	7
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	13915452, 2386110X	2011-2021	Homepage How to publish in this journal bjcp@ymail.com


SCOPE

This is the only journal of child health in Sri Lanka. It is designed to publish original research articles and scholarly articles by recognized authorities on paediatric subjects. It is distributed widely in Sri Lanka and bears the ISSN number 1391-5452 for the print issues and e-ISSN 2386-110x for the electronic version in the internet. The journal is published quarterly and the articles are reviewed by both local and foreign peers. The Journal is the primary organ of Continuing Paediatric Medical Education in Sri Lanka.

 $\ensuremath{\bigcirc}$ Join the conversation about this journal





Metrics based on Scopus® data as of April 2022




Home	About Contac	t Content		Q Log in Register				
	-	tion on serum evels in children om severe acute , A. V. y,	 - 2018 Volume 47 - Issue 1 - 2018 Volume 47 - Issue 1 - 2018 Volume 46 - Issue 4 - 2017 Volume 46 - Issue 3 - 2017 Volume 46 - Issue 3 - 2017 					
	Association be anthropometri and carotid in thickness in o adolescents Nur Aisiyah Wi Aji Prihaningty Herdiana Hani Roedi Irawan, Retno Handaja G. Ugrasena	ric parameters tima-media bese idjaja, Rendi yas, Meta ndita,	(*) ** 05 Mar 2021 50(1):28-31	- 2017 Volume 46 - Issue 1 - 2017 Volume 45 - Issue 4 - 2016 Volume 45 - Issue 3 - 2016 Volume 45 - Issue 2 - 2016 Volume 45 - Issue 1 - 2016				
	Indian childre Idiopathic Art sectional stud	, Ajanta Haldar,	• * 05 Mar 2021 50(1):32-37	Volume 44 - Issue 4 - 2015 Volume 44 - Issue 3 - 2015 Volume 44 - Issue 2 - 2015 Volume 44 - Issue 1				
	Study of glyca and microalbu children with mellitus Ujjwala Manth Shrikiran Aroo G. Kini, Leslie Shravan Kanap Ramesh Bhat	uminuria in type 1 diabetes a, or, Pushpa Edward Lewis,	(*************************************	Volume 43 - Issue 4 - 2015 Volume 43 - Issue 4 - 2014 Volume 43 - Issue 3 - 2014 Volume 43 - Issue 2 - 2014 Volume 43 - Issue 1 - 2014				
	The agreemer cytomegalovi serology exan CMV polymera reaction of liv infants with cl Bagus Setyobo	rus (CMV) nination and ase chain er tissue in holestasis	(*) ** 05 Mar 2021 50(1):43-48	- 2014 Volume 42 - Issue 4 - 2013 Volume 42 - Issue 3 - 2013 Volume 42 - Issue 2 - 2013				


Home	About Contact Content		Q Log in Register
	Dwiyanti Puspitasari, Rendi Aji Prihaningtyas		Volume 41 - Issue 4 - 2012
	'Don't judge by outer	۲	Volume 41 - Issue 3 - 2012
	promotion to children through food packages in the	05 Mar 2021 50(1):49-55	Volume 41 - Issue 2 - 2012
	Sri Lankan market Manori Gamage		Volume 41 - Issue 1 - 2012
	Baby oral health promotion	۲	Volume 40 - Issue 4 - 2011
	centre at a tertiary care hospital in India: A successful model for prevention of early	05 Mar 2021 50(1):56-62	Volume 40 - Issue 3 - 2011
childhood cari Indira Mysore I D. Narayanappa Nandlal Bhojraj	childhood caries Indira Mysore Devraj,		Volume 40 - Issue 2 - 2011
	D. Narayanappa, Nandlal Bhojraj		Volume 40 - Issue 1 - 2011
	Complementary feeding	۲	Volume 39 - Issue 4 - 2010
	implementation in Pintu Padang Health Centre, South Tapanuli District, Indonesia:	05 Mar 2021 50(1):63-68	Volume 39 - Issue 3 - 2010
	a qualitative study Cecep Triwibowo,		Volume 39 - Issue 2 - 2010
	Rusdi Rangkuti, . Ridesman, Niniek Lely Pratiwi, . Tinah,		Volume 39 - Issue 1 - 2010
	Lita Heni Kusumawardani, Yulia Fauziyah		Volume 38 - Issue 4 - 2009
	Urinary tract infections and vesolution of renal pelvic	۲	Volume 38 - Issue 3 - 2009
	dilatation in infants Sajida Abdulla, Peter Vazhayil,	05 Mar 2021 50(1):69-74	Volume 38 - Issue 2 - 2009
	Hariprasad Pallippurath Gopalakrishna		Volume 38 - Issue 1 - 2009
	Pillai, Manjula Pallippurath Gopalakrishna Divak		Volume 37 - Issue 4 - 2008
	The association of iron	۲	Volume 37 - Issue 3 - 2008
	deficiency anaemia with simple febrile seizures in	05 Mar 2021 50(1):75-82	Volume 37 - Issue 2 - 2008
	children less than 5 years age: A single centre,		Volume 37 - Issue 1 - 2008
	prospective, case-control study		Volume 36 - Issue 4 - 2007









Home	About	Contact	Content		Q	Log in	Register
	O RIAL BOAL surnames):	<u>RD MEMBERS</u> :	6 (in alphat	oetical order			
MS, F	RCS.	<u>Abeygunaseka</u> ssed as Dr. Sa					
Specia	alist Consult	hotmail.com tant Paediatri al for Childrer	0,	•			
💿 of	RCID:						
		archer with 09 ic journals, as	-	-			
Hepar Hypo comp incon Mitro (prev	tobiliary dis spadias and lex, Surgica tinence (Aug fanoff proce ention and e	and expertise: ease Esp Bilia bladder extro l managemen gmentation cy edures), Paedi early intervent ka Forum of N					
MSc I		<u>a Milani Atap</u> diff) MD (Colo (UK)					
Comr	nonly addre	ssed as Profe s	ssor Piyush	a Atapattu			
Profe	ssor in Phys	tapattu@yah iology, Facult ombo, Sri Lan	y of Medici	ne,			
💿 of	RCID: 0000-	0002-8252-	5446				
19 pu		archer and Me peer-reviewe 22.					





Home	About	Contact	Content		Q	Log in	Register
2022.					•		
with a breast and yo Neona care, I Infant of hig	special int feeding an oung child atal behavio nfant body developm h-risk neo er, Sri Lank	•	erm nutrit ng challeng responsive ent, Develo n, evelopmen	tion, ges, infant feeding, opmental tal outcome			
-		<u>h Mettananda</u> on), FRCP(Ed		-			
Metta E-mai Chair I Univer Consu	nanda ll: sachithm Professor of rsity of Kela	ssed as Profe ettananda@{ f Paediatrics, niya, Sri Lanl atrician, Color a, Sri Lanka	gmail.com Faculty of <i>I</i> ka and Hon	Medicine, orary			
© or	CID: 0000-	0002-0760-	0418				
		rcher with 65 c journals as	-	-			
-		and expertises tology, Rare I		liatrics,			
Memb	er, Sri Lank	a Forum of M	Iedical Edit	ors (SLFME)			
-	essor Heman 1), FRCPsyc	<u>mali Niranjalo</u> h.	<u>ı Perera</u> MB	BS, MD			
Comm	only addre	ssed as Profe	ssor Hemai	mali Perera			



Home About Contact Content	Q Log in Register
Professor in Community Medicine, Faculty of Medicine, University of Colombo, Sri Lanka Member, Sri Lanka Forum of Medical Editors (SI	
ORCID: 0000-0002-0760-0418	
A renowned researcher with over 70 publications peer-reviewed scientific journals as of Septembe 2022.	
Special interests and expertise: Maternal and Ch Health, Public Health Nutrition, Medical Statistic	
Member, Sri Lanka Forum of Medical Editors (SI	LFME)

PANEL OF INTERNATIONAL ADVISORS (in alphabetical order of the surnames):	
1. Professor Zulfiqar Bhutta E-mail: zulfiqar.bhutta@aku.edu Husein Laljee Dewraj Professor and the Foundin Chair of the Division of Women and Child Health Khan University, Karachi, Pakistan.	•
2. Dr. Timothy L Chambers E-mail: TimothyLChambers@physicians.ie Consultant Paediatrician, Bristol, UK.	
3. Professor P. T. Chandrasoma E-mail: ptchandr@usc.edu Professor of Pathology, Keck School of Medicine University of Southern California, Los Angeles, California, USA.	2,
4. Professor E. A. S. Nelson E-mail: tony-nelson@cuhk.edu.hk Professor (Clinical), Department of Paediatrics, ' Chinese University of Hong Kong.	The
5. Professor Sir Terence Stephenson E-mail: tstephenson@gmc-uk.org Nuffield Professor of Child Health, Institute of C Health, UCL Current Chair, General Medical Council, UK	hild

Home	About	Contact	Content			Q	Log in	Register
	resident, R 2009-201	oyal College o 12	of Paediatric	s and Child			I	
E-ISSN: 23	86-110X	Published by	Sri Lanka Col	lege of Paediatri	cians	Privad	cy Policy	

The agreement of cytomegalovirus (CMV) serology examination and CMV polymerase chain reaction of liver tissue in infants with cholestasis

*Bagus Setyoboedi¹, Reny Widayanti¹, Sjamsul Arief¹, Dwiyanti Puspitasari¹, Rendi Aji Prihaningtyas¹

Sri Lanka Journal of Child Health, 2021; 50(1): 43-48

Abstract

Introduction: The incidence of cholestasis in infancy is associated with congenital or viral infection. Commonest cause is *Cytomegalovirus* (CMV) infection. In Indonesia, diagnosis of cholestasis due to CMV infection is still not optimal because the support of virological examination is still limited, so serological examination is an option.

Objective: Analysing the agreement of CMV IgG or IgM serological examination and CMV polymerase chain reaction (PCR) examination of liver tissue in infants with cholestasis

Method: This is an analytic observational research using cross sectional design. Samples were infants with cholestasis, 1-12 months of age, in the Department of Child Health, Dr. Soetomo General Hospital, who met the inclusion criteria. Samples were taken from December 2017- May 2018. Liver biopsy was done in Department of Child Health, Dr. Soetomo Hospital. CMV PCR examination of liver tissue was carried out in the Tropical Disease Laboratory at Airlangga University. Data were analysed by calculating the kappa coefficient CMV IgG or IgM serology examination and CMV PCR examination of liver tissue.

Results: There were 30 infants with cholestasis, consisting of 16 (53%) boys and 14 (47%) girls. Positive CMV IgG was found in 28 subjects and CMV IgM was found in 15 subjects. CMV PCR was positive in 20 (67%) subjects. The most common age range with positive CMV IgG and IgM as well as CMV PCR was 1-3 months. Agreement of CMV

¹Airlangga University, Indonesia *Correspondence: bagus.setyoboedi@fk.unair.ac.id

D https://orcid.org/0000-0002-3923-6913

(Received on 26 February 2020: Accepted after revision on 20 March 2020)

The authors declare that there are no conflicts of interest

Personal funding was used for the project.

Open Access Article published under the Creative

Commons Attribution CC-BY Commons Attribution

IgG compared to CMV PCR of liver tissue showed weak result with kappa value of 0.25. The agreement of CMV IgM compared to CMV PCR of liver tissue was not found with kappa value 0.00.

Conclusions: In infants with cholestasis, there is no agreement for CMV serological examination with CMV PCR of liver tissue. CMV serologic examination cannot replace CMV PCR of liver tissue to determine the presence of CMV virus in infant with cholestasis.

DOI: http://dx.doi.org/10.4038/sljch.v50i1.9401

(Key words: cholestasis, CMV serology, CMV PCR liver tissue, agreement, infant).

Introduction

Cholestasis is a decrease or obstruction of bile flow at each level to the extrahepatic biliary tract and duodenum¹. Cholestasis is found in 1 in 2500 births². Incidence of cholestasis in infancy is associated with congenital or viral infection. Commonest cause is Cytomegalovirus (CMV) infection. Research conducted in Brazil on 76 infants with cholestasis showed 29.4% positive for CMV serology in intrahepatic cholestasis and 28.5% positive for CMV serology in extrahepatic cholestasis³. CMV infection is diagnosed in neonates by isolating the virus from urine, detecting IgM in blood, detecting CMV antigen in blood and identifying CMV-DNA by polymerase chain reaction (PCR)⁴. CMV-DNA can be taken from blood or tissue⁵.

Research conducted in Brazil showed a positive frequency of CMV-DNA of 34.3% of 35 samples of liver tissue examined⁶. Research in Poland in nine children with cholestasis showed 9 CMV IgG positive, 4 CMV IgM positive, and 9 CMV DNA positive in the blood but on immunohistochemistry and molecular examination in liver biopsy tissue, CMV was not found⁷. Research in Brazil showed that in comparison with PCR, serology had a low sensitivity (33.3%) and a low positive predictive value (28.6%)⁸. In Indonesia, diagnosis of cholestasis due to CMV infection is still not optimal because the support of virological examination is an option.

Objectives

Analysing the agreement of CMV IgG or IgM serological examination and CMV PCR examination of liver tissue in infants with cholestasis

Method

This is an analytic observational research using cross sectional design. Subjects were 1-12 months old infants with cholestasis in the Department of Child Health, Dr. Soetomo General Hospital, who met the inclusion criteria. Samples were taken during a 6 month period from December 2017-May 2018.

Ethical issues: Ethical approval was obtained from the Ethics Committee of Dr. Soetomo General Hospital (ERC No. 729/Panke. KKE/XII/2017). Parental written informed consent was obtained prior to data collection.

Total sampling was performed in this study. Inclusion criteria were infants with cholestasis 1-12 months of age. The exclusion criteria were septicaemia, HIV infection, miliary tuberculosis, severe malnutrition, thrombocytopenia, prolonged clotting time, ascites and a history of using immunosuppressants including and steroids cytostatic agents. Liver biopsy was carried out in the Department of Child Health, Dr. Soetomo General Hospital. Serology examination was carried out in the clinical pathology laboratory in the Dr. Soetomo General Hospital. CMV PCR examination of liver tissue was carried out in the Tropical Disease Laboratory at Airlangga University. Descriptive statistics and calculating the kappa coefficient CMV IgG or IgM serology examination and CMV PCR examination of liver tissue were done.

Results


During the 6 month period, 33 subjects were sampled, 30 subjects met the inclusion criteria and 3 subjects were excluded, 2 with severe infection (sepsis) and 1 with ascites. The basic characteristics of the sample are shown in Table 1. The average age in this study was 3.6 months.

Based on the results of CMV serology, out of 30 study subjects, CMV IgG was positive in 28 patients, and CMV IgM was positive in 15 patients. Age distribution based on CMV serology results shows that the most common age range with CMV IgG and CMV IgM positive is at 1-3 months of age (Figure 1).


Table 1: Basic characteristics of the sample (n=30)

Variable	Result
Sex - n (%)	
Male	16 (53)
Female	14 (47)
Age in months - n (%)	
1-3	20 (66.7)
4-6	06 (20.0)
7-9	02 (06.7)
10-12	02 (06.7)
AST in IU – Mean (SD)	221.7 (153.3)
ALT in IU - Mean (SD)	161.4 (86.5)
Direct bilirubin in mg/dl Mean (SD)	8.16 (4.51)
Total bilirubin in mg/dl Mean (SD)	11.11 (5.77)

AST: Aspartate transaminase, ALT: Alanine transaminase



Based on CMV PCR results of liver tissue, CMV PCR was positive in 20 (67%) and CMV PCR was negative in 10 (33%). The most positive CMV PCR was found in the 1-3 month age range in 15 (75%) subjects (Figure 2).



From CMV serologic examination compared with CMV PCR liver tissue was obtained, from 28 subjects with positive CMV IgG, CMV PCR positive liver tissue as much as 20 (71.4%). Based on CMV IgM results compared to CMV PCR liver tissue, from 15 subjects with positive CMV IgM, positive CMV PCR of liver tissue as many as 10 (66.7%) subjects (Table 2).

	Variable	PCR	Total	
		Negative	Positive	
I-C CMV	Negative	02	0	02
IgG CMV	Positive	08	20	28
I-M CMV	Negative	05	10	15
IgM CMV	Positive	05	10	15

Table 2: Results between CMV serology and CMV PCR liver tissue

The agreement of CMV IgG compared to CMV PCR of liver tissue obtained a weak agreement with kappa value of 0.25. The agreement of CMV IgM

compared to CMV PCR of liver tissue was not in agreement with a kappa value of 0.00 (Table 3).

Table 3: Results o	f kappa suitability	between CMV	serology and CMV	PCR liver tissue based on age

Age	IgG CMV with PCR CMV	IgM CMV with PCR CMV
	Карра	Карра
1-3 months	0.273	0.053
1-4 months	0.362	0.015
1-6 months	0.272	0.031
1-12 months	0.250	0.000

In terms of age, at 1-3 months and 1-4 months of age, CMV IgG compared to CMV PCR liver tissue had a weak agreement with kappa values of 0.273 and 0.362. At the age of 1-6 months also CMV IgG compared to CMV PCR liver tissue had a weak agreement with kappa value 0.272 (Table 3).

Based on the sensitivity test and specificity test IgG CMV against CMV PCR, sensitivity was 100%, specificity was 20%, positive predictions were 71.4%, negative predictions were 100% and accuracy was 73.3%. Whereas for IgM CMV against CMV PCR, sensitivity was 50%, specificity was 50%, positive predictions were 66.7%, negative predictions were 33.4% and accuracy was 50%.

Discussion

From this study, the age range in as many as 20 (66.7%) subjects, was 1-3 months. The mean age of the study population was 3.6 months. In research conducted in Brazil, the median age of subjects at the first visit was 82.5 days $(\pm 3 \text{ months})^6$. In a study conducted in India, in 101 infants with neonatal cholestasis, the mean age of the subjects was 2.8 \pm 1.7 months⁹. In another study, the average age at time of liver biopsy was 2.5 ± 1 month¹⁰. In this study there were 53% males. Studies by Liberek (2002), Brandao (2009) and Goel (2018) also showed greater male prevalence^{8,10,11}. In this study, the mean AST and ALT values were 221.7 mg/dl and 161.4 mg/dl respectively. A study by Liberek (2002) had AST 156 mg/dl / ALT 169 mg/dl, a study by Talachian (2014) had AST 280 mg/dl / ALT 238 mg/dl and a study by Sira (2016) had AST 134 mg/dl / ALT 230 mg/dl^{11,12,13}.

From the 30 study subjects, CMV IgG was positive in 28 (93%) and CMV IgM was positive in 15

(50%). Research in Brazil on 76 subjects with neonatal cholestasis was divided into 2 groups, intrahepatic cholestasis (34) and extrahepatic cholestasis (42); in the intrahepatic cholestasis group, CMV IgM was positive in 29.4%, while in the extrahepatic cholestasis group, CMV IgM was positive in 28.5%³. In the study conducted in Poland, all 9 subjects with cholestasis were CMV IgG positive and 4 were CMV IgM positive⁷. In an African study of 78 infants with prolonged jaundice, 72 (92.3%) were CMV IgG positive and 31 (39.7%) were CMV IgM positive; the 92.3% positive CMV IgG is likely due to maternal IgG transfer via the placenta or IgG production because the baby is infected, even though 39.7% positive for IgM in the baby, presumably because there was a possibility of infection in the uterus¹⁴. CMV IgG is useful to determine whether the patient has been infected with CMV in the past, whilst IgM antibodies indicate an acute condition or new infection⁵.

In the current study, the highest number of positive CMV IgG and IgM subjects were found in the 1-3 month age range. Research in China showed that primary infection with CMV in children occured in the age range 1-3.5 months and that there was very high anti-CMV IgG seroprevalence in young infants (<6 months) due to persistent maternal antibodies¹⁵. Generally, moderate to high levels of IgM antibodies can be detected at 1-3 months after the onset of infection, after which IgM levels begin to fall but remain positive for more than 9 months after primary infection¹⁶.

In our study of 30 subjects, 20 (67%) were liver tissue CMV PCR positive. These results indicate high CMV frequency in infants with cholestasis. In a study in Taiwan, 26 liver tissue subjects with biliary atresia were positive for HCMV DNA in liver tissue with PCR of 2 $(7.6\%)^{17}$. In research from Brazil, from 33 liver tissue in subjects with extrahepatic cholestasis, 9 (27.3%) subjects with PCR were positive for HCMV DNA⁶. Research in India, from 31 subjects with neonatal cholestasis, found positive CMV PCR liver tissue with PCR in 16 $(52\%)^{10}$. Variations in CMV PCR results in liver tissue may be explained because of the few patients involved in study as well as the geographic variation of subjects¹⁸.

CMV PCR positive liver tissue was mostly found in the 1-3-month age range. Research in India on 31 infants with neonatal cholestasis, where the average age at the time of liver biopsy was 2.5 ± 1.0 months, positive CMV PCR was 16 (52%)¹⁰. CMV serological examination compared with CMV PCR of liver tissue, from 28 subjects with positive CMV IgG, 20 positive PCR (71.4%) subjects. CMV IgG compared to CMV PCR obtained a weak agreement with a kappa value of 0.25. Research conducted in India 26 CMV IgG positive subjects, 15 (58%) positive CMV PCR liver tissue with a kappa 0.2 value¹⁰. In the Brazilian study, 9 CMV IgG positive subjects, 3 subjects were positive for HCMV DNA. CMV IgG represents the presence of transplacental maternal IgG⁶.

CMV IgM results compared to CMV PCR of liver tissue, from 15 subjects with positive CMV IgM, 66.7% positive PCR subjects. In this study, CMV IgM compared with CMV PCR did not obtain agreement with kappa value 0.00. Research in Poland, in 9 children with cholestasis, obtained a weak correlation between positive serological results and virus detection in liver tissue⁷. Other studies have shown a positive association between CMV IgM and CMV PCR in liver tissue even though the relationship was very weak. It is suspected that CMV IgM is a poor indicator of the presence or absence of CMV in liver tissue¹⁰. In infants with CMV serology positive but liver tissue PCR negative, possibly at the time of the study, subjects passed the acute phase of hepatitis due to CMV where the CMV virus was eliminated from liver tissue by the host immune system or CMV infection may be outside liver tissue or hepatitis caused by other aetiologies. Infants with positive HCMV DNA with PCR on liver tissue but negative CMV serology may not produce enough antibodies for some unknown reason¹⁷.

This study showed that at the age of 1-3 months, 1-4 months, 1-6 months and 1-12 months, CMV IgG compared to CMV PCR liver tissue, obtained a weak agreement. There was no agreement obtained in CMV IgM in that age range. Other studies have

shown that serological examination has weak accuracy to detect active CMV virus infections⁶. There was weak correlation between positive serology and detection of CMV virus in liver tissue⁷. CMV blood serology or PCR is not an accurate marker for CMV infection in liver tissue¹⁰. CMV IgM and CMV IgG serology examination is not recommended for diagnosing CMV infection in infants. CMV IgG positivity reflects IgG antibodies obtained from the mother through the placenta and not diagnosis of CMV infection. Maternal IgG antibodies can last up to 18 months in infants. IgG antibodies that remain in infants cannot distinguish congenital, natal or postnatal infections¹⁹.

In this study, CMV IgM sensitivity to CMV PCR liver tissue was 50%, specificity 50%, accuracy 50%, positive predictive value 66.7% and negative predictive value 33.4%. Other studies also showed CMV IgM sensitivity to CMV PCR liver tissue was 69% and specificity was 61%¹⁰. Research in Brazil showed CMV PCR of liver tissue compared to serology with sensitivity of 54%, specificity of 61% and accuracy of 59%⁶. Compared to PCR, serology has a high accuracy of 82.4%, sensitivity of 33.3%, specificity of 88.9%, positive predictive value of 28.6%, and negative predictive value of 90.01%8. Examination of CMV IgM antibodies for primary infection has less specificity, because IgM can be positive in active CMV infection⁵. In patients with positive CMV IgM but negative DNA PCR, results are likely to be due to false positive results from IgM. False positives can be secondary to the presence of rheumatoid factors (infant IgM against maternal IgG) or cross-reaction with herpesviruses. In patients with IgM negative but positive PCR DNA, results are likely to be due to false negative. False negative results are likely secondary due to competition between high maternal IgG antibody levels and relatively low infant IgM levels⁶.

One limitation of this study is that the biopsy sample was taken with a special needle biopsy on the liver tissue so that the sample obtained was not as good as if the sample was taken during surgery. Ideally, samples should be taken from the bile duct tissue during Kasai surgery. Another drawback of this study is not to examine the types of intrahepatic or extrahepatic cholestasis. The results of this study found no agreement of CMV serological examination and CMV PCR liver tissue in infant cholestasis.

Conclusions

In infants with cholestasis, there was no agreement for CMV IgM examination with CMV PCR of liver tissue. However, for CMV IgG with CMV PCR of liver tissue, there was low agreement. Serologic examination of CMV IgG or IgM is not sensitive and specific, and cannot replace CMV PCR of liver tissue to detect the presence of CMV virus in infant cholestasis.

Acknowledgements

The authors thank to Dr. Soetomo General Hospital, Surabaya for supporting this study.

References

- Girard M, Lacaille F. Diagnosis of neonatal cholestasis. *Annales Nestle* 2008; 66: 109-20. https://doi.org/10.1159/000147408
- Suchy FJ. Neonatal cholestasis. *Pediatrics* in Review 2004; 25(11): 388-96. https://doi.org/10.1542/pir.25-11-388 PMid: 15520084
- Oliviera NL, Kanawaty FR, Costa S, Hessel G. Infection by cytomegalovirus in patients with neonatal cholestasis. *Arquivos de Gastroenterologia* 2002; **39**(2): 132-6. https://doi.org/10.1590/S00042803200200 0200012 PMid: 12612719
- Bhatia P, Narang A, Minz RW.. Neonatal cytomegalovirus infection: diagnostic modalities available for early disease detection. *Indian Journal of Pediatrics* 2010; 77: 77-9. https://doi.org/10.1007/s12098-009-0255-2
 PMid: 19936660
- Ross S, Novak Z, Pati S, Boppana S. Diagnosis of cytomegalovirus infections. *Infectious Disorders Drug Targets* 2011; 11: 466-74. https://doi.org/10.2174/187152611797636 703 PMid: 21827433 PMCid: PMC3730495
- De Tommaso AM, Andrade PD, Costa SC, Escanhoela CA, Hessel G. High frequency of human cytomegalovirus DNA in the liver of infants with extrahepatic neonatal cholestasis. *BMC Infectious Diseases* 2005; 5: 108. https://doi.org/10.1186/1471-2334-5-108 PMid: 16321152 PMCid: PMC1315325
- 7. Pawłowska JE, Dorota G, Irena J, Przemysław K, Bożena C, Katarzyna D..

The role of cytomegalovirus infection in pathogenesis of neonatal cholestasis. *Experimental and Clinical Hepatology* 2010; **6**: 25-9.

- Brandao MA, Andrade PD, Costa S, Escanhoela C, Vassallo J, Porta G, *et al.*. Cytomegalovirus frequency in neonatal intrahepatic cholestasis determined by serology, histology, immunohistochemistry, and PCR. *World Journal of Gastroenterology* 2009; 15: 3411-6. https://doi.org/10.3748/wjg.15.3411 PMid: 19610143 PMCid: PMC2712903
- Poddar U, Thapa BR, Das A, Bhattacharya A, Rao KLN, dan Singh K. Neonatal cholestasis: differentiation of biliary atresia from neonatal hepatitis in a developing country. *Acta Pædiatrica* 2009; **98**: 1260-4. https://doi.org/10.1111/j.16512227.2009.0 1338.x

PMid: 19469771

- Goel A, Chaudhari S, Sutar J, dan Bhonde G. Detection of cytomegalovirus in liver tissue in infants with neonatal cholestasis. *Pediatric Infectious Disease Journal* 2018; **37**(7): 632-6. https://doi.org/10.1097/INF.0000000000 01889
 - PMid: 29389827
- 11. Liberek A, Rytlewska M, Sidorkiewicz AS. Cytomegalovirus disease in neonates and infants - clinical presentation, diagnostic and therapeutic problems - own experience. *Medical Science Monitor* 2002; **8**: 815-20.
- Talacian E, Bidari A, Mehrazma M, Nickkhah N. Biopsy-driven diagnosis in infants with cholestatic jaundice in Iran. *World Journal of Gastroenterology* 2014; 20(4): 1048-53. https://doi.org/10.3748/wjg.v20.i4.1048 PMid: 24574777 PMCid: PMC3921528
- Sira MM, Sira AM, Elhenawy IA, Khalil FO. Prevalence of serological markers of TORCH infections in biliary atresia and other neonatal cholestatic disorders. Available from: https://www.peertechz.com/articles/OJPC H-2-110.php
- 14. Goedhals D, Kriel J, Hertzog MI, Janse van Rensburg MN. Human cytomegalovirus

infection in infants with prolonged neonatal jaundice. *Journal of Clinical Virology* 2008; **43**: 216-8. https://doi.org/10.1016/j.jcv.2008.06.001 PMid: 18619898

- Chen J, Hu L, Wu M, Zhong T, Zhou Y-H, Hu Y. Kinetics of IgG antibody to cytomegalovirus (CMV) after birth and seroprevalence of anti-CMV IgG in Chinese children. *Virology Journal* 2012;
 Article number 304(2012). https://doi.org/10.1186/1743-422X-9-304 PMid: 23228149 PMCid: PMC3544651
- 16. Revello MG, Gerna G. Diagnosis and management of human cytomegalovirus infection in the mother, fetus, and newborn infant. *Clinical Microbiology Reviews* 2002; 15: 680-715. https://doi.org/10.1128/CMR.15.4.680-715.2002
 PMid: 12364375 PMCid: PMC126858

- Chang MH, Huang HH, Huang ES, Kao CL, Hsu HY, Lee CY. Polymerase chain reaction to detect human cytomegalovirus in livers of infants with neonatal hepatitis. *Gastroenterology* 1992; 103: 1022-5. https://doi.org/10.1016/00165085(92)9003 8-Z
- El-Araby HA, Ghoneim EM, El-Rahman MH. Cytomegalovirus frequency in neonates with biliary atresia. *Egyptian Journal of Medical Microbiology* 2010; 19(2), 25-32.
- Stehel EK, Sánchez PJ. Cytomegalovirus infection in the fetus and neonate. *NeoReviews* 2005; 6: e38-e45. https://doi.org/10.1542/neo.6-1-e38