

18. Pediatric Biliary Atresia Prenatal and Postnatal Risk Factors

by Dina Aristiya Sumarno

Submission date: 08-May-2023 10:25AM (UTC+0800)

Submission ID: 2087008684

File name: ediatric_Biliary_Atresia_Prenatal_and_Postnatal_Risk_Factors.pdf (726.28K)

Word count: 2868

Character count: 15141

Pediatric Biliary Atresia: Prenatal and Postnatal Risk Factors

Dina Aristiya Sumarno¹, Sjamsul Arief², Bagus Setyoedi³

¹Resident in Department of Child Health, Faculty of Medicine, Universitas Airlangga/ Dr. Soetomo General Hospital, Surabaya, Indonesia, ²Lecturer and Consultant in Paediatric Hepatology Division Department of Child Health, Faculty of Medicine, Universitas Airlangga/ Dr. Soetomo General Hospital, Surabaya, Indonesia, ³Lecturer and Consultant in Paediatric Hepatology Division Department of Child Health, Faculty of Medicine, Universitas Airlangga/ Dr. Soetomo General Hospital, Surabaya, Indonesia

Abstract

Background: Biliary atresia is the most common cause of cholestasis in infants caused by intrauterine and neonatal inflammatory process leads to progressive biliary tract obstruction and fibrosis. Risk factors are expected to contribute in the disease type. Many studies have been done to determine the risk factors of biliary atresia. Yet, there is still limited information about risk factors in Indonesia. This study aims to determine the risk factors contributing to biliary atresia in East Java, Indonesia.

Methods: This cross-sectional study collected 219 medical record data of cholestasis patients during January 2010 to April 2017 that met the inclusion criteria. 85 of them were diagnosed with biliary atresia. Selected prenatal and postnatal risk factor were evaluated and analyzed.

Result: There were significant results in maternal age on pregnancy ($p= 0.009$), parity ($p= 0.035$), and gestational age ($p= 0.005$) among 85 patients which were diagnosed with biliary atresia. The regression test showed significant result on those 3 factors.

Conclusion: Maternal age on pregnancy, parity, and gestational age are the prenatal and postnatal risk factors that contribute to the biliary atresia incidence. These results are expected to be used as one of the providing information regarding prenatal and postnatal risk factors for pediatric biliary atresia in East Java, Indonesia

Keyword Biliary Atresia, Risk Factor, Prenatal, Postnatal

Background

Biliary atresia is well known as a progressive obstructive cholangiopathy affecting biliary tract, both in the intrahepatic and extrahepatic tract, resulting in cirrhosis, liver failure, and death if it is not well treated. It also becomes the most common cause

of pediatric cholestasis besides infection and α -1 antitrypsin deficiency¹. Despite the poor prognosis and complications in delayed diagnosis, the clear etiology and pathophysiology are remained unknown. The incidences of biliary atresia are reported around the world. Biliary atresia seems to be more commonly occurred in Asia than Europe or America, with the incidence approximately 1:8000 in Asia compared to 1: 15,000-20,000 in Europe or America^{2,3}. Another study also stated that Asia-pacific region (1:3300) has higher biliary atresia incidence than Europe (1:18000)⁴.

The initial findings of this disease are quite similar to other condition like physiological neonatal icteric, thus the diagnosis is likely to be delayed and the

Corresponding author:

Bagus Setyoedi, MD

Lecturer and Consultant in Paediatric Hepatology Division Department of Child Health, Faculty of Medicine, Universitas Airlangga/ Dr. Soetomo General Hospital, Surabaya, Indonesia

prognosis becomes worse. There are two major forms of biliary atresia, embryonic form, which is related to other congenital disorders, and perinatal form, where normal bile duct underwent fibro-obliteration during perinatal period³. Some risk factors, including environment, prenatal, and postnatal risk factors, play significant role to determine the type and form of biliary atresia.

Many researches around the world have been done to describe the risk factors of biliary atresia incidences. Previous study reported that genetical factor or chromosome alteration, viral-induced immune dysregulation, maternal disease including gestational diabetes, are several factors that affect the incidence of biliary atresia³. Another older study suggests that maternal age, parity and gravidity, gestational age, Intrauterine Growth Retardation (IUGR), birth weight, and race/ethnic as the potential risk factors for pediatric biliary atresia⁵. The clear However, in Indonesia, data showing the biliary atresia risk factors is still limited. The authors aim to describe both the prenatal and postnatal risk factors in children with biliary atresia.

Methods and Materials

This was a cross-sectional study conducted in April 2017 using medical record data of children aged a-month-old to 72-months-old that were diagnosed with cholestasis in pediatric hepatology outpatient ward in Dr. Soetomo Hospital Surabaya during January 2010 to April 2017. Any incomplete data in medical record were excluded.

The diagnosis of biliary atresia was decided based on clinical manifestation (icteric, acholic stool), laboratory finding (direct bilirubin > 2 mg/dL if total bilirubin < 5 mg/dL, or 20% of total bilirubin if its value > 5 mg/dL), and histological findings of liver biopsy (ductular proliferation, periportal fibrosis, and bile plug).

We evaluated the maternal age on pregnancy, maternal history on pregnancy, and maternal parity as the prenatal risk factors and Birth weight, gestational age, gender, plurality, anti CMV serology as postnatal risk factors. The data were collected and analyzed by chi square and prevalence ratio was evaluated for subsequently. All statistical analysis was conducted using SPSS version 25.

Result

There were total 219 patients with cholestasis that met the inclusion criteria. Among those patients, 85 patients (38.8%) were diagnosed with biliary atresia and 134 (61.2 %) were not biliary atresia. The characteristics of the subjects were depicted in table 1.

Table 1. Data Characteristic Including Prenatal and Postnatal Risk Factors of Biliary Atresia

Characteristic Profile	Biliary Atresia (n=85)	Non-Biliary Atresia (n=134)
Age		
0-6 m.o	51 (41.5 %)	72 (58.5 %)
6 m.o – 12 m.o	28 (35.0 %)	52 (65.0 %)
12 m.o – 36 m.o	5 (50 %)	5 (50.0 %)
36 m.o – 72 m.o	1 (16.7 %)	5 (83.3 %)
Gender		
Girl	49 (44.5 %)	61 (55.5 %)
Boy	36 (33.0 %)	73 (67.0 %)
Domicile		
Surabaya	34 (38.2 %)	55 (61.8 %)
Outside Surabaya	51 (39.2 %)	79 (60.8 %)

Cont... Table 1. Data Characteristic Including Prenatal and Postnatal Risk Factors of Biliary Atresia

Maternal Age on Pregnancy < 35 y.o. ≥ 35 y.o.	38 (31.1 %) 47 (48.5 %)	84 (68.9 %) 50 (51.5 %)
Maternal history on pregnancy Hypertension Diabetes Mellitus Healthy	19 (48.7 %) 14 (43.8 %) 52 (35.1 %)	20 (51.3 %) 18 (56.3 %) 96 (64.9 %)
Maternal Parity < 5 times ≥ 5 times	41 (32.5 %) 44 (47.3 %)	85 (67.5 %) 49 (52.7 %)
Birth Weight < 2500 g ≥ 2500 g	35 (39.8 %) 50 (38.2 %)	53 (60.2 %) 81 (61.8 %)
Gestational Age Premature Aterm	51 (48.6 %) 34 (29.8 %)	54 (51.4 %) 80 (70.2 %)
Plurality Single Multiple	81 (38.2 %) 4 (57.1 %)	131 (61.8 %) 3 (42.9 %)
Ig G and Ig M anti-CMV IGM + IgG + Ig G and Ig M +	11 (28.2 %) 42 (37.5 %) 32 (47.1 %)	28 (71.8 %) 70 (62.5 %) 36 (52.9 %)

In Table 2, the Chi square bivariate analysis showed that maternal age on pregnancy ($p= 0.009$, PR= 2.07) and parity (0.035) differed significantly, in contrast with the maternal history on pregnancy ($p= 0.247$, PR= 1.86). Furthermore, Table 3 presented the analysis result on postnatal risk factors, where gestational age ($p= 0.005$, PR= 1.27-3.86) was the only risk factor in postnatal that differed significantly.

Table 2. Prenatal Risk Factors in Pediatric Biliary Atresia

Variable	n (%)	Prevalence Ratio (PR)	CI 95%	P
Maternal age on pregnancy < 35 y.o. ≥ 35 y.o.	38 (44.7) 47 (55.3)	2.07	1.19-3.61	0.009
Maternal history on pregnancy Hypertension Diabetes Mellitus Healthy	19 (22.4) 14 (16.5) 52 (61.2)	-	-	-
Maternal parity < 5 times ≥ 5 times	41 (48.2) 44 (51.8)	1.86	1.07-3.23	0.035

Table 3. Postnatal Risk Factors in Pediatric Biliary Atresia

Variable	N (%)	PR	CI 95 %	P
Gender				
Girl	49 (57.6)	-	-	0.096
Boy	36 (42.4)			
Birth Weight				
< 2500 g	35 (41.2)	-	-	0.888
≥ 2500 g	50 (58.8)			
Gestational Age		2.22	1.27-3.86	0.005
Premature	51 (60.0)			
Aterm	34 (40.0)			
IgG and IgM anti CMV				
IgM +	11 (12.9)	-	-	0.144
IgG +	42 (49.4)			
IgG and IgM +	32 (47.1)			
Plurality				
Single	81 (95.3)	-	-	0.435
Multiple	4 (4.7)			

The logistic regression analysis showed that maternal age on pregnancy (p= 0.002), parity (p= 0.015), and gestational age (p= 0.002) are 3 most influential factors on the biliary atresia incidence. The data is depicted in table 4.

Table 4. Multivariate Analysis on Prenatal and Postnatal Risk Factors in Pediatric Biliary Atresia

Risk Factors	Exp(B)	CI 95%	P
Maternal age on pregnancy	0.383	0.21 – 0.69	0.002
Parity	0.482	0.26 – 0.86	0.015
Gestational Age	0.386	0.21 – 0.70	0.002
Gender	0.594	0.33 – 1.06	0.081
Constant	1.972		

Discussion

Among 219 patients, 85 patients were diagnosed with biliary atresia. The diagnosis is based on clinical manifestation, laboratory finding, and histological finding after liver biopsy. Liver biopsy procedure has been known as a predictor and differentiator between biliary atresia and non-biliary atresia in cholestasis

patient^{6,7}. There are more non-biliary atresia patients than biliary atresia in this study. Some other studies stated differently that biliary atresia is the most cause of cholestasis in children^{5,8}.

The advanced maternal age (≥ 35 y.o) on pregnancy is associated with unfavorable outcome of delivery (premature, low birth weight, other complications)⁹. A

study done in Sweden showed the similar result with this study where a mother whose age ≥ 35 years old has higher risk to have a children with biliary atresia⁵. Old maternal age on pregnancy is a fetomaternal factor that contribute to biliary atresia incidence. Older mother is associated with oocyte aging, history of prior disease, and chromosome abnormalities toward the baby^{10,11}.

Maternal history on pregnancy, including asthma, diabetes mellitus, epilepsy, urinary tract infection, respiratory infection, gastrointestinal infection, and healthy. A study stated that there is no significant association between maternal history on pregnancy with the biliary atresia incidence¹². Another study stated that diabetes mellitus and hypertension are included to risk factors of biliary atresia incidence. In diabetes mellitus mother, vascular dysfunction stimulate the hypoxia-induced ischemia in fetal liver vascularization¹³. Furthermore, gestational hypertension can cause several complications and poor outcome, including premature delivery, intrauterine growth retardation (IUGR), and death. High blood pressure decreases the placental blood flow and creates circulation disruption and nutrition transfer delay that lead to premature delivery. Direct association between gestational hypertension and biliary atresia incidence has not been established, but it may affect the biliary atresia incidence through the premature delivery incidence³.

High parity rate is defined as maternal delivery event more than 5 times with gestational age in each pregnancy is more than 20 weeks¹⁴. The parity data finding in this study has similar result in other studies^{5,14}. A study assumed that higher parity leads to a potential risk factor of having fetal growth abnormalities, including low birth weight and premature delivery, but the mechanism is still unknown¹⁴.

¹⁰ Birth weight is classified into two groups, low birth weight (< 2500 gram) and normal birth weight (≥ 2500 gram). a low birth weight is associated with sequences of public health problem consequences¹⁵. Previous study showed that there is no significant association between low birth weight and biliary atresia incidence, but premature delivery has significant association with this disease⁵. Prematurity has been reported as a risk factor for biliary atresia incidence despite of unclear mechanism. Some previous studies have reported the

significant association between biliary atresia and prematurity¹⁶⁻¹⁸.

Gender has been suggested as one of the risk factors of biliary atresia. Many studies have been evaluating how gender differences affect the biliary atresia incidence with the different results. An epidemiological study in Korea discovered that female infant has higher risk in biliary atresia incidence that male infant¹⁹. Other studies also showed similar result^{20,21}. On the other hand, a study in Sweden stated differently that higher biliary atresia prevalence found in male infants than female⁵.

Plurality defined as the number of the children born in a delivery. A study in Texas reported a biliary atresia incidence in monozygotic/identical twins. A recent meta-analysis study of 12 articles stated that biliary atresia is found in almost half of twin subjects with monozygotic and dizygotic twins share the same proportion²².

Conclusion

Among several prenatal factors that have been established, the presence of maternal age ≥ 35 years old on pregnancy and maternal parity (≥ 5 times) are two times higher to develop a biliary atresia. Whereas in postnatal factor, presence of prematurity risks twice higher to develop a biliary atresia. This study result may be a providing information to determine the risk factors of biliary atresia in Indonesia. **Conflict of interest :** None declared.

Source of Funding : The authors received no specific grants from any funding agency in the public, commercial, or not-for-profit sectors.

Ethical Clearance : Approved by researched ethical committee Dr. Soetomo General Hospital Surabaya No: 1787/KEPK/I/2020.

References

1. Fischler B, Lamireau T. Cholestasis in the newborn and infant. *Clin Res Hepatol Gastroenterol*. 2014;38(3):263-267. doi:10.1016/j.clinre.2014.03.010
2. Mezina A, Karpen SJ. Genetic Contributors and Modifiers of Biliary Atresia. *Dig Dis*. 2015:408-414. doi:10.1159/000371694
3. Mack CL, Sokol RJ. Unraveling the Pathogenesis

- and Etiology of Biliary Atresia. *Pediatr Res.* 2005;57(5):87-94.
4. Chiang LW, Lee CY, Krishnaswamy G, et al. Seventeen years of Kasai portoenterostomy for biliary atresia in a single Southeast Asian paediatric centre. *J Paediatr Child Health.* 2016;(September). doi:10.1111/jpc.13379
 5. Fischler B, D P, Haglund B, D P, Hjern A, D P. A population-based study on the incidence and possible pre- and perinatal etiologic risk factors of biliary atresia. *J Pediatr.* 2002;217-222. doi:10.1067/mpd.2002.126001
 6. Rashed YK. Histopathological features and accuracy for diagnosing biliary atresia by prelaparotomy liver biopsy in Egypt. *Egypt Pediatr Assoc Gaz.* 2013;61(1):42-45. doi:10.1016/j.epag.2013.05.001
 7. Venigalla S, Gourley GR. Neonatal Cholestasis. *Semin Perinatol.* 2004. doi:10.1053/j.semperi.2004.09.008
 8. Zabiegaj-Zwick C, Nel E, Moore SW. Problems related to cytomegalovirus infection and biliary atresia. *South African Med J.* 2012;102(11):890-892. doi:10.7196/SAMJ.6163
 9. Cleary-goldman J, Malone FD, Vidaver J, et al. Impact of Maternal Age on Obstetric Outcome. *Am Coll Obstet Gynecol.* 2005;105(5):983-990. doi:10.1097/01.AOG.0000158118.75532.51
 10. Huang L, Sauve R, Birkett N, Fergusson D, van Walraven C. Maternal Age and Risk of Stillbirth: a systematic review. *Can Med Assoc J.* 2008;178(2).
 11. Hunt PA, Hassold TJ. Human female meiosis : what makes a good egg go bad ? *Trends Genet.* 2008;(January). doi:10.1016/j.tig.2007.11.010
 12. The NS, Honein MA, Caton AR, et al. Risk Factors for Isolated Biliary Atresia , National Birth Defects Prevention Study , 1997 – 2002. *Am J Med Genet Part A.* 2007;2284:2274-2284. doi:10.1002/ajmg.a
 13. Xavier L, Fratta S, Regina G, et al. Hypoxic-ischemic gene expression profile in the isolated variant of biliary atresia. *J Hepatobiliary Pancreat Sci.* 2015:846-854. doi:10.1002/jhbp.297
 14. Aliyu MH, Salihu HM, Keith LG, Ehiri JE, Islam MA, Jolly PE. High Parity and Fetal Morbidity Outcomes. *Am Coll Obstet Gynecol.* 2005;105(5):1045-1051. doi:10.1097/01.AOG.0000157444.74674.75
 15. World Health Organization. *WHA Global Nutrition Targets 2025 : Low Birth Weight Policy Brief;* 2014.
 16. Durkin N, Deheragoda M, Davenport M. Prematurity and biliary atresia : a 30-year observational study. *Pediatr Surg Int.* 2017;0(0):0. doi:10.1007/s00383-017-4193-1
 17. Chiu C, Chen P, Chan C, et al. Biliary Atresia in Preterm Infants in Taiwan: A Nationwide Survey. *J Pediatr.* 2013:1-5. doi:10.1016/j.jpeds.2012.12.085
 18. Van Wessel DB, Boere T, Hulzebos C V, de Kleine RH, Verkade HJ, Hulscher JB. Preterm Infants with Biliary Atresia, a Nationwide Cohort Analysis from The Netherlands. *J Pediatr Gastroenterol Nutr.* 2017. doi:10.1097/MPG.0000000000001692
 19. Lee KJ, Kim JW, Moon JS, Ko JS. Epidemiology of biliary atresia in Korea. *J Korean Med Sci.* 2017;32(4):656-660. doi:10.3346/jkms.2017.32.4.656
 20. Wada H, Muraji T, Yokoi A, et al. Insignificant seasonal and geographical variation in incidence of biliary atresia in Japan: a regional survey of over 20 years. *J Pediatr Surg.* 2007;42(12):2090-2092. doi:10.1016/j.jpedsurg.2007.08.035
 21. Utterson EC, Shepherd RW, Sokol RJ, et al. Biliary atresia: Clinical profiles, risk factors, and outcomes of 755 patients listed for liver transplantation. *J Pediatr.* 2005;147(2):180-185. doi:10.1016/j.jpeds.2005.04.073
 22. Xu X, Zhan J. Biliary atresia in twins: a systematic review and meta-analysis. *Pediatr Surg Int.* 2020;(0123456789):1-6. doi:10.1007/s00383-020-04690-4

18. Pediatric Biliary Atresia Prenatal and Postnatal Risk Factors

ORIGINALITY REPORT

12%

SIMILARITY INDEX

8%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	mts.intechopen.com Internet Source	1%
2	core.ac.uk Internet Source	1%
3	repository.unar.ac.id Internet Source	1%
4	journal2.unusa.ac.id Internet Source	1%
5	www.ncbi.nlm.nih.gov Internet Source	1%
6	Anya Mezina, Saul J. Karpen. "Genetic Contributors and Modifiers of Biliary Atresia", <i>Digestive Diseases</i> , 2015 Publication	1%
7	1library.net Internet Source	1%
8	biology.ejournals.ca Internet Source	1%

9	Jesse K. Sandberg, Yinghua Sun, Zhaoru Ju, Shaoling Liu et al. "Ultrasound shear wave elastography: does it add value to gray-scale ultrasound imaging in differentiating biliary atresia from other causes of neonatal jaundice?", <i>Pediatric Radiology</i> , 2021 Publication	1 %
10	e-journal.unair.ac.id Internet Source	1 %
11	Swati Antala, Sarah A. Taylor. "Biliary Atresia in Children", <i>Clinics in Liver Disease</i> , 2022 Publication	<1 %
12	"The Impact of Kidney Development on the Life Course: A Consensus Document for Action", <i>Nephron</i> , 2017 Publication	<1 %
13	Andrea Catzola, Pietro Vajro. "Management options for cholestatic liver disease in children", <i>Expert Review of Gastroenterology & Hepatology</i> , 2017 Publication	<1 %
14	<i>An Illustrated Guide to Pediatric Surgery</i> , 2014. Publication	<1 %
15	Jee Hyun Lee, Hyeong Sik Ahn, Seungjin Han, Heather S. Swan, Yoon Lee, Hyun Jung Kim. "Nationwide population - based study showed	<1 %

that the rotavirus vaccination had no impact on the incidence of biliary atresia in Korea", *Acta Paediatrica*, 2019

Publication

16

Ries Lu, Brandy, and Ronald J. Sokol. "Special Considerations in Pediatric Liver Transplantation", *Medical Care of the Liver Transplant Patient Clavien/Medical Care of the Liver Transplant Patient*, 2012.

Publication

17

Vonita Chawla, Miran Rhee Anagnost, Alaa-Eldin Eldemerdash, Dahna Reyes et al. "A Novel Case of Biliary Atresia in a Premature Neonate With 1p36 Deletion Syndrome", *Journal of Investigative Medicine High Impact Case Reports*, 2018

Publication

18

Yahya Al-Farsi, Brooks, Werler, Howard Cabral, Al-Shafei, Wallenburg. "Effect of high parity on occurrence of some fetal growth indices: a cohort study", *International Journal of Women's Health*, 2012

Publication

19

eprints.qut.edu.au

Internet Source

20

f6publishing.blob.core.windows.net

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

21

"1-B1: Respiratory Infections (Non-Tuberculosis) 1 : Poster Sessions",
Respirology, 2013.

Publication

<1 %

22

"Pediatric Hepatology and Liver Transplantation", Springer Science and Business Media LLC, 2019

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

18. Pediatric Biliary Atresia Prenatal and Postnatal Risk Factors

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
