

Home / Editorial Team

## **Editorial Team**

#### **Editor in Chief:**

Prof. Harsono Salimo

SCOPUS ID <u>57205023539</u>.

Department of Pediatrics, Dr. Moewardi Hospital/ Universitas Sebelas Maret, Surakarta, Indonesia

#### **Editor:**

1. Prof. Ana Diez Roux

SCOPUS ID <u>7004701264</u>, ORCID ID <u>0000-0001-9658-7050</u>, Dornsife School of Public Health, Drexel University, Philadelphia, PA, United States

- 2. Prof. Jessica Gipson
  - SCOPUS ID <u>16028551300</u>, ORCID ID <u>0000-0001-8719-2081</u>, UCLA Fielding School of Public Health, Los Angeles, United States
- 3. Prof. Linda S. Adair
  - SCOPUS ID <u>7005651928</u>, ORCID ID <u>0000-0002-3670-8073</u>, Department of Nutrition, University of North Carolina School of Public Health, Chapel Hill, United States
- 4. Prof. Diana Kuh
  - SCOPUS ID <u>8329745200</u>, ORCID ID <u>0000-0001-7386-2857</u>, MRC Unit for Lifelong Health and Ageing, University College London, London, United Kingdom
- 5. Prof. Adang Bachtiar
  - SCOPUS ID <u>56683183900</u>, ORCID ID <u>0000-0003-1008-3906</u>, SINTA ID <u>6027980</u>, Faculty of Public Health, Universitas Indonesia
- 6. <u>Dr. Rita Benya Adriani</u> SCOPUS ID <u>57274125200</u>, ORCID ID <u>0000-0001-7386-2857</u>, SINTA ID <u>5995045></u>,

School of Health Polytechnics, Surakarta, Indonesia

#### **Reviewer:**

1. Prof. Mai Do

SCOPUS ID <u>13204304000</u>, ORCID ID <u>0000-0002-2965-4812</u>, Tulane University School of Public Health and Tropical Medicine, New Orleans, United States

2. Prof. Michael Dibley

SCOPUS ID <u>26665373100</u>, ORCID ID <u>0000-0002-1554-5180</u>, The University of Sydney School of Public Health, Sydney, Australian

- 3. Prof. Nicola Reavley
  - SCOPUS ID <u>6507792925</u>, ORCID ID <u>0000-0001-5513-8291</u>, Melbourne School of Population and Global Health, Melbourne, Australia
- 4. <u>Dr. Uki Retno Budihastuti</u>, SCOPUS ID <u>57212480740</u>, ORCID ID <u>0000-0003-0611-7464</u>, SINTA ID <u>6689455</u>, Department of Obstetrics and Gynecology, Dr. Moewardi Hospital/ Universitas Sebelas Maret, Surakarta, Indonesia
- 5. <u>Dr. Eti Poncorini Pamungkasari</u>, SCOPUS ID <u>57195310832</u>, ORCID ID <u>0000-0002-4197-3226</u>, SINTA ID <u>6657185</u>, Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia
- 6. <u>Dr. CSP. Wekadigunawan</u>, ORCID ID <u>0000-0001-9354-7133</u>, SINTA ID <u>6680156</u>, Faculty of Health Sciences, Universitas Esa Unggul, Jakarta, Indonesia
- 7. <u>Dr. Noer Saudah</u>, ORCID ID <u>0000-0002-6638-0996</u>, SINTA ID <u>6657185</u>, School of Health Sciences Bina Sehat PPNI, Mojokerto, Indonesia

## **Managing Editor:**

1. Aris Widiyanto

SCOPUS ID <u>58046530800</u>, ORCID ID <u>0000-0002-3234-3265</u>,SINTA ID <u>6673181</u>, School of Health Sciences Mamba'ul Ulum, Surakarta, Indonesia

2. <u>Ika Yuliayuningrum</u>

SCOPUS ID <u>57205018824</u>, ORCID ID <u>0000-0003-0851-1131</u>, Universitas Sebelas Maret, Surakarta, Indonesia

#### **Assistant Editors:**

- 1. Siti Mar'atul Munawaroh
- 2. Amallia Wijiwinarsih
- 3. Nindita Arum Veibiani

#### Web Designer:

Abdush Shomad

#### **Editorial Office:**

Journal of Maternal and Child Health

## Masters Program in Public Health,

Graduate School, Universitas Sebelas Maret,

Jl. Ir. Sutami 36A, Surakarta 57126, Indonesia.

Telephone: (62 271) 632450 ext 208.

Email: jurnals2ikm@gmail.com

Website: www.thejmch.com.



Home / Archives / Vol. 8 No. 2 (2023)

## Vol. 8 No. 2 (2023)

PUBLISHED: 2023-03-16

### **ARTICLES**

Meta-Analysis: Effect of Bullying on the Risk of Suicide in Adolescents

Sholikha Dela Aprilia, Hanung Prasetya, Bhisma Murti 138-147



## Relationship Between Family Social Support with Anxiety of Postpartum

Zakia Jihan Salsa Bila, Budi Santoso, Muhammad Ilham Aldika Akbar  $^{148\text{-}153}$ 



Meta Analysis the Effects of Maternal Education, Residence, and Birth Delivery Place, on Exclusive Breastfeeding

Ade Amalia, Eti Poncorini Pamungkasari, Rita Benya Adriani 154-168



	Relationship stfeeding Moth		Anxiety	and	Smoothness	Breast	Milkexpenditu	ires in
Annis	a Nur Rohma, Ma 75	artono Tri Uto	omo, Risa Et	ika, Wo	oro Setia Ningtyas	3		
	PDF							
	ysis of Using lemic Based or			ication	n for Pregnai	nt Wom	en During CO	)VID-19
Nurar 176-18	ni Zulfa Zakiya, M 87	Iuhammad Ar	dian Cahya,	Samsri	yaningsih Handa	yani		
B	PDF							
The (	Correlation of	Knowledge,	Mother					
Anggr 188-2	raini Dian Prastic	ha, Mahendra	Tri Arif San	npurna	, Linda Dewanti			
	PDF							
	_				ntrol Among P	regnant \	Women with As	sthma
Winda 203-2	a Aenah, Arief Ba 209	khtiar, Endyk	a Eyre Frety					
B	PDF							
	Role of Ferritin		vel as Pred	lictor (	Sepsis Mortali	ty on Chi	ldren in Dr. Me	oewardi
Agus \$	Sucianto, Pudjias 16	tuti Pudjiastu	ti, Diah Linta	ang Ka	wuryan			
B	PDF							

Yemen, 2015-2016
Ahmed Hamood Al-Shahethi, Rafdzah Ahmad Zaki, Abdulwahed Abdulgabar Al-Serouri, Awang Bulgiba 217-226
₽DF
Social Determinants in Association with Postpartum Blues during the Transition Period of COVID-19 Pandemic
Salwa Tsabitah Althaf Mujab, Ria Maria Theresa, Arman Yurisaldi Saleh, Soroy Lardo 227-236
₽DF
Meta-Analysis the Effect of Exclusive Breastfeeding on Anemia in Children Under Five Fikri Dian Dinu Azizah, Eti Poncorini Pamungkasari, Bhisma Murti 237-246
₽DF
The Relationship Between History of COVID-19 Disease with Anxiety and Symptoms of Obsessive Compulsive Disorder (OCD) in Adolescents Age 20-24 Years
Ayu Novia Christanti, Woro Setia Ningyas, Azimatul Karimah, Pudji Lestari 247-254
PDF



#### **INFORMATION**

Information for Author

Licenses and Copyright

**Publication Ethics for Authors** 

**Article Processing Charge** 

**Author Guidelines** 

Template for Author

**Originality Statement** 

Information for Reviewer

**Publication Ethics for Reviewers** 

**Reviewers Guidelines** 

**Reviewer Statement** 

**Reviewer Assessment Form** 

**Publication Ethics for Editor** 



School, Universitas Sebelas Maret. Jl. Ir. Sutami 36A, Surakarta 57126, Central Java, Indonesia. This journal is licensed under the Creative Commons Attribution 4.0 International (CC BY 4.0).



Pages 190 710 Online 3 Vis. today 149





JOURNAL OF MATERNAL AND CHILD HEALTH

UNIVERSITAS SEBELAS MARET

♣ P-ISSN: <> E-ISSN: 25490257
Subject Area: Health







2025

Google Scholar

**▶** <u>Garuda</u> **᠍ Website** 

**Editor URL** 

**History Accreditation** 

2018 2019 2020 2021 2022 2023 2024

<u>Garuda</u> Google Scholar

Meta-Analysis: The Effect of Physical Activity in Reducing Excessive Gestational Weight Gain and Gestational Diabetes Mellitus among Overweight and Obese Pregnant Women

Journal of Maternal and Child Health

■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 78-90

<u> ■ DOI: -</u> **2023** O Accred: Sinta 3

Increased plasma caspase-3 in children with Down syndrome is associated with an increasing risk of pulmonary hypertension

Journal of Maternal and Child Health ■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 23-32

**2**023 <u> ■ DOI: -</u> O Accred: Sinta 3

**The Differences Women** 

Journal of Maternal and Child Health ■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 70-77

**2**023 <u> ■ DOI: -</u> O Accred: Sinta 3

Effect of Stunting Education Counseling with Booklet Media on Parental Knowledge and Attitude As an Effort to Overcome Stunting

■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 105-Journal of Maternal and Child Health <u>115</u>

**2**023 O Accred: Sinta 3

The Effect of Vitamin D Deficiency on Anemia in Children and Adolescents: Meta-Analysis Journal of Maternal and Child Health ■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 12-22

<u>2023</u> <u> ■ DOI: -</u> O Accred: Sinta 3

Epidemiological Aspects and Factors Associated with Early Neonatal Death From 2018 to 2020 in the Maternity of the Sav

Journal of Maternal and Child Health Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 91-<u>104</u>


<u>2023</u>

<u> БОІ: -</u>

O Accred: Sinta 3

Get More with SINTA Insight

Go to Insight



Journal By Google Scholar						
All	Since 2018					
1645	1619					
18	18					
47	45					
	All 1645 18					

<u>Do Massage and Relaxation Music Therapy Increase Breast Milk Volume? A Meta-Analysis</u> ■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 33-47 Journal of Maternal and Child Health

**2023** 

<u> БОІ: -</u>

O Accred: Sinta 3

Relationships between Exclusive Breastfeeding, and History of Illness, and Stunting in **Children Under Five** 

Journal of Maternal and Child Health ■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 116-<u>124</u>

<u>2023</u>

<u> ■ DOI: -</u>

O Accred: Sinta 3

Meta-Analysis: The Effect of Anemia in Pregnant Women on the Risk of Postpartum Bleeding and Low Birth Weight

Journal of Maternal and Child Health ■ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 58-<u>69</u>

<u> 2023</u> <u> ББ DOI: -</u>

O Accred: Sinta 3

Comparative Study of Feto-Maternal Outcome of Ante-natal Cases as per the Standard WHO Guidelines Vs Revised Antenatal Visit Schedule during Covid19 Pandemic

Journal of Maternal and Child Health ▶ Journal of Maternal and Child Health Vol. 8 No. 1 (2023) 125-<u>137</u>

O Accred: Sinta 3



Get More with SINTA Insight

Go to Insight

Citation Per Year By Google Scholar

	Journal By Google Scholar	
	All	Since 2018
Citation	1645	1619

h-index 18 18 i10-index 47 45



## Relationship between Anxiety and Breast Milk Expulsion in Lactating Mothers

Annisa Nur Rohma<sup>1)</sup>, Martono Tri Utomo<sup>2)</sup>, Risa Etika<sup>2)</sup>, Woro Setia Ningtyas<sup>3)</sup>

<sup>1)</sup>Midwifery Program Study, Faculty of Medicine, Airlangga University, Surabaya, Indonesia <sup>2)</sup>Departement of Pediatrics, Dr.Soetomo Hospital, Airlangga University, Surabaya, Indonesia <sup>3)</sup>Midwifery Program Study, Faculty of Medicine, Airlangga University, Surabaya, Indonesia

#### **ABSTRACT**

**Background:** Breastfeeding is the process of giving mother's milk to babies from birth to two years. Exclusive breastfeeding can protect babies and children against dangerous diseases and strengthens the affection (bonding) between mother and child. However, the exclusive breastfeeding coverage rate is still low due to failure to breastfeed. One of the factors that influence failure in the breastfeeding process can be caused by not releasing milk. The smoothness of breastfeeding is strongly influenced by psychological factors, namely anxiety. This study aims to analyze the relationship between anxiety and the smoothness of breastfeeding in breastfeeding mothers in Slahung District, Ponorogo Regency.

**Subjects and Method:** A cross-sectional study was conducted in Slahung District, Ponorogo Regency in November 2022. A total of 83 breastfeeding mothers were selected for this study. The dependent variable is the smooth expulsion of breast milk. The independent variable is anxiety. The instrument used is a questionnaire. Research data were analyzed using the Chi-Square test and odds ratio (OR).

**Results:** Anxiety affects the smooth release of breast milk in nursing mothers. Breastfeeding mothers who experience anxiety are at risk of experiencing a milk ejection process that is not smooth as much as 26.18 times compared to breastfeeding mothers who do not experience anxiety, and these results are statistically significant (OR= 26.18; 95% CI= 3.20 to 214.12; p<0.001).

**Conclusion:** Anxiety affects the smooth release of breast milk in nursing mothers.

**Keywords:** anxiety, smooth milk production.

#### **Correspondence:**

Annisa Nur Rohma, Midwifery Program Study, Faculty of Medicine, Universotas Airlangga, Indonesia. Jl. Mayjen Prof. Dr. Moestopo No. 47, Pacar Kembang, Tambaksari, Surabaya, East Java, 60132. Email: annisa.nur.rohma-2019@fk.unair.ac.id. Mobile: +6285708790065.

#### Cite this as:

Rohma AN, Utomo MT, Etika R, Ningtyas WS (2023). The Relationship Between Anxietyand Smoothness Breast Milkexpenditures in Breastfeeding Mothers. J Matern Child Health. 08(02): 169-175. https://doi.org/10.26911/thejmch.2023.08.02.04.



Journal of Maternal and Child Health is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

#### **BACKGROUND**

Breast milk is the main food for babies which contains a complete nutritional composition and is needed for their growth and development (Liliek Fauziah, 2020). Breast milk is a natural source of nutrition for babies. Human breast milk contains special com-

ponents that prevent malnutrition and offer optimal growth for infants (Martini et al., 2020). Breast milk is produced by the mammary glands, namely the alveoli cells. The milk that has been produced will flow into the lactiferous sinuses and gather in the lactiferous duct organs, then the breasts expe-

e-ISSN: 2549-1172

rience pressure and reflex release of milk (let down reflex) through both mother's breasts (Putri, 2020).

The WHO (World Health Organization) and the United Nations International Children's Emergency Fund (UNICEF) recommend that babies should only be given breast milk (ASI) for at least 6 months and continued breastfeeding until the child is 2 vears old (Korompis, 2019). The government has regulated breastfeeding in Law Number 33 of 2012 to support exclusive breastfeeding mothers. However, even though breastfeeding is mandatory, it turns out that many mothers choose not to breastfeed their babies for several reasons. According to Aprilia 2011 (in Korompis, 2019) In the process of breastfeeding there are two important processes, namely the process of forming breast milk (milk production reflex) and the process of expressing breast milk (reflex let down), both of which are influenced by hormones regulated by the hypothalamus. several factors, namely the baby's weight at birth, frequency of breastfeeding, exposure to cigarette smoke, gestational age at birth, age and parity of the mother, stress and acute illness, IMD (Early Breastfeeding Initiation), alcohol consumption, breast care, contraceptive use, and nutritional status (Hastuti and Wijayanti, 2017). In premature babies, there is Immunoglobulin A and breast milk is the main source of Ig A but levels decrease with age (Handayani et al., 2012).

Breastfeeding mothers are mothers who give breast milk to babies from birth to two years of age from the breast. The results of research on the benefits of breastfeeding for the physical health of mothers include reducing blood pressure, reducing the risk of breast cancer and ovarian cancer as well as for their psychological health, namely reducing stress and sleep which is increasing (Figueiredo et al., 2013). Breastfeeding is associated with improved infant health and

immune development, fewer incidences of gastrointestinal disease and lower mortality than formula-fed infants. As well as providing basic nutrition for baby growth. Breast milk is a source of commensal bacteria which further improves infant health by preventing adhesion of pathogens and promoting beneficial microbial colonization of the gut. Breast milk is produced by the mammary glands i.e. in alveoli cells. The milk that has been produced will flow into the lactiferous sinuses and gather in the lactiferous duct organs, then the breasts experience pressure and reflex release of milk (let down reflex) through both mother's breasts (Putri, 2020). The calm mental and emotional condition of the mother greatly influences the smooth flow of milk. If the mother experiences stress, depressed thoughts, uneasy, anxious, sad, and tense it will affect the smoothness of breastfeeding (Riksani, 2012).

Anxious mothers will secrete less milk than mothers who are not anxious. Based on the results of the study (Iin Febrina, 2011) said that there is a relationship between the level of anxiety and the smoothness of breastfeeding in primiparous postpartum mothers. Efforts to keep breastfeeding running smoothly start from a mother's strong desire to provide the best nutrition, namely breast milk for her baby. Strong motivation will affect the physical and emotional mother to produce breast milk. Mothers who have strong and sincere desires and high affection, breast milk production can be stimulated. This study aimed to analyze the relationship between anxiety and the smooth release of breast milk in breastfeeding mothers in Slahung, Ponorogo Regency.

#### SUBJECTS AND METHOD

#### 1. Study Design

This study uses an observational analytic study design with a cross sectional approach. The location of this research is in Sla-

hung District, Ponorogo Regency, East Java, and will be held in November 2022.

#### 2. Population and Sample

The population in this study were all breastfeeding mothers in Slahung District, Ponorogo Regency. The sample size in this study was 83 samples using a simple random sampling technique.

#### 3. Study Variable

The dependent variable was the smooth expulsion of breast milk. The independent was anxiety.

4. Operational Definition of Variables Expenditure of breast milk is the process of releasing breast milk as measured by several indicators, namely scheduled or unscheduled breastfeeding time, condition of the mother's breast milk seeping or not, sensation in the mother, namely feeling tingly, swallowing sound and feeling tired, the baby's response remains fussy or not after being breastfed, BAK normal or abnormal, normal or abnormal bowel movements. Code 1 for breastfeeding is not smooth. Code 2 is for breastfeeding smoothly.

**Anxiety** is a feeling of anxiety, tension, fear, insomnia, intellectual, depressed feelings, somatic symptoms in muscles and sensory, cardiovascular symptoms, respiratory, gastrointestinal, autonomic symptoms and also behavior shown by breastfeeding mothers. Code 1 for anxiety. Code 2 for no anxiety.

#### 5. Study Instrument

The instruments used in data collection were anxiety questionnaires and breastfeeding expenditure questionnaires which were distributed offline to breastfeeding mothers who were research subjects or samples. Data management techniques include editing, coding, data entry, cleaning and tabulating techniques. Questions in the questionnaire included 14 questions about anxiety and 24 questions about breastfeeding.

#### 6. Data Analysis

The data analysis method of this study uses

univariate and bivariate analysis. Univariate analysis in the form of frequency distribution and percentage of each variable and bivariate analysis using the chi-square statistical test with a significance level of 5% (p =0.05) to determine the relationship between anxiety and the smoothness of breastfeeding.

#### 7. Research Ethics

Research ethics namely with informed consent, anonymity, confidentiality. A research ethics permit approval letter was obtained from the Health Research Ethics Committee, Faculty of Medicine, Airlangga University, Surabaya, Indonesia, No.187/EC/KEPK/FKUA/2022, on October 3, 2022.

#### **RESULTS**

#### 1. Characteristics of Sample

Table 1 shows that of the 83 subjects, the majority of breastfeeding mothers aged 20-35 years were 75 (90.4%), and 56 (67.5%) had normal deliveries. The gestational age at birth in this study was mostly in the term group of 66 (79.5%) people. Most of the subjects were mothers who did IMD as many as 49 (59.0%) people. The majority of subjects with maternal parity who had 1 child were 49 (59.4%) people, and used nonhormonal contraception as many as 52 (67.2%) people. Most of the subjects were mothers who did not have dietary restrictions as many as 51 (61.4%) people and their family members did not smoke as many as 49 (59.0%) people. Most of the babies at the time of study were aged < 6 months as many as 62 (74.7%) babies, with babies weighing 2500-4000 grams as many as 65 (78.3%) babies.

#### 2. Univariate Analysis

Univariate analysis in this study included anxiety and the smoothness of expressing breast milk which can be seen in Table 2. Table 2 shows that the majority of breastfeeding mothers experienced anxiety as many as 13 (15.7%) breastfeeding mothers, and 34

(41.0%) experienced a process of ejecting milk that was not smooth. breastfeeding mothers.

#### 3. Bivariate Analysis

Bivariate analysis was carried out to see the relationship between anxiety and the smoothness of expressing milk in nursing mothers. Table 3 shows that anxiety has an effect on the smooth release of milk in nursing mothers. Breastfeeding mothers who experience anxiety are at risk of experiencing a milk ejection process that is not smooth as much as 26.18 times compared to breastfeeding mothers who do not experience anxiety, and these results are statistically significant (OR= 26.18; 95% CI= 3.20 to 214.12; p<0.001).

Tabel 1. Study subject characteristics

Characteristics Category Frequency (n) Percentage (%)						
Maternal Age						
Maternal Age	20-35 Years	75	90.4			
	>35 Years	8	9.6			
Baby's age	< 6 months	62	74.7			
	> 6 months	21	25.3			
Type of Childbirth	Normal paravaginal	56	67.5			
Type of emidsh th	SC	27	32.5			
Birth Weight	<2500 gram	18	21.7			
Ditti Weight	2500-4000 gram	65	78.3			
Gestational Age at Birth	Preterm	17	20.5			
oestational rige at Difti	Aterm	66	79.5			
Early Breastfeeding	Yes	49	59.0			
Initiation	No	34	41.0			
Parity	1 child	49	59.4			
	2-4 children	34	41.0			
Contraception	Non-Hormonal	52	62.7			
	Hormonal	31	37.3			
<b>Eating Abstinence</b>	No	51	61.4			
Lating Absumence	Yes	32	38.6			
Smoking Family	No	49	59.0			
Sinoking Family	Yes	34	41.0			

**Tabel 2. Univariate analysis** 

Variable	Category	Frequency (n)	Percentage (%)
Anxiety	Yes	13	15.7
	No	70	84.3
Smooth milk production	Smooth	49	59.0
	Not smooth	34	41.0

Table 3. Bivariate results between anxiety and fluency in expressing milk in nursing mothers

Variable	Smooth milk production							
	No		Yes		Total		OR (95% CI)	p
Anxiety	N	%	N	%	N	%		_
Yes	12	92.3	1	7.7	13	100	26.18	<0.001
No	22	31.4	48	68.6	70	100	(3.20 to 214.12)	

#### **DISCUSSION**

# 1. The relationship between anxiety and smooth milk expulsion

Based on the results of this study showed that most (84.3%) of the subjects had no anxiety. The results of this study also showed that the majority (59.0%) of the subjects had smooth breastfeeding. Anxiety is an emotion, a feeling that arises as an initial response to psychological stress and threats to values that are meaningful to individuals. Anxiety can be described as a feeling of uncertainty, doubt and helplessness, restlessness, worry, unrest which is often accompanied by physical complaints (Lilik, 2016). Anxiety is a major public health problem. Anxiety, like any class of mental disorders, is also associated with various medical conditions, exacerbating symptoms, when anxiety symptoms do not reach the criteria for a disorder, they can cause severe distress. (Christian Grillon, 2019)

Milk production is strongly influenced by psychological factors (Prasetyono, 2009). Postpartum stress conditions are experienced by 80% of women after giving birth. Feelings of sadness or gloom that hit the mother arise within two days to two weeks after delivery (Danuatmaja and Meiliasari, 2013). The condition of mothers who are easily anxious and stressed can interfere with lactation so that it can affect milk production. This is because stress can inhibit milk production (Kodrat, 2010). The higher the level of emotional disturbance, the less stimulation of the hormone prolactin is given to produce breast milk (Prasetvono, 2009). The mental and psychological factors of nursing mothers greatly influence the breastfeeding process and the smooth production of breast milk. Feelings of stress, pressure, and discomfort experienced by a mother can inhibit the amount of milk that comes out (Bahayatun, 2009).

Breastfeeding mothers must think

positively and relax so they don't experience anxiety and the mother's psychological condition becomes good, a good psychological condition can trigger the work of hormones that produce breast milk. The results of this study are in accordance with the results of research conducted by (Kamariyah et al. 2014) that there is a relationship between the psychological condition of the mother and the smooth production of breast milk, a good psychological state of the mother will motivate her to breastfeed her baby so that the hormones that play a role in milk production will increase because Breast milk production starts from the breastfeeding process and will stimulate milk production. Anxious conditions in postpartum mothers affect milk expenditure. Breastfeeding mothers should relax and think positively. A good mother's psychological condition can stimulate hormone work to produce breast milk (Aprilia and Krisnawati, 2017; Destri et al., 2022)

Factors that can affect breastfeeding failure due to not releasing milk are psychological factors (Mardjun, 2019). If the mother experiences anxiety, sadness, stress, depressed thoughts, unrest, or tension, it can affect the smooth flow of milk. Mothers who experience anxiety will secrete less milk compared to mothers who do not experience anxiety (Arfiah, 2018; Friscila et al., 2022).

#### **AUTHOR CONTRIBUTION**

In this study, Martono Tri Utomo and Annisa Nur Rohma collaborated to create a conceptual framework and research methodology. Annisa Nur Rohma collected data. Risa Etika, Annisa Nur Rohma, and Woro Setia Ningtyas collaborated to analyze the data.

#### FUNDING AND SPONSORSHIP

The study was self-funded.

#### **CONFLICT OF INTERESTS**

There is no conflict of interest in this study.

#### **ACKNOWLEDGMENT**

The researcher expresses his deepest gratitude to the Slahung District Health Center and the Posyandu of Caluk Village and Slahung Village who have granted research permission.

#### REFERENCES

- Aprilia D, Krisnawati AM (2017). Faktor-faktor yang mempengaruhi kelancaran pengeluaran asi pada ibu post partum (Factors affecting the smooth expulsion of breast milk in post partum mothers). Jurnal Kebidanan. 6(1): 11-17. Retrieved from: https://jurnal.stikes-williambooth.ac.id/index.php/Keb/article/view/199
- Arfiah A. (2018). Pengaruh pemenuhan nutrisi dan tingkat kecemasan terhadap pengeluaran ASI pada ibu post partum primipara (The effect of fulfillment of nutrition and levels of anxiety on expenditure of breast milk in post partum primipara mothers). Jurnal Kebidanan. DOI:10.33486/jk.v8i2.60. Volume 8 Nomor 2 November 2018
- Bahayatun. (2009). Buku ajar kebidanan asuhan nifas normal (Normal postpartum care midwifery textbook). EGC.
- Christian Grillon OJ (2019). Modeling anxiety in healty humans: a key intermediate bridge between basic and clinical sciences. Neuropsychopharmacology, 12. doi: 10.1038/s41386-019-0445-1 2019 Nov;44(12):1999-2010.
- Danuatmaja, Meiliasari (2013). Persalinan, sakit terkuat sepanjang hidup dalam: persalinan normal tanpa rasa sakit (Childbirth, the strongest pain throughout life in: normal childbirth without pain). Puspa Swara.
- Destri U, Sarkiah S, Friscila I (2022).

- Efektivitas pemberian aromaterapi lavender terhadap produksi asi pada ibu nifas normal: literatur review (The effectiveness of giving lavender aromatherapy on breast milk production in normal postpartum mothers: literature review). Proceeding of Sari Mulia Univercity Midwifery National Seminar. doi: https://doi.org/10.33859/psmumns.v3i1.813 Published Feb 24,2022 Vol.3 No.1 (2021):
- Friscila I, Hasanah SN, Hidayah N, Sari SP, Nabila S, Fitriani A, Fonna L, Dashilva NA (2022). Education preparation for exclusive breast milk at sungai andai integrated services post. Prosiding Seminar Nasional PKM: Masyarakat Tangguh. 1(1). Prosiding seminar nasional PKM: Masyarakat Tangguh https://orcid.org/0000-0001-8912-1367
- Handayani KD, Etika R, Alami N, Wijayanti D, Harianto A (2012). Gain velocity and iga secretory fecal between preterm. 11(5): 108-121 https://doi.org/10.34-302/erpjfst/2019.11.5.16
- Korompis, Widad AS, Adisti A, Rumayar, Grace EC (2019). Hubungan antara pengetahuan dan sikap ibu dengan pemberian asi eksklusif di Wilayah Kerja Puskesmas Tuminting Kota Manado (The relationship between mother's knowledge and attitudes and exclusive breastfeeding in the Work Area of the Tuminting Health Center in Manado City). Laporan Penelitian. 8(7): 298–304.
- Kamariyah N, Anggasari Y, Muflihah S (2014). Buku ajar kehamilan untuk mahasiswa dan praktisi keperawatan serta kebidanan (Pregnancy textbook for nursing and midwifery students and practitioners). Salemba Medika.
- Lilik AM, Zainuri I, Akbar A (2016). Buku ajar keperawatan jiwa teori dan ap-

- likasi klinik (Textbook of psychiatric nursing theory and clinical application). Indomedika Pustaka.
- Manggabarani S, Hadi AJ, Said I,Bunga S (2018). Hubungan status gizi, pola makan, pantangan makanan dengan kelancaran produksi asi pada ibu menyusui di Kota Makassar (Correlation between nutritional status, diet, food abstinence and smooth milk production in breastfeeding mothers in Makassar City). Jurnal Dunia Gizi, 1(1): 1. https://doi.org/10.33085/jdg.-v1i1.2902
- Mardjun Z (2019). Hubungan kecemasan dengan kelancaran pengeluaran ASI pada ibu post partum selama dirawat di Rumah Sakit Ibu dan Anak Kasih Ibu Manado (The relationship between

- anxiety and smooth expulsion of breast milk in post partum mothers during treatment at Mother and Child Care Hospital Manado). E-Journal Keperawatan (e-Kp). 7(1). doi: https://doi.org/10.35790/jkp.v7i1.22901
- Martini, Irwanto, Irawan R, Widjaja NA (2020). Breastmilk macronutrient levels and infant growth during the first three months: A cohort study. Siriraj Medical Journal. 72(1):10–17. https://doi.org/10.33192/Smj.2020.02
- Putri IM, Utami FS (2020). ASI dan menyusui (ASI and breastfeeding). In Buku. UNISA Press. https://doi.org/-10.26699/jnk.v
- Riksani R (2012). Keajaiban ASI (air susu ibu) (The miracle of ASI (breast milk).

  Jakarta Timur: Dunia Sehat