



Source details

Pharmacognosy Journal

Scopus coverage years: from 2009 to Present

Publisher: Pharmacognosy Network Worldwide

ISSN: 0975-3575

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmacology

Pharmacology, Toxicology and Pharmaceutics: Drug Discovery

Source type: Journal

CiteScore 2022

1.9



SJR 2022

0.229



SNIP 2022

0.484



[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology



CiteScore 2022 counts the citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022. [Learn more >](#)

CiteScore 2022 ▼

$$1.9 = \frac{1,691 \text{ Citations } 2019 - 2022}{878 \text{ Documents } 2019 - 2022}$$

Calculated on 05 May, 2023

CiteScoreTracker 2023 ⓘ

$$1.7 = \frac{1,267 \text{ Citations to date}}{741 \text{ Documents to date}}$$

Last updated on 05 July, 2023 • Updated monthly

CiteScore rank 2022 ⓘ

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#219/301	27th
Pharmacology		
Pharmacology, Toxicology and Pharmaceutics	#121/156	22nd
Drug Discovery		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Pharmacognosy Journal

COUNTRY

India



Universities and research institutions in India



Media Ranking in India

SUBJECT AREA AND CATEGORY

Pharmacology,
Toxicology and
Pharmaceutics
Drug Discovery
Pharmacology

PUBLISHER

EManuscript Technologies

H-INDEX

30

PUBLICATION TYPE

Journals

ISSN

09753575

COVERAGE


2009-2022


INFORMATION

[Homepage](#)[How to publish in this journal](#)editor@phcogj.com

SCOPE

Pharmacognosy Journal (Phcog J.) covers different topics in natural product drug discovery, and also publishes manuscripts that describe pharmacognostic investigations, evaluation reports, methods, techniques and applications of all forms of medicinal plant research









 Join the conversation about this journal

 [Quartiles](#)



FIND SIMILAR JOURNALS ?


<p>1 Oriental Pharmacy and Experimental Medicine USA</p> <p>82% similarity</p>	<p>2 Pharmacognosy Research IND</p> <p>81% similarity</p>	<p>3 Indian Journal of Natural Products and Resources IND</p> <p>77% similarity</p>	<p>4 Journal of H Medicinal Pl USA</p> <p>7 s</p>
--	---	---	---



Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```



Metrics based on Scopus® data as of April 2023

E **edi utomo** 8 months ago

I want to get information about drug poison crimes, is there a journal about this?

reply



Melanie Ortiz 8 months ago

SCImago Team

Dear Edi,

Thank you for contacting us.

We suggest you use the Journal Rankings' Search Tool shown on our website, as the journals can be filtered by Subject Area/ Category:

<https://www.scimagojr.com/journalrank.php>

Best Regards, SCImago Team

K **Kori Yati** 1 year ago

I want to submit an article, can I send a link?

reply



Melanie Ortiz 1 year ago

SCImago Team

Dear Kori, thank you very much for your comment, we suggest you look for the author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

Home (/) About Journal ▾ Editorial Board (/editorial-board) For Authors ▾
 Contact Us (/contact-us)

Pharmacognosy Journal (/)

An Open Access, Peer Reviewed Journal in the field of
 Pharmacognosy

Enter terms then hit Search...



Articles In Press (/inpress) Current Issue (/v15/i3) Archives (/archives)

RSS Feeds (/rss.xml) Submit Article (https://www.phcogj.info)

HOME (/) / PHARMACOGNOSY JOURNAL,
 VOL 15, ISSUE 2, MAR-APR, 2023

Pharmacognosy Journal, Vol 15, Issue 2, Mar-Apr, 2023

RECENT ARTICLES



Original Article

Acute and Sub-acute Oral Toxicity Profile of Root Bark Methanol Extract of *Carissa Edulis* Vahl (/article/1988)

Jane Wanja Mbiri, Kenneth Ogila, Patrick Kisangau, Michael Gicheru

Pharmacognosy Journal, 15(2):253-258

DOI: 10.5530/pj.2023.15.36

Published: Wed, 26-Apr-2023


[Read More \(/article/1988\)](/article/1988)

Original Article

An Analysis of Local Knowledge of Phit-Samdaeng in Yang Sisurat District, Maha Sarakham Province and The Medicinal Plants Used to Relieve Its Symptoms (/article/1990)



**Hatthapan Wipanso,Bhanubong Bongcheewin,Suppachai
Tiyaworanant,Suksalin Booranasubkajorn,Natchagorn Lumlerdkij,Pravit
Akarasereenont**



Pharmacognosy Journal,15(2):259-265

DOI: 10.5530/pj.2023.15.37

Published: Wed, 26-Apr-2023

[Read More \(/article/1990\)](#)

Original Article

**Development and Evaluation of Bio fabricated Silver
Nanoparticles from Blumea lacera for In-vitro
Antibacterial, Antioxidant and Anti-inflammatory Activity
(/article/1991)**

Tarkeshwar Dubey,Kancharla Bhanukiran,Kuna Das,Siva Hemalatha



(/article/1991)

Pharmacognosy Journal,15(2):266-278

DOI: 10.5530/pj.2023.15.38


Published: Wed, 26-Apr-2023

[Read More \(/article/1991\)](#)

Original Article

**Antibacterial and Anti-quorum Sensing Activities of
Erianthemum dregei`s Leaf Extract and Molecular
Docking (/article/1992)**

**Tsolanku Sidney Maliehe,Tlou Nelson Selepe,Nokuthula Nomusa
Mthembu,Jabulani Siyabonga Shandu**



Pharmacognosy Journal,15(2):279-285

DOI: 10.5530/pj.2023.15.39

Published: Wed, 26-Apr-2023

[Read More \(/article/1992\)](#)

Original Article

**The Effect of Sapodilla Leaf Extract (Manilkara zapota L.)
on Lipid Profiles of Alloxan-Induced Diabetic Mice (/article
/1994)**

**Tridiganita Intan Solikhah,Tania Anggreani Wijaya, ,Daffa Alice Pavita,Ragil
Kusnandar Miftakhurrozaq,Hartanto Mulyo Raharjo,Maya Nurwartant
Yunita,Faisal Fikri**



Pharmacognosy Journal,15(2):286-289

DOI: 10.5530/pj.2023.15.40

Published: Wed, 26-Apr-2023

[Read More \(/article/1994\)](#)

Original Article

The Effect of Preconception Moringa Leaf Extract Supplementation on Cotinine Serum Levels in Pregnant Women Exposed to Tobacco Smoke (/article/1995)

Haerani Harun,Anwar Daud,Ridwan Amiruddin,M Sabir,Veni Hadju,Anwar Mallongi, , Tamrin Talebe,M. Faris

**Pharmacognosy Journal**,15(2):290-293

DOI: 10.5530/pj.2023.15.41


Published: Wed, 26-Apr-2023

[Read More \(/article/1995\)](#)

Original Article

Relationship Between Individual Characteristics and the Risk of Exposure to Heat Stress in Indonesian Fishermen (/article/1996)

Putri Ayuni Alayyannur,Dani Nasirul Haqi,Fatimatuz Zahroh,Tazkia Azrielda Munib,Moh. Malik Alhakim,Diah Pramesthi Ningrum

**Pharmacognosy Journal**,15(2):294-297

DOI: 10.5530/pj.2023.15.42


Published: Wed, 26-Apr-2023

[Read More \(/article/1996\)](#)

Original Article

Another Mode of Action of Temephos Against Aedes aegypti Larvae: A Stomach Poison Investigation (/article/1997)

Hebert Adrianto,Sri Subekti,Heny Arwati,Etha Rambung,Hanna Tabita Has Silitonga,Etik Ainun Rohmah

**Pharmacognosy Journal**,15(2):298-303

DOI: 10.5530/pj.2023.15.43

Published: Wed, 26-Apr-2023

[Read More \(/article/1997\)](#)

Original Article


The Aqueous Leaf Extract of Moringa Oleifera had Immunomodulatory Effects on Sheep that had Been Infected by Fasciola Gigantica, Clostridium Novyi Naturally and Impact to Fasciola gigantica Non-Embryonated (/article/1998)

Maslichah Mafruchati,Wan Irjani Wan Ismail

Pharmacognosy Journal,15(2):304-309

DOI: 10.5530/pj.2023.15.44

Published: Wed, 26-Apr-2023

[Read More \(/article/1998\)](#)

Original Article

The Effect of Moringa Leaf Extract Intervention Since Preconception Period on the Prevention of Oxidative Stress in Pregnant Women and Adverse Pregnancy Outcomes (/article/1999)

,Veni Hadju,Arsunan Arsin,Aminuddin Syam,Anwar Mallongi,Abd. Farid Lewa,Haerani Harun,

Pharmacognosy Journal,15(2):310-314

DOI: 10.5530/pj.2023.15.45

Published: Wed, 26-Apr-2023

[Read More \(/article/1999\)](#)

Original Article

Antihypertensive Choices during Pregnancy in Limited Setting (/article/2000)

, ,Salsabila Nabilah Rifdah,Agus Sulistyono

Pharmacognosy Journal,15(2):315-318


DOI: 10.5530/pj.2023.15.46

Published: Wed, 26-Apr-2023

[Read More \(/article/2000\)](#)

Original Article

The Influence of Job Satisfaction and Organizational Commitment on Nurse Performance with Work Motivation as a Mediating Factor at I Lagaligo East Luwu



Hospital In 2022 (/article/2001)

Maria Ulfha Ibrahim, Alimin Maidin, Indahwaty Sidin, Fridawaty Rivai, Khalid Shaleh

Pharmacognosy Journal, 15(2):319-324

DOI: 10.5530/pj.2023.15.47

Published: Wed, 26-Apr-2023

[Read More \(/article/2001\)](#)

Original Article

The Antidepressant Effect of Fluoxetine and Mozart K448 Combination Therapy on Hippocampal Serotonin and BDNF Levels (/article/2002)

Lisa Pangemanan, Irwanto Irwanto, Margarita M. Maramis

Pharmacognosy Journal, 15(2):325-328

DOI: 10.5530/pj.2023.15.48

Published: Wed, 26-Apr-2023

[Read More \(/article/2002\)](#)

Research Article

Pharmacognostic Characteristics and Antioxidant Activity of Gendola Stem (Basella Rubra L.) Ethanol Extract from South Kalimantan (/article/2003)

Arnida Arnida, Dian Kurnia, Sutomo Sutomo

Pharmacognosy Journal, 15(2):329-332

DOI: 10.5530/pj.2023.15.49

Published: Wed, 26-Apr-2023

[Read More \(/article/2003\)](#)

Research Article

Analysis of Deferred Blood Donor Candidates at Dr. Sardjito Hospital, Yogyakarta, Indonesia (/article/2004)




Teguh Triyono, Tsaniatul Afifah, Usi Sukorini

Pharmacognosy Journal, 15(2):333-337

DOI: 10.5530/pj.2023.15.50

Published: Wed, 26-Apr-2023

Pharmacognosy Journal
(/article/2001)



Home (/) About Journal (v) Editorial Board (/editorial-board) For Authors (v)

Contact Us (/contact-us)

Pharmacognosy Journal (/)

An Open Access, Peer Reviewed Journal in the field of Pharmacognosy

Enter terms then hit Search...



Articles In Press (/inpress) Current Issue (/v15/i3) Archives (/archives)

RSS Feeds (/rss.xml) Submit Article (https://www.phcogj.info)



(http://www.phcogj.com/#/)

HOME (/) / EDITORIAL BOARD

View (/editorial-board)

What links here (/node/25/backlinks)

Time to read
1 minute

Editorial Board

Share



(https://www.facebook.com/)

/sharer.php?u=https%3A%2Fwww.phcogj.com%2Feditorial-board&t=Editorial+Board

Editors & Editorial Board Members (2021)

Dr.Djemli Samir

Department of Biology , Applied Neuroendocrinology Laboratory
Bach Mektar Annaba University
Algeria

Dr. Raghava Naidu, Ph.D

Department of Human Oncology,
University of Wisconsin,
1111, Highland Ave, Madison,
Wisconsin 53705, USA

Dr.Karim Raafat

Associate Professor of Pharmacognosy and Phytochemistry,
Pharmaceutical Sciences Department,
Faculty of Pharmacy,
Beirut Arab University (BAU),
Beirut 115020, Lebanon

Ourlad Alzeus Tantengco, MD-PhD Molecular Medicine

College of Medicine, University of the Philippines Manila
EDSA Shrine Road, Alabang, Manila, Philippines, 1000

Janib Achmad

Lecturer of Faculty of Fisheries and Marine Science,
University of Khairun Ternate
Kampus 2 Jalan Pertamina, Kelurahan Gambesi,
Ternate Selatan



Print

a+ a-

Muammar Fawwaz, Ph.D

Department of Pharmaceutical Chemistry
Faculty of Pharmacy
Universitas Muslim Indonesia
Makassar 90231, South Sulawesi, Indonesia

Hany Ezzat Khalil

Associate Professor,
College of Clinical Pharmacy,
King Faisal University,
KSA

Emad Yousif

Department of Chemistry
College of Science
Al-Nahrain University
Baghdad,Iraq

Sughosh Upasani

R.C Patel Institute of pharmacy,
Shirpur,Dist-Dhule,Maharashtra,
India.

Gurusiddaiah suresh kumar

Scientist

Dept of biochemistry
CSIR-CFTRI
Mysore, Karnataka, INDIA

Arjun Patra

Assistant Professor
School of Pharmaceutical Sciences
Guru Ghasidas Central University
Koni, Bilaspur - 495 009
Chattisgarh, India

Francis O. Atanu, Ph.D

Department of Biochemistry
Faculty of Natural Sciences
Kogi State University
Anyigba, Nigeria.

Vijay Kumar Chattu

Faculty of Medical Sciences
University of the West Indies
St. Augustine, Trinidad & Tobago.

Dr.Kunle Okaiyeto, PhD

Applied and Environmental Microbiology Research Group (AEMREG)
Department of Biochemistry and Microbiology
University of Fort Hare
Alice campus
5700, Alice
South Africa.

Dr. Srisailam Keshetti, Ph.D

Principal, University College of Pharmaceutical Sciences, Satavahana University
Karimnagar 505001
Telangana
INDIA

Dr. Gayathri M Rao

Associate Professor
Department of Biochemistry
Kasturba Medical Collge, Mangaluru.

Shuge Tian

Experimental Teaching Demonstration Center of TCM in Xinjiang Medical
University
Department of traditional medicine ,TCM
Xinjiang Medical University
Xinjiang CHINA 830054

Dr. Ramachandra Setty Siddamsetty,

Professor, Govt College of Pharmacy,
Mission Road, Bengaluru, INDIA

Dr. (Mrs.) Sayyada Khatoon

HOD, Pharmacognosy Division
CSIR-National Botanical Research Institute,
Rana Pratap Marg, Post Box 436,
Lucknow-226001 (U.P.) India

Dr. A. Sajeli Begum

Department of Pharmacy
Birla Institute of Technology & Science
Hyderabad, India

Olga Silva

Department of Pharmacological Sciences,
Faculdade de Farmácia,
Universidade de Lisboa, Portugal

Xinwen Wang

Department of Clinical Pharmacy
University of Michigan
USA

Roman Lysiuk

Department of Pharmacognosy and Botany,
Danylo Halytsky Lviv National Medical University,
Pekarska,69., Lviv 79010, Ukraine

Arif Nur Muhammad Ansori

Universitas Airlangga
Indonesia

Prabhu Manickam Natarajan

Specialist in Periodontics, College of Dentistry
Ajman University of Science and Technology, UAE

Neeta Mahesh Deshpande

Rural Institute of Ayurved Research Centre And Hospital

Mayani, Satara, Maharashtra, India

103752 reads

About

Pharmacognosy Journal (Phcog J.) covers different topics in natural product drug discovery, and also publishes manuscripts that describe pharmacognostic investigations, evaluation reports, methods, techniques and applications of all forms of medicinal plant research

Distinctions: The most widely read, cited, and known Pharmacognosy journal and website is well browsed with all the articles published. More than 50,000 readers in nearly every country in the world each month

ISSN : 0975-3575 ; Frequency : Rapid at a time publication (6 issues/year)

Indexed and Abstracted in : SCOPUS, Scimago Journal Ranking, Chemical Abstracts, Excerpta Medica / EMBASE, Google Scholar, CABI Full Text, Index Copernicus, Ulrich's International Periodical Directory, ProQuest, Journalseek & Genamics, PhcogBase, EBSCOHost, Academic Search Complete, Open J-Gate, SciACCESS.

Rapid publication: Average time from submission to first decision is 30 days and from acceptance to In Press online publication is 45 days.

Open Access Journal: Pharmacognosy Journal is an open access journal, which allows authors to fund their article to be open access from publication.

Submit your Next Article


- Online submission
- Highly indexed and abstracted
- 10 years of successful publishing
- Wider visibility though open access
- Higher impact with wider visibility
- Prompt review

Submit your next article to Phcog J

and be a part of many successful authors.

Create free account (<http://phcogj.info>) / Login (<https://www.phcogj.com/submissions/index.php/phcogj/index>)

Copyright © 2020 Pharmacogn J. All rights reserved.



Pharmacognosy Journal and its contents are licensed under a Creative Commons Attribution-Non Commercial-No Derivs 4.0 License. Permissions beyond the scope of this license may be available with editor@phcogj.com (<mailto:editor@phcogj.com>)

[Home \(/\)](#)

[Advertise with us \(/\)](#)

[Privacy Statement \(/\)](#)

The Antidepressant Effect of Fluoxetine and Mozart K448 Combination Therapy on Hippocampal Serotonin and BDNF Levels

Lisa Pangemanan^{1,2}, Irwanto Irwanto^{3,*}, Margarita M. Maramis⁴

Lisa Pangemanan^{1,2}, Irwanto Irwanto^{3,*}, Margarita M. Maramis⁴

¹Doctoral Program of Medical Science, Faculty of Medicine, Airlangga University, Surabaya, INDONESIA.

²Department of Child Health, Faculty of Medicine, Widya Mandala Catholic University, Surabaya, INDONESIA.

³Department of Child Health, Faculty of Medicine, Airlangga University, Surabaya, INDONESIA.

⁴Department of Psychiatry, Faculty of Medicine, Airlangga University, Surabaya, INDONESIA.

Correspondence

Irwanto Irwanto

Department of Child Health, Faculty of Medicine, Airlangga University, Surabaya, INDONESIA.

E-mail: irwanto@fk.unair.ac.id

History

- Submission Date: 12-02-2023;
- Review completed: 02-04-2023;
- Accepted Date: 06-04-2023.

DOI : 10.5530/pj.2023.15.48

Article Available online

<http://www.phcogj.com/v15/i6>

Copyright

© 2023 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

ABSTRACT

Despite the use of fluoxetine as a first-line therapy, some patients do not show a good therapeutic effect. Effective antidepressant therapy will reverse the low serotonin and BDNF levels found in depression. Mozart K. 448 was reported to yield a good therapeutic effect for depression. Based on findings, the combination of Mozart K. 448 and fluoxetine as a therapy for depression is very rare. Therefore, this study aimed to determine the effect of the combined therapy of fluoxetine and Mozart K. 448 on hippocampal serotonin and BDNF levels in an animal model of depression under CUMS conditions. In this study, the animal model of depression was administered three different treatments, i.e. fluoxetine, Mozart, or fluoxetine-Mozart combined therapy, respectively. Hippocampal serotonin and BDNF levels were assessed after 21 days of treatment. Statistical analysis was then carried out using T-test or Mann-Whitney test and ANOVA or Kruskal-Wallis. The fluoxetine-Mozart group has higher BDNF levels, but lower serotonin levels compared to other groups with values of $1,694 \pm 0.215$ and $44,533 \pm 3,275$, respectively.

Key words: Fluoxetine, Mozart, Serotonin, BDNF, Hippocampus.

INTRODUCTION

Fluoxetine is a first-line antidepressant.^{1,2} Fluoxetine can increase the low serotonin levels as well as low BDNF levels found in depression.^{3,4} Despite its usefulness, some patients did not respond well to this treatment.^{5,6} Music is one of the alternative and complementary therapies available for those patients.^{7,8} Slow rhythm music can increase BDNF levels in the brain.⁹ Mozart K448 is one of the music that can increase serotonin in certain brain areas.¹⁰ Mozart's music is one of depression's alternative therapy, and it acts differently on several areas of the brain to improve mood.⁹⁻¹³

Depression can show widely varied manifestations.^{14,15} The animal model used to evaluate the therapeutic effects of certain treatments should be able to mimic various pathogenesis and symptoms of depression in humans.¹⁶⁻¹⁸ An animal model of depression shows decreased serotonin and BDNF levels compared to the normal control.¹⁹⁻²² These changes can be reversed using effective management.^{19,23}

A combination of music and antidepressant therapy can be designed to obtain a better outcome in depressive patients.²⁴ Therefore, this study aims to determine the combined effects of music and fluoxetine therapy on hippocampal serotonin and BDNF levels.

METHODS

This study was carried out from January to March 2022 in the experimental animal laboratory (LPHC), Faculty of Medicine, Brawijaya University, Malang, Indonesia. The protocol was reviewed and approved by the Animal Care and Use Committee, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia, with reference number 2.KE.120.10.2021. Minimal

numbers of animals were used and all efforts were done to minimize suffering. This study was carried out to fulfill the requirements for a doctoral program.

Animals and study design

Male Wistar rats obtained from PT. Indoanilab (Bogor, Indonesia) undergoes one week of acclimatization. Later, they were randomly divided into two groups, namely control (no CUMS given) and CUMS (CUMS given). The CUMS group received psychological dominant CUMS protocols (cold swimming, foot shock, forced swimming, tail pierced, immobilization, no bedding, bright light, tail tied, isolation in a narrow dark space, predator exposure, wet bedding, and continuous light) by randomly giving 1 or 2 unexpected treatments daily for 21 days.²⁵ During the process, the same stressor was not given for two consecutive days.²⁶⁻²⁸ A sucrose preference test was carried out before and after the use of CUMS to determine the successful creation of an animal model of depression. The CUMS procedure was carried out continuously in the treatment period for another 21 days.

All animals were group-housed with two rats per cage, and they were separated by a wire mesh. The conditions in the room included a temperature of $23 \pm 2^\circ\text{C}$ and humidity of 40 to 70% with 12 hours/12 hours light/dark cycle (lights on at 6:00 AM). The samples were fed ad libitum with food, water, and sucrose 1.5%. The food given was weighed every morning before feeding, and the weight of the leftover was obtained the next day. Water and sucrose solutions were freshly prepared daily in separate bottles.

The inclusion criteria included male rats aged fourteen weeks old, which were exhibiting depressive-like behavior in the treatment group, as assessed by SPT. Meanwhile, the exclusion criteria were physical illness and disability, which were

Cite this article: Pangemanan L, Irwanto I, Maramis MM. The Antidepressant Effect of Fluoxetine and Mozart K448 Combination Therapy on Hippocampal Serotonin and BDNF Levels. *Pharmacogn J.* 2023;15(2): 325-328.

assessed by a veterinarian. The animals in the treatment group were randomly assigned to four groups, namely CUMS (CUMS + water), fluoxetine (CUMS + fluoxetine), Mozart (CUMS + Mozart K. 448 + water), and fluoxetine + Mozart (CUMS + Mozart K. 448 + fluoxetine) group. The control was given standard care and water using gavage feeding. Fluoxetine was dissolved in water at 2 ml/kg and given in a dose of 10 mg/kg body weight through gavage feeding. Dose selection for fluoxetine was carried out based on a previous study.²⁹ Mozart K. 448 was administered at a dosage of 60 – 80 dB from 06.00 PM – 06.00 AM. Water was also given in a dose of 2 ml/kg body weight through gavage feeding. A separate room was provided for groups without exposure to music, and all treatment lasted for 21 days. The design of the study is summarized in Figure 1.

Outcome measure

Weekly body weight measurement, sucrose preference test, and forced swimming test were done and published elsewhere. All rats were euthanized after the treatment period. The hippocampus was retrieved




Figure 1: Study's design

to measure serotonin levels and BDNF levels using ELISA (E0866Ra and E0476Ra respectively). Hippocampal tissue was taken, rinsed in cold PBS (pH 7.4), and weighed before homogenization. The sample was homogenized in PBS (tissue mass (g): PBS volume (mL) = 1:9) using a glass homogenizer on ice, followed by sonication. Centrifugation was carried out for 15 minutes at 12,000 RPM (4°C), and the supernatant was collected to measure the hippocampal serotonin levels and BDNF levels based on the manufacturer's instructions.

Statistical analysis

Data were presented as mean ± SD, and the difference between groups was analyzed using ANOVA or Kruskal-Wallis test, depending on the homogeneity. Furthermore, differences between the CUMS and fluoxetine-Mozart groups were analyzed with a t-test or Mann-Whitney test, depending on whether the data distribution was normally distributed or not (Kolmogorov-Smirnov test). A p-value of <0.05 was considered statistically significant in this study. Data analysis was carried out using IBM SPSS Statistics software version 23.0 (IBM Corp., Armonk, NY, USA).

RESULTS

A total of 39 rats with body weights of 92-159 g were included in this study (33 and nine rats in CUMS and control groups, respectively). Serotonin levels and BDNF levels between groups showed a significant difference, as shown in Table 1. Based on the results, the fluoxetine + Mozart group had the lowest serotonin level and the highest BDNF levels compared to the fluoxetine group and the CUMS group. Both the Mozart group and the fluoxetine + Mozart group show no significant difference compared to the control group, as shown in Table 2.

DISCUSSION

Despite the high efficacy and relatively well-tolerated antidepressant drug,^{30,31} some patients show a low response to the drug.³⁰ Additional treatment with other modalities needs to be considered to optimize the treatment's result.³¹ Mozart K. 448 is one of the music that can be used as a complementary therapy to improve depressive management.³²⁻³⁴ This study revealed that the combination of fluoxetine and Mozart can be used as an option for depression management, as it yields a similar response to control in terms of hippocampal BDNF levels (p>0.05).

The CUMS is a method commonly used to obtain a valid animal model of depression.³⁵ The modification in this study was done to mimic a human stressor that is psychologically dominant.²⁵ The CUMS protocol consists of chronic and unpredictable mild stressor exposure for 21

Table 1: Serotonin and BDNF between groups.

	ELISA level (ng/mL), mean ± SD					p value
	CUMS (n=8)	CUMS+F (n=7)	CUMS+F+M (n=7)	CUMS+M (n=8)	Control (n=9)	
Serotonin	50,242 ± 4,411	45,079 ± 4,432	44,533 ± 3,275	46,579 ± 4,047	50,672 ± 3,458	0.007**a
BDNF	1,757 ± 0.125	1,510 ± 0.229	1,694 ± 0.215	1,646 ± 0.090	1,759 ± 0.130	0.031*a

aANOVA*significant, p value <0.05 **significant, p value <0.01.

Table 2: Significance comparison between groups.

	Comparison between groups									
	CUMS Vs control	CUMS+F Vs control	CUMS+F+M Vs control	CUMS +M Vs control	CUMS Vs CUMS+F	CUMS Vs CUMS +F+M	CUMS Vs CUMS +M	CUMS+F Vs CUMS+F+M	CUMS+F Vs CUMS+M	CUMS+F+M Vs CUMS+M
Serotonin	0.825 ^a	0.013* ^a	0.003* ^a	0.040* ^a	0.042* ^a	0.015* ^a	0.105 ^a	0.798 ^a	0.505 ^a	0.306 ^a
BDNF	0.984 ^a	0.030* ^a	0.471 ^a	0.059 ^a	0.031* ^a	0.493 ^a	0.060 ^a	0.146 ^a	0.180 ^a	0.569 ^a

aT-test *Significant, p value <0.05 ** Significant, p value <0.01.

days. The sucrose preference test was used to evaluate the successful creation of an animal model of depression, as it represents anhedonia.³⁶ This study used psychologically dominant CUMS modification for 21 days to create an animal model of depression. The CUMS protocols were still given during the treatment periods.

Lower serotonin levels and BDNF levels were found in the depression compared to the normal control.³⁴ A decrease in brain serotonin levels alone is not sufficient to cause symptoms of depression.³⁷ A combination of serotonin pathway disruption and neurobiological vulnerability plays a role in the pathophysiology of depression.³⁸ Improvements in serotonin pathways contribute to depression improvements.³⁷ Lower BDNF levels that can be reversed in a well-tolerated treatment is found in an animal model of depression,²³ as well as in human.³⁹ This study found that the CUMS group indeed showed lower serotonin levels compared to the control group. While it did not achieve a significant difference statistically ($p > 0.05$), it did show a trend that fits with the depression paradigm. This study also found that the CUMS group had lower serotonin levels than the control group, although the difference was not significant statistically ($p > 0.05$). Marzban *et al.* found that Mozart K. 448 can indeed increase hippocampal BDNF levels.¹¹ Moraes *et al.* found that music exposure can increase serotonin.¹⁰ Among the treatments group, the fluoxetine + Mozart group had the highest BDNF levels (1,6940.215), while the Mozart group had the highest serotonin levels (46,579±4,047) in this study

Serotonin and BDNF affect each other.⁴⁰ Decreased BDNF levels will affect serotonin sensitivity.⁴⁰ Subsequently, this will reduce resilience to the environmental stressor and cause phenotype alteration that will last for life.^{37,40} Although fluoxetine increases serotonin levels, the response will not be optimal if the BDNF level is decreased.^{40,41} This study found that the fluoxetine + Mozart group had the highest BDNF levels with the lowest serotonin levels, although it did not reach statistical significance.

This study provided additional data related to the impact of fluoxetine and Mozart K. 448 combined therapy on serotonin levels and BDNF levels. This study has several limitations. First, hippocampal serotonin levels and hippocampal BDNF levels measurement were done after the treatment was given only. Second, the number of cells as well as neuron complexity in the hippocampus was not evaluated. Third, the serotonin and BDNF receptors were not assessed in this study.

CONCLUSION

The group treated with the combined therapy of fluoxetine and Mozart for 21 days had higher BDNF levels, but lower serotonin levels compared to other groups. This study found that therapy in addition to standard therapy did not necessarily provide a better outcome in all aspects. The addition of Mozart to fluoxetine treatment resulted in higher BDNF levels compared to either Mozart or fluoxetine alone.

ACKNOWLEDGMENT

The authors are thankful to the LPHC and Physiology Laboratory, Faculty of Medicine, Brawijaya University, Malang, Indonesia for providing the necessary facilities. This study received funding from Widya Mandala Catholic University Surabaya, Indonesia, a university that sent LP for the doctoral program at the Faculty of Medicine, Airlangga University, Surabaya, Indonesia. This study did not receive any funding.

CONFLICTS OF INTEREST

There is no competing interest.

AUTHOR'S CONTRIBUTION

LP, II, and MMM: designed the research, methodology, validation, formal analysis and review, and editing. LP: data collection, and

writing original draft preparation. All authors have read, reviewed, and approved the final manuscript.

RECOMMENDATIONS


The addition of Mozart to fluoxetine treatment can be used as it resulted in higher BDNF levels compared to either Mozart or fluoxetine alone. Further research needs to be done to unveil the pathways of depression improvement in Mozart and fluoxetine combined therapy.

REFERENCES

- Algristian H, Bintarti TW, Baroroh RNM, Leila Q, Ulfa R, Amelia K, *et al.* Protective effect of lavender essential oils on depression and multi-organ stress. *Bali Med J.* 2022;11(3):1357-63.
- Faisal FO, Algristian H, Azizah N. Anticipating suicide act of patient with borderline personality disorder and history of severe depression. *Bali Med J.* 2022;11(2):910-2.
- Blardi P, De Lalla A, Leo A, Auteri A, Iapichino S, Di Muro A, *et al.* Serotonin and Fluoxetine Levels in Plasma and Platelets After Fluoxetine Treatment in Depressive Patients. *J Clin Pharmacol.* 2002;22(2):131-6.
- Gupta K, Gupta R, Bhatia MS, Tripathi AK, Gupta LK. Effect of Agomelatine and Fluoxetine on HAM-D Score, Serum Brain-Derived Neurotrophic Factor, and Tumor Necrosis Factor- α Level in Patients with Major Depressive Disorder with Severe Depression. *J Clin Pharmacol.* 2017;57(12):1519-26.
- Zohar AH, Eilat T, Amitai M, Taler M, Bari R, Chen A, *et al.* An exploratory study of adolescent response to fluoxetine using psychological and biological predictors. *PeerJ.* 2018;2018(1):1-13.
- Maramis MM, Pantouw JG, Lesmana CBJ. Depression screening in Surabaya Indonesia: Urgent need for better mental health care for high-risk communities and suicide prevention for men. *Int J Soc Psychiatry.* 2021;67(5):421-31.
- Paul N, Lotter C, van Staden W. Patient Reflections on Individual Music Therapy for a Major Depressive Disorder or Acute Phase Schizophrenia Spectrum Disorder. *J Music Ther.* 2020;57(2):168-92.
- Ardani Y, Putranto R, Yunir E, Koesnoe S, Muhadi, Shatri H, *et al.* Effects of virtual music therapy on Burnout Syndrome in health workers in Tertiary Hospitals. *IJBS.* 2023;17(1):2302-906.
- Angelucci F, Ricci E, Padua L, Sabino A, Tonali PA. Music exposure differentially alters the levels of brain-derived neurotrophic factor and nerve growth factor in the mouse hypothalamus. *Neurosci Lett.* 2007;429(2-3):152-5.
- Moraes MM, Rabelo PCR, Pinto VA, Pires W, Wanner SP, Szawka RE, *et al.* Auditory stimulation by exposure to melodic music increases dopamine and serotonin activities in rat forebrain areas linked to reward and motor control. *Neurosci Lett.* 2018;673:73-8.
- Marzban M, Shahbazi A, Tondar M, Soleimani M, Bakhshayesh M, Moshkforoush A, *et al.* Effect of Mozart Music on Hippocampal Content of BDNF in Postnatal Rats. *Basic Clin Neurosci.* 2011;2(3):21-6.
- Putra AANK, Irwanto I, I'tishom R, Setyo-boedi B, Mustakim MRD. Mozart music stimulation effect on wistar rats' neurogenesis. *Bali Med J.* 2023;12(1):921-5.
- Amigo TAE, Mariati M. Music and video music therapy are effective in reducing stress among elderly at Yogyakarta Social Service Center of Tresna Werdha, Abiyoso Pakem Unit, Sleman regency. *Bali Med J.* 2020;9(1):211-5.
- Nardi B, Francesconi G, Catena-Dell'Osso M, Bellantuono C. Adolescent depression: Clinical features and therapeutic strategies. *Eur Rev Med Pharmacol Sci.* 2013;17(11):1546-51.
- Setiawati Y, Wahyuhadi J, Joestandari F, Maramis MM, Atika A. Anxiety and Resilience of Healthcare Workers During COVID-19 Pandemic in Indonesia. *J Multidiscip Healthc.* 2021;14(1):1-8.
- Zhang Y, Wang Y, Lei H, Wang L, Xue L, Wang X, *et al.* Optimized animal model to mimic the reality of stress-induced depression in the clinic. *BMC Psychiatry.* 2017;17(1):171.

17. Katz RJ, Roth KA, Schmaltz K. Amphetamine and tranylcypromine in an animal model of depression: pharmacological specificity of the reversal effect. *Neurosci Biobehav Rev.* 1981;5(2):259-64.
18. Becker M, Pinhasov A, Ornoy A. Animal Models of Depression: What Can They Teach Us about the Human Disease? *Diagnostics (Basel).* 2021;11(1):123.
19. Sen S, Duman R, Sanacora G. Serum BDNF, Depression and Anti-Depressant Medications: Meta-Analyses and Implications. *Biol Psychiatry.* 2008;64(6):527-32.
20. De Morais H, de Souza CP, da Silva LM, Ferreira DM, Baggio CH, Vanvossen AC, *et al.* Anandamide reverses depressive-like behavior, neurochemical abnormalities and oxidative-stress parameters in streptozotocin-diabetic rats: Role of CB1 receptors. *Eur Neuropsychopharmacol.* 2016;26(10):1590-600.
21. Fitrikasari A, Wardani ND, Sumekar TA, Saktini F, Asikin HG, Sulchan M. The role of psychosocial stressors, carbohydrate and protein intake on serum serotonin and cortisol levels in patients with depression: a preliminary evaluation. *Bali Med J.* 2021;10(1):137-41.
22. Purnawati S, Wrasati LP, Jaya Lesmana CB, Megantara S, Lesmana R. A study of molecular docking of l-tryptophan ligand as a compound in pineapples and bananas binding with the human serotonin transporter (SERT). *Bali Med J.* 2022;11(3):1243-9.
23. Jin X, Liu P, Yang F, Zhang YH, Miao D. Rosmarinic acid ameliorates depressive-like behaviors in a rat model of CUS and up-regulates BDNF levels in the hippocampus and hippocampal-derived astrocytes. *Neurochem Res.* 2013;38(9):1828-37.
24. Bulaj G. Combining non-pharmacological treatments with pharmacotherapies for neurological disorders: A unique interface of the brain, drug-device, and intellectual property. *Front Neurol.* 2014;5(1):126.
25. Pangemanan L, Irwanto I, Maramis MM. Psychological dominant stressor modification to an animal model of depression with chronic unpredictable mild stress. *Vet World.* 2023;16(3):595-600.
26. Maramis MM, Mahajudin MS, Khotib J. Impaired Cognitive Flexibility and Working Memory Precedes Depression: A Rat Model to Study Depression. *Neuropsychobiology.* 2021;80(3):225-33.
27. Pandey DK, Pati D, Joshi A, Mahesh R. Chronic Unpredictable Stress: Possible Animal Model of Comorbid Depression. *Int J Preclin Pharm Res.* 2010;1(1):54-63.
28. López López AL, Escobar Villanueva MC, Brianza Padilla M, Bonilla Jaime H, Alarcón Aguilar FJ. Chronic unpredictable mild stress progressively disturbs glucose metabolism and appetite hormones in rats. *Acta Endocrinol (Buchar).* 2018;14(1):16-23.
29. Klomp A, Václavů L, Meerhoff GF, Reneman L, Lucassen PJ. Effects of chronic fluoxetine treatment on neurogenesis and tryptophan hydroxylase expression in adolescent and adult rats. *PLoS One.* 2014;9(5):e97603.
30. Zhou X, Teng T, Zhang Y, Del Giovane C, Furukawa TA, Weisz JR, *et al.* Comparative efficacy and acceptability of antidepressants, psychotherapies, and their combination for acute treatment of children and adolescents with depressive disorder: a systematic review and network meta-analysis. *Lancet Psychiatry.* 2020;7(7):581-601.
31. Cipriani A, Zhou X, Del Giovane C, Hetrick SE, Qin B, Whittington C, *et al.* Comparative efficacy and tolerability of antidepressants for major depressive disorder in children and adolescents: a network meta-analysis. *Lancet.* 2016;388(10047):881-90.
32. Verrusio W, Moscucci F, Cacciafesta M, Gueli N. Mozart Effect and Its Clinical Applications: A Review. *Br J Med Res.* 2015;8(8):639-50.
33. Papadakakis A, Sidiropoulou K, Panagis G. Music exposure attenuates anxiety- and depression-like behaviors and increases hippocampal spine density in male rats. *Behav Brain Res.* 2019;372:112023.
34. Aalbers S, Fusar-Poli L, Freeman RE, Spreen M, Ket JC, Vink AC, *et al.* Music therapy for depression. *Cochrane Database Syst Rev.* 2017;11(11):CD004517.
35. Willner P. Reliability of the chronic mild stress model of depression: A user survey. *Neurobiol Stress.* 2017;6(1):68-77.
36. Willner P, Towell A, Sampson D, Sophokleous S, Muscat R. Reduction of sucrose preference by chronic unpredictable mild stress, and its restoration by a tricyclic antidepressant. *Psychopharmacology (Berl).* 1987;93(3):358-64.
37. Jauhar S, Cowen PJ, Browning M. Fifty years on: Serotonin and depression. *J Psychopharmacol.* 2023;026988112311618.
38. Fusar-Poli P, Allen P, McGuire P, Placentino A, Cortesi M, Perez J. Neuroimaging and electrophysiological studies of the effects of acute tryptophan depletion: A systematic review of the literature. *Psychopharmacology.* 2006;188(2):131-43.
39. Yeh SH, Lin LW, Chuang YK, Liu CL, Tsai LJ, Tsuei FS, *et al.* Effects of music aerobic exercise on depression and brain-derived neurotrophic factor levels in community dwelling women. *Biomed Res Int.* 2015;2015:135893.
40. Homberg JR, Molteni R, Calabrese F, Riva MA. The serotonin-BDNF duo: Developmental implications for the vulnerability to psychopathology. *Neurosci Biobehav Rev.* 2014;43(1):35-47.
41. Alboni S, van Dijk RM, Poggini S, Miliot G, Perrotta M, Drenth T, *et al.* Fluoxetine effects on molecular, cellular and behavioral endophenotypes of depression are driven by the living environment. *Mol Psychiatry.* 2017;22(4):552-61.

GRAPHICAL ABSTRACT



Cite this article: Pangemanan L, Irwanto I, Maramis MM. The Antidepressant Effect of Fluoxetine and Mozart K448 Combination Therapy on Hippocampal Serotonin and BDNF Levels. *Pharmacogn J.* 2023;15(2): 325-328.