

# NeuroQuantology


An Interdisciplinary Journal of Neuroscience and Quantum Physics



ISSN 1303 5150

## Quantum Physics and its Relation to the Nervous System

A  
n  
K  
a  
  
P  
u  
b  
l  
i  
s  
h  
e  
r



Sultan Tarlacı



## Volume 20 No 21 (2022)

### RADIO SQUARE DIFFERENCE D-DISTANCE NUMBER OF SNAKE GRAPHS

 Vishma George G, John Bosco K

DOI: 10.48047/NQ.2022.20.21.NQ99001

Pages: 1 - 6

### An Optimized HR Analytics Approach to Impact Decision Making Process in Banking Sector to increase employee performance

 Tanya Nagpal, Mridula Mishra

DOI: 10.48047/NQ.2022.20.21.NQ99003

Pages: 17 - 33

### Study of Public Transport Service Level of 34 'Friend Bus' Mamminasata Route Corridor 3

 Mulyadi Nur , Fatmawati Sabur

DOI: 10.48047/NQ.2022.20.21.NQ99004

Pages: 34 - 60

### Composition of Plankton in Waters and Digestive of Vanname Shrimp (*Litopenaeus vannamei*) with Semi-biofloc System of Cultivation

 Irsyah Rahmi , Diana Arfiati , Muhammad Musa

DOI: 10.48047/NQ.2022.20.21.NQ99005

Pages: 41 - 49

### ANALYSING THE PREVALENCE OF ACADEMIC RESILIENCE IN COLLEGE GOING STUDENTS


 L Jyoti Reddy

DOI: 10.48047/NQ.2022.20.21.NQ99008

DOI: 10.48049/NQ.2022.20.21.NQ99059

Pages: 494 - 502

### Comparative evaluation of Canthal distances and Canthal index in males and females

 Khushboo Rajput1\*, Dr. Vimal Modi, Dr. Kuldeep Kumar, Dr. Chandan Kumar Yadav, Dr. Shivani Dhingra

DOI: 10.48049/NQ.2022.20.21.NQ99060

Pages: 503 - 507


### PROPOSAL MODEL: THE INFLUENCE OF SPORTS ACTIVITIES ON WORKPLACE PRODUCTIVITY IN VIETNAM: THE MEDIATING ROLE OF STRESS MANAGEMENT AND WORKLIFE BALANCE

 Thi Diep Uyen Doan, Thi Diep Uyen Doan

DOI: 10.48049/NQ.2022.20.21.NQ99061

Pages: 508 - 516

### The Impact of Laparoscopic Gastric Sleeve on blood glucose and HbA1c: A Retrospective Cohort Study

 Aysha Saqer Mustafa, Aysha Saqer Mustafa, Dr. Zumin Shi, Dr. Marios Hadjicharalambous, Prof. Eiman Aleem, Mohammad Mollazehi

DOI: 10.48049/NQ.2022.20.21.NQ99061

Pages: 508 - 518

### ADAPTIVE RADIOTHERAPY WITH SURFACEGUIDED BREATHING FOR BREAST CANCER PATIENTS IN VIETNAM

 Thi Hue Bui, Anh Kiet Duong, Thanh Binh Nguyen

DOI: 10.48047/NQ.2022.20.21.NQ99062

Pages: 517 - 522


## Hashtag Emotional Study using Hidden Linguistic Extraction for Textual Characterization of Internet Abusive Behaviour

 Pawan Kuma

DOI: 10.48049/NQ.2022.20.21.NQ99063

Pages: 523 - 529

## Research scholars information seeking anxiety based on time spent for accessing resources from the Library and the Internet

 A.E.K. SELVAN DEVADOSS, Dr. K. KANNAN, Dr.P. SARAVANAN

DOI: 10.48049/NQ.2022.20.21.NQ99064

Pages: 530 - 535

## Analysis of Research Scholars' Information Literacy based on their Demographic Factors

 A.E.K. SELVAN DEVADOSS, Dr. K. KANNAN, Dr.P. SARAVANAN

DOI: 10.48049/NQ.2022.20.21.NQ99065

Pages: 536 - 541

## IMPACT OF ETHICAL LEADERSHIP STYLE ON GREEN HRM

 V VARISHA BEGUM, Dr. M. KAVITHA

DOI: 10.48049/NQ.2022.20.21.NQ99066

Pages: 542 - 545

## Expression of B-Cell Lymphoma 2 (BCL-2) Protein and Regeneration of Incision Wound Skin Surface Epithel Cells in Infiltration of Ropivacaine (Laboratory Experimental Study on Wistar Rats)

 Catur Zannuar Abidin, Prananda S. Airlangga, Imam Susilo

DOI: 10.48049/NQ.2022.20.21.NQ99067

Pages: 546 - 561

## Identifying the Best ML Algorithm to Predict Real-Time Data Classification Problems:A Non-ML to ML Algorithm Transition

 Dr. RohitaYamaganti, Dr.P.N.SivaJyothi, NelliSreevidya

DOI: 10.48049/NQ.2022.20.21.NQ99068

Pages: 553 - 563

### Correlation Between Degrees of Stress and Hair Cortisol Levels in Anesthesiology Residents Caring for Critically Ill Patients at Dr Soetomo Hospital Surabaya

 Zakaria, Prananda Surya Airlangga, Yunias Setiawati

DOI: 10.48049/NQ.2022.20.21.NQ99069

Pages: 581 - 595

### SOCIO-ECONOMIC ASPECTS OF DEVELOPMENT OF INLAND FISHERIES IN KERALA

 Anu B, Dr.D. Deepha Jeya Merlin

DOI: 10.48049/NQ.2022.20.21.NQ99070

Pages: 596 - 601

### EFFECT OF DETERMINANTS ON PURCHASE INTENTION TOWARDS ENERGY-EFFICIENT HOUSEHOLD APPLIANCES

 Dr.A.Michael John SJ, M. Siva Bharathy,

DOI: 10.48049/NQ.2022.20.21.NQ99071

Pages: 602 - 607

### Evaluation of System Reliability: CNG Kit

 KUNDAN MISHRA

DOI: 10.48049/NQ.2022.20.21.NQ99073

Pages: 625 - 630

### A Critical Study of Scientific value and Psychological values implied in Science Textbook of NCERT of Class 7th

 Tripty Saini, Dr. Vibha KaushiK

DOI: 10.48049/NQ.2022.20.21.NQ99074

Pages: 631 - 635

### AN ALLELIC LOSS AT CHROMOSOME 9p21 TO DETERMINE

# Neuroquantology



## Editorial Board

Editor-In-Chief, Founder

Prof.Dr.Riyaz Ahmed Abdul Khan, Editor in Chief (Neuroquantology)

### Advisory and Editorial Board

Mohammadreza Khoshbin-e-Khoshnazar, Physics Department, Curriculum Development Center Organization & Educational Planning

Dr. Mansi Babbar, Postdoctoral Research Fellow, Laboratory of Molecular Gerontology, National Institute on Aging/National Institute of Health, 251 Bayview Blvd, Baltimore, MD 21224., United States

Dirk K.F. Meijer, University of Groningen, Netherlands

Attila Grandpierre, Konkoly Observatory, Budapest, Hungary; Chapman University, Orange, California, United States

Alfredo Pereira, Institute of Biosciences UNESP - Botucatu, Brazil

Daegene Song, Chungbuk National University, Korea, Republic of

Donald Mender, Lecturer in Psychiatry, Yale University, New Haven CT USA, United States

Diana Gasparyan, National Research University - Higher School of Economics (NRU-HSE), Associate Professor, Russian

### Federation

Gelengul Kocaslan, Department of Economics, Faculty of Economics, Istanbul University, Turkey

Greg P. Hodes, PhD, University of Kansas, United States

Gustav Bernroider, Univ. Prof. University of Salzburg, Dept. of Ecology & Evolution, Austria

Kemal Koç, Baskent University, Dept of Physics, Ankara, Turkey

Michael B. Mensky, P. N. Lebedev Physics Institute, Moscow, Russian Federation

Fred H. Thaheld, Independent Researcher, Atomic, Molecular and Quantum Physics, United States

Michel Planat, Institut FEMTO-ST, Departement LPMO, 32 Avenue de l'observatoire, Cedex, France

Panar, Askdar University

Pierre Uzan, Fondation Santa des Etudiants de France, University Paris, France

Dr Pravin Thomas, University Hospitals Birmingham NHS Foundation Trust, United Kingdom

Sam Anderson, Singapore

Subhash Kak, School of Electrical and Computer Engineering, Stillwater, United States

Sinan Canan, Yadaram Beyazat University, Dept of Physiology, Turkey

Tatyana Petrovna Lifintseva, Professor of the Department of History of Philosophy Faculty of Philosophy National Research University Higher School of Economics, Russian Federation

Aner Tan, Cukurova University, Department of Physiology, Turkey

Włodzisław Duch, Nicholas Copernicus University, Dept of

Physics, Poland

Dr Jiapei Dai, Wuhan Institute for Neuroscience and Neuroengineering Director of Chinese Brain Bank Center South-Central University for Nationalities Minzhu Dadao Wuhan, Hubei China, China

Dr. Richard Allen Sieb, Independent Researcher Edmonton, Alberta, Canada, Canada

Dr John Gardiner, Freelance Scientist, Glebe 2037, Australia, Australia

Dr Paul Bessler, Department of French, University of Toronto 50 St. Joseph Street, Toronto, Ontario, Canada, Canada

Michele Caponigro, Italy

Dr. Contzen Pereira, Independent Researcher, Mumbai, India, India

Dr Lex Neale, Integral Research Center, JFK University, CA, United States, United States

Dr Dorina Lauritano, Department of Medicine and Surgery., Italy

Dr Germano Resconi, Catholic University, Brescia, Italy, Italy

Quantum Dynamical Psychology

Mark Germine, Psychoscience, Mount Shasta, California, United States

Mladen Pecujlija, Serbia

Quantum Paradigms of Psychopathology

Mladen Pecujlija, Serbia

Prof. Massimo Cocchi, Professor of Nutrition Biochemistry Department of Veterinary Medical Sciences University of

Bologna, Via Tolara di Sopra, Italy

Mansoor Malik, Howard University Hospital, Washington, United States

Altered States of Consciousness

Dr Raul Valverde, Concordia University

Hideyuki Kokubo, Meiji University, Japan

Alan S. Haas, Harvard University Department of Psychology, United States

Bruce Greyson, Carlson Prof. of Psychiatry & Neurobehavioral Sciences, Director, Division of Perceptual Studies University of Virginia Health System, United States

Cheryl Fracasso, Saybrook University, State of Washington, Department of Social & Health Services, Developmental Disability Administration, United States

Harris Friedman, Psychology at the University of Florida, United States

Stanley Krippner, Saybrook University, Oakland, CA, USA, United States

Vernon M Neppe, Director, Pacific Neuropsychiatric Institute, Professor, Dept Neurology and Psychiatry, United States

Philosophy of Consciousness and Mind

Diana Gasparyan, National Research University - Higher School of Economics (NRU-HSE), Associate Professor, Russian Federation

Tatyana Petrovna Lifintseva, Professor of the Department of History of Philosophy Faculty of Philosophy National Research University Higher School of Economics, Russian Federation

Dr. Sergey Korotaev, Geoelectromagnetic Research Centre of Schmidt Institute of Physics of the Earth Russian Academy of Sciences; Physics Department of Bauman Moscow State Technical University Moscow, Russia, Russian Federation

Social Neuroscience

Gelengul Kocaslan, Department of Economics, Faculty of Economics, Istanbul University, Turkey

Stephanie Cacioppo, University of Geneva and Syracuse University, United States

Basics of Quantum Physics

L.A. Glinka, Poland

Paolo Di Sia, University of Padova, Italy

Nagadi Tidjani, Department de physique, Faculty des Sciences Exactes et Appliquees, University d'Oran1, Algeria

Antonio Manzalini, Telecom Italia, Italy

Prof. Carlos Eduardo Maldonado, Universidad del Rosario, Colombia

Teruaki Nakagomi, Professor Emeritus, Kochi University, Department of Information Science, Japan

Michael B. Mensky, P. N. Lebedev Physics Institute, Moscow, Russian Federation

Basics of Neuroscience

Burak Yulug, Dept of Neurology, Assoc. Prof. Private Hospital, Swaziland

James Giordano, Professor, Integrative Physiology and

Interdisciplinary Neurosciences, United States

Aner Tan, Cukurova University, Department of Physiology, Turkey

Behavioral Neuroscience

Associate Professor Dr Mohd Normani Zakaria, Universiti Sains Malaysia, Malaysia

Mohammad Hossien Delshad, PhD of Health Education and Health Promotion Department, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran, Islamic Republic of

Dimensional Biopsychophysics

Vernon M Neppe, Director, Pacific Neuropsychiatric Institute, Professor, Dept Neurology and Psychiatry, United States

Dr. Anabela Cardoso, Licentiate in Germanic Philology, Faculty of Letters, Classical University of Lisbon Doctor of Public Service (Honoris Causa), Roger Williams University, Bristol, R. I., USA., United States

Neuropsychology and Mental Health

Fernando Fajardo, Department of Psychology, University of Extremadura, Spain

Child and Adolescent Psychiatry

Dr. Katayoon Razjouyan, Associate Professor of Child and Adolescent Psychiatry, Shahid Beheshti University of Medical Sciences, Tehran, Iran



Neurology Disorders

Angelo Lavano, University Magna Graecia Catanzaro,  
Department of Neurosurgery, Italy

Honorary Retired Members

Michael Persinger, Died August 14, 2018. Laurentian University,  
Canada

Ivan Godfroid, Belgium

Dr. Florin Gaiseanu, Independent Senior Researcher 1-st Degree  
(Professor), Ph. D. (Physics), Information Science and  
Technology., Romania

Alex Kaivarainen, Finland

Brian Josephson, United Kingdom

Danko D. Georgiev, Bulgaria

Stuart Hameroff, United States

English Editors

Alec Rylands, Turkey

Dr. Stanley Grant, University of Melbourne, Australia

Dr. Andrew Gavin, Universität München, Germany

Joseph Charlie, Universiteit Utrecht, Netherlands

Dr. Lucas Jacob, Monash University, Australia

Aiden Jackson, Massachusetts Institute of Technology, United  
States

Derya Turan, Turkey

Yeliz Turan, Turkey

**Dr Atul Viraj Wadagale**


(MSc, PhD)

Assistant Professor(Statistics)

Department of Community Medicine

Vilasrao Deshmukh Government Medical College, Latur,  
Maharashtra, INDIA.

**Journal Cover**



This block contains a dashboard or sidebar with several key metrics and navigation options. At the top left, there is a large orange square with the text 'Q3' in white, indicating the journal's quartile ranking. To its right, the text 'Atomic and Molecular Physics, and Optics' is displayed, followed by 'best quartile'. Below this, the journal title 'Neuroquantology' is shown in a light grey font. The 'SJR 2021' index is prominently displayed as '0.29' in a large, bold, black font. A Creative Commons Attribution-NonCommercial license icon (CC BY-NC) is visible. A line graph with an orange line and a red dot at the end is partially visible. Below the graph, the text 'powered by scimagojr.com' is written in a bold, black font. At the bottom of this section, there is a copyright notice: 'copyright © 2023 Neuroquantology. All Rights Reserved.' The main navigation area includes a 'SUBMIT PAPER' button with a blue icon of a document and a checkmark. Below this are two sections: 'Indexed In' with a link to 'Indexing', and 'Useful Links' with links to 'Author Instruction', 'Publication Ethics', and 'Open Access Statement'.



# Expression of B-Cell Lymphoma 2 (BCL-2) Protein and Regeneration of Incision Wound Skin Surface Epithel Cells in Infiltration of Ropivacaine (Laboratory Experimental Study on Wistar Rats)

Catur Zannuar Abidin<sup>1\*</sup>, Prananda S. Airlangga<sup>2</sup>, Imam Susilo<sup>3</sup>

Corresponding Author: Catur Zannuar Abidin

<sup>1</sup>Anesthesiology and Reanimation Department of Medical Faculty of Universitas Airlangga / Soetomo General Hospital, Surabaya, Indonesia.

<sup>2</sup>Anesthesiology and Reanimation Department of Medical Faculty of Universitas Airlangga / Soetomo General Hospital, Surabaya, Indonesia.

<sup>3</sup>Anatomical Pathology Department of Medical Faculty of Universitas Airlangga/ Soetomo General Hospital, Surabaya, Indonesia.

<sup>1</sup>\*E-mail: [zanncatur@gmail.com](mailto:zanncatur@gmail.com)

## Abstract

**Background:** Inadequate pain management will result in disruption in the healing process. Cell regeneration and proliferation of epithelial cells is an important part of the wound healing process. Bcl-2 is one of the regulators in apoptosis and cell proliferation. **Materials and Method:** The treatment group was given ropivacaine infiltration while the control group was not given any treatment. Each group Thirty-two male Wistar white rats were divided into 2 treatment and control groups, then given a superficial wound (incision) on the back of each rat which was divided into sub groups, each terminated on the 3rd and 7th day. Histological evaluation to assess the process of cell regeneration and proliferation as well as BCL-2 protein expression at the edges of the wound using immunohistochemistry. The data obtained will then be processed using SPSS. **Result:** The results showed that there were differences in cell regeneration and Bcl-2 expression at the wound edges between the group that was given ropivacaine infiltration and the group that was not given infiltration. **Conclusion:** Administration of ropivacaine by infiltration has the effect of increasing the proliferation of epithelial cells in the superficial wound healing of wistar rats.

**Key Words:** Ropivacaine, Cell Regeneration, Bcl-2

**DOI Number:** 10.48047/NQ.2022.20.21.NQ99067

**Neuro Quantology 2022; 20(21): 546-561**

## Introduction

The skin was the body's outermost organ and serves crucial roles in maintaining life by controlling body temperature, regulating water and electrolyte balance, and protecting the body from microorganisms. However, the function was no longer adequately performed when the skin disturbed for any reason (chronic wound, burn, neoplasm, or trauma). Consequently, it was crucial to ensure its integrity as quickly as feasible (Enoch & Price, 2004).

Any wound causes a loss of skin integrity resulting in an imbalance in function, anatomy and possibly accompanied by disability or even death. Skin wound repair through a well-organized multiphase process (inflammatory stage, cell proliferation, matrix deposition to remodeling phase) between the interactions of various cell types, growth factors and cytokines that led to the wound area was aimed to seal the skin and produce tissue repair (Cohen & Mercandetti, 2016; Sunarto et al., 2019).



Multiple tissue and cell types, including inflammatory cells, fibroblasts, keratinocytes, endothelial cells, and macrophages, should work together to heal a wound. Cytokines, growth factors, and extracellular matrix (ECM) components all exert strong control over these cells. Numerous growth factors, including platelet-derived growth factor (PDGF), epidermal growth factor (EGF), fibroblast growth factor (FGF), transforming growth factor (TGF), and nerve growth factor (NGF), were indeed produced and released by inflammatory cells, keratinocytes, and fibroblasts in the wound and border areas (J.-C. Chen et al., 2014).

Wounds on the skin of adults could be regenerated by repair or regeneration, and there were major differences between the two processes. After a loss of function, repair consists of physiological modifications that create continuity without seeking to restore the original tissue. Regeneration, on the other hand, tries to rebuild damaged tissue with identical replicas in order to restore tissue form and function. Mature mammalian skin does not spontaneously regenerate, but rather heals through scar formation (Borena et al., 2015).

However, surgical wounds cause acute pain and may result in a variety of negative outcomes due to inefficient patient care, including anxiety, hemodynamic compromise, increased morbidity, reduced physical performance and quality of life, delayed recovery, prolonged use of opioids during and after hospitalization, and increased maintenance costs (Gan, 2017).

In this instance, transmission of nociceptive impulses from the periphery to the CNS owing to pain leads to a neuroendocrine stress response, a combination of local inflammatory substances (e.g., cytokines, prostaglandins, leukotrienes, TNF- $\alpha$  (tumor necrosis factor- $\alpha$ )) and systemic neuroendocrine mediators. The dominant neuroendocrine response to pain involves interactions between hypothalamic-pituitary-adrenocortical and sympathoadrenal suprasedgmental reflexes. These reflexes result in increased sympathetic tone, increased catecholamine and catabolic hormone secretion (e.g., cortisol, adrenocorticotropic hormone,

antidiuretics), glucagon, aldosterone, renin, and angiotensin (Hurley et al., 2015).

The function of a cellular protein from the lymphoma family 2 B-cell (Bcl-2) was to regulate the initiation of the intrinsic (mitochondrial) and extrinsic pathways of apoptosis (Abdurrahman et al., 2022). Recent studies have shown that Bcl-2 family proteins also contributed to the regulation of other intracellular pathways that have a significant impact on cell survival, including antiapoptosis, autophagy, endoplasmic reticulum (ER) stress response, intracellular calcium dynamics, mitochondrial dynamics and energy metabolism, and cell cycle development (Hatok & Racay, 2016; Hurley et al., 2015).

The inflammatory stimulus activates intracellular signaling pathways which in turn activate the production of inflammatory mediators. Primary inflammatory stimuli, including microbial products and cytokines such as interleukin-1 $\beta$  (IL-1 $\beta$ ), interleukin-6 (IL-6), and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), mediate inflammation through interaction with the TLR, IL-1 receptor (IL-1R), IL-6 receptor (IL-6R), and TNF receptor (TNFR) involved in cell regeneration and cell death through the extrinsic pathway (Chen et al., 2018).

Apoptosis or programmed cell death has various purposes, including the elimination of damaged or dangerous cells, the creation of tissues during embryonic development, and the maintenance and regeneration of tissue homeostasis. The intrinsic and extrinsic pathways were two distinct classical apoptotic processes that have been identified. The activation of the pathways occurs in a variety of distinct ways due to the utilization of specific cellular signal components. The extrinsic pathway was dependent on death receptors of the TNF family such as Fas ligand (CD95) or TNFR-1 and was activated upon binding of their ligands. Upon activation, adapter proteins such as the Fas-associated death domain (FADD) or TNFR-1-associated death domain (TRADD) were recruited and activate the caspase cascade. Conversely, intrinsic or mitochondrial apoptotic pathways were activated in response to different types of stress signals such as loss of growth factor (trophic factor) or DNA damage. It

was primarily controlled by the B-cell lymphoma protein family 2 (Bcl-2). The Bcl-2 protein was characterized by up to four homological domains Bcl-2 (BH), BH1-4, and uses pro- or antiapoptosis functions.

Members of the antiapoptotic Bcl-2 family (that is, Bcl-2, Bcl-xL, Mcl-1, and A1) protected cells from apoptotic stimuli by binding to and inactivating proapoptotic antagonists. Caspase-8 also cuts BH3-only protein BH3 - interacting-domain death agonist (Bid) protein to produce tBid fragments which then bind Bcl-2 in the outer mitochondrial membrane which leads to the formation of Bak/Bax pores and release of cytochrome c into the cytosol, and activation of the intrinsic apoptotic pathway (Lodish et al., 2016).

Following the initial hemostatic event, early wound healing was characterized by the invasion of neutrophils, macrophages, and lymphocytes, which serve as a source of inflammatory and growth-promoting cytokines. Fibroblasts migrate, proliferate, and produce extracellular matrix components, hence contributed to the creation of granulation tissue. Infiltration and proliferation of cells should be sufficient for the initial stages of normal wound healing. The early decrease in apoptosis made possible this rapid surge in cell proliferation. Subsequently, as the inflammatory response begins to subside with wound healing and scar formation, there was a substantial decrease in cell number, which has been demonstrated to be mediated by increased apoptotic cell death (Bhan et al., 2013).

Infiltration of a local anesthetic reduces the severity of pain by suppressing pain impulse transmission pathways. This results in a reduction in the secretion of the hormone glucocorticoid, which in turn eliminates one of the elements that inhibits the healing of wounds (Hancı et al., 2012). Giving a local anesthetic injection around the wound was thought to change the inflammatory response and speed up wound healing (Utariani et al., 2020).

Subcutaneous administration has shown bacteriostatic and bactericidal results (Bhaskar, 2015). Infiltration of local anesthetics could

reduce the release of inflammatory mediators from neutrophils, reduce oxidant formation, has an anti-inflammatory effects by binding to G proteins (inhibiting adhesion of polymorphonuclear leukocytes, macrophages, and monocytes), increasing glutamate release, and interfering with the activity of several intracellular signaling pathways (Abrão et al., 2020).

Previous animal and human studies have shown that ropivacaine has less toxic effects than bupivacaine. From the description above, there were no studies that have directly observed the effect of local anesthetic infiltration around the wound on Bcl-2 gene expression and the regeneration process of skin epithelial cells. As such, we were encouraged to perform research on the effect of ropivacaine infiltration as a local anesthetic drug through inflammatory inhibition processes on changes in bcl-2 expression and the process of epithelial cell regeneration which plays an important role in the process of wound healing. This research was an experimental study on experimental rats wistar. Based on the background, this study aimed to analyze the effect of infiltration ropivacaine around the wound on bcl-2 expression and epithelial cell regeneration on the healing process of Wistar rat incision wounds.

## Literature Review

### Wound

Damage to the integrity of the skin's epithelium was what constitutes a wound, which may also be followed by disturbances to the structure and function of the normal tissue that lies beneath the damaged area. Extensive tissue damage may happen from the exact disruption of tissue with a surgeon's knife (incision) (eg major trauma, burns). Likewise, the wound may be the result of a bruise, hematoma, laceration, or abrasion (Enoch & Sumartono, 1987).

The healing of skin wounds seems to be a complex physiological process involving the collaboration of numerous cell types and biochemical mediators in a progressive and overlapping process. The healing of wound-



induced lesions begins early in the inflammatory phase. In the end, wound healing ends in repair, which consists of the replacement of specialized structures due to collagen deposition, and regeneration, which corresponds to processes of cell proliferation and differentiation via preexisting tissue cells and/or stem cells. These methods were not mutually exclusive; – in other words, regeneration and repair may occur following a skin lesion in the same tissue depending on the cell strain affected by the injury (Cristina and Gonzalez, 2016).

### *Apoptosis*

A form of programmed cell death known as apoptosis occurs when a cell responds to various stimuli, either physiological or pathological, by undergoing a sequence of structural alterations. Apoptosis was characterized by cellular shrinkage, membrane blebbing, chromatin condensation, and nuclear disintegration. Caspases, a class of protease enzymes that could degrade the cytoskeleton and protein metabolites through proteolysis, were activated to produce this picture. These enzymes include poly (adenosine-5 disphosphate-ribose) polymerase (PARP), DNA-dependent protein kinase, lamins, proteins kinases, and actin. Extrinsic or death receptor route and extrinsic or mitochondrial pathway were the two primary apoptosis pathways (Lodish et al., 2016).

### *The Role of Apoptosis in Wound Healing*

Neutrophils were the first cells to arrive at the trauma area because they perform an important defense function against invading organisms. Neutrophil activity was also involved in local and systemic tissue damage through the release of oxygen free radicals and proteases. Most neutrophils enter to eliminate microorganisms, undergo apoptosis, and thereafter were rapidly and efficiently phagocytosed by macrophages in a process that does not cause further inflammation (Rai et al., 2005).

Macrophages also undergo apoptosis although the mechanism was less known. Tidball and St. Pierre reported that macrophages predominate

in the inflammatory response and disappear rapidly through apoptosis. However, the mechanism was suspected to be related to transforming growth factors that affect the downregulation of inflammation. In the wound remodeling phase, different cell populations need to be removed. Fibroblasts need to be downregulated along with decreasing concomitant vascularization. Preliminary studies showed that endothelial cells undergo apoptosis followed by loss of myofibroblasts. In vitro studies showed that c-myc was involved in fibroblast apoptosis in a process involving fas and fasL interactions on the cell membrane. Growth factors, especially insulin-like growth factor-1 to increase fibroblast proliferation. Conversely, another class of growth factors may be involved in decreasing fibroblast activity in wounds (Rai et al., 2005).

### *BCL-2*

Bcl-2 was B-cell lymphoma / leukemia-2 and the second of several proteins found in lymphoma. This gene was discovered due to its participation in B-cell malignancy, which involves chromosomal translocations that activate the most of genes in non-Hodgkin's B-cell follicular lymphoma. The Bcl-2 gene has over 230 kb of DNA and consists of three exons of which exon 2 and a small part of exon 3 encode proteins. Bcl-2 encodes 2 mRNAs, namely Bcl-2 $\alpha$  and Bcl-2 $\beta$ , of which only Bcl-2 $\alpha$  seemed to be biological relevance. The Bcl-2 protein was a membrane protein that has a molecular weight of 26 kDa and was located in the cytosolic portion of the nuclear envelope, endoplasmic reticulum and outer mitochondrial and cytoplasmic membranes (Borner, 2003; Hatok & Racay, 2016).


### *Ropivacaine*

Ropivacaine was a local anesthetic with a lengthy duration of action that was structurally similar to bupivacaine. Ropivacaine was a pure S(-) enantiomer, unlike bupivacaine which was a racemix. The S(-) enantiomer has lower toxicity and increases sensory block relative to motor block. The physicochemical properties of both enantiomers were the same, but they both has different affinities in places that might cause side

effects. The R(+) and S(-) enantiomers of local anesthetics has different affinities for sodium, potassium, and calcium ion channels. This resulted in a decrease in the toxicity of S(-) enantiomer towards the central nervous system and heart compared to R(+) enantiomer. Ropivacaine was slightly less potent than bupivacaine. Ropivacaine was suitable for use in epidural and regional anesthesia of similar duration to bupivacaine (Maheshwari & Naguib, 2015).

Ropivacaine inhibits conduction or decreases or prevents the typical transient rise in sodium permeability of excitable membranes caused by membrane depolarization. This effect was due to the interaction of ropivacaine with sodium channels (voltage-gated sodium channels). As ropivacaine acts on neurons, the threshold for electrical excitability increases, action potentials decrease, and impulse conduction slowed down. As a result, nerve conduction was obstructed (Maheshwari & Naguib, 2015).

In addition to inhibiting conduction in neurons in the peripheral nervous system, ropivacaine also affects the function of all organs where conduction or transmission of impulses occurs. Thus, the effect also occurs in the central nervous system, autonomic ganglia, neuromuscular junctions, and all muscles. The danger of side effects that occur depends on the circulating concentration of ropivacaine. In general, local anesthetics with the S-enantiomer were less toxic than the R-enantiomer (Kuthiala & Chaudhary, 2011).



**Figure 1:** Conceptual framework

Information :

- : Next arrow or influenced
- ▭ : Variable under study
- ⊥ : Inhibit the process

**Hypothesis:**

- 1) There is a difference in the expression of the Bcl-2 protein which was higher in the infiltration of ropivacaine around the wound during the healing process of incisions in Wistar rats.
- 2) There is a difference in the higher regeneration of skin surface epithelial cells when infiltrated with ropivacaine around the wound during the healing process of incisions in Wistar rats.

**Conceptual Framework**

**Materials and Methods**

**Research Design**

This research was an experimental study with a "randomized post test only control group design" using pure strain Wistar rats as research objects. The treatment given was infiltration of the local anesthetic ropivacaine with evaluation of Bcl-2 expression and the level of regeneration of



epithelial cells on the surface of the incision wound in the tissue. Preparations were made by HE and thricrom staining and then examined under a microscope.

### Research Sample

The experimental animals were Wistar rats obtained from the Experimental Animal Care Unit (UPHP) of the Department of Pharmacology, Faculty of Medicine, Universitas Airlangga.

#### Inclusion criteria :

- 1) Pure ancestry
- 2) Age two to two and a half months
- 3) Body weight 250-300 grams.
- 4) No anatomical abnormalities were seen

#### Exclusion criteria:

- 1) Sick during the 7 day adaptation period.
- 2) Has a previous scar.
- 3) There were anatomical abnormalities in mice.
- 4) There was a 10% weight loss during the adaptation period

The size of the Wistar rat sample was determined by the formula

$$n = \frac{2\sigma^2(Z_{1-\alpha/2} + Z_{1-\beta})^2}{(\mu_1 - \mu_2)^2}$$

- N : Total number of samples  
 $\alpha$  : 0,05  $\rightarrow$   $Z_{1/2\alpha} = 1,96$   
 $\beta$  : 0,10  $\rightarrow$   $Z_b = 1,28$   
 $\mu_1$  : Average Bcl-2 expression in previous studies  
 $\mu_2$  : Average Bcl-2 expression in the control group  
 $\sigma$  : SD in previous studies

Because there was no data in previous studies, based on the formula, the number of samples needed was 7,2 and rounded up to 8 per treatment group.

### Research Location

The selection of experimental animals was carried out at the Department of Pharmacology, Faculty of Medicine, Universitas Airlangga. Preparations were made at the Department of

Pharmacology, Faculty of Medicine, Universitas Airlangga and microscopic observations were made at the Department of Anatomical Pathology FK UNAIR/Dr. Sutomo Surabaya.

### Treatment Material

The experimental animals were Wistar rats aged 2 to 2,5 months and weighing 250-300 grams. The Wistar rat was one of hundreds of strains, originating from the American continent. Widely used as experimental animals in research in the fields of medicine and veterinary medicine. The Wistar rat were obtained from the Experimental Animal Care Unit at Universitas Airlangga, Surabaya. During the experiment, the experimental animals were placed in cages and given standard feed and sufficient drinking.

a) Materials and tools for injection and wound creation (Minor surgical devices in the form of a scalpel knife, chirurgis tweezers, Tegaderm and sterile doek).

b) Materials and means for infiltration (The tools used include: 1 ml disposable syringe, 0,2% ropivacaine solution)

c) Materials and tools for making histopathological preparations (The tools and materials needed include: 10% formalin buffer, alcohol (50%, 70%, 80%, 96%, 100%), xylol, liquid paraffin (Histoplast), Van Gieson stain, and Canadian balsam).


### Research procedure

- 1) The selection of Wistar rat was carried out at the Corba Animal Care Unit, Department of Pharmacology, Faculty of Medicine, Universitas Airlangga. The selection was carried out by taking into account the inclusion and exclusion criteria, so that 24 male rats were selected to be used in the study.
- 2) The next step was acclimatization to give the rats time to adapt to the research environment. Acclimatization was carried out for one week. Acclimatization was done by equalizing food, drink, and measuring the weight of the rats.
- 3) Wistar rat that have gone through the



- acclimatization period could be used for research. These rat were then grouped using a simple random method, grouped into 4 experimental groups. Each group contained 8 rats which were divided into 4 cages with the size of each cage 30x20x7 cm.
- 4) A total of 32 rats were divided into 4 groups which were randomly assigned. Each consisted of 8 rat for the Control-1 and Control-2 groups (K1 and K2) with 0,9% of NaCl infiltration, then 8 rat each for the treatment group (P1 and P2) with ropivacaine infiltration 0,2 %.
  - 5) All rats were anesthetized using ketamine-xylazine at a dose of 75-100 mg/kg + 5-10 mg/kg intraperitoneally for 10-30 minutes. Afterwards, the hair on the rat's back was shaved. A 2 cm incision was made on the rat's back.
  - 6) K1 and K2 rat group were given infiltration of 0,9% NaCl as much as 1 ml around the wound approximately 0,5 cm around the wound, then the wound was closed with

- 7) On day 3, 12 rats from groups K1 and P1 were anesthetized using ketamine. After the rats were anesthetized, a biopsy excision was performed on the scar tissue, which was sliced 3 cm square with a depth to the subcutis.
- 8) Histological preparations were made and microscopically assessed.
- 9) On the 7th day, 16 rats from groups K2 and P2 were anesthetized using ketamine. After the rats were anesthetized, a biopsy excision was performed on the scar tissue, which was sliced 3 cm square with a depth to the subcutis.
- 10) Histological preparations were made and microscopically assessed.



**Figure 2:** Operational framework

**Data Analysis**

The collected data were processed with the help of SPSS statistical software, then analyzed descriptive statistics in the form of percentages,

graphic tables or pictures. To answer the specific objective, a parametric statistical test is used if the data curve was normally distributed, and if the data curve was skewed, a non-parametric statistical test was used.



## Results and Discussion

### Demographic Characteristics of Research Subjects

This study was an experimental study using pure strain Wistar rats as research objects as many as 32. The sample was then divided into 4 groups, namely 2 treatment groups were given 0,2% infiltration and examined on the 3rd day (P1) and 7th day group (P2), and the other 2 groups were given 0,9% NaCl infiltration and also examined on the 3rd day (K1) and 7th day group (K2).

Parameters observed were the expression of Bcl-2 protein in basal layer epithelial cells in the area around the wound and cell regeneration which was estimated from the proliferation affinity of epithelial cells in the area of basal epithelial cells. Observation of Bcl-2 protein expression was carried out using immunohistochemical staining examination using specific monoclonal antibodies from paraffin block preparations. Calculation of Bcl-2 protein expression using immunohistochemical staining and measurement using the Allred Scoring Guideline which was the sum of the proportion score and intensity score of epithelial cells expressing Bcl-2 protein (Maae et al., 2011).

### Bcl-2 Expression and Cell Regeneration in Treatment

Tables 1 and 2 showed the characteristics of Bcl-2 expression and descriptions of cell regeneration in each study group. In this case, the highest average expression of Bcl-2 and the highest cell regeneration figure was P2 with values of 7,13 & 0,641 and 2 respectively 2,88 & 0,354. The median Bcl-2 expression and the

highest cell regeneration features were P2 with values of 7,00 (6-8) and 3,00 (2-3), respectively.


**Table 1:** Characteristics of Bcl-2 expression after administration of wound incisions in Wistar rats

| Group | Average | Standard Deviation | Median | Minimum | Maximum |
|-------|---------|--------------------|--------|---------|---------|
| K1    | 4,63    | 0,518              | 5,00   | 4       | 5       |
| P1    | 5,88    | 0,354              | 6,00   | 5       | 6       |
| K2    | 5,75    | 0,463              | 6,00   | 5       | 6       |
| P2    | 7,13    | 0,641              | 7,00   | 6       | 8       |
| Total | 5,84    | 1,019              | 6,00   | 4       | 8       |

**Table 2:** Characteristics of cell regeneration after infiltration of 0,2% ropivacaine in wound incisions of Wistar rats

| Group | Average | Standard Deviation | Median | Minimum | Maximum |
|-------|---------|--------------------|--------|---------|---------|
| K1    | 0,75    | 0,463              | 1,00   | 0       | 1       |
| P1    | 1,63    | 0,518              | 2,00   | 1       | 2       |
| K2    | 1,88    | 0,641              | 2,00   | 1       | 3       |
| P2    | 2,88    | 0,354              | 3,00   | 2       | 3       |
| Total | 1,78    | 0,906              | 2,00   | 0       | 3       |


On the 3rd day of observation (Figure 3) it was seen that Bcl-2 expression was higher in the treatment group (P1) than in the control group (K1) while on the 7th day of observation (Figure 4) Bcl-2 expression in the group (P2) was seen to be higher when compared to the control group (K2). In the treatment group Bcl-2 expression increased on the 3rd and 7th day of observation. Likewise, the control group also experienced an increase on the 3rd and 7th day of observation. The highest Bcl-2 expression was found in the P2 treatment group on the 7th day of observation.



**Figure 3.** Expression of Bcl-2 (green arrows) on day 3 by Immunohistochemical staining (X100) in the control group (A) and the treatment group (B)

(C)

(D)



**Figure 4:** Bcl-2 expression (green arrows) on day 7 with Immunohistochemical staining (X100) in the control group (C) and the treatment group (D)

*Effect of Ropivacaine Infiltration around the Wound on Bcl-2 Expression in the Healing Process of Incision Wounds in Wistar Rats*

**Table 3:** The normality test results for Bcl-2 expression in each study group

| Group | p-value |
|-------|---------|
| K1    | 0,0001  |
| P1    | 0,0001  |
| K2    | 0,0001  |
| P2    | 0,037   |

\*Shapiro-Wilk test of normality

Based on the results of the normality test, Bcl-2 expression in each study group was not normally distributed ( $p < 0,05$ ). The results of the different tests showed that there were significant differences between variables, except for the P1 and K2 groups which showed that there was no significant difference in the average Bcl-2 expression between the P1 and K2 groups (see Table 4).

**Table 4:** The results of different Bcl-2 expression tests for each study group

| Bcl-2 expression different test results | Group    | p-value |
|---|----------|---------|
| Research Group                          | K1       | 0,0001* |
|   | P1       |         |
|   | K2       |         |
|   | P2       |         |
| Group K1 and P1                         | K1<br>P1 | 0,001*  |
| Group K1 and K2                         | K1<br>K2 | 0,002*  |

|                  |    |        |
|------------------|----|--------|
| Group K1 and P2  | K1 | 0,001* |
|                  | P2 |        |
| Groups P1 and K2 | P1 | 0,535* |
|                  | K2 |        |
| Group P1 and P2  | P1 | 0,001* |
|                  | P2 |        |
| Group K2 and P2  | K2 | 0,001* |
|                  | P2 |        |

\*Mann-Whitney test

*Effect of Ropivacaine Infiltration around the wound on the description of Epithelial Cell Regeneration in the Wound Healing Process of Wistar Rats*

**Table 5:** Normality test results depict epithelial cell regeneration

| Group | p-value* |
|-------|----------|
| K1    | 0,0001   |
| P1    | 0,0001   |
| K2    | 0,037    |
| P2    | 0,0001   |

\*Shapiro-Wilk test of normality

Based on the results of the normality test, the description of epithelial cell regeneration in each study group was not normally distributed ( $p < 0,05$ ). The results of the different tests showed that there were significant differences between variables, except for groups P1 and K2, that is, there were no significant differences in the average appearance of epithelial cell regeneration between groups P1 and K2 (Table 6).

**Table 6:** Test results of different epithelial cell regeneration description in each research group

| Test results of different epithelial cell regeneration description | Group | p-value |
|--|-------|---------|
| Research group   | K1    | 0,0001* |
|  | P1    |         |
|  | K2    |         |
|  | P2    |         |
| Group K1 and P1  | K1    | 0,007*  |
|  | P1    |         |
| Group K1 and K2  | K1    | 0,003*  |
|  | K2    |         |
| Group K1 and P2  | K1    | 0,0001* |
|  | P2    |         |
| Groups P1 and K2   | P1    | 0,424*  |
|  | K2    |         |
| Group P1 and P2  | P1    | 0,001*  |
|  | P2    |         |
| Group K2 and P2  | K2    | 0,004*  |
|  | P2    |         |

### Discussion

Wound healing was indeed a very complex process and it was influenced by a variety of factors, including several cellular and paracellular pathways (Eming et al., 2014; Janis & Harrison, 2014; Reinke & Sorg, 2012; Rousselle et al., 2019). Wound healing has four overlapping phases, namely hemostasis, inflammation, proliferation, and remodeling (Enoch & Leaper, 2005). Any disturbance of the four phases may cause a disruption in wound healing and vice versa (Cullen et al., 2002). The cellular apoptotic mechanisms were responsible for the strict regulation and control of the processes that determine wound healing and other tissue repair processes (Rodrigues et al., 2019).

Ropivacaine, one of the long duration local anesthetics, played a role in enhancing wound healing through two mechanisms, namely anti-inflammatory and analgesic agents. Wounds in the process may cause stress and then inflammation appears, the role of neutrophils, and stimulates anti-apoptosis. Wounds also cause pain which could stimulate immune

release and cortisol thereby stimulating anti-apoptosis. The role of ropivacaine in inhibiting the inflammatory process has a positive impact on wound healing, for example shortening the inflammatory process in the wound healing process (Sgonc & Gruber, 2013). Ropivacaine which acts as an analgesic might maximize wound healing through complex mechanisms, including inhibiting cortisol and the immune response (Kiecolt-Glaser et al., 1995). Previous studies using experimental models and cultured fibroblasts in wounds investigated the effect of local anesthetics, such as bupivacaine, prilocaine, and lidocaine on wound healing (Dere et al., 2009; Drucker et al., 1998; Fedder et al., 2010; Vasseur et al., 1984; Waite et al., 2010). However, the function of ropivacaine in the process of wound healing was not yet completely recognized.

The process and signaling pathways of ropivacaine in wound healing were still not fully understood, including its effects on gene expression, apoptosis, and cell regeneration. Recent studies have shown that 0,2% ropivacaine infiltration, a local anesthetic, has the best efficacy and safety (Bianconi et al., 2003; Labaille et al., 2002; Niyama et al., 2016). In addition, local anesthetics were believed to be an easy, cost-effective, and effective modality for many surgical methods (Gupta, 2010). Hence, this study analyzed the expression of Bcl-2 protein, anti-apoptotic protein, and regeneration of the incision wound epithelium of Wistar rats after infiltration with ropivacaine. The experimental study was chosen since the pharmacokinetic and pharmacodynamic characteristics of ropivacaine are comparable in animals and humans (Arthur et al., 1988; Yi et al., 2009). Incisions on Wistar may also demonstrated the process of primary wound healing.

### *Effect of Ropivacaine Infiltration around the Wound on Bcl-2 Expression in the Healing Process of Incision Wounds in Wistar Rats*

In this study, there were significant differences in Bcl-2 expression between variables. This shows that there was an increase in Bcl-2 expression after administration of 0,2% ropivacaine infiltration. Our data support the





concept that 0,2% ropivacaine can affect the Bcl-2 gene and confer a cell survival advantage. Following this idea, the method of wound healing could be utilized using this information. Besides, the results of this study were supported by previous studies (Özkiriş et al., 2012; Pramono et al., 2016). Initially, Bcl-2 was found in follicular lymphoma associated with a t chromosome translocation (14:18) (Adachi et al., 1990). The translocation causes overexpression of Bcl-2 which leads to inhibition of apoptosis (Hockenbery et al., 1990). On the other hand, overexpression of Bcl-2 protein has been reported in many types of human cancer, including leukemia, lymphoma, and carcinoma (Cimmino et al., 2005). Still, inhibition of apoptosis also has the potential to be involved in the wound healing process. Following injury and the onset of injury, Bcl-2 expression increases, to allow cellular proliferation to occur (Appleton et al., 1996). Expression of apoptosis-related markers such as Bcl-2 provides a novel way to regulate the pattern of apoptosis in wound healing mouse models.

Bcl-2 expression increased with increasing days, for example Bcl-2 expression was higher on day 3 compared to day 7. Even though Bcl-2 was naturally produced during the wound healing process, administration of 0,2% ropivacaine increased Bcl-2 expression. This result was supported by previous study that Bcl-2 expression in the 0,2% ropivacaine group on day 3 was the same as the group without ropivacaine on day 7. Bcl-2 increased on the second day after the appearance of the sores (Azimian et al., 2015). Some studies has shown improved regulation of Bcl-2 expression after wound healing. Studies in rats showed that immediately after injury, Bcl-2 increases to allow for cell proliferation necessary for tissue repair (Kane & Greenhalgh, 2000). Similarly, another study also showed that the low expression of Bcl-2 protein might cause deregulation of the wound healing process (Bhan et al., 2013).

The mechanism between Bcl-2 and wound healing was still not fully understood. Bcl-2 has been described as an anti-apoptotic protein. Pro-apoptotic members of the Bcl-2 family of proteins, such as Bcl-2-associated X protein

(BAX) induce mitochondrial outer membrane permeabilization to cause cytochrome-c release, whereas anti-apoptotic members such as Bcl-2 act as outer membrane protectors and guard against integrity by opposing the BAX function (Furlong et al., 2013). Bcl-2 serves as a protective outer membrane and maintains its integrity (Korsmeyer et al., 2000).

#### *Effect of Ropivacaine Infiltration Around the Wound on the Description of Epithelial Cell Regeneration in the Wound Healing Process of Wistar Rats*

Wound healing studies focus on identifying molecular level target genes that might be enhanced to accelerate the wound healing process. The hedgehog signaling pathway, for example, has been used in various studies because of its role in epithelial-mesenchymal interactions in the wound healing process (Park et al., 2012). Epithelialization or the process of epithelial cell regeneration was a crucial component of wound healing which was used as a determining parameter for successful wound closure (Leoni et al., 2015). Wounds cannot be considered healed without epithelial cell regeneration. If this process does not work, the wound may develop into a chronic wound (Pastar et al., 2014). Several regulators such as growth factors and cytokines, integrins, keratin, MMPs, chemokines, and the extracellular matrix modulate epithelialization during wound closure (Pastar et al., 2014). Epithelial cells were considered to be most important in maintaining the structural integrity and function of the skin's chemical barrier (Martin, 1997). Hence, epithelial cell regeneration was a positive representation of the wound healing process.

Local anesthetics were a class of medications routinely used to alleviate post-operative pain. Long-acting local anesthetics, such as ropivacaine, were widely being used postoperatively by surgical infiltration of the incision area (Stamenkovic et al., 2021). Since the direct interaction between the local anesthetic and the surgical wound, there has been increasing interest in the possible wound healing benefits of ropivacaine. Ropivacaine was considered one of the safest long-acting local anesthetics as of its low toxicity to the



cardiovascular and central nervous systems and was most widely used in infiltration of postoperative incisions (B.-L. Li et al., 2020; K. Li et al., 2020; Wang et al., 2020). Previously, 0,2% ropivacaine increased fibroblast expression (Kurniawan et al., 2020). This was in accordance with the observations in this study, namely 0,2% ropivacaine may contribute to the healing of incision wounds in Wistar rats.

In this study, there were significant differences in the description of epithelial cell regeneration between the variables. This shows that there was an increase in the pattern of epithelial cell regeneration after administration of 0,2% ropivacaine infiltration giving an illustration of epithelial cell regeneration increasing with increasing days. In this study, it was also observed that there was no significant difference in the appearance of epithelial cell regeneration between the P1 and K2 groups. The description of epithelial cell regeneration after infiltration of 0,2% ropivacaine on day 3 was the same as that of epithelial cell regeneration without ropivacaine on day 7.

In this study, there was no significant difference in the appearance of epithelial cell regeneration between the P1 and K2 groups. The description of epithelial cell regeneration after infiltration of 0,2% ropivacaine on day 3 was the same as epithelial cell regeneration without ropivacaine on day 7. Similar research on the role of 0,2% ropivacaine in epithelial regeneration and wound healing was still limited. In addition, similar studies also stated that apoptosis decreased in the ropivacaine infiltration group when compared to the control group and decreased even more on the 7th day (Abdurrahman et al., 2022). On the other hand, previous findings also revealed that on the third day after the incision, Wistar rats that had received infiltration injections of ropivacaine around the wound experienced faster and more complete recovery from superficial wounds (Utariani et al., 2021). This might pave the way for studies on the impact of ropivacaine and other local anesthetics at concentrations of 0,2% on the healing process of wounds. To that end, our research focuses on epithelial regeneration, a critical element of the wound healing process.

The effect of local anesthetics on this process has been investigated in a number of animal studies. Although lidocaine has been reported to enhance wound re-epithelialization in rats, the effect of local anesthetics on wound healing has been a source of debate (Chvapil et al., 1979; Dere et al., 2009; Drucker et al., 1998; Madhuchandra et al., 1991). Previous studies reported that local anesthetic drugs used for infiltration anesthesia showed neutral, negative, and positive effects on wound healing (Dere et al., 2009; Nietgen et al., 1997; Vasseur et al., 1984; Waite et al., 2010). Another studies report that bupivacaine and lidocaine cause no negative or positive changes in wound healing on day 3 (Waite et al., 2010). In addition, several studies have also proven that ropivacaine exhibits anti-proliferative and anti-migratory activity in many cells (Qin et al., 2020; Yin et al., 2020; Zhang et al., 2018). The dose of ropivacaine has an impact on the role that ropivacaine plays (Scherb et al., 2009). Research in Indonesia demonstrated an increase in collagen after infiltration of 0,2% ropivacaine in rats after the 3rd and 7th day (Pramono et al., 2016). This was consistent with the theory that re-epithelialization begins after fibroblasts begin laying down proteoglycans, including new collagen and glycosaminoglycans on day 7 (Wallace et al., 2021).

However, there were a number of issues with this research. First, this research was an experimental study on experimental animals so that clinical application in humans needs to be done. Second, this study did not compare the doses and administration techniques of ropivacaine. The dosage and technique used in this study were safe and effective doses according to previous studies (Pramono et al., 2016). Third, this study did not examine the long-term effects of ropivacaine infiltration. However, according to the literature, the 3rd to 7th day was the optimal time to observe Bcl-2 expression and epithelial cell regeneration.

## Conclusions

In light of the findings of this research, we concluded that there were differences in the expression of Bcl-2 in the wound healing process of the Wistar rat incisions on the 3rd and 7th day after infiltrating ropivacaine. Besides, there were differences in the description of epithelial cell regeneration in the wound healing process of Wistar rat incisions on the 3rd and 7th day after administration of ropivacaine infiltration. In order to reap the benefits of ropivacaine infiltration on the wound healing process, the researchers suggest to consider a retrospective human study on the effect of ropivacaine infiltration in wound healing. Likewise, performing experimental animal experiments with ropivacaine at various doses and procedures as well as exploring the long-term effects of ropivacaine on wound healing.

## References

- Abdurrahman, A., Airlangga, P. S., & Susilo, I. (2022). Effect Ropivacaine Infiltration in Incisional Wound on Caspase-3 Expression and Apoptosis Squamous Cells Wound Healing. *International Journal of Health Sciences*, 6(S4), 1312–1320. <https://doi.org/https://doi.org/10.53730/ijhs.v6nS4.6148>
- Abrão, J., Antunes, M., & Garcia, L. V. (2020). Local Anesthetics Infiltration and Wound Healing Process. *Topics in Local Anesthetics*, 119.
- Adachi, M., Tefferi, A., Greipp, P. R., Kipps, T. J., & Tsujimoto, Y. (1990). Preferential linkage of bcl-2 to immunoglobulin light chain gene in chronic lymphocytic leukemia. *The Journal of Experimental Medicine*, 171(2), 559–564.
- Appleton, I., Brown, N. J., & Willoughby, D. A. (1996). Apoptosis, necrosis, and proliferation: possible implications in the etiology of keloids. *The American Journal of Pathology*, 149(5), 1441.
- Arthur, G. R., Feldman, H. S., & Covino, B. G. (1988). Comparative pharmacokinetics of bupivacaine and ropivacaine, a new amide local anesthetic. *Anesthesia and Analgesia*, 67(11), 1053–1058.
- Azimian, H., Bahreyni-Toossi, M. T., Rezaei, A. R., Rafatpanah, H., Hamzehloei, T., & Fardid, R. (2015). Up-regulation of Bcl-2 expression in cultured human lymphocytes after exposure to low doses of gamma radiation. *Journal of Medical Physics/Association of Medical Physicists of India*, 40(1), 38.
- Bhan, S., Mitra, R., Arya, A. K., Pandey, H. P., & Tripathi, K. (2013). A study on evaluation of apoptosis and expression of bcl-2-related marker in wound healing of streptozotocin-induced diabetic rats. *International Scholarly Research Notices*, 2013.
- Bhaskar, S. B. (2015). Case for local infiltration analgesia: Is all the evidence in black and white? *Indian Journal of Anaesthesia*, 59(1), 1.
- Bianconi, M., Ferraro, L., Traina, G. C., Zanolli, G., Antonelli, T., Guberti, A., Ricci, R., & Massari, L. (2003). Pharmacokinetics and efficacy of ropivacaine continuous wound instillation after joint replacement surgery. *British Journal of Anaesthesia*, 91(6), 830–835.
- Borena, B. M., Martens, A., Broeckx, S. Y., Meyer, E., Chiers, K., Duchateau, L., & Spaas, J. H. (2015). Regenerative skin wound healing in mammals: state-of-the-art on growth factor and stem cell based treatments. *Cellular Physiology and Biochemistry*, 36(1), 1–23.
- Borner, C. (2003). The Bcl-2 protein family: sensors and checkpoints for life-or-death decisions. *Molecular Immunology*, 39(11), 615–647.
- Chen, C.-B., Wang, C.-W., Yang, L.-Y., Ho, H.-C., Hung, S.-I., Yang, C.-H., Chang, C.-J., Su, S.-C., Hui, R. C.-Y., & Chin, S.-W. (2018). Randomized, controlled trial of TNF- $\alpha$  antagonist in CTL-mediated severe cutaneous adverse reactions. *The Journal of Clinical Investigation*, 128(3), 985–996.
- Chen, J.-C., Lin, B.-B., Hu, H.-W., Lin, C., Jin, W.-Y., Zhang, F.-B., Zhu, Y.-A., Lu, C.-J., Wei, X.-J., & Chen, R.-J. (2014). NGF accelerates cutaneous wound healing by promoting the migration of dermal fibroblasts via the PI3K/Akt-Rac1-JNK and ERK pathways. *BioMed Research International*, 2014.
- Chvapil, M., Hameroff, S. R., O’Dea, K., & Peacock Jr, E. E. (1979). Local anesthetics and wound healing. *Journal of Surgical Research*, 27(6), 367–371.
- Cimmino, A., Calin, G. A., Fabbri, M., Iorio, M. V., Ferracin, M., Shimizu, M., Wojcik, S. E., Aqeilan, R. I., Zupo, S., & Dono, M. (2005). miR-15 and miR-16 induce apoptosis by targeting BCL2. *Proceedings of the National Academy of Sciences*, 102(39), 13944–13949.
- Cohen, A., & Mercandetti, M. (2016). *Wound*



- Healing, Healing and Repair*. Emedicine.
- Cullen, B., Smith, R., Mcculloch, E., Silcock, D., & Morrison, L. (2002). Mechanism of action of PROMOGRAN, a protease modulating matrix, for the treatment of diabetic foot ulcers. *Wound Repair and Regeneration*, 10(1), 16–25.
- Dere, K., Sen, H., Teksoz, E., Ozkan, S., Dagli, G., Sucullu, I., Filiz, A. I., Ipcioglu, O. M., & Kucukodaci, Z. (2009). The comparison of the effects of different doses of levobupivacaine infiltration on wound healing. *Journal of Investigative Surgery*, 22(2), 112–116.
- Drucker, M., Cardenas, E., Arizti, P., Valenzuela, A., & Gamboa, A. (1998). Experimental studies on the effect of lidocaine on wound healing. *World Journal of Surgery*, 22(4), 394–398.
- Eming, S. A., Martin, P., & Tomic-Canic, M. (2014). Wound repair and regeneration: mechanisms, signaling, and translation. *Science Translational Medicine*, 6(265), 265sr6-265sr6.
- Enoch, S., & Leaper, D. J. (2005). Basic science of wound healing. *Surgery (Oxford)*, 23(2), 37–42.
- Enoch, S., & Price, P. (2004). Cellular, molecular and biochemical differences in the pathophysiology of healing between acute wounds, chronic wounds and wounds in the aged. *World Wide Wounds*, 13, 1–17.
- Enoch, & Sumartono. (1987). Knowledge of Food Consumption. *Center for Nutrition Research and Development (P3G), Bogor*.
- Fedder, C., Beck-Schimmer, B., Aguirre, J., Hasler, M., Roth-Z'graggen, B., Urner, M., Kalberer, S., Schlicker, A., Votta-Velis, G., & Bonvini, J. M. (2010). In vitro exposure of human fibroblasts to local anaesthetics impairs cell growth. *Clinical & Experimental Immunology*, 162(2), 280–288.
- Furlong, H., Mothersill, C., Lyng, F. M., & Howe, O. (2013). Apoptosis is signalled early by low doses of ionising radiation in a radiation-induced bystander effect. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 741, 35–43.
- Gan, T. J. (2017). Poorly controlled postoperative pain: prevalence, consequences, and prevention. *Journal of Pain Research*, 10, 2287.
- Gupta, A. (2010). Wound infiltration with local anaesthetics in ambulatory surgery. *Current Opinion in Anesthesiology*, 23(6), 708–713.
- Hancı, V., Hakimoğlu, S., Özçmak, H., Bektaş, S., Özçmak, H. S., Özdamar, Ş. O., Yurtlu, S., & Turan, I. Ö. (2012). Comparison of the effects of bupivacaine, lidocaine, and tramadol infiltration on wound healing in rats. *Revista Brasileira de Anestesiologia*, 62, 804–810.
- Hatok, J., & Racay, P. (2016). Bcl-2 family proteins: master regulators of cell survival. *Biomolecular Concepts*, 7(4), 259–270.
- Hockenbery, D., Nuñez, G., Milliman, C., Schreiber, R. D., & Korsmeyer, S. J. (1990). Bcl-2 is an inner mitochondrial membrane protein that blocks programmed cell death. *Nature*, 348(6299), 334–336.
- Hurley, R. W., Murphy, J. D., & Wu, C. L. (2015). Chapter 98, acute postoperative pain. *Miller's Anesthesia. 8th Ed. Philadelphia: Elsevier*, 2974–2998.
- Janis, J. E., & Harrison, B. (2014). Wound Healing: Part I. Basic Science. *Plastic and Reconstructive Surgery*, 138(3 Suppl), 9S-17S.
- Kane, C. D., & Greenhalgh, D. G. (2000). Expression and localization of p53 and bcl-2 in healing wounds in diabetic and nondiabetic mice. *Wound Repair and Regeneration*, 8(1), 45–58.
- Kiecolt-Glaser, J. K., Marucha, P. T., Mercado, A. M., Malarkey, W. B., & Glaser, R. (1995). Slowing of wound healing by psychological stress. *The Lancet*, 346(8984), 1194–1196.
- Korsmeyer, S. J., Wei, M. C., Saito, M. T., Weiler, S., Oh, K. J., & Schlesinger, P. H. (2000). Pro-apoptotic cascade activates BID, which oligomerizes BAK or BAX into pores that result in the release of cytochrome c. *Cell Death & Differentiation*, 7(12), 1166–1173.
- Kurniawan, R., Rahaju, A. S., & Djojodimedjo, T. (2020). Profile of Patients with Urinary Tract Stone at Urology Department of Soetomo General Hospital Surabaya in January 2016-December 2016. *Indonesian Journal of Urology*, 27(1), 22–25.
- Kuthiala, G., & Chaudhary, G. (2011). Ropivacaine: A review of its pharmacology and clinical use. *Indian Journal of Anaesthesia*, 55(2), 104.
- Labaille, T., Mazoit, J. X., Paqueron, X., Franco, D., & Benhamou, D. (2002). The clinical efficacy and pharmacokinetics of intraperitoneal ropivacaine for laparoscopic cholecystectomy. *Anesthesia & Analgesia*,





- 94(1), 100–105.
- Leoni, G., Neumann, P. A., Sumagin, R., Denning, T. L., & Nusrat, A. (2015). Wound repair: role of immune–epithelial interactions. *Mucosal Immunology*, 8(5), 959–968.
- Li, B.-L., Liu, X., Cui, L., Zhang, W., Pang, H., Wang, M., & Wang, H.-Q. (2020). Local infiltration analgesia with ropivacaine improves postoperative pain control in ankle fracture patients: a retrospective cohort study. *Pain Research and Management*, 2020.
- Li, K., Li, H., Luo, D., Feng, H., Ji, C., Yang, K., Liu, J., Zhang, H., & Xu, H. (2020). Efficacy of local infiltration analgesia with ropivacaine for postoperative pain management in cervical laminoplasty: a retrospective study. *Scientific Reports*, 10(1), 1–8.
- Lodish, H., Berk, A., Kaiser, C. A., Krieger, M., Bretscher, A., Ploegh, H., Amon, A., & Martin, K. C. (2016). *Loose-leaf Version for Molecular Cell Biology*. Macmillan Higher Education.
- Maae, E., Nielsen, M., Steffensen, K. D., Jakobsen, E. H., Jakobsen, A., & Sørensen, F. B. (2011). Estimation of immunohistochemical expression of VEGF in ductal carcinomas of the breast. *Journal of Histochemistry & Cytochemistry*, 59(8), 750–760.
- Madhuchandra, S. P., Bhat, M. P., & Ramesh, K. V. (1991). Wound healing profile of topical xylocaine preparations in rodents. *Indian Journal of Experimental Biology*, 29(9), 877–878.
- Maheshwari, K., & Naguib, M. A. (2015). Stoelting's pharmacology & physiology in anesthetic practice. In: Flood, P., J.P. Rathmell and S. Shafer. (Editors). *Stoelting's Pharmacology & Physiology in Anesthetic Practice*, 5th Ed. Wolters Kluwer Health. 282–308.
- Martin, P. (1997). Wound healing--aiming for perfect skin regeneration. *Science*, 276(5309), 75–81.
- Nietgen, G. W., Chan, C. K., & Durieux, M. E. (1997). Inhibition of lysophosphatidate signaling by lidocaine and bupivacaine. *The Journal of the American Society of Anesthesiologists*, 86(5), 1112–1119.
- Niiyama, Y., Yotsuyanagi, T., & Yamakage, M. (2016). Continuous wound infiltration with 0.2% ropivacaine versus a single intercostal nerve block with 0.75% ropivacaine for postoperative pain management after reconstructive surgery for microtia. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 69(10), 1445–1449.
- Özkiriş, M., Kapusuz, Z., & Saydam, L. (2012). Comparison of ropivacaine, bupivacaine and lidocaine in the management of post-tonsillectomy pain. *International Journal of Pediatric Otorhinolaryngology*, 76(12), 1831–1834.
- Park, H.-J., Lee, J., Kim, M.-J., Kang, T. J., Jeong, Y., Um, S. H., & Cho, S.-W. (2012). Sonic hedgehog intradermal gene therapy using a biodegradable poly (β-amino esters) nanoparticle to enhance wound healing. *Biomaterials*, 33(35), 9148–9156.
- Pastar, I., Stojadinovic, O., Yin, N. C., Ramirez, H., Nusbaum, A. G., Sawaya, A., Patel, S. B., Khalid, L., Isseroff, R. R., & Tomic-Canic, M. (2014). Epithelialization in wound healing: a comprehensive review. *Advances in Wound Care*, 3(7), 445–464.
- Pramono, W. B., Leksana, E., & Satoto, H. H. (2016). Pengaruh pemberian ropivacain infiltrasi terhadap tampilan kolagen di sekitar luka insisi pada tikus Wistar. *JAI (Jurnal Anestesiologi Indonesia)*, 8(1), 1–10.
- Qin, A., Liu, Q., & Wang, J. (2020). Ropivacaine inhibits proliferation, invasion, migration and promotes apoptosis of papillary thyroid cancer cells via regulating ITGA2 expression. *Drug Development Research*, 81(6), 700–707.
- Rai, N. K., Tripathi, K., Sharma, D., & Shukla, V. K. (2005). Apoptosis: a basic physiologic process in wound healing. *The International Journal of Lower Extremity Wounds*, 4(3), 138–144.
- Reinke, J. M., & Sorg, H. (2012). Wound repair and regeneration. *European Surgical Research*, 49(1), 35–43.
- Rodrigues, M., Kosaric, N., Bonham, C. A., & Gurtner, G. C. (2019). Wound healing: a cellular perspective. *Physiological Reviews*, 99(1), 665–706.
- Rousselle, P., Braye, F., & Dayan, G. (2019). Re-epithelialization of adult skin wounds: Cellular mechanisms and therapeutic strategies. *Advanced Drug Delivery Reviews*, 146, 344–365. <https://doi.org/doi:10.1016/j.addr.2018.06.019>
- Scherb, M. B., Han, S.-H., Courneya, J.-P., Guyton, G. P., & Schon, L. C. (2009). Effect of bupivacaine on cultured tenocytes. *Orthopedics (Online)*, 32(1), 26.
- Sgonc, R., & Gruber, J. (2013). Age-related



aspects of cutaneous wound healing: a mini-review. *Gerontology*, 59(2), 159–164.

- Stamenkovic, D. M., Bezmarevic, M., Bojic, S., Unic-Stojanovic, D., Stojkovic, D., Slavkovic, D. Z., Bancevic, V., Maric, N., & Karanikolas, M. (2021). Updates on Wound Infiltration Use for Postoperative Pain Management: A Narrative Review. *Journal of Clinical Medicine*, 10(20), 4659.
- Sunarto, O. I., Bajamal, A. H., & Abdurachman. (2019). Outcome of Brain Injury Patients After Decompressive Craniectomy in Tertiary Referral Hospital in East Java Indonesia. *Research Journal of Pharmacy and Technology*, 12(12), 6057–6061. <https://doi.org/10.5958/0974-360X.2019.01051.5>
- Utariani, A., Rahardjo, E., & Perdanakusuma, D. S. (2020). Effects of albumin infusion on serum levels of albumin, proinflammatory cytokines (TNF- $\alpha$ , IL-1, and IL-6), CRP, and MMP-8; Tissue expression of EGRF, ERK1, ERK2, TGF- $\beta$ , collagen, and MMP-8; and wound healing in Sprague Dawley rats. *International Journal of Inflammation*, 2020, 1–13. <https://doi.org/https://doi.org/10.1155/2020/3254017>
- Utariani, A., Sumartono, C., Semedi, B. P., & Susilo, I. (2021). Ekspresi Interleukin 1 dan Tumor Necrosis Factor Alpha pada Pemberian Ropivacain Di Sekitar Luka terhadap Proses Penyembuhan Luka. *Jurnal Syntax Transformation*, 2(2), 157–165.
- Vasseur, P. B., Paul, H. A., Dybdal, N., & Crumley, L. (1984). Effects of local anesthetics on healing of abdominal wounds in rabbits. *American Journal of Veterinary Research*, 45(11), 2385–2388.
- Waite, A., Gilliver, S. C., Masterson, G. R., Hardman, M. J., & Ashcroft, G. S. (2010). Clinically relevant doses of lidocaine and bupivacaine do not impair cutaneous wound healing in mice. *British Journal of Anaesthesia*, 104(6), 768–773.
- Wang, B., Yan, T., Kong, X., Sun, L., Zheng, H., & Zhang, G. (2020). Ropivacaine infiltration analgesia of the drainage exit site enhanced analgesic effects after breast Cancer surgery: a randomized controlled trial. *BMC Anesthesiology*, 20(1), 1–7.
- Yi, J.-W., Lee, B.-J., Kim, D.-O., Park, S.-W., Choi, Y.-K., Chang, H.-K., Kim, C.-J., & Park, J.-H. (2009). Effects of bupivacaine and ropivacaine on field potential in rat hippocampal slices. *British Journal of Anaesthesia*, 102(5), 673–679.
- Yin, D., Liu, L., Shi, Z., Zhang, L., & Yang, Y. (2020). Ropivacaine inhibits cell proliferation, migration and invasion, whereas induces oxidative stress and cell apoptosis by circSCAF11/miR-145-5p axis in glioma. *Cancer Management and Research*, 12, 11145.
- Zhang, Y., Peng, X., & Zheng, Q. (2018). Ropivacaine inhibits the migration of esophageal cancer cells via sodium-channel-independent but prenylation-dependent inhibition of Rac1/JNK/paxillin/FAK. *Biochemical and Biophysical Research Communications*, 501(4), 1074–1079.

