

Effects of Pomegranate Extract Supplementation (*Punica granatum* L.) on Clinical Manifestations of Pulmonary Arterial Hypertension in Children with Acyanotic Congenital Heart Disease

by Budi Utomo

Submission date: 07-Aug-2023 02:46PM (UTC+0800)

Submission ID: 2142530345

File name: Artikel_Tambahan_5.pdf (404.62K)

Word count: 4165

Character count: 23195

Effects of Pomegranate Extract Supplementation (*Punica granatum* L.) on Clinical Manifestations of Pulmonary Arterial Hypertension in Children with Acyanotic Congenital Heart Disease

Audyliya Hartono¹, Mahrus A. Rahman^{1,*}, Budi Utomo²

Audyliya Hartono¹, Mahrus A. Rahman^{1,*}, Budi Utomo²

¹Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, INDONESIA.

²Department of Public Health, Faculty of Medicine, Airlangga University, Surabaya, East Java, Indonesia.

Correspondence

Mahrus A. Rahman

Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya, East Java, INDONESIA.

E-mail: mahrus.a@fk.unair.ac.id

History

- Submission Date: 11-11-2022;
- Review completed: 15-12-2022;
- Accepted Date: 17-12-2022.

DOI : 10.5530/pj.2022.14.210

Article Available online

<http://www.phcogj.com/v14/i6>

Copyright

© 2022 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

ABSTRACT

Background: Pulmonary arterial hypertension (PAH) in congenital heart disease (CHD) mechanism includes vasoconstriction and vascular remodeling. In Indonesia, sildenafil as a vasodilator is the only standard treatment for PAH. Research on pomegranate extract showed its potential as an anti-inflammatory and anti-proliferative agent that could work synergically with sildenafil in reducing pulmonary arterial pressure (PAP). **Objective:** To investigate the effect of pomegranate supplementation in the clinical manifestation of PAH in children with acyanotic CHD. **Methods:** This study was a randomized controlled clinical trial that compared the clinical manifestation of PAH in children with acyanotic CHD who received standard therapy for PAH and pomegranate extract supplementation with the placebo group. Observations were carried out over three months, and pre-and post-test evaluations were conducted to assess changes in PAP, functional classification of PAH, body weight, systolic blood pressure, heart rate, respiratory rate, and oxygen saturation in both groups. Statistical analysis was undertaken using the Wilcoxon, Mann-Whitney, and Spearman correlation tests. **Results:** There was no significant difference in PAP reduction between the pomegranate and placebo groups ($p = .44$) and no significant difference in the functional classification of PAH in the placebo and pomegranate groups ($p = .55$). There was a considerable reduction in the respiratory rate in the pomegranate group ($p = .017$). Respiratory rate was positively correlated with the functional classification of PAH ($p = .011$). **Conclusion:** There was a significant reduction in the respiratory rate in the pomegranate group that correlated with the functional classification of PAH. **Key words:** Heart disease, Pomegranate, Pulmonary hypertension, Sildenafil.

INTRODUCTION

In children with acyanotic CHD, PAH is a pulmonary vascular disease that causes severe morbidity and mortality. It is a problem for clinicians to manage as part of CHD management. Cardiovascular surgery and catheterization to repair the underlying anatomic defect are the definitive treatments for PAH caused by CHD. Due to inadequate resources for cardiac surgery and catheterization, managing PAH in developing countries remains difficult. Thus, pharmacological therapy is essential to reduce the mortality and morbidity of the disease.¹

According to the National Registries of the United Kingdom, the incidence of PAH is 48 percent per 1 million children each year. In cases of CHD, 4-15 percent progress to PAH, and 5-10% progress to Eisenmenger's syndrome.² The Registry to Evaluate Early and Long-Term PAH (REVEAL) estimates a survival rate of 1.3, and the 5-year survival rate in PAH patients is predicted to be 96 +/- 4%, 84 +/- 5%, and 74 +/- 6%. Survival rates were significantly worse in CHD without shunts and in PAH that persisted after surgery.³

PAH etiology is characterized by two mechanisms: vasoconstriction and pulmonary vascular remodeling. In individuals with left-to-right shunt

CHD, increasing flow and pressure to the pulmonary blood arteries decreases vasodilators such as NO-cGMP and prostacyclin (PGI₂) while boosting vasoconstrictors such as endothelin (ET-1), Rho GTPases, and thromboxane. Endothelial damage promotes matrix metalloproteinase breakdown and the release of FGF and TGF- β . These factors cause smooth muscle cell hypertrophy and proliferation. Endothelial injury also activates pro-inflammatory cytokines and macrophages (e.g., IL-6, IL-1 β , MCP-1, and TNF- α), which promote remodeling and the creation of the neointima layer. As a result, effective pharmacological therapy of PAH CHD must include vasodilators as well as anti-remodeling medicines.⁴

Currently, the standard pharmacological therapy for children with PAH CHD in Indonesia is a class of medications known as phosphodiesterase-5 [PDE-5] inhibitors (sildenafil), which acts as a vasodilator in the pulmonary arteries. As a result, anti-remodeling medicines that work in tandem with vasodilators to achieve appropriate pulmonary vascular pressure are still required.⁵

Pomegranate has been demonstrated in studies to be an anti-inflammatory and anti-proliferative agent that can lower oxidative stress, lipid peroxidase, and foam cell formation.⁶ According to Rahman MA's 2016 research, pomegranate extract showed an anti-remodeling impact by decreasing the advancement

Cite this article: Hartono A, Rahman MA, Utomo B. Effects of Pomegranate Extract Supplementation (*Punica granatum* L.) on Clinical Manifestations of Pulmonary Arterial Hypertension in Children with Acyanotic Congenital Heart Disease. Pharmacogn J. 2022;14(6) Suppl: 1061-1066.




Figure 1: Research flowchart.

of fibrosis in the tunica adventitia of the pulmonary arteries in PAH model rats by inhibiting the expression of TGF-1, type I collagen, and an increase in the MMP-1 /TIMP-1 ratio.⁷ Previous research on the use of pomegranate extract in the treatment of PAH CHD was still in the process of proving its efficacy in an animal model. However, additional research is needed to assess the effect of pomegranate extract supplementation as a therapy for PAH CHD in children.

METHODS

This study was a double-blind and randomized controlled trial with a pre-and post-test control group, which compared the clinical manifestation of PAH in children with acyanotic CHD who received sildenafil and supplementation of pomegranate extract. The samples were children aged one month to 10 years old with PAH CHD at Dr. Soetomo General Hospital's Pediatric Cardiology Outpatient Clinic from February 2022 to October 2022. The Health Research Ethics Committee of Dr. Soetomo General Hospital Surabaya approved this research with ethics certificate 0353/KEPK/I/2022.

Selection of participants

The inclusion criteria were children aged one month to 10 years with CHD with left-to-right shunts (ASD, VSD, PDA, or a combination) who received standard therapy for PAH and parents with completed consent forms. The exclusion criteria included those who had undergone heart surgery or catheterization, those with chronic lung parenchymal disease, autoimmune diseases, impaired thyroid function, connective tissue diseases, chromosomal abnormalities, neuromuscular disorders, and those who could not be contacted to come during the observation.

Pomegranate

The pomegranate extract used in this study resulted from the whole pomegranate fruit (*Punica granatum L.*), which contains 40 percent

ellagic acid. The dose was 120 mg/kg/day with an ellagic acid content of 48 mg/kg/day.^{7,11-13}

Intervention and measurement

Patients who fulfilled the requirements criteria completed the informed consent and had measurements of their height, weight, systolic blood pressure, pulse, respiratory rate, peripheral oxygen saturation, and echocardiogram to determine the estimated PAP. Afterward, patients were randomly assigned to the pomegranate or placebo groups. We gave sildenafil 0.5-2mg/kg/dose 3-4 times daily and 120 mg/kg/day of pomegranate extract to the patients in the intervention group. Sildenafil and placebo capsules were given to patients in the control group. After three months of observation, evaluations of body weight, systolic blood pressure, heart rate, respiratory rate, peripheral oxygen saturation, and echocardiographic examination were carried out. The measurements were carried out blindly without knowing whether the study subjects were included in the intervention or control groups.

Statistical analysis

The Kolmogorov-Smirnov test was used to determine the normality of all data. The Wilcoxon statistical test was used to compare changes before and after treatment in each test group. The Mann-Whitney test was used to compare the clinical symptoms of the intervention and control groups. The Spearman correlation test was used to assess the relationship between clinical manifestation of PAH and functional classification of PAH.

RESULTS

Of the forty patients who met the inclusion criteria during the study period, only 30 completed the post-test evaluation at the end of the research. Four patients in the intervention group and three in the control group were excluded because these patients had heart defects closed.

Table 1 shows that there were no significant differences in the age range, gender proportion, nutritional status, mean systolic blood pressure, heart rate, respiratory rate, peripheral oxygen saturation, type of heart defect, and size of heart defect, as well as severity degree of PAH in both test groups before the intervention with a $p > 0.05$.

Table 2 compares echocardiographic features, decreasing PAP, and functional classification of PAH before and after three months of intervention. This study showed no significant difference in pre-test and post-test PAP reduction ($p = 0.44$). There was no significant difference in the functional classification of PAH between the study subjects who received standard therapy and pomegranate compared to those who received standard therapy and placebo ($p = 0.55$).

Table 3 compares changes in clinical manifestations, including body weight, systolic blood pressure, heart rate, respiratory rate, and oxygen saturation, in both test groups. In this table, there was a significant increase in the body weight of the research subjects in the pomegranate group ($p = 0.012$) and the placebo group ($p = 0.005$). This table also shows a significant decrease in the average respiratory rate in the pomegranate group ($p = 0.017$). In contrast, in the placebo group, there was no significant decrease in the average respiratory rate ($p = 0.361$). Respiratory rate was positively correlated with the functional classification of PAH ($p = 0.011$).

Out of fifteen study samples, no side effects occurred in the placebo group during the three months of observation. In the pomegranate

Table 1: The characteristic of the participants.

	Pomegranate n (%)	Placebo n (%)	P
Age			
0-5 months	1 (6.7)	1 (6.7)	0.29
6-11 months	1 (6.7)	2 (13.3)	
1-2 years	2 (13.3)	3 (20)	
<2 - 5 years	4 (26.7)	5 (33.3)	
>5 - 10 years	7 (46.7)	4 (26.7)	
Mean age	4.86	4.12	0.91
Gender			
Male	7 (46.7)	7 (46.7)	1
Female	8 (53.3)	8 (53.3)	
Nutritional status			
Normal	4(26.7)	8(53.5)	0.91
Moderate malnutrition	11(73.3)	2(13.3)	
Severe malnutrition	0(0)	5(33.3)	
Systolic blood pressure (mmHg)	85.4 ± 6.8	87.47 ± 14.0	0.61
Heart rate (x/minutes)	113.4 ± 18.1	104.1 ± 17.1	0.16
Respiratory rate (x/minutes)	31.6 (20-60)	32 (20-60)	0.98
Oxygen saturation (%)	98 (86-99)	98 (90-99)	0.52
Type of congenital heart defect			
ASD	4 (26.7)	5 (33.3)	0.81
VSD	7 (46.7)	6 (40.0)	
PDA	2 (13.3)	2 (13.3)	
ASD + VSD	0 (0.0)	1 (6.7)	
ASD + PDA	2 (13.3)	0 (0)	
VSD + PDA	0 (0)	0 (0)	
ASD + VSD + PDA	0 (0)	1 (6.7)	
Size of the heart defect			
Small	2 (13.3)	1 (6.7)	0.87
Moderate	5 (33.3)	6 (40.0)	
Large	8(53.3)	8(53.3)	
Degree of PAH			
Mild	7(46.7)	1 (6.7)	0.078
Moderate	2(13.3)	5 (33.3)	
Severe	6(40)	9(60)	

Table 2: Comparison of the effect of pomegranate supplementation vs placebo.

	Pomegranate n (%)	Placebo n (%)	p
Degree of PAH			
Normal	3 (20.0)	3 (20)	0.25
Mild	5 (33.3)	1 (6.7)	
Moderate	3 (20.0)	4 (26.7)	
Severe	3(26.7)	7 (53.3)	
Change in degree of PAH			
Improved	5(33.3)	7(46.7)	0.71
Not improved	10(66.7)	8(53.3)	
Reduction of PAP (mmHg)	-12.02± 32	-14 ± 19.46	0.44
Functional classification of PAH			
I	4 (26.7)	6 (40)	0.55
II	9 (60.0)	7 (46.7)	
IIIa	2 (13.3)	2 (13.3)	
IIIb	0 (0)	0 (0)	
Change in functional classification of PAH			
Improved	3(20)	2(13.3)	1
Not improved	12(80)	13(86.7)	

group, out of fifteen study samples, one patient was found with complaints of diarrhea, and two patients complained of vomiting during the first month of the study. In the three patients, dosage adjustments and medication schedules were adjusted to minimize side effects. No severe complication of pomegranate supplementation, including hypotension, was reported in this study.

DISCUSSION

There was no significant difference in pre-test and post-test PAP reduction between the pomegranate and placebo groups. Moreover, there was no significant difference in the functional classification of PAH after three months of pomegranate supplementation.

Pomegranate was proven by Shao *et al.* to produce pulmonary vascular vasodilation by raising levels of acetylcholine acting on endothelial receptors and correcting endothelial dysfunction via the NO-cGMP signaling pathway. This study also showed that punicalagin exhibited antioxidant and anti-inflammatory properties, as seen by lower levels of MMP-9, TNF-a, and VEGF.¹⁴ Previous research revealed that pomegranate could decrease the fibrosis process that occurs in the right ventricle and pulmonary arteries during PAH.⁷ According to Rahman MA's study, pomegranate extract could slow the advancement of fibrosis in the tunica adventitia of the pulmonary arteries in PAH model rats by inhibiting the expression of TGF-1, type I collagen, and increasing the MMP-1/TIMP-1 ratio.^{13,21}

The average age in this study was 4.5 years old. According to Egitto *et al.*, which included 26 children with PAH who underwent open lung biopsy as well as intracardiac defect closure surgery, patients who had thickening of the tunica media consisting of vascular smooth muscle cells, as well as thickening Collagen layer in the tunica adventitia, had persistent PAH after surgical closure of the defect. The thickening of this tunica media and adventitia were indicators of an irreversible state in pulmonary vascular remodeling. Collagen layer thickening in the tunica adventitia was discovered as early as 12 months of life and was most common in patients aged two years. The likelihood of irreversible tunica media and pulmonary vascular adventitia remodeling increases with age.⁹

In this trial, the pomegranate and sildenafil intervention lasted three months. In 2014, Barst *et al.* conducted a research on children aged

Table 3: Comparison of clinical manifestation after pomegranate supplementation vs. placebo.

	Pomegranate		p	Placebo		p
	Pre-test	Post-test		Pre-test	Post-test	
Body weight (kg)	14.6 ± 8.5	15.5 ± 8.3	0.012	11.9 ± 7.3	12.5 ± 7.1	0.005
Systolic blood pressure (mmHg)	85 (71-97)	88 (74-119)	0.308	87 (67-115)	90 (70-120)	0.335
Heart rate (x/minute)	110 (92-150)	110 (84-144)	0.77	110 (76-130)	98 (60-140)	0.353
Respiratory rate (x/minute)	31.6 ± 9.8	28.3 ± 5.8	0.017	32.4 ± 10.8	29.9 ± 7.7	0.361
Oxygen saturation (%)	96.8 ± 3	94.4 ± 8	0.86	97.4 ± 3.3	96.8 ± 5	0.529

Table 4: Correlation analysis of clinical manifestation of PAH and functional classification of PAH.

	p
Body weight	0.055
Systolic blood pressure	0.182
Heart rate	0.504
Respiratory rate	0.011
Oxygen saturation	0.97
Side effect	0.681

1-17 years with PAH, demonstrating that sildenafil successfully reduced PAP after 16 weeks of sildenafil medication. Clinical improvement and exercise capacity, on the other hand, improved within 12 months of treatment.¹⁵ Pomegranate extract was administered to male Sprague-Dawley rats for two weeks in previous studies to establish a vasculoprotective effect.^{7,11,12,16} According to Sengupta's prior research from 2013, one year in human life was nearly similar to two weeks in mouse life (13.8 days).¹⁷ Based on this literature, it could be concluded that in human research subjects, pomegranate extract showed a vasculoprotective effect after administration for one year.

This study showed a significant decrease in respiratory rate after administering pomegranate extract compared to the respiratory rate in the placebo group. From the correlation test analysis, the respiratory rate was positively correlated with the functional classification of pulmonary hypertension, which means that the higher the respiratory rate in the study subjects, the more severe the functional classification of PAH.

The standard therapy for pulmonary arterial hypertension, the phosphodiesterase-5 [PDE-5] inhibitor class (sildenafil), acted as a vasodilator in the pulmonary vessels. Pomegranate extract had the potential to act as an anti-inflammatory and anti-proliferative agent by lowering oxidative stress, lipid peroxidase, and the development of foam cells, all of which functioned as anti-remodeling agents. Pomegranate extract supplementation may improve the efficacy of PAH management. Pomegranate and sildenafil may work together to reduce PAP. In comparison to the group receiving standard therapy and placebo, the group receiving standard therapy plus pomegranate supplementation had a superior functional classification of PAH.^{6,19,22}

This was the first study to investigate at the effect of pomegranate supplementation on pediatric patients with PAH caused by acyanotic CHD. After three months of pomegranate consumption, there were no serious side effects such as hypotension or anaphylaxis. There was no prior evidence on the combination of anti-failure medicines and standard therapy for pulmonary arterial hypertension with pomegranate supplements that could be utilized to provide information about the rules for using these drugs. More research with age grouping, frequency of administration, dosage, and longer duration of pomegranate supplementation is required to provide data on the proper dose and duration of pomegranate administration in children to achieve the desired anti-remodeling effect. More research is also needed to investigate the interaction of pomegranate extract with sildenafil and other anti-failure medicines taken by study participants.

CONCLUSION

A significant reduction in the respiratory rate in the pomegranate group is correlated with the functional classification of PAH in children with PAH caused by acyanotic CHD. There are no severe side effects, such as hypotension or anaphylaxis, after consuming pomegranate for three months.

REGISTRATION

This research has been reviewed and approved by the Thai Clinical Trials Registry (TCTR) with the identification number: TCTR20221006005. The entire clinical trial protocol can be accessed at <http://www.thaiclinicaltrials.org/show/TCTR20221006005>

ACKNOWLEDGMENTS

The authors express their gratitude to Dr. Muhammad Faizi, Head of the Department of Child Health, Dr. Joni Wahyuhadi, Director of Dr. Soetomo General Academic Teaching Hospital, Surabaya, and Professor Doctor Budi Santoso, Dean of the Faculty of Medicine, Universitas Airlangga. We appreciate the help of nurses and our supervisors during the research period. We also want to thank our colleagues and the pediatric residents for supporting our efforts to gather the data.

DISCLOSURE


The author reports no conflicts of interest in this work.

REFERENCES

- Hopper RK, Abman SH, Ivy DD. Persistent Challenges in Pediatric Pulmonary Hypertension. *Chest*. 2016;150(1):226-36.
- Ivy D. Pulmonary Hypertension in Children. *Cardiol Clin*. 2016;34(4):451-72.
- Duffels MGJ, Engelfriet PM, Berger RMF, van Loon RLE, Hoendermis E, Vriend JWJ, et al. Pulmonary arterial hypertension in congenital heart disease: An epidemiologic perspective from a Dutch registry. *Int J Cardiol*. 2007;120(2):198-204.
- Van Der Feen DE, Bartelds B, De Boer RA, Berger RMF. Assessment of reversibility in pulmonary arterial hypertension and congenital heart disease. *Heart*. 2019;105(4):276-82.
- Benedict N, Seybert A, Mathier MA, Benedict N. Evidence-Based Pharmacologic Management of Pulmonary Arterial Hypertension. *Clin Ther*. 2007;29(10):2134-53.
- Wang D, özen C, Abu-Reidah IM, Chigurupati S, Patra JK, Horbanczuk JO, et al. Vasculoprotective effects of pomegranate (*Punica granatum* L.). *Front Pharmacol*. 2018;9:544.
- Hidayat T, Ontoseno T, A. Rahman M, Utamayasa IKA. Peran MMP-1, TIMP-1, rasio MMP-1/TIMP-1 dan kolagen pada hambatan fibrosis otot jantung ventrikel kanan hewan coba model hipertensi arteri pulmonal yang mendapatkan ekstrak buah delima (*Punica granatum* L.). *Univ Airlangga, Surabaya*. 2018;2017.
- Hansmann G, Koestenberger M, Alastalo TP, Apitz C, Austin ED, Bonnet D, et al. 2019 updated consensus statement on the

- diagnosis and treatment of pediatric pulmonary hypertension: The European Pediatric Pulmonary Vascular Disease Network (EPPVDN), endorsed by AEPC, ESPR and ISHLT. *J Hear Lung Transplant.* 2019;38(9):879-901.
9. Egito EST, Aiello VD, Bosisio IBJ, Lichtenfels AJ, Horta ALM, Saldiva PHN, *et al.* Vascular remodelling process in reversibility of pulmonary arterial hypertension secondary to congenital heart disease. *Pathol Res Pract.* 2003;199(8):521-32.
 10. Lasmono SF, Rahman MA, Ontoseno T. Efikasi Dan Safety Beraprost Dalam Menurunkan Tekanan Arteri Pulmonalis Pada Hipertensi Arteri Pulmonalis Terkait Penyakit Jantung Bawaan Pirau Kiri Ke Kanan. Tesis. 2018.
 11. Rais R, Budiono A, Ontoseno T, A. Rahman M, Utamayasa IKA. Pengaruh ekstrak buah delima terhadap ekspresi endotelin A, p38-mitogen activated protein kinase, dan penebalan otot polos arteri pulmonalis pada tikus model hipertensi arteri pulmonal. *Univ Airlangga, Surabaya.* 2003;1-21.
 12. Pramudiawan IB, Ontoseno T, Rahman MA, Utamayasa IKA, Hidayat T, Budiono. Ekspresi Tgf- β 1 Dan Kolagen Tipe I Pada Ventrikel Kanan Tikus Model Hipertensi Arteri Pulmonal Yang Mendapat Ekstrak Buah Delima. 2016;7-43.
 13. Pramudiawan, Bondan I, Hidayat, Taufiq, Utamayasa, Alit IK, *et al.* Ekspresi Tgf- β 1 Dan Kolagen Tipe I Pada Ventrikel Kanan Tikus Model Hipertensi Arteri Pulmonal Yang Mendapat Ekstrak Buah Delima. *Univ Airlangga, Surabaya.* 2016;7-43.
 14. Shao J, Wang P, Liu A, Du X, Bai J, Chen M. Punicalagin Prevents Hypoxic Pulmonary Hypertension via Antioxidant Effects in Rats. *Am J Chinese Med.* 2016;44(4):1-17.
 15. Barst RJ, Ivy DD, Gaitan G, Szatmari A, Rudzinski A, Garcia AE, *et al.* A randomized, double-blind, placebo-controlled, dose-ranging study of oral sildenafil citrate in treatment-naive children with pulmonary arterial hypertension. *Circulation.* 2012;125(2):324-34.
 16. Putri CG, Ontoseno T, A. Rahman M, Ganesha, Citra, Taufiq, *et al.* Pengaruh ekstrak buah delima terhadap ekspresi endothelin-1, inducible nitric oxide synthase dan ketebalan otot polos pada tunika media arteri pulmonalis. *Univ Airlangga, Surabaya.* 2019;1-157.
 17. Sengupta P. The laboratory rat: Relating its age with human's. *Int J Prev Med.* 2013;4(6):624-30.
 18. Viladomiu M, Hontecillas R, Lu P, Bassaganya-Riera J. Preventive and prophylactic mechanisms of action of pomegranate bioactive constituents. *Evidence-based Complement Altern Med.* 2013;2013:789764.
 19. Barst RJ, Beghetti M, Pulido T, Layton G, Konourina I, Zhang M, *et al.* STARTS-2: Long-term survival with oral sildenafil monotherapy in treatment-naive pediatric pulmonary arterial hypertension. *Circulation.* 2014;129(19):1914-23.
 20. Paller CJ, Ye X, Wozniak PJ, Gillespie BK, Sieber PR, Greengold RH, *et al.* A randomized phase II study of pomegranate extract for men with rising PSA following initial therapy for localized prostate cancer. *Prostate Cancer Prostatic Dis.* 2013;16(1):50-5.
 21. Hidayat T, Utamayasa IK, Rahman MA. Effects of Pomegranate Extracts on MMP-1, TIMP-1, MMP-1/TIMP-1 Ratio, and Type I Collagen to Inhibit Right Ventricular Fibrosis in Animal Models. *Carpathian Journal of Food Science and Technology.* 2019; 11: 78-82.
 22. Lasmono SF, Rahman MA, Utamayasa MA. Beraprost as initial pharmacologic treatment for pulmonary hypertension related to left to right shunt congenital heart disease. *Current Pediatric Research.* 2021; 25 :347-349.

GRAPHICAL ABSTRACT



ABOUT AUTHORS



Audyia Hartono is a pediatric resident of Department of Child Health, Faculty of Medicine, Dr. Soetomo General Academic Teaching Hospital, Surabaya. Research interest in pediatric cardiology, pediatric congenital heart disease, and pediatric pulmonary hypertension.



Dr. Mahrus A. Rahman, dr., Sp.A(K) is a lecturer and staff at the Pediatric Cardiology Department, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia. He completed a Bachelor of Medical Doctor at Universitas Airlangga back in 1987. He continued his studies in the Pediatric Department, Faculty of Medicine, Universitas Airlangga, graduating in 1995. After that, he took a fellowship training in Pediatric Cardiology at UN Malaysia in 2001. In 2005, he took a consultant in pediatric cardiology at the Faculty of Medicine, Universitas Indonesia. He earned his doctorate degree in 2016.



Dr. dr. Budi Utomo, MPH is a staff at Department of Public Health - Preventive Medicine, Universitas Airlangga, Surabaya, East Java, Indonesia.

Cite this article: Hartono A, Rahman MA, Utomo B. Effects of Pomegranate Extract Supplementation (*Punica granatum* L.) on Clinical Manifestations of Pulmonary Arterial Hypertension in Children with Acyanotic Congenital Heart Disease. *Pharmacogn J.* 2022;14(6)Suppl: 1061-1066.

Effects of Pomegranate Extract Supplementation (*Punica granatum* L.) on Clinical Manifestations of Pulmonary Arterial Hypertension in Children with Acyanotic Congenital Heart Disease

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

12%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Padjadjaran University Student Paper	1%
2	assets.researchsquare.com Internet Source	1%
3	Submitted to Curtin University of Technology Student Paper	1%
4	journals.plos.org Internet Source	1%
5	dokumen.pub Internet Source	1%
6	www.scielo.br Internet Source	1%
7	sinta3.ristekdikti.go.id Internet Source	1%
8	Amirhossein Sahebkar, Camelia Gurban, Alexandru Serban, Florina Andrica, Maria-	1%

Corina Serban. "Effects of supplementation with pomegranate juice on plasma C-reactive protein concentrations: A systematic review and meta-analysis of randomized controlled trials", Phytomedicine, 2016

Publication

9

Maria Elisabetta Clementi, Giovambattista Pani, Beatrice Sampaolese, Giuseppe Tringali. " Punicalagin reduces H O -induced cytotoxicity and apoptosis in PC12 cells by modulating the levels of reactive oxygen species ", Nutritional Neuroscience, 2017

Publication

1 %

10

ia800700.us.archive.org

Internet Source

1 %

11

Submitted to iGroup

Student Paper

1 %

12

Miftakhur Rohmah, Nurwinda Saputri, Justitia Bahari. "Effectiveness Of Use Of Nesting On Body Weight, Oxygen Saturation Stability, And Breath Frequency In Prematures In Nicu Room Gambiran Hospital Kediri City", STRADA Jurnal Ilmiah Kesehatan, 2020

Publication

1 %

13

ijcbnm.sums.ac.ir

Internet Source

1 %

14

ouci.dntb.gov.ua

Internet Source

1 %

15

Anang Endaryanto, Arlina Dewi, Kusbaryanto, Ricardo Adrian Nugraha. "Trend in the admissions of patients with non-COVID-19 respiratory symptoms during COVID-19 pandemic and its impact on hospital finances in Surabaya, Indonesia", Heliyon, 2023

Publication

<1 %

16

www.alliedacademies.org

Internet Source

<1 %

17

Submitted to Angeles University Foundation

Student Paper

<1 %

18

pediatriccardiology.conferenceseries.com

Internet Source

<1 %

19

koreascience.or.kr

Internet Source

<1 %

20

www.dovepress.com

Internet Source

<1 %

21

www.visual-paradigm.com

Internet Source

<1 %

22

apallergy.org

Internet Source

<1 %

23

f1000researchdata.s3.amazonaws.com

Internet Source

<1 %

24

www.cialis-pills.buy-cheap-medications.net

Internet Source

<1 %

25

www.pjbs.org

Internet Source

<1 %

26

Natalia D. Kalalo, Vivekenanda Pateda, Praevillia Salendu. "Gambaran pertumbuhan pada anak dengan penyakit jantung bawaan di RSUP Prof. Dr. R. D. Kandou Manado", e-CliniC, 2016

Publication

<1 %

27

dergipark.org.tr

Internet Source

<1 %

28

www.science.org

Internet Source

<1 %

29

Sharon M. Moe, Neal X. Chen. " The Role of the Synovium and Cartilage in the Pathogenesis of β - Microglobulin Amyloidosis ", Seminars in Dialysis, 2008

Publication

<1 %

30

Xiaohui, Li, Du Junbao, Shi Lin, Li Jian, Tang Xiuying, Qi Jianguang, Wei Bing, Jin Hongfang, and Tang Chaoshu. "Down-Regulation of Endogenous Hydrogen Sulfide Pathway in Pulmonary Hypertension and Pulmonary Vascular Structural Remodeling Induced by

<1 %

High Pulmonary Blood Flow in Rats", Circulation Journal, 2005.

Publication

31

[covid-19.elsevier.health](https://www.covid-19.elsevier.health)

Internet Source

<1 %

32

[sciencescholar.us](https://www.sciencescholar.us)

Internet Source

<1 %

33

theses.gla.ac.uk

Internet Source

<1 %

34

www.chemie.de

Internet Source

<1 %

35

"APASL Abstracts, Bali 2005", Liver
International, 12/2005

Publication

<1 %

36

[recentscientific.com](https://www.recentscientific.com)

Internet Source

<1 %

37

static.frontiersin.org

Internet Source

<1 %

38

www.medicopublication.com

Internet Source

<1 %

39

www.sciencegate.app

Internet Source

<1 %

40

Jiaxing Xie, Yin Xi, Qingling Zhang, Kefang Lai,
Nanshan Zhong. "Impact of short term forced
oral breathing induced by nasal occlusion on

<1 %

respiratory function in mice", Respiratory
Physiology & Neurobiology, 2015

Publication

41

Omar Tamimi, Mohammed H. A. Mohammed.
"Pulmonary Vascular Resistance
Measurement Remains Keystone in
Congenital Heart Disease Management",
Frontiers in Cardiovascular Medicine, 2021

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Effects of Pomegranate Extract Supplementation (Punica granatum L.) on Clinical Manifestations of Pulmonary Arterial Hypertension in Children with Acyanotic Congenital Heart Disease

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
