

Effects of Golden Sea Cucumber Extract (*Stichopus hermanni*) on Hyphae, Neutrophils and TNF- α in BALB/c Mice Inoculated with *C. albicans* Intravaginally

by Budi Utomo

Submission date: 07-Aug-2023 02:48PM (UTC+0800)

Submission ID: 2142531130

File name: Artikel_Tambahan_7.pdf (830.44K)

Word count: 6617

Character count: 35324

Effects of Golden Sea Cucumber Extract (*Stichopus hermanni*) on Hyphae, Neutrophils and TNF- α in BALB/c Mice Inoculated with *C. albicans* Intravaginally

Varidianto Yudo¹, Widjiati², Harianto Notopuro³, Yulianto Listiawan², Budi Utomo², Purwo Sri Rejeki², Prawesty Diah Utami², Aryati^{2,*}

Varidianto Yudo¹, Widjiati², Harianto Notopuro², Yulianto Listiawan², Budi Utomo², Purwo Sri Rejeki², Prawesty Diah Utami², Aryati^{2,*}

¹Medical Faculty, Hong Tuah University, Surabaya, Indonesia - Affiliated Doctoral Program of Medical Science, Medical Faculty, Airlangga University, Surabaya, INDONESIA.

²Department of Doctoral Program, Medical Faculty, Airlangga University, Surabaya, INDONESIA.

³Medical Faculty, Hong Tuah University, Surabaya, INDONESIA.

Correspondence

Aryati

Department of Doctoral Program, Medical Faculty, Airlangga University, Surabaya, INDONESIA.

E-mail: aryati@k.unair.ac.id

History

- Submission Date: 31-03-2022;
- Review completed: 14-05-2022;
- Accepted Date: 02-06-2022.

DOI : 10.5530/pj.2022.14.97

Article Available online

<http://www.phcogj.com/v14/i3>

Copyright

© 2022 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



ABSTRACT

Introduction: *Candidal vaginitis* is an inflammatory disease that caused mainly by *Candida albicans*. Yeast transitions to filamentous hyphae considered the most important virulence factor. Neutrophils are the first line of defense of the immune system, but in patients with *Candidal vaginitis* the recruitment of neutrophils into the vaginal lumen is positively correlated with symptoms of the disease. This is supported by the release of proinflammatory cytokines such as TNF- α . Standard treatment is considered less effective in relieving symptoms, so other alternative/adjunctive treatments are needed. Golden sea cucumber (*Stichopus hermanni*) extract has been widely studied, especially for anti-fungal and anti-inflammatory. This study aims to analyze the mechanism of decreasing number of hyphae and neutrophils, and pro-inflammatory cytokine TNF- α in BALB/c mice inoculated intravaginally with *C. albicans* after administration of golden sea cucumber extract (*S. hermanni*). **Methods:** Experimental research uses a post-test only control group design. The experimental unit consisted of 36 BALB/c mice that were inoculated intravaginally with *C. albicans* and divided into 4 groups, group that did not receive treatment (K-), group that received standard treatment fluconazole (K+), group that received treatment with golden sea cucumber extract (*S. hermanni*) (P1) and group that received standard treatment with fluconazole plus extract of golden sea cucumber (*S. hermanni*) (P2). The hyphae and neutrophils number were seen microscopically on vaginal mucosal tissue. Cytokine levels of TNF- α were seen from the ELISA blood samples. **Results:** Results showed from the vaginal mucosal tissue of mice, there was significant difference in the number of hyphae ($p = 0.001$) between groups and no significant difference in the number of neutrophils ($p = 0.070$) between groups. From the blood serum of mice, there were significant differences in TNF- α levels ($p = 0.001$) between groups. From the path analysis obtained a significant relationship from the number of hyphae to the number of neutrophils ($p = 0.034$) and the number of neutrophils to TNF- α levels ($p = 0.021$). The strength of the pathway from number of hyphae to number of neutrophils ($\beta = 0.354$) and number of neutrophils to TNF- α levels ($\beta = 0.382$) with positive interactions all. **Conclusion:** In summary, the administration of *S. hermanni* extract was able to reduce the number of hyphae, neutrophils and TNF- α levels through the hyphae, neutrophil and TNF- α pathway.

Key words: *Stichopus hermanni* extract, *Candida vaginitis*, Hyphae, Neutrophils, TNF- α .

INTRODUCTION

Worldwide, recurrent vulvovaginal candidiasis (RVVC) has a global annual prevalence around 4 per 100,000 women and at least 400 million women are affected by recurrent vulvovaginal candidiasis during their lifetime.^{1,3} Abnormal vaginal discharge is the main feature and the first sign of VVC that often causes women to seek health care from a gynecologist.^{4,5} Various risk factors for VVC include douching, antibiotic therapy, elevated estrogen and diabetes. Many factors that increase estrogen levels are risk factors for the development of this disease. The association between increased estrogen and the incidence of *C. albicans* colonization or vaginal disease is due to the effects of estrogen on the host, on fungal cells, or both.^{6,7}

The morphological transition of the budding yeast to the form of filamentous hyphae is considered to represent the most important virulence factor of *C. albicans*.^{8,9} Hyphae are more invasive and contribute to host tissue damage. Pathogen recognition by the host immune system broadly involves pattern recognition receptors (PRR). Vaginal epithelial

cells "detect" the danger signal generated by *C. albicans* and respond by activating immune cells, secreting inflammatory immune mediators and eliciting an immune system response.^{10,11}

Neutrophils are the main innate immune cells that are recruited at the site of infection in response to epithelial cells that secrete immune mediators. Neutrophils further release TNF- α which consequently regulates TLR4 expression in epithelial cells.^{12,13} The recruitment of polymorphonuclear neutrophils (PMNs) into the vagina is strongly associated with symptoms of vaginal inflammation. This vaginal inflammation was observed to decrease with a decrease in PMN.^{2,14,15} Neutrophils do not always play a protective role, as uncontrolled influx of neutrophils in the vagina provokes and amplifies pathogenic inflammation.^{10,16,17} Damage to the vaginal epithelium occurs by a series of mechanisms as above as well as other factors such as hyphae elongation and the simultaneous release of fungal peptide toxin secretion (Candidalysin).^{18,19}

Most women with RVVC (71%) require long-term antifungal treatment as maintenance therapy to

Cite this article: Yudo V, Widjiati, Notopuro H, Listiawan Y, Utomo B, Rejeki PS, et al. Effects of Golden Sea Cucumber Extract (*Stichopus hermanni*) on Hyphae, Neutrophils and TNF- α in BALB/c Mice Inoculated with *C. albicans* Intravaginally. Pharmacogn J. 2022;14(4): 278-285.

control symptoms.⁵ Standard treatment is considered less effective and causes many side effects, for example, topical antifungals cause local hypersensitivity reactions (itching or burning), oral antifungals cause systemic side effects, especially gastrointestinal effects and drug interaction toxicity.²⁰ With standard treatment, especially the azole group, relapses often occur and treatment is only a temporary solution for patients with RVVC.²¹ In addition to specific antimicrobial treatment, anti-inflammatory therapy, both systemic and topical (vaginal douche), is usually added. Most used are non-steroidal anti-inflammatory drugs (NSAIDs). Side effects that often occur are the occurrence of platelet aggregation and digestive tract disorders.²² Problems associated with the management of *Candida* infections require discovery and innovation, where the need for herbal medicines is increasing and natural plant-derived products may offer more amelioration of the existing infections.²³

Sea cucumbers in the form of extracts have been shown to have inhibitory activity against pathogenic fungi.²⁴ The most abundant and most important secondary metabolites of sea cucumbers are triterpene glycosides (saponins).²⁵ Saponin compounds are able to inhibit the formation of biofilms from fungi, increase hyphal permeability and inhibit hyphal growth.^{26,27} Sea cucumbers reduce levels of inflammatory markers interleukin- β , interleukin-6, nitric oxide, and matrix metalloproteinase 9 in cancer cells.²⁴ A study of liver anti-tumor and immunomodulatory effects showed that sea cucumbers decreased serum levels of ALT, AST, GGT and TNF- α and increased serum IL-2.²⁹ Thus, the tested extracts could be explored as new marine sources for novel anti-fungal and anti-inflammatory agents.³⁰ The purpose of this study was to analyze the effects of golden sea cucumber (*S. hermanni*) extract on the number of hyphae and neutrophils, and pro-inflammatory (TNF- α) levels in BALB/c mice inoculated with *C. albicans* intravaginally.

MATERIALS AND METHODS

The design of this study was a true experimental study using a post-test only control group design to prove the relationship between effect on number of hyphae and neutrophils, and pro-inflammatory cytokine (TNF- α) on BALB/c mice inoculated with *C. albicans* intravaginally after administration of golden sea cucumber (*S. hermanni*) extract (ethanolic extract).

The units of examination in this study were 36 female BALB/c mice aged 6-8 weeks that were inoculated with *C. albicans* intravaginally. The experimental unit will be divided into 4 groups as follows: the first group was a group that received standard feed and drink and inoculated with *C. albicans* but did not receive gold sea cucumber extract (K(-) / negative control); the second group was a group that received standard feed and drink and was inoculated with *C. albicans* and received 0.25 mg/kgBW of azole drug (fluconazole) orally (K(+)/ positive control);³¹ the third group was a group that received standard feed and drink and was inoculated with *C. albicans* and received treatment with 17 mg/kgBW orally administered gold sea cucumber extract (P1 / treatment 1);^{32,34} the fourth group was a group that received standard feed and drink and was inoculated with *C. albicans* and received treatment with 17 mg/kgBW orally administered gold sea cucumber extract along with azole class drugs (fluconazole) 0.25 mg/day. kgBB orally (P2 / treatment 2).^{31,34}

The study was started by observing *C. albicans* colonization (hyphae) and neutrophil infiltration in the infected group (on the fourth day). Then the golden sea cucumber (*S. hermanni*) extract was administered to the treatment group (P1 and P2) and azole (fluconazole) to the control group (K+ and P2) for seven days. The parameters studied were examined then, which were the number of hyphae, neutrophils, and TNF- α levels.

Extraction of *Stichopus hermanni*

S. hermanni weighing 100-250 grams was taken from the sea waters of Sumenep in Madura, Indonesia. The sea cucumbers were cleaned, cut into pieces with a size of 3-10 cm, weighed by wet weight after drying on a solar drying rack for samples until they looked dry (3-4 days) to reduce the moisture content. The sea cucumber samples were dried, cut into \pm 1 cm pieces, crushed in a blender. The extraction process was carried out by a maceration process, by immersing 250 grams of dry sample in 500 mL of fine methanol solvent until all samples were submerged and left at room temperature for 24 hours. After filtering with filter paper to separate the filtrate and residue, then soaked again with 500 mL of methanol solvent for 24 hours. After being filtered with filter paper to separate the filtrate and residue, the filtrate will be obtained with a sample ratio of 250 grams / 1000 ml of solvent (1: 4 w/v). The methanol filtrate (polar) was homogenized with hexane solvent (non-polar) and 1,000 mL was performed by partitioning a separating funnel, then each layer of the methanol filtrate solvent and hexane solvent was separated. The methanol filtrate was homogenized again with chloroform solvent (semi-polar) and 1,000 mL was carried out with a separate funnel partition, then each layer of the methanol filtrate solvent and chloroform solvent was separated. Each filtrate was then separated from the solvent using a rotary evaporator to obtain the extract.³²

Administration of *Stichopus hermanni* extract

Treatment groups 1 and 2 (P1 and P2) were given a golden sea cucumber (*S. hermanni*) extract at a dose of 17 mg/kgBW using a feeding tube.^{33,34} Treatment group 2 and positive control (P2 and K+) were given fluconazole 0.25 mg/kgBW using a feeding tube.³¹ On the 7th day after the treatment, the mice were killed.³²

Vaginal inoculation of *C. albicans*

Mice were held to expose the stomach so that 100 ml of sesame oil containing 0.1-0.5 mg estradiol could be injected intraperitoneally (three days before inoculation). The needle is inserted about 5 to 10 mm lateral to the skin to minimize leakage from the injection site. The injections were repeated once a week during the study period. Inoculum was prepared by adding a full circle of *C. albicans* blastoconidia subcultured on Sabouraud-Dextrose Agar (SDA) into 10 ml of Phytone-peptone medium supplemented with 0.1% glucose. The Phytone-peptone medium mixture containing *C. albicans* was incubated for 18 h at 25°C in a vibrating water bath. After incubation, culture medium was collected into 15 ml conical tubes and centrifuged at 800 x G for 5 min. The pellets were washed twice using sterile PBS. Blastoconidia were counted using a hemocytometer. The cell concentration was adjusted to 2.5 x 10⁶/ml (or desired inoculum concentration) in sterile PBS. Mice were stabilized by holding the base of the tail with two fingers and lifting the hips up so that the vagina was facing the examiner. The inoculum suspension was taken using a pipette as much as 20 L (or the desired volume did not exceed 20 L). The inoculum suspension was inserted by inserting the tip of the pipette about 5 mm into the vaginal lumen.³⁶

Mice vaginal fluid sampling for examination of hyphae and neutrophils (PMN)

Mice were anesthetized using Ketamine. The mice were held at the base of the tail with two fingers so that the vaginal opening was opened. The vaginal lumen was rinsed using 100 l of sterile PBS and repeated aspiration was performed with the tip of the pipette. The rinsing fluid was collected into a microcentrifuge tube.³⁶

Hyphae and Neutrophil (PMN) count examination

Wet preparations were made by transferring 10 μ L of vaginal rinse fluid to a glass slide. Cell and nuclear morphology examination (hyphae and neutrophils/PMN) were carried out by staining the cellular fraction of the rinsing fluid. Hyphae and neutrophil count were observed with a light microscope at 400-1000x magnification.²⁶

Serum sampling for examination of TNF- α levels

The mice were anesthetized using Ketamine. To take blood, surgery is done first. Then the needle is inserted directly into the heart and aspirated slowly. The blood obtained was used for the purposes of examining TNF- α levels. The mice were then euthanized.²⁷

Measurement of TNF- α levels using ELISA technique

TNF- α levels were examined quantitatively using ELISA with the following examination principles: reagents, standard solutions, and sample solutions were prepared according to the manufacturer's instructions. All reagents were placed at room temperature before use. This test was carried out at room temperature. The number of strips required is determined for the test. The strip is inserted in the appropriate place. Unused strips should be stored at 2-8 $^{\circ}$ C. 50 μ L standard solution was introduced into standard wells. A 40 μ L sample solution was added to the sample well and then antibody was added according to the desired test (TNF- α), 10 μ L was added to the sample well, 50 μ L Streptavidin-HRP was added to the sample well and standard well. The plate was closed with a sealer and incubated for 60 minutes at 37 $^{\circ}$ C. The sealer was removed and the plate was washed 5 times with wash buffer. Well soaked with 0.35 ml of wash buffer for 30 seconds to 1 minute for each wash. The plate is dried with blotting paper/paper towels or other absorbent material. 50 μ L of substrate solution A was added to each well and then L of substrate solution B was added to each well. The plate was covered with a new sealer for 10 minutes at 37 $^{\circ}$ C in a dark room. Stop Solution 50 μ L was added to each well until the blue color turned yellow. The color intensity was calculated by ELISA Reader at an optical density (OD value) of 450 nm for 30 minutes.²⁸

Data analysis

The selection of statistical tests referred to the type of data from the independent variables. Data analysis was carried out on the number of hyphae, neutrophils, and TNF- α levels in each group by analyzing the mean and standard deviation, then testing for the normality of the distribution in all groups. The homogeneity of variance test between groups was carried out. The Anova comparison test for data with normal distribution and Kruskal Wallis test for data that were not normally distributed.

RESULTS

Mean value and standard deviation of the examination of variables in each group

The results of the mean and standard deviation of the examination of the variables (hyphae count, neutrophils count and TNF- α levels) are presented in table 1.

Research data normality test

The results of the normality distribution of examination data for variables (hyphae count, neutrophils count, and TNF- α levels) are presented in table 2.

The results of the Shapiro-Wilk test showed that the data on the number of hyphae (K- and K+ groups) and the number of neutrophils (K-group) was not normally distributed ($p < \alpha$; $\alpha = 0.05$), so that the differences in hyphae and neutrophil number between groups were analyzed using Kruskal Wallis test. Data on TNF- levels in all groups were normally distributed ($p > \alpha$; $\alpha = 0.05$), so differences between groups were tested using analysis of variance.

Hyphae examination in the vaginal tissue of Balb/c mice

The results of microscopic examination of the number of *C. albicans* hyphae in the vaginal tissue of Balb/c mice are presented in table 3.

The results of the Kruskal Wallis test showed that there was a significant difference in the number of hyphae between groups ($p = 0.001$; $p < \alpha$; $\alpha = 0.05$), so further tests were needed to determine which groups

Table 1: The results of the mean and standard deviation (SD) of the examination of variables for each group.

No.	Variable	Groups							
		K-		K+		P1		P2	
		Means	SD	Means	SD	Means	SD	Means	SD
1.	Hyphae Count	2.78	0.23	1.69	0.63	1.8	0.87	1.18	0.78
2.	Neutrophils Count	2.13	0.52	1.78	0.67	1.38	1.07	1.11	1.07
3.	TNF- α levels	357.11	129.56	575.04	137.9	120.65	142.08	199.66	179.66

Table 2: Test results for the normality distribution of research data.

No.	Variable	p Value			
		K- (n = 9)	K+ (n = 9)	P1 (n = 9)	P2 (n = 9)
1.	Hyphae Count	0.041	0.026	0.220	0.303
2.	Neutrophils Count	0.029	0.069	0.261	0.058
3.	TNF- α levels	0.247	0.290	0.412	0.052

Table 3: Number of *C. albicans* hyphae in the vaginal tissue of Balb/c mice of each group.

No.	Groups	n	Median (min - maks)	p Value
1.	K-	9	2.8 (2.4 - 3) ^a	< 0.001
2.	K+	9	1.8 (1 - 2.4) ^b	
3.	P1	9	2 (0.8 - 3) ^b	
4.	P2	9	1.2 (0.2 - 2.2) ^b	

24

were different. The results of the Mann Whitney test showed that the number of hyphae in the K- group was significantly different from the other three groups, while the number of hyphae in the K+, P1 and P2 groups was not significantly different.

Figure 1. shows a significant decrease in the number of hyphae between the K- group (without treatment) compared to the K+ (fluconazole), P1 (*S. hermanni* extract) and P2 (fluconazole + *S. hermanni* extract) groups. Figure 2. shows images comparison of hyphal elements on the vaginal mucosa in several treatment groups.

Neutrophil examination in the vaginal tissue of Balb/c mice

The results of microscopic examination of the number of neutrophils in the vaginal tissue of Balb/c mice are presented in table 4.

The results of the Kruskal Wallis test showed that there was no significant difference in the number of neutrophils between groups ($p=0.070$; $p>\alpha$; $\alpha=0.05$). Figure 3. shows a decrease in the number of neutrophils in the K+ (fluconazole), P1 (*S. hermanni* extract) and P2 (fluconazole + *S. hermanni* extract) groups compared to the K- group (without treatment). Figure 4. shows images comparison of vaginal mucosal inflammation (neutrophils infiltration) in several treatment groups.

TNF- α ELISA examination in the serum of Balb/c mice

The results of the ELISA examination of TNF- α levels in the blood serum of Balb/c mice can be seen in table 5.




Figure 1: Average number of hyphae in vaginal tissue of Balb/c mice for each group.




Figure 2: Comparison of images of hyphal elements on the vaginal mucosa in several treatment groups. (arrow showing hyphae)




Figure 3: Average number of neutrophils in vaginal tissue of Balb/c mice for each group.




Figure 4: Comparison of vaginal mucosal inflammation (neutrophils infiltration) in several treatment groups. (arrow denotes neutrophils). (Hematoxylin –Eosin : 400x magnification).

Table 4: Number of neutrophils in the vaginal tissue of Balb/c mice of each group.

No.	Groups	n	Median (min – maks)	p Value
1.	K-	9	1.80 (1.6 – 3)	0.07
2.	K+	9	2 (0.6 – 2.4)	
3.	P1	9	1 (0.2 – 3.4)	
4.	P2	9	0.6 (0.2 – 3.4)	

Table 5: TNF- α levels in the blood serum of Balb/c mice of each group.

No.	Groups	n	Median (min – maks)	p Value
1.	K-	9	357.11 \pm 129.564 ^a	< 0.001
2.	K+	9	575.04 \pm 137.900 ^b	
3.	P1	9	120.65 \pm 64.280 ^c	
4.	P2	9	199.66 \pm 179.659 ^c	

Analysis of variance showed that there were significant differences in TNF- α levels between groups ($p<0.001$; $p<\alpha$; $\alpha=0.05$), so further testing was necessary to determine which groups were different. The results of the LSD test showed that the levels of TNF- α in the K- and K+ groups were significantly different from those in the P1 and P2 groups, while the P1 group was not significantly different from the P2 group. Figure 5, shows TNF- α levels in the blood serum of Balb/c mice in all groups. The K+ group (fluconazole) had the highest levels of TNF- α compared to all groups. The P1 group (*S. hermanni* extract) had the lowest levels




Figure 5: Average levels of TNF- α in blood serum of Balb/c mice of each group.




Figure 6: Regression analysis model to see interactions between number of hyphae and neutrophils, and TNF- α levels.

of TNF- α compared to all groups. The P2 group (fluconazole + *S. hermanni* extract) had TNF- α levels between the K+ and P1 groups, but lower than the K- group (without treatment).

Causalistic relationship between the variables studied

Regression analysis was conducted to determine the causal relationship between research variables and the causes of changes in research variables. The pattern of causal relationships of variables can be described in a hypothetical path analysis model as shown in Figure 6.

The regression analysis show significant relationship results were obtained from the path of hyphae number to neutrophils number ($p = 0.034$) and neutrophils number to TNF- α levels ($p = 0.021$). From the path analysis, the path coefficients power is moderate (0.354 - 0.382). The strength of the pathway of hyphae number to neutrophils number ($\beta = 0.354$) and the number of neutrophils to TNF- α levels ($\beta = 0.382$) with all positive interactions.

DISCUSSION

Activity of *Stichopus hermanni* extract against *Candida albicans* hyphae

The pathogenicity of *C. albicans* in causing *Candidal vaginitis* is caused by a change in the shape of yeast cells into hyphae. The formation of these hyphae will trigger an inflammatory reaction that causes symptoms. So that efforts to inhibit the formation of candida hyphae will relieve symptoms.^{4,4-10} From Figure 1, the administration of *S. hermanni* extract reduced the number of hyphae significantly ($p=0.011$; $p<\alpha$; $\alpha=0.05$) compared to the K- group (without treatment). Administration of *S. hermanni* extract reduced the number of *C. albicans* hyphae almost as much as fluconazole alone, where statistically the difference between administration of *S. hermanni* extract and administration of fluconazole was not significant ($p=0.894$; $p>\alpha$; $\alpha=0.05$). When given concurrently with fluconazole, it was very effective in reducing hyphae of *C. albicans*, which is more hyphae reduction than the administration of *S. hermanni* extract and fluconazole alone, although the difference was not statistically significant.

Sea cucumbers are rich in glycosides, especially triterpene glycosides (saponins) which have been shown to have antifungal activity. *S.*

hermanni also produces triterpene glycosides (saponins) both holostane (Stichlorosides, Stichoposides and Holotoxins) and non-holostane.^{25,30} The biological activity of holothurian saponins occurs through their membranolytic function after a certain threshold concentration is reached. Triterpene glycosides cause membrane disruption; alter membrane permeability, loss of barrier function, and rupture of cell membranes. The interaction of glycosides (aglycone moiety) with membrane D5(6)-sterols (ergosterol in fungi) creates a glycoside-sterol complex in the membrane, modifying microviscosity, ion permeability and activity of membrane proteins. The strong membranolytic function of D5-sterol-containing biological membranes due to the formation of single ion channels and larger pores is the basis of the antifungal features of this substance. The glycoside activity of sea cucumbers in sub-cytotoxic doses disrupts specific membrane transport proteins in fungal cells and alters their activity.²⁵

Activity of *Stichopus hermanni* extract on neutrophil count

Neutrophils are the most abundant circulating leukocyte, and have been considered the first line of defense of the immune system. Neutrophils also act as inflammatory mediators.³⁹ However, in patients with *Candidal vaginitis*, as well as in some studies using mice, recruitment of neutrophils into the vaginal lumen is positively correlated with disease symptoms. Many studies have targeted neutrophil removal as a treatment for *Candidal vaginitis*.⁷ From Figure 3 and Table 4, the administration of *S. hermanni* extract decreased the number of neutrophils more than the K- group (without treatment) and the K+ group (fluconazole) but slightly higher than the P2 group (fluconazole + *S. hermanni* extract), although this difference in decrease was not significant in statistics ($p=0.070$; $p>\alpha$; $\alpha=0.05$). The administration of *S. hermanni* extract decreased the number of neutrophils more than the administration of fluconazole alone. And when combined with fluconazole further increase the effectiveness of reducing the number of neutrophils.

This decrease in the number of neutrophils may because of stichopus group contains fucosylated chondroitin sulfate (FucCS) substances. This polysaccharide isolated from sea cucumbers is a strong inhibitor of P- and L-selectin. FucCS reduces neutrophil recruitment to inflamed tissues.^{40,41} Activation, migration, and aggregation of neutrophils can also be inhibited by PGE2 where PGE2 can be produced by host cells or from *Candida* cells themselves.^{42,43}

Activity of *Stichopus hermanni* extract on TNF- α levels

Tumor necrosis factor α (TNF- α) exhibits a large number of functions in host defense mechanisms, whereas excessive release of TNF- α in inflammation promotes tissue damage. Cytokines released in the inflammatory environment can influence the development of microorganisms either by promoting their growth or exhibiting antimicrobial activity. Excessive release of TNF- α during inflammation is associated with the development of pain, cell infiltration, and tissue structural damage.⁴⁴

From figure 5 and table 5, the administration of *S. hermanni* extract decreased TNF- α levels compared to the K- group (without treatment), this decrease was statistically significant ($p=0.001$; $p<\alpha$; $\alpha=0.05$). The administration of *S. hermanni* extract also significantly reduced TNF- α levels compared to the administration of fluconazole alone ($p=0.001$; $p<\alpha$; $\alpha=0.05$). Likewise, the combined administration of *S. hermanni* extract and fluconazole also reduced TNF- α levels, where the decrease in TNF- α levels was not significantly different from the administration of *S. hermanni* extract alone ($p=0.221$; $p>\alpha$; $\alpha=0.05$). Some of the things that can reduce TNF- α levels from some types of sea cucumbers may be cerebroside content,⁴⁵ and all things that reduce neutrophils including fucosylated chondroitin sulfate and PGE2.⁴⁰⁻⁴³

This study analyzed the effect of *S. hermanni* extract on the pathway of *C. albicans* hyphae, neutrophil recruitment and production of cytokine TNF- α . This study did not see the relationship with other immune cells, especially dendritic, macrophage and lymphocyte cells.

CONCLUSION

In summary, the administration of *S. hermanni* extract was able to reduce the number of *C. albicans* hyphae. The combination of fluconazole – *S. hermanni* extract decreased the most of *C. albicans* hyphae. The administration of *S. hermanni* extract was able to reduce the number of neutrophils. The combination of fluconazole – *S. hermanni* extract decreased the most neutrophil counts. The administration of *S. hermanni* extract was able to reduce TNF- α levels. The decrease in TNF- α due to the administration of *S. hermanni* extract alone was the largest.

ACKNOWLEDGEMENTS


We thank EJA for editing the manuscript.

REFERENCES

- Denning DW, Kneale M, Sobel JD, Richardson RR. Global burden of recurrent vulvovaginal candidiasis: A systematic review. *Lancet Infect Dis*. 2018;18(11):e339-47.
- Swidergall M. *Candida albicans* at Host Barrier Sites: Pattern Recognition Receptors and Beyond. *Pathogens*. 2019;8(1):40.
- Rosati D, Bruno M, Jaeger M, Oever JT, Netea MG. Recurrent Vulvovaginal Candidiasis: An Immunological Perspective. *Microorganisms*. 2020;8(2):144.
- Kalia N, Singh J, Kaur M. Immunopathology of Recurrent Vulvovaginal Infections: New Aspects and Research Directions. *Front Immunol*. 2019;10:2034.
- Yano J, Sobel JD, Nyirjesy P, Sobel R, Williams VL, Yu Q, et al. Current patient perspectives of vulvovaginal candidiasis: incidence, symptoms, management and post-treatment outcomes. *BMC Women's Health*. 2019;19(1):48.
- Cessone A. Vulvovaginal *Candida albicans* infections: pathogenesis, immunity and vaccine prospects. *BJOG*. 2014;122(6):785-94.
- Willems HME, Ahmed SS, Liu J, Xu Z, Peters BM. Vulvovaginal Candidiasis: A Current Understanding and Burning Questions. *J Fungi*. 2020;6(1):27.
- Mayer FL, Wilson D, Hube B. *Candida albicans* pathogenicity mechanisms. *Virulence*. 2013;4(2):119-28.
- Yang W, Yan L, Wu C, Zhao X, Tang J. Fungal invasion of epithelial cells. *Microbiol Res*. 2014;169(11):803-10.
- Hofs S, Mogavero S, Hube B. Interaction of *Candida albicans* with host cells: virulence factors, host defense, escape strategies, and the microbiota. *J Microbiol*. 2016;54(3):149-69.
- Mengesha BG, Conti HR. The Role of IL-17 in Protection against Mucosal *Candida* Infections. *J fungi*. 2017;3(4):52.
- Husen SA, Syadzha MF, Setyawan MF, Pudjiastuti P, Ansori ANM, Susilo RJK, et al. Evaluation of the Combination of *Sargassum Duplicatum*, *Sargassum Illicifolium*, *Abelmoschus Esculentus*, and *Garcinia Mangostana* Extracts for Open Wound Healing in Diabetic Mice. *Sys Rev Pharm*. 2020;11(9):888-92.
- Husen SA, Setyawan MF, Syadzha MF, Susilo RJ, Hayaza S, Ansori AN, et al. A Novel Therapeutic effect of *Sargassum illicifolium* Algininate and Okra (*Abelmoschus esculentus*) Pods extracts on Open wound healing process in Diabetic Mice. *Res J Pharm Technol*. 2020;13(6):2764-70.
- Peters BM, Palmer GE, Nash AK, Lilly EA, Fidel PL Jr, Noverr MC. Fungal morphogenetic pathways are required for the hallmark inflammatory response during *Candida albicans* vaginitis. *Infect Immun*. 2014;82(2):532-43.
- Kumar KP, Nicholls AJ, Wong C. Partners in crime: neutrophils and monocytes/macrophages in inflammation and disease. *Cell Tissue Res*. 2018;371(3):551-65.
- Yano J, Noverr MC, Fidel PL Jr. Cytokines in the host response to *Candida* vaginitis: Identifying a role for non-classical immune mediators, S100 alarmins. *Cytokine*. 2012;58(1):118-28.
- Yano J, Noverr MC, Fidel PL Jr. Vaginal Heparan Sulfate Linked to Neutrophil Dysfunction in the Acute Inflammatory Response Associated with Experimental Vulvovaginal Candidiasis. *Bio*. 2017;8(2):e00211-17.
- Moyes DL, Wilson D, Richardson JP, Mogavero S, Tang SX, Wernecke J, et al. Candidalysin is a fungal peptide toxin critical for mucosal infection. *Nature*. 2016;532(7597):64-8.
- Wilson D, Naglik JR, Hube B. The Missing Link between *Candida albicans* Hyphal Morphogenesis and Host Cell Damage. *PLoS Pathog*. 2016;12(10):e01005867.
- Paladine HL, Desai UA. Vaginitis: diagnosis and treatment. *Am Fam Physician*. 2018;97(5):321-9.
- Crouss T, Sobel JD, Smith K, Nyirjesy P. Long-Term Outcomes of Women with Recurrent Vulvovaginal Candidiasis After a Course of Maintenance Antifungal Therapy. *J Low Genit Tract Dis*. 2018;22(4):382-6.
- Milani M, Iacobelli P. Vaginal use of Ibuprofen isobutanolammonium (Iginerom): efficacy, tolerability, and pharmacokinetic data: a review of available data. *ISRN Obstet Gynecol*. 2012;2012:673131.
- Ahangari F, Farshbaf-Khalili A, Javadzadeh Y, Adibpour M, Oskouei BS. Comparing the effectiveness of *Salvia officinalis*, clotrimazole and their combination on vulvovaginal candidiasis: A randomized, controlled clinical trial. *J Obstet Gynaecol Res*. 2019;45(4):897-907.
- Adibpour N, Nasr F, Nematpour F, Shakouri A, Ameri A. Antibacterial and antifungal activity of *Holothuria leucospilota* isolated from Persian gulf and Orman sea. *Jundishapur J Microbiol*. 2014;7(1):e8708.
- Bahrami Y, Franco CM. Acetylated Triterpene Glycosides and Their Biological Activity from *Holothuroidea* Reported in the Past Six Decades. *Mar Drugs*. 2016;14(8):147.
- Coleman JJ, Okoli I, Tegos GP, Holson EB, Wagner FF, Hamblin MR, et al. Characterization of plant-derived saponin natural products against *Candida albicans*. *ACS Chem Biol*. 2010;5(3):321-32.
- Yano A, Abe A, Aizawa F, Yamada H, Minami K, Matsui M, et al. The effect of eating sea cucumber jelly on *Candida* load in the oral cavity of elderly individuals in a nursing home. *Mar Drugs*. 2013;11(12):4993-5007.
- Kareh M, El Nahas R, Al-Aaraj L, Al-Ghadban S, Naser Al Deen N, Saliba N, et al. Anti-proliferative and anti-inflammatory activities of the sea cucumber *Holothuria polii* aqueous extract. *SAGE Open Med*. 2018;6:2050312118809541.
- Song Y, Jin SJ, Cui LH, Ji XJ, Yang FG. Immunomodulatory effect of *Stichopus japonicus* acid mucopolysaccharide on experimental hepatocellular carcinoma in rats. *Molecules*. 2013;18(6):7179-93.
- Widyananda MH, Pratama SK, Samoedra RS, Sari FN, Dhea V, Kharisma AN, et al. Molecular docking study of sea urchin (*Arbacia lixula*) peptides as multi-target inhibitor for non-small cell lung cancer (NSCLC) associated proteins. *J Pharm Pharmacogn Res*. 2021;9(4):484-96.
- Graybill JR, Najjar LK, Holmberg JD, Correa A, Luther MF. Fluconazole treatment of *Candida albicans* infection in mice: does in vitro susceptibility predict in vivo response? *Antimicrob Agents Chemother*. 1995;39(10):2197-200.
- Revianti S, Parisihni K. Potency of *Stichopus hermanni* extract

- as oral candidiasis treatment on epithelial rat tongue. Dent J. 2016;49(1):10-6
33. Parishni K. Mekanisme molekuler ekstrak etanol teripang (*Stichopus hermanni*) sebagai imunomodulator pada periodontitis kronis yang diinduksi *Porphyromonas gingivalis*. Disertasi, Universitas Airlangga. 2015.
 34. Yudo V, Utami PD, Setianingsih H. Studi Eksperimental: Penurunan Profil Netrofil Dan TNF- α Setelah Pemberian Ekstrak Teripang Emas (*Stichopus hermanni*). Surabaya Biomed J. 2022;1(2):101-9.
 35. Olu S, Pandarangga P, Ndacong N. Persembuhan luka insisi kulit mencit (*Mus musculus*) dengan pemberian ekstrak etanol teripang getah (*Holothuria leucospilota*). J Veteriner Nusantara. 2019;2(1):60-9.
 36. Yano J, Fidel PL Jr. Protocols for vaginal inoculation and sample collection in the experimental mouse model of *Candida* vaginitis. J Vis Exp. 2011;58:3382.
 37. Nugroho RA. Mengenal Mencit Sebagai Hewan Laboratorium. Mulawarman University Press. Samarinda. 2018;76.
 38. Setianingsih H. Mekanisme Perbaikan Disfungsi Endotel Pasca Pemberian Oksigen Hiperbarik Pada Hewan Coba Terpapar Diet Aterogenik. Disertasi, Universitas Airlangga. 2018.
 39. Kharisma VD, Agatha A, Ansori AN, Widyananda MH, Rizky WC, Dings TG, et al. Herbal combination from *Moringa oleifera* Lam. and *Curcuma longa* L. as SARS-CoV-2 antiviral via dual inhibitor pathway: A viroinformatics approach. J Pharm Pharmacogn Res. 2022;10(1):138-46.
 40. Borsig L, Wang L, Cavalcante MCM, Cardio-Reis L, Ferreira PL, Mourao PAS, et al. Selectin Blocking Activity of a Fucosylated Chondroitin Sulfate Glycosaminoglycan from Sea Cucumber: Effect on Tumor Metastasis And Neutrophil Recruitment. J Biol Chem. 2007;282(20):14984-91.
 41. Myron P, Siddiquee S, Al Azad S. Fucosylated chondroitin sulfate diversity in sea cucumbers: a review. Carbohydr Polym. 2014;112:173-8.
 42. Agard M, Asakrah S, Morici LA. PGE(2) suppression of innate immunity during mucosal bacterial infection. Front Cell Infect Microbiol. 2013;3:45.
 43. Lucas RC, Taglialegna R, França CN, Segato F, Cazzaniga RA, Da Fonseca CK, et al. Immune Response, Detection of IgE and PGE2 during Vaginal Candidiasis in Mice. Am J Immunol. 2016;12(2):29-36.
 44. Padmi H, Ansori AN, Probojati RT, Murtadlo AA, Sunarwidhi AL, Hernawen A, et al. Anti-inflammatory potential of λ -carrageenan by inhibition of IL-6 receptor: in silico study. IOP Confr Ser: Earth Environ Sci. 2021;913:012106.
 45. Zhang L, Zhang T, Ding L, Xu J, Xue C, Yanagita T, et al. The Protective Activities of Dietary Sea Cucumber Cerebrosides against Atherosclerosis through Regulating Inflammation and Cholesterol Metabolism in Male Mice. Mol Nutr Food Res. 2018;62(16):e1800315.

GRAPHICAL ABSTRACT



ABOUT AUTHORS



Varidianto Yudo: He is a doctoral student from Faculty of Medicine, Universitas Airlangga. He is also a lecturer at Universitas Hang Tuah. Recently, his research topic about the effect of *Stichopus hermanni* extract on candidal vaginitis.



Widjiati: She is Professor at the Faculty of Veterinary Medicine, Universitas Airlangga. The research topic is currently being carried out is related to the reproduction of livestock.



Harianto Notopuro: He is Professor at Faculty of medicine, Universitas Airlangga. His main topic research is related to biochemistry and biomolecular in human.



Yulianto Listiawan: He is a Dermatologist and Venereologist who graduated from Universitas Airlangga in 1998. He later obtained the title of Consultant Skin and Venereal Health from the Collegium of Dermatology and Venereology in 2007. In 2011, he succeeded in completing the Doctoral Program in Medical Sciences, which he also did at Airlangga University. He works in Expression of TNF- α in skin tissue.



Budi Utomo: He is senior lecturer in Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga. Experience with data management, epidemiology research and biostatistics. Strong background in research project management and data management.



Purwo Sri Rejeki: She is staff member of Department of Physiology, Faculty of Medicine, Universitas Airlangga. She is a lecturer of physiology department faculty of medicine Universitas Airlangga. She is also interest in metabolism and obesity field research. Recently, she conducts some research for preventive obesity about. She is also a general practitioner with esthetic medicine interested



Prawesty Diah Utami: She is a lecturer at Universitas Hang Tuah, Parasitology laboratory. Recently, her research topic about Hyperbaric Oxygen Exposure Reduces ICAM-1 And HIF-1 α Expression in Brain Endothelial Cells from Experimental Cerebral Malaria Mice.



Aryati: She is Professor of Clinical Pathology at Faculty of Medicine, Universitas Airlangga. Her expertise is in Immunology and molecular biology.

Cite this article: Yudo V, Widjiati, Notopuro H, Listiawan Y, Utomo B, Rejeki PS, et al. Effects of Golden Sea Cucumber Extract (*Stichopus hermanni*) on Hyphae, Neutrophils and TNF- α in BALB/c Mice Inoculated with *C. albicans* Intravaginally. *Pharmacogn J.* 2022;14(4): 278-285.

Effects of Golden Sea Cucumber Extract (*Stichopus hermanni*) on Hyphae, Neutrophils and TNF- α in BALB/c Mice Inoculated with *C. albicans* Intravaginally

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

15%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to University of Alabama, Huntsville Student Paper	<1 %
2	coek.info Internet Source	<1 %
3	d-scholarship.pitt.edu Internet Source	<1 %
4	jjmicrobiol.com Internet Source	<1 %
5	repository.up.ac.za Internet Source	<1 %
6	rjptonline.org Internet Source	<1 %
7	"Sunday, 3 September 2006", European Heart Journal, 08/02/2006 Publication	<1 %

8

Evandro Ribeiro, Sibelle Faleiro, Cerise Campos, Maria Breseguelo et al. "Oral Candida albicans in Patients in the ICU of a Brazilian Hospital School and in vitro Susceptibility of Isolated Yeasts to Extra-virgin Coconut Oil", International Journal of TROPICAL DISEASE & Health, 2017

Publication

<1 %

9

armaghanj.yums.ac.ir

Internet Source

<1 %

10

eprints.whiterose.ac.uk

Internet Source

<1 %

11

Eva Pericolini, Stefano Perito, Anna Castagnoli, Elena Gabrielli et al. "Epitope unmasking in vulvovaginal candidiasis is associated with hyphal growth and neutrophilic infiltration", PLOS ONE, 2018

Publication

<1 %

12

Meng-Yuan Bao, Ming Li, Qing-Ru Bu, Yue Yang, Hang Song, Chang-Zhong Wang, Tian-Ming Wang, Ning Li. "The effect of herbal medicine in innate immunity to Candida albicans", Frontiers in Immunology, 2023

Publication

<1 %

13

Submitted to Universitas Andalas

Student Paper

<1 %

14	Internet Source	<1 %
15	bmcmicrobiol.biomedcentral.com Internet Source	<1 %
16	www.recentmedicalfindings.com Internet Source	<1 %
17	Juliana Lírio, Paulo Cesar Giraldo, Rose Luce Amaral, Ayane Cristine Alves Sarmiento et al. "Antifungal (oral and vaginal) therapy for recurrent vulvovaginal candidiasis: a systematic review protocol", BMJ Open, 2019 Publication	<1 %
18	benthamscience.com Internet Source	<1 %
19	www.escavador.com Internet Source	<1 %
20	Submitted to University College London Student Paper	<1 %
21	Xiaofei Shao, Xiaobo Wang, Kaidi Zhu, Yongjun Dang, Biao Yu. "Synthesis of Sea Cucumber Saponins with Antitumor Activities", The Journal of Organic Chemistry, 2020 Publication	<1 %
22	Submitted to UNITEC Institute of Technology Student Paper	<1 %

23

ejurnal.undana.ac.id

Internet Source

<1 %

24

Inggita Kusumastuty, Dian Handayani, Shafira Hanifa, Melinda Lisan, Etik Sulistyowati. "The Effect of Brown Rice on Superoxide Dismutase Level and Non-alcoholic Fatty Liver in an Sprague–Dawley Rat Model of High-fat High-fructose Diet-induced Obesity", Open Access Macedonian Journal of Medical Sciences, 2020

Publication

<1 %

25

Waqas Nasir, Alejandro Gomez Toledo, Fredrik Noborn, Jonas Nilsson, Mingxun Wang, Nuno Bandeira, Göran Larson. " : A Bioinformatics Workflow for Glycopeptide MS/MS Spectral Analysis ", Journal of Proteome Research, 2016

Publication

<1 %

26

elea.unisa.it

Internet Source

<1 %

27

pdfs.semanticscholar.org

Internet Source

<1 %

28

bmrat.org

Internet Source

<1 %

29

text-id.123dok.com

Internet Source

<1 %

30

unsworks.unsw.edu.au

Internet Source

<1 %

31

f1000research.com

Internet Source

<1 %

32

www.e-repository.unsyiah.ac.id

Internet Source

<1 %

33

www.sciencegate.app

Internet Source

<1 %

34

www.scilit.net

Internet Source

<1 %

35

"Abstracts of the 37th Annual Scientific Meeting of the Indonesian Urological Association : Abstracts", BJU International, 2014.

Publication

<1 %

36

aeronline.org

Internet Source

<1 %

37

bmcinfectdis.biomedcentral.com

Internet Source

<1 %

38

jnanobiotechnology.biomedcentral.com

Internet Source

<1 %

39

repository.ub.ac.id

Internet Source

<1 %

40

Submitted to University of Leicester

Student Paper

<1 %

41	cyberleninka.org Internet Source	<1 %
42	doaj.org Internet Source	<1 %
43	news.unair.ac.id Internet Source	<1 %
44	Bambang Purwanto, Sundari Indah Wiyasihati, Putri Ayu Masyitha, Kristanti Wanito Wigati, Irfiansyah Irwadi. "Golden sea cucumber extract revives glucose transporter-4 and interleukin-6 protein level in diabetic mouse muscle", Veterinary World, 2019 Publication	<1 %
45	eprints.umm.ac.id Internet Source	<1 %
46	Bemnet Mengesha, Heather Conti. "The Role of IL-17 in Protection against Mucosal Candida Infections", Journal of Fungi, 2017 Publication	<1 %
47	Submitted to University of South Alabama Student Paper	<1 %
48	Submitted to Universitas Brawijaya Student Paper	<1 %
49	ajp.mums.ac.ir Internet Source	<1 %

50 journal.uwks.ac.id <1 %
Internet Source

51 www.cell.com <1 %
Internet Source

52 www.veterinaryworld.org <1 %
Internet Source

53 Bruno R.A. Pelozin, Ursula Paula Reno Soci, João L. P. Gomes, Edilamar Menezes Oliveira, Tiago Fernandes. "mTOR signaling-related microRNAs as cardiac hypertrophy modulators in high - volume endurance training", Journal of Applied Physiology, 2021
Publication

54 journal.ipb.ac.id <1 %
Internet Source

55 Adewale Oluwasogo Olalemi, Abiola Stephen Okunade. "Effect of some physicochemical parameters on the decay rate of enteric bacteria isolated from a surface water", Research Square Platform LLC, 2023
Publication

56 jppres.com <1 %
Internet Source

57 new.kjo.or.kr <1 %
Internet Source

tndt.org

58

Internet Source

<1 %

59

"Mechanisms Underlying Host-Microbiome Interactions in Pathophysiology of Human Diseases", Springer Science and Business Media LLC, 2018

Publication

<1 %

60

Elizabeth A. Lilly, Junko Yano, Shannon K. Esher, Emily Hardie, Paul L. Fidel, Mairi C. Noverr. " Spectrum of Trained Innate Immunity Induced by Low-Virulence Species against Lethal Polymicrobial Intra-abdominal Infection ", Infection and Immunity, 2019

Publication

<1 %

61

L. Cárdenas-Freytag. "Partial protection against experimental vaginal candidiasis after mucosal vaccination with heat-killed *Candida albicans* and the mucosal adjuvant LT(R192G)", Medical Mycology, 6/1/2002

Publication

<1 %

62

digital.library.unt.edu

Internet Source

<1 %

63

digitalcommons.unomaha.edu

Internet Source

<1 %

64

jnsbm.org

Internet Source

<1 %

-
- 65 repositorio.unesp.br Internet Source <1 %
-
- 66 thesis.library.adelaide.edu.au Internet Source <1 %
-
- 67 www.oncotarget.com Internet Source <1 %
-
- 68 www.sysrevpharm.org Internet Source <1 %
-
- 69 Yenni Andriana, Agung Dwi Wahyu Widodo, Pepy Dwi Endraswari. "A Correlation between the Number of Airborne Bacteria and Fungi using the Settle Plate Method with Temperature and Relative Humidity at the Clinical Microbiology Laboratory of Dr. Soetomo Regional General Hospital Surabaya, Indonesia", *Journal of Pure and Applied Microbiology*, 2023
Publication <1 %
-
- 70 Yudi Her Oktaviono, Ferry Sandra, Achmad Lefi, Christian Pramudita Budianto. "Secretome And Ramiprilat Effects On Endothelial Progenitor Cells Proliferation In Chronic Coronary Syndrome Patient", 2020 7th International Conference on Biomedical and Bioinformatics Engineering, 2020
Publication <1 %
-

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Effects of Golden Sea Cucumber Extract (*Stichopus hermanni*) on Hyphae, Neutrophils and TNF- α in BALB/c Mice Inoculated with *C. albicans* Intravaginally

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
