


Journal of
Pakistan Association of Dermatologists

JPAD

www.jp.ad.com.pk

SPECIAL FOCUS;

Tinea corporis caused by *microsporum canis* in children with HIV



A Publication of

PAKISTAN ASSOCIATION OF DERMATOLOGISTS

About the Journal

Editor-in-Chief

Prof. Ijaz Hussain

Editor

Prof. Zahida Rani

Associate Editors

Prof. Faria Asad

Prof. Shehla Shaukat

Assistant Editors

Dr. Ghazala Butt
Dr. Bushra Bashir
Dr. Sadia Jabeen
Dr. Rutaba Kiran

Dr. Mahwish Zahoor
Dr. Umara Siddique
Dr. Wajieha Saeed

Dr. Amina Afzal
Dr. Zareen Saqib
Dr. Nabeela Shehzadi

JPAD, the official journal of Pakistan Association of Dermatologists is published quarterly, four issues per volume and one volume per year (ISSN 1560-9014, print; ISSN 2413-7650, online). The journal is recognized by Pakistan Medical Commission, Higher Education Commission of Pakistan and is indexed in College of Physicians and Surgeons Pakistan MEDLIP; Ulrich's International Periodical Directory, USA; ExtraMED, London; EMBASE/Excerpta Medica, PakMediNet, The Netherlands; and Index Medicus, WHO Alexandria, Egypt.



Published: 2022-12-08

Editorial

The importance of dermatological lesions in the agile diagnosis of Monkeypox

Gonzalo Dominguez Alvarado, Angie Marcela Arboleda Roca , Aymer Daniel Bolívar Cabrera , Gabrielle Yolmari Curicó Martínez , Laura Camila Millán Guerrero
662-664

[PDF](#)

Original Articles

The Spectrum of Dermatoses among Prisoners; A Retrospective Analysis

Hira Tariq
665-668

[PDF](#)

Basil Leaf Extract and Eugenol against Isolates of Candida sp. Causing Oral Candidiasis in HIV/AIDS

Dwi Murtiastutik, Emma Hidayati Sasmito, Diah Mira Indramaya, Sawitri Sawitri, Rahmadewi Rahmadewi, Afif Nurul Hidayati, Pepy Dwi Endraswari, Sudjarwo Sudjarwo, Budi Utomo
669-675

[PDF](#)

Can Ki-67 immunohistochemistry marker differentiate mycosis fungoides from cutaneous lichen planus?

Mazaher Ramezani, Kambiz Kamyab Hesari, Negar Ramezanizoorab, Masoud Sadeghi, Mansour Rezaei, Sedigheh Khazaei
676-682

PDF

Correlation between quality of life and clinical severity of melasma in Pakistani women

Maryam Qayyum, Saadiya Siddiqui, Mohsina Awais, Maimona Ilyas, Atif Shahzad, Nadia Ali Azfar
683-689

PDF

Association of Pruritus Visual Analogue Scale and Risk Factors in Adolescence Pediculosis Capitis in Two Public Boarding Schools, West Java

Arlene Rainamira, Firman Parrol, Yari Castiliani, Githa Rahmayunita, Mochamad Helmi Aziz, Kusmarinah Bramono, Sandra Widaty
690-695

PDF

Scalp defects and their reconstruction in patients at Allama Iqbal Memorial Teaching Hospital, Sialkot

Sarfraz Ahmad, Raffad, Natasha Rehman, Zahida Rani, Atika Malik
696-700

PDF

Novel trends in cutaneous manifestations of geriatric dermatoses in a tertiary care hospital, South India

Latest trends in geriatric dermatosis

Hasini budeda, Indira B, Darsan S, V. V. V. Satyanarayana, Suruthi Purushothaman
701-706

PDF

Effect of Lactobacillus plantarum IS-10506 supplementation with oral metronidazole for the treatment of bacterial vaginosis : a randomized placebo-controlled clinical trial

Afif Nurul Hidayati, Ridha Ramadina Widiatma, Dwi Murtiastutik, Ingrid Suryanti Surono, Cita Rosita Sigit Prakoeswa, Lunardi Bintanjoyo
707-714

PDF

The Platelet Rich Plasma (PRP) Preparation and Standardization of time interval for scalp hair loss

Munir Alam
715-718

PDF

Review Articles

Zinc as an essential element for normal immune reactions and as a therapeutic agent for autoimmune diseases

Inas K. Sharquie, Khalifa E. Sharquie , Wasan W. Al-bassam

719-725

PDF

Review of treatment modalities in the management of Genital Warts

Review of treatment modalities in the management of Genital Warts

Jai Singh Solanki, Swagat Waghmare, Manjyot Gautam, Praneet P. Awake

726-733

PDF

Case Reports

Kaposi's sarcoma in a male with human immunodeficiency virus and condyloma acuminata: A case report

Minna Hasniah, Budi Eko Prasetyorini, Endra Yustin Ellistasari

734-740

PDF

Tinea Corporis Caused by Microsporum Canis in Children with HIV

Sri Mariyani

741-744

PDF

Recurrent herpes simplex in elderly with vitiligo vulgaris: A dilemmatic case

Novita Suprpto, dr. Lili Legiawati, dr. Rinadewi Astriningrum, dr. Shannaz Nadia Yusharyahya

745-747

PDF

Basil leaf extract and eugenol against isolates of *Candida* sp. causing oral candidiasis in HIV/AIDS

Dwi Murtiastutik¹, Emma Hidayati Sasmito¹, Afif Nurul Hidayati^{1,2}, Rahmadewi¹, Sawitri¹, Budi Utomo³, Sudjarwo⁴, Pepy Dwi Endraswari^{5,6}, Diah Mira Indramaya¹

¹Department of Dermatology and Venereology, Dr. Soetomo General Academic Teaching Hospital, Surabaya, Indonesia.

²Department of Dermatology and Venereology, Universitas Airlangga Teaching Hospital, Surabaya, Indonesia.

³Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

⁴Department of Pharmaceutical Science, Faculty of Pharmacy, Universitas Airlangga, Surabaya, Indonesia.

⁵Department of Clinical Microbiology, Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Academic Teaching Hospital, Surabaya, Indonesia.

⁶Department of Clinical Microbiology, Universitas Airlangga Teaching Hospital, Surabaya, Indonesia.

Abstract

Background Oral candidiasis is an opportunistic infection of the oral mucosa caused by *Candida* sp., frequently found in HIV/AIDS patients. Basil leaf extract (*Ocimum sanctum* Linn.), which contains eugenol, a compound believed to inhibit the growth of *Candida* sp.

Objective To evaluate the antifungal effect of basil leaf extract (*Ocimum sanctum* Linn.) and eugenol compared to fluconazole against *Candida* sp. isolates.

Methods Basil leaf extract (*Ocimum sanctum* Linn.) with doses equivalent to eugenol 800 µg/mL and 400 µg/mL, eugenol doses at 800 µg/mL and 400 µg/mL, which was then compared with fluconazole 25 µg/mL against 40 *Candida* sp. isolated stored from the oral cavity of HIV/AIDS patients.

Results The average inhibition zone of fluconazole was 21.81 mm, the mean inhibition zone of eugenol with doses of 800 µg/mL and 400 µg/mL were 17.07 mm and 15.89 mm, and the mean inhibition zone of basil leaf extract (*Ocimum sanctum* Linn.) with doses equivalent to eugenol 800 µg/mL and 400 µg/mL were 14.87 mm and 14.01 mm ($p = 0.001$ and $p < 0.05$).

Conclusion Fluconazole had a significantly higher inhibition zone against *Candida albicans* and *Candida non-albicans* isolates than basil leaf extract (*Ocimum sanctum* Linn.) and eugenol.

Key words

Basil leaf extract (*Ocimum sanctum* Linn.), eugenol, fluconazole, antifungal susceptibility testing, oral candidiasis, HIV, AIDS.

Introduction

Oral candidiasis is a fungal infection of the oral mucosa caused by *Candida* sp.¹ *Candida* sp. is a commensal fungus in healthy people, but under certain conditions it can cause an opportunistic infection. Opportunistic infections can occur as a

consequence of infection with the human immunodeficiency virus (HIV) and the acquired immunodeficiency syndrome (AIDS). During the course of their illness, nearly 90% of HIV/AIDS patients develop oral candidiasis.^{2,3}

Oral candidiasis can be treated with the

antifungal drug nystatin oral suspension and/ or fluconazole as a systemic antifungal.^{4,5} In a test of *Candida* sp. resistance to fluconazole, isolates resistant to fluconazole were found to be 48.6%. *Candida non-albicans* was the most resistant isolate, accounting for 72.2% of the total.² Antifungal drug biofilm formation and resistance are two issues that can be addressed by seeking for alternative antifungal drugs.⁶

Basil (*Ocimum sanctum* Linn.) is a tropical plant that can be found in abundance in Indonesia.⁷ The main component of basil leaf (*Ocimum sanctum* Linn.) is eugenol. This compound is effective against the adaptive mechanism of *Candida albicans* biofilm resistance to fluconazole.⁸ Based on these findings, the researchers proposed to hold an in vitro test of basil leaf extract (*Ocimum sanctum* Linn.) and eugenol against stored isolates of *Candida* sp. and compare them to standard oral candidiasis treatment fluconazole, which was now showing signs of resistance.

Methods

The research design used in this study was an experimental laboratory with the aim of evaluating the comparison of the antifungal activity of basil leaf extract (*Ocimum sanctum* Linn.) with doses equivalent to eugenol 800 µg/mL and 400 µg/mL, eugenol at doses 800 µg/mL and 400 µg/mL was then compared with fluconazole 25 µg/mL against 40 isolates stored *Candida* sp. which consisted of 20 *Candida albicans* and 20 *Candida non-albicans* isolated

Address for correspondence

Dwi Murtiastutik,
Department of Dermatology and Venerology
Faculty of Medicine, Universitas Airlangga/ Dr.
Soetomo General Academic Teaching Hospital/
Universitas Airlangga Teaching Hospital,
Surabaya, Jl. Mayjen Prof. Dr. Moestopo No. 6-8
Surabaya 60131, Indonesia.
Ph: (031) 5501609
Email: dwimurtiastutik@yahoo.co.id

from the oral cavity of HIV/AIDS patients who were hospitalized at the Infectious Disease Intermediate Treatment Unit (UPIPI) RSUD Dr. Soetomo Surabaya for the period April 2019 to July 2019 which was reactivated.

The antifungal activity was evaluated using the disk diffusion method with paper discs or blank discs. These data were entered into a data collection sheet and analysed with SPSS (Statistical Package for Social Sciences). This research has obtained ethical approval from the Ethics Committee of Dr. Soetomo General Academic Teaching Hospital Surabaya (0522/LOE/301.4.2/VII/2021).

Results

In this study, using the disc diffusion method, the average results of the inhibition zone test of basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to 800 µg/mL eugenol, basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to 400 µg/mL eugenol, eugenol 800 µg/mL, and eugenol 400 µg/mL will be tested and compared with the standard antifungal treatment fluconazole against *Candida* sp.

The average fluconazole zone for all *Candida* sp. was 21.81 mm, eugenol 800 µg/mL was 17.07 mm, eugenol 400 µg/mL was 15.89 mm, basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to eugenol 800 µg/mL was 14.87 mm, and basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to eugenol 400 µg/mL was 14.01 mm. **Table 1** shows a comparison of the average inhibition zone results for all *Candida* sp.

The average fluconazole inhibition zone in *Candida albicans* species was 22.75 mm, eugenol 800 µg/mL was 16.76 mm, eugenol 400 µg/mL was 15.52 mm, basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to

Table 1 Comparison of the mean diameter of the inhibition zone of basil leaf extract (*Ocimum sanctum* Linn.), eugenol and fluconazole in all species.

No.	Antifungal type	Number of Isolates	Average Inhibition Zone Diameter (mm)	P value
1	Fluconazole 25 µg/mL	40	21.81	0.001
2	Eugenol 800 µg/mL	40	17.07	
3	Eugenol 400 µg/mL	40	15.89	
4	Basil leaf extract equivalent to eugenol 800 µg/mL	40	14.87	
5	Basil leaf extract equivalent to eugenol 400 µg/mL	40	14.01	

Table 2 Comparison of the mean diameter of the inhibition zone of basil leaf extract (*Ocimum sanctum* Linn.), eugenol and fluconazole in *Candida albicans*.

No.	Antifungal type	Number of Isolates	Average Inhibition Zone Diameter (mm)	P value
1	Fluconazole 25 µg/mL	40	22.75	0.001
2	Eugenol 800 µg/mL	40	16.76	
3	Eugenol 400 µg/mL	40	15.52	
4	Basil leaf extract equivalent to eugenol 800 µg/mL	40	14.41	
5	Basil leaf extract equivalent to eugenol 400 µg/mL	40	13.62	

Table 3 Comparison of the mean diameter of the inhibition zone of basil leaf extract (*Ocimum sanctum* Linn.), eugenol and fluconazole in *Candida non-albicans*.

No.	Antifungal type	Number of Isolates	Average Inhibition Zone Diameter (mm)	P value
1	Fluconazole 25 µg/mL	40	20.86	0.001
2	Eugenol 800 µg/mL	40	17.38	
3	Eugenol 400 µg/mL	40	16.25	
4	Basil leaf extract equivalent to eugenol 800 µg/mL	40	15.34	
5	Basil leaf extract equivalent to eugenol 400 µg/mL	40	14.39	

eugenol 800 µg/mL was 14.41 mm, and basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to eugenol 400 µg/mL was 13.62 mm. **Table 2** shows a comparison of the average inhibition zone results against *Candida albicans*.

In *Candida non-albicans*, the average fluconazole inhibition zone was 20.86 mm, eugenol 800 µg/mL was 17.38 mm, eugenol 400 µg/mL was 16.25 mm, basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to eugenol 800 µg/mL was 15.34 mm, and basil leaf extract (*Ocimum sanctum* Linn.) was equivalent to eugenol 400 g/mL was 14.39 mm. The comparison of the average inhibition zone results against *Candida non-albicans* could be observed in **Table 3**.

Because the data were not normally distributed and homogeneous, non-parametric statistical

methods (Mann Whitney Test) was used to analyze it. The results of the non-parametric statistical test showed that the data significance value was 0.001, so <0.05, which means that there was a significant difference between the mean inhibition zone of eugenol 800 µg/mL, eugenol 400 µg/mL, and basil leaf extract (*Ocimum sanctum* Linn.) equivalent to eugenol 800 µg/mL, and basil leaf extract (*Ocimum sanctum* Linn.) equivalent to eugenol 400 µg/mL compared to fluconazole which is the standard antifungal drug against the growth of *Candida* in all species, *Candida albicans* and *Candida non-albicans*.

Discussion

In this study, the average inhibition zone of fluconazole was greater than the average inhibition zone of eugenol and basil leaf extract

(*Ocimum sanctum* Linn.). The inhibition zones of both eugenol and basil leaf extract (*Ocimum sanctum* Linn.) are not better than fluconazole as a standard antifungal drug, but both eugenol and basil leaf extract (*Ocimum sanctum* Linn.) have antifungal effects capable of inhibiting the growth of *Candida* sp., according to the mean comparison.

In a similar study published in 2010, Ahmad and colleagues discovered that fluconazole had a better antifungal effect than eugenol, but some *Candida* sp. Have become resistant to fluconazole. Subsequently a study was conducted with combining fluconazole and eugenol which concluded that a combination of the two were superior to inhibit *Candida* sp. than fluconazole alone.⁹ In a similar study published in 2021, Godil and colleagues discovered that basil leaf (*Ocimum sanctum* Linn.) had antifungal activity by inhibiting the growth of *Candida* sp., but in an in vitro test, the fluconazole inhibition zone was larger than that of basil (*Ocimum sanctum* Linn.) against *Candida*.¹⁰

In a study by Khan and colleagues in 2014, fluconazole revealed an MIC value of 256 µg/mL, capable of inhibiting *Candida* sp., whereas MIC of eugenol were 400µg/mL, showing that MIC of fluconazole were inferior to MIC of eugenol. This indicates that fluconazole is more sensitive in inhibiting *Candida* sp.¹¹ Basil leaf (*Ocimum sanctum* Linn.) have been shown to inhibit *Candida* sp. in previous studies, but the antifungal activity of basil leaf (*Ocimum sanctum* Linn.) is considered small due to the high MIC. Eugenol is a constituent of basil leaf (*Ocimum sanctum* Linn.) that has been shown to be effective in inhibiting *Candida* sp.⁹

The fluconazole inhibition zone was also significantly higher than the eugenol and basil

leaf extract (*Ocimum sanctum* Linn.) inhibition zones. Fluconazole works by inhibiting the enzyme lanosterol 14-demethylase, a microsomal cytochrome P450 enzyme found in fungal cell membranes that prevents lanosterol from being converted to ergosterol. Disturbances in these enzymes cause the integrity of the fungal membrane to be disrupted, causing fungal growth to be inhibited and thus having a fungistatic effect.^{4,12}

In a study published in 2020 by Sharifzadeh and Shokri, it was discovered that eugenol can inhibit the growth of *Candida* sp.¹³ Eugenol has an antifungal effect by inhibiting the biosynthesis of ergosterol, a key component of fungal cell membranes, causing damage to the membranes and a reduction in function. Damage to cell membranes disrupts the transport of nutrients (compounds and ions) through the cell membrane, preventing fungal cells from growing and resulting in cell lysis. Eugenol also contains lipophilic compounds, which can penetrate the lipid bilayer membrane, which is made up of fatty acid chains, altering the fluidity and permeability of the cell membrane, causing the cell to lose its structure and function, resulting in cell lysis.¹⁴ Eugenol can also prevent the formation of pre-formed biofilms and *Candida albicans* biofilms. This substance inhibits *Candida albicans* biofilm resistance to fluconazole through an adaptive mechanism.¹⁵

In several studies investigating the antifungal effect of eugenol, the mechanism by which eugenol induces *Candida* cell death has not been fully understood. The inactivation of ergosterol synthesis and the production of free radicals, which can be an antifungal effect of eugenol, are the mechanisms of action of eugenol against *Candida*. However, the mechanism of action of eugenol is probably not related to the degradation of *Candida* fungal cell walls.¹⁵ According to Jawetz and colleagues, the

mechanism by which antifungal active substances inhibit fungal growth is through membrane damage. The integrity of the cellular components will be compromised if the cell membrane is damaged, and the fungal respiration process will be disrupted. In the end, there is insufficient energy for active substance transport, causing fungal growth to be disrupted.¹⁶

The results of this study showed that by giving basil leaf extract (*Ocimum sanctum* Linn.) to both *Candida albicans* and *Candida non-albicans*, all isolates showed inhibition zones for the growth of *Candida* sp. Basil leaf (*Ocimum sanctum* Linn.) are a tropical plant that can be found throughout Indonesia. Research by De Ornaty *et al.* in 2017 and Desmara *et al.* in 2017, found antifungal activity from basil leaf extract (*Ocimum sanctum* Linn.) on the in vitro growth of *Candida albicans*.^{7,8} Eugenol is the main component of basil leaf (*Ocimum sanctum* Linn.). Several in vitro and in vivo studies show that eugenol inhibits the growth of the *Candida albicans* fungus, which causes candidiasis.¹⁷

Khan and colleagues found that basil leaf (*Ocimum sanctum* Linn.) inhibited the transition of yeast formation into hyphae, inhibited fungal proteinase enzymes, and inhibited the expression levels of HWP1, SAP1, and PLB2 genes, which are pathogenesis crucially expressed during a *Candida albicans* infection, in a study published in 2014. As a therapeutic effect of the content of basil leaf (*Ocimum sanctum* Linn.), the content of basil leaf (*Ocimum sanctum* Linn.) can cause *Candida* apoptosis by performing programmed cell death (PCD) on fungi.¹¹

The main component of eugenol, which is highly volatile, may have caused the inhibition zone of basil leaf extract (*Ocimum sanctum* Linn.) to be inferior to fluconazole in the study.¹⁸ The antifungal activity test method used can also

have an impact; in this study, the diffusion method was used to test antifungal activity, which is a common method. The diffusion method has the advantage of being a simple and quick way to see the antifungal effect by measuring the diameter of the inhibition zone formed; however, the disc diffusion method has a disadvantage in that not all substances are absorbed on the paper disc, potentially affecting the inhibition zone formed.¹⁶

The concentration of the extract, the content of antifungal compounds, the type of fungus inhibited, and the diffusion power all affect the antifungal activity of a compound. The concentration of the extract can also influence the inhibition zone formed, with the higher the concentration, the larger the clear zone. The more active compounds present, the more focused the concentration, affecting the diameter of the inhibition zone formed on fungal growth.¹⁹

Several other studies produced very encouraging results. According to Ahmad and colleagues 2010 research, eugenol is an antifungal agent with antifungal activity in vitro against *Candida albicans* and *Candida non-albicans*, that are intrinsically resistant to fluconazole. The interaction of eugenol and fluconazole was found to have a high level of synergism. In the strains that were tested, no antagonistic interactions were found. Fluconazole in combination with eugenol improves efficacy while lowering the fluconazole minimum effective dose.⁹

In a study conducted by Jafri and colleagues in 2020, it was discovered that administering fluconazole and eugenol to *Candida* isolates worked synergistically, with eugenol disrupting cell membrane integrity and allowing the entry of standard fluconazole drugs into mold cells. Combining antifungal drugs with eugenol has

several benefits, including increased potency, reduced drug doses, and reduced toxicity, all of which help to inhibit or overcome biofilms and antifungal drug resistance.²⁰

Given the growing treatment failure and antifungal resistance in *Candida* sp., more in vivo studies are needed to assess the potential of these compounds for therapeutic applications and suggest ways to treat resistant *Candida* infections using a combination drug approach. The synergistic interactions between eugenol and antifungal drugs must also be evaluated, with additional studies in animal models needed to assess therapeutic efficacy, topical formulation and toxicity.^{9,20}

References

1. Quindós G, Gil-Alonso S, Marcos-Arias C, Sevillano E, Mateo E, Jauregizar N, dan Eraso E. Therapeutic tools for oral candidiasis: Current and new antifungal drugs. *Med Oral Patol Oral Cir Bucal*. 2019;**24(2)**:172-80.
2. Murtiastutik D, Maharani CS, Rahmadewi R, dan Listiawan MY. Nystatin profile on *Candida* species in HIV/AIDS patients with oral candidiasis: a phenomenology study. *J Pure Appl Microbiol*. 2019;**13(4)**:2013-19.
3. Patil S, Majumdar B, Sarode SC, Sarode GS, dan Awan KH. Oropharyngeal candidosis in HIV-Infected patients—an update. *Front Microbiol*. 2018;**1(1)**:1-9.
4. Kundu RV, dan Garg A. 2012. ‘Yeast Infections: Candidiasis, tinea (pityriasis) versicolor, and malassezia (pityrosporum) folliculitis’. In: Goldsmith LA, Katz SI, Gilchrist BA, Paller AS, Leffel DJ, Wolff K, editors. *Fitzpatrick’s dermatology in general medicine*. 8th ed. New York: McGraw Hill. 2298-301.
5. Lydiawati E, Listiawan MY, Murtiastutik D, Rahmadewi R, Prakoeswa CRS, Avanti C, Fitriani E, Astha ET, Astari L, dan Zulkarnain I. In vitro antifungal susceptibility testing of tea tree oil (TTO) 5% compared with nystatin against *Candida* sp. as important agent of oral candidiasis in HIV/AIDS patients. *BIKKK*. 2020;**32(3)**: 189-94.
6. Rauseo AM, Coler-Reilly A, Larson L, dan Spec A. Hope on the horizon: novel fungal treatments in development. *Open Forum Infect Dis*. 2020;**7(2)**:5-23.
7. Desmara S, Rezeki S, dan Sunnati S. Konsentrasi hambat minimum dan konsentrasi bunuh minimum ekstrak daun kemangi (*O. sanctum* L.) terhadap pertumbuhan *Candida albicans*. *J Caninus Denst*. 2017;**2(1)**: 31-9.
8. De Ornay AK, Prehananto H, dan Dewi ASS. Growth inhibition of *Candida albicans* and power kill *Candida albicans* extract basil leave. *Jurnal Wiyata*. 2017;**4(1)**:78-83.
9. Ahmad A, Khan A, dan Yousuf S. Proton translocating ATPase mediated fungicidal activity of eugenol and thymol. *Fitoterapia*. 2010;**81(1)**:1157-62.
10. Godil AZ, Bhagat D, Das P, Kazi AI, Dugal, R, dan Satpute S. Fluconazole and Ocimum sanctum Oil in soft denture liners to treat biofilms of *Candida albicans* associated with denture stomatitis. *Dentistry*. 2021;**9(1)**:1-11.
11. Khan A, Ahmad A, Khan LA, dan Manzoor N. *Ocimum sanctum* (L.) essential oil and its lead molecules induce apoptosis in *Candida albicans*. *Res Microbiol*. 2014;**165(6)**:411-9.
12. Ghannoum M, Salem I, dan Christensen L. 2019. ‘Antifungals’. In: Kang, S., Amagai, M., Bruckner, A.L., Enk, A.H., Margolis, D.J., McMichael, A.J., Orringers, J.S., editors. *Fitzpatrick’s dermatology*. 9th ed. New York: McGraw Hill, 3436-45.
13. Sharifzadeh A, dan Shokri H. In vitro synergy of eugenol on the antifungal effects of voriconazole against *Candida tropicalis* and *Candida krusei* strains isolated from the genital tract of mares. *Equine Vet J*. 2020;**3(1)**:94-101.
14. Brajawikalpa RS. dan Ramzy AN. Uji efektivitas antijamur minyak atsiri daun cengkeh (*Syzygium aromaticum* L.) terhadap pertumbuhan *Malassezia furfur*. *Jurnal Kedokteran dan Kesehatan*. 2018;**4(1)**:49-51.
15. Silva ICG, Santos HBP, Cavalcanti YW, Nonaka CFW, Sousa SA, dan Castro RD. Antifungal activity of eugenol and its association with nystatin on *Candida albicans*. *Pesquisa Brasileira em Odontopediatria e Clínica Integrada*. 2017;**17(1)**:1-8.
16. Jawetz, Melnick, dan Adelberg. 2007. ‘Mikrobiologi Kedokteran’. 23rd ed. Jakarta: ISBN 978-979- 448-859-1. 273-5.

17. Núñez IC, Arranz JC, Rivas CB, Mendonça PM, Perez K, Sánchez CD, Cortinhas LB, Silva CF, Carvalho M, dan Queiroz M. Chemical composition and toxicity of *O. sanctum* L. var. *cubensis* essential oil up-growing in the Eastern of Cuba. *Phytopathology*. 2017;**9(7)**:1021-8.
18. De Paula SB, Bartelli TF, Di Raimo V, Santos JP, Morey AT, Bosini MA, Nakamura CV, Yamauchi LM, Yamada-Ogatta SF. Effect of eugenol on cell surface hydrophobicity, adhesion, and biofilm of *Candida tropicalis* and *Candida dubliniensis* isolated from oral cavity of HIV-Infected patients. *Evid Based Complement Alternat Med*. 2014;**1(1)**:1-8.
19. Andayani A, Susilowati A, dan Pangastuti A. AntiCandida minyak atsiri lengkuas putih (*Alpinia galanga*) terhadap *Candida albicans* penyebab candidiasis secara invitro. *El Vivo*. 2014;**2(2)**:1-9.
20. Jafri H, Banerjee G, dan Khan MSA. Synergistic interaction of eugenol and antimicrobial drugs in eradication of single and mixed biofilms of *Candida albicans* and *Streptococcus mutans*. *AMB Expr*. 2020;**1(10)**:185.